

THE JOHNS HOPKINS UNIVERSITY SCHOOL OF HYGIENE AND PUBLIC HEALTH, LLOYD E. ROZEBOOM MOSQUITO COLLECTION¹

E. L. PEYTON,² JAMES E. PECOR,^{2,3} THOMAS V. GAFFIGAN,² MILAN TRPIS,⁴ LEOPOLDO M. RUEDA^{2,5}
AND RICHARD C. WILKERSON²

ABSTRACT. The Lloyd E. Rozeboom mosquito collection from the Johns Hopkins University School of Hygiene and Public Health, Baltimore, MD, has been transferred to the National Museum of Natural History, Smithsonian Institution, Washington, DC. This historically and taxonomically important collection has specimens representing 32 genera, 78 subgenera, and 763 species from most mosquito faunal areas of the world. The collection consists of 12,015 individually pinned and labeled adults, and 7,179 microscope slide preparations of immature stages and male and female genitalia. A tabulation of genera, subgenera, species, countries, faunal areas, and number of specimens for each type of preparation is provided.

KEY WORDS Collection, Culicidae, Diptera, Rozeboom

INTRODUCTION

During 1998, the Johns Hopkins mosquito collection, also widely known for its primary organizer and curator, Lloyd E. Rozeboom, was donated and transferred to the National Museum of Natural History (NMNH), Smithsonian Institution, Washington, DC. From 1961 to 1981, specimens from the collection were studied and extensively cited by researchers from the Mosquitoes of Middle America Project (University of California at Los Angeles), the Southeast Asia Mosquito Project, and the Medical Entomology Project (both located at the NMNH). Numerous new species were described from this material, including 25 new taxa authored or coauthored by Doctor Rozeboom, 11 of which are listed here. Primary type specimens of these were previously donated to the NMNH. We present here an inventory of the collection annotated by faunal area and country, following the practice established by Harbach et al. (1990).

Lloyd E. Rozeboom was awarded his Sc.D. in 1934 from the Johns Hopkins University School of Hygiene and Public Health, where he worked under Professor Francis M. Root. In 1939 he returned from Oklahoma A&M College to Johns Hopkins University, to replace his former professor. After retiring in 1975, he continued working for 2 years on a part-time basis to curate the insect collection and to organize a tropical medicine center for the

university (Kitzmiller 1982:470). He was succeeded by Professor Milan Trpis, who cared for the collection and was instrumental in facilitating its transfer to the Smithsonian Institution.

Professor Rozeboom began with a modest collection of specimens from several different sources. The collection had its origins in the early 1920s with collections from the West Indies and South America made by F. M. Root. Other early small, but notable, contributions also were made from South America by O. R. Causey and from Thailand by W. A. Hoffman. Doctor Rozeboom added to these holdings with extensive accessions from 3 main geographical areas. The 1st included collections from Panama made while he was at the Gorgas Memorial Laboratory from 1934 to 1937, followed by Pacific Islands collections (mostly from the Philippines, New Guinea, New Hebrides, and Molucca Islands, Indonesia), made while he was serving in the U.S. Navy during 1944-45, and the 3rd included collections from an arbovirus study site sponsored by the Rockefeller Foundation in Meta, Villavicencio, Colombia, ca. 1947-48. These were augmented by numerous smaller personal collections and donations from colleagues. The World War II Pacific islands collections are widely known as the L. E. Rozeboom, K. L. Knight, and J. Laffoon collections. The Smithsonian Institution acquired the K. L. Knight portion of these collections in the early days (1965-66) of the Southeast Asia Mosquito project, an organization at the NMNH funded by the U.S. Army.

The Rozeboom collection consists of specimens that represent most major faunal regions of the world. Figure 1 (from Belkin 1962) shows the world mosquito faunal areas that correspond to the areas listed in Tables 1 and 2. Table 1 reflects specimen labeling and is divided into countries and island groups. The locality numbers listed in Table 1 are used in Table 2 to indicate where each species was collected. Because specific island names have often been overlooked in previous publications, we believe it is more useful to cite them when known. For example, the Philippines, Indonesia, and Solo-

¹ The views of the authors do not purport to reflect the views of the Department of the Army or the Department of Defense.

² Department of Entomology, Division of Communicable Diseases and Immunology, Walter Reed Army Institute of Research, Washington, DC 20307-5100.


³ Author to whom correspondence and reprint requests should be addressed.

⁴ Department of Molecular Microbiology and Immunology, Johns Hopkins School of Hygiene and Public Health, 615 N Wolfe Street, Baltimore, MD 21205.

⁵ Present address: Public Health Pest Management Section, Division of Environmental Health, Department of Environment and Natural Resources, PO Box 29593, Raleigh, NC 27626-0593.

WORLD MOSQUITO

FAUNAL AREAS



- ① South Pacific
- ④ Papuan
- ⑦ Oriental
- ⑩ Palaeartic
- ⑬ Neotropical
- ② South Australian
- ⑤ Western Pacific Is.
- ⑧ Malagasy
- ⑪ Nearctic
- ⑭ Patagonian-So. Chilean
- ③ North Australian
- ⑥ Indomalayan
- ⑨ Ethiopian
- ⑫ Amer. Medit.

Fig. 1. World mosquito faunal areas used in Tables 1 and 2 (from Belkin 1962).

Table 1. Geopolitical divisions. The majority of names are taken from Knight and Stone (1977); others are from maps and gazetteers.

Country	Faunal area
1. Afghanistan	10
2. Argentina	13, 14
3. Australia	2, 3
4. Belize	12
5. Bolivia	13
6. Brazil	13
7. Canada	11
8. China	7-10
a. Hong Kong	
9. Colombia	12, 13
10. Costa Rica	12
11. Ecuador	12, 13
12. El Salvador	12
13. England (United Kingdom)	10
14. Egypt	10
15. Ethiopia	9
16. France	10
a. Corsica	
17. Germany	10
18. Ghana	9
19. Greece	10
20. Guatemala	12
21. Guyana	12
22. Honduras	12
23. India	7
24. Indonesia	4, 6

Table 1. Continued.

Country	Faunal area
a. Irian Jaya (New Guinea)	4
b. Java	6
c. Moluccas	4
d. Sumatra	6
e. Sulawesi [Celebes]	6
f. Morotai	4
g. Rau [Raoe]	4
25. Italy	
a. Sardinia	10
26. Iraq	10
27. Japan	10
a. Honshu	10
b. Ryukyus Archipelago	7
c. Okinawa	7
28. Kenya	9
29. Lebanon	10
30. Liberia	9
31. Malaysia	6
a. East (N. Borneo)	
b. West (Peninsular)	
32. Malawi	9
33. Mauritania	9, 10
34. Mexico	11, 12
35. Mozambique	9
36. Nepal	1
37. Netherlands	10
38. New Zealand	1
39. Nicaragua	12
40. Nigeria	9
41. Panama	12

Table 1. Continued.

Country	Faunal area
42. Pakistan	7, 10
43. Paraguay	13
44. Papua New Guinea	4
45. Peru	13
46. Philippines	6
a. Balabac	
b. Basilan	
c. Busuanga	
d. Culion	
e. Jolo	
f. Leyte	
g. Luzon	
h. Mindanao	
i. Negros	
j. Palawan	
k. Samar	
47. Rhodesia (Zambia/Zimbabwe)	9
48. Sierra Leone	9
49. Singapore	6
50. South Africa	9
51. Sudan	9, 10
52. Suriname	10, 13
53. Syria	10
54. Taiwan (Republic of China)	7
55. Tanzania	9
56. Thailand	6, 7
57. Uganda	9
58. United States	11, 12
a. Alaska	
59. Venezuela	12, 13
60. Zaire	9
61. Zimbabwe	9
Pacific Island groups	
62. Caroline Islands	5
a. Truk	
b. Moen	
63. Fiji Islands	1
64. Gilbert and Ellice Islands	1
65. Loyalty Islands	1
66. Marianas Islands	5
a. Guam	
b. Rota	
67. Marquesas Islands	1
68. New Caledonia Islands	3
69. New Hebrides Islands	1
a. Santo	
b. Malekoula	
c. Malo	
70. Samoa Islands	1
a. American	
b. Western	
c. Swains	
d. Tajuha	
e. Tutuila	
f. Mangpasago	
g. Fajatago	

Table 1. Continued.

Country	Faunal area
71. Santa Cruz Islands	1
a. Nupani	
72. Society Islands	1
a. Tahiti	
73. Solomon Islands	1
a. Bellona	
b. Bougainville	
c. Gizo	
d. Guadalcanal	
e. Malaita	
f. Rennell	
g. Tulaghi	
h. Uki Ni Masi	
i. Mbanika	
74. South Pacific (island unknown)	1
75. Tokelau Islands	1
76. Tuamotu Archipelago	1
a. Taiao	
77. Tonga Islands	1
a. Tafahi	
b. Tongatapu	
78. Wallis Islands	1
Indian Ocean	
79. Mascarene Islands	8
a. Mauritius	
West Indian islands and countries	
80. Aruba	12
81. Bahamas	12
82. Barbados	12
83. Caicos	12
84. Carricou	12
85. Curacao	12
86. Dominica	12
87. Grenada	12
88. Guadaloupe	12
89. Hispaniola	12
a. Dominican Republic	
b. Haiti	
90. Jamaica	12
91. Marie Galante	12
92. Martinique	12
93. Montserrat	12
94. Mayreau	12
95. Nevis	12
96. Puerto Rico	12
97. Saint Croix	12
98. Saint Kitts	12
99. Saint Lucia	12
100. Saint Martin	12
101. Saint Vincent	12
102. Tobago	12
103. Trinidad	12
104. Turks	12
105. Union	12

Table 2. Summary of the Johns Hopkins L. E. Rozeboom mosquito collection.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Aedeomyia</i> (<i>Aedeomyia</i>)				
<i>catantacta</i> Knab	12	—	6	46(t)
<i>squamipennis</i> (Lynch Arribalzaga)	31 ²	47	12, 13	6, 9, 41
<i>venustipes</i> (Skuse)	1 ²	—	6	56
<i>Aedes</i> (<i>Aedes</i>)				
<i>cinereus</i> Meigen	3 ²	21	11	7, 58
<i>Aedes</i> (<i>Aedimorphus</i>)				
<i>alboscutellatus</i> (Theobald)	1	7	6	46(h)
<i>apicoannulatus</i> (Edwards)	—	1	9	40
<i>dentatus</i> (Theobald)	—	5	9	50
<i>irritans</i> (Theobald)	—	3	9	18, 40
<i>oakleyi</i> Stone	13	4	5	66(a)
<i>pampangensis</i> (Ludlow)	12	4	6	46(f)
<i>vexans</i> (Meigen)	51 ²	211	1, 5, 6, 9-11	8, 46(g, k), 50, 56, 58, 64(a), 66(a), 69, 71
<i>vexans nipponii</i> (Theobald)	—	4	10	27
<i>vittatus</i> (Bigot)	1	8	6, 9	40, 48, 55, 56
<i>Aedes</i> (<i>Edwardsaedes</i>)				
<i>imprimens</i> (Walker)	2 ²	6	1, 4	44, 46, 73(d)
<i>Aedes</i> (<i>Finlaya</i>)				
<i>albilabris</i> Edwards	19 ²	15	1	73(a, d, e, h)
<i>ananae</i> Knight and Laffoon	1	1	6	46(k)
<i>argenteitarsis</i> Brug	1	14	4	24(a)
<i>argyronotum</i> Belkin	3 ²	—	1	73
<i>aureostriatus</i> (Doleschall)	1	—	6	46(d)
<i>aureostriatus okinawanus</i> Bohart	—	1	10	27
<i>croceus</i> Knight and Laffoon	1	1	6	46(f)
<i>dobodurus</i> King and Hoogstraal	—	1	4	44
<i>elsiae</i> (Barraud)	7 ²	—	6, 7	54, 56
<i>fijiensis</i> Marks	7	—	1	63
<i>flavipennis</i> (Giles)	—	7	6	46
<i>hatorii</i> Yamada	6	—	7, 10	27, 54
<i>hollandius</i> King and Hoogstraal	1	2	4	24(a)
<i>ingrami</i> Edwards	14	15	9	30, 57
<i>japonicus</i> (Theobald)	9	—	7, 10	27(a), 54
<i>kochi</i> (Doenitz)	3	—	4	24(a)
<i>longipalpis</i> (Grunberg)	3	5	9	30, 40, 57
<i>melanopterus</i> (Giles)	14	1	6	46(f)
<i>niveus</i> (Ludlow)	—	1	6	46
<i>notoscriptus</i> (Skuse)	2 ²	4	4	24(a)
<i>novalbitarsis</i> King and Hoogstraal	3	17	4	24(a)

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>papueus</i> (Taylor)	1	17	4	24(a)
<i>poicilius</i> (Theobald)	4 ²	26	6	46(a, b)
<i>rizali</i> (Banks)	6 ²	1	6	46(l)
<i>samoanus</i> (Grunberg)	—	2	1	71(a)
<i>saperoi</i> Knight	1 ²	1	6	46(a)
<i>saxicola</i> Edwards	2 ²	4	6	46(j)
<i>solomonis</i> Stone and Bohart	7 ²	8	1	73(d)
<i>subalbataris</i> King and Hoogstraal	2	4	4	24(a)
<i>togoi</i> (Theobald)	8 ²	8	7, 10	27(a, c), 54
<i>wallacei</i> Edwards	1	2	4	24(a)
<i>Aedes</i> (<i>Geoskusea</i>)	—	1	1	73(d)
<i>longiforceps</i> Edwards	1 ²	5	1	73(d)
<i>perryi</i> Belkin	—	—	—	—
<i>Aedes</i> (<i>Gynometopa</i>)	—	—	—	—
<i>mediovitatus</i> (Coquillett)	13 ³	9	12	89(b), 90, 96, 97
<i>Aedes</i> (<i>Howardina</i>)	—	—	—	—
<i>albonotatus</i> (Coquillett)	2 ²	8	12	59, 89(b)
<i>busckii</i> (Coquillett)	12 ²	—	12	86, 87, 92, 99
<i>fulvithorax</i> (Lutz)	—	4	12, 13	6, 103
<i>ioliota</i> Dyar and Knab	3	2	12	59
<i>pseudodominicii</i> Komp	2 ²	5	12	9
<i>quadrivittatus</i> (Coquillett)	—	3	12	10
<i>septemstriatus</i> Dyar and Knab	10	18	12	9
<i>sexlineatus</i> (Theobald)	10	13	12	9, 59
<i>walkeri</i> (Theobald)	—	1	12	90
<i>whitmorei</i> Dunn	3 ²	1	12	9
<i>Aedes</i> (<i>Kenknighita</i>)	—	—	—	—
<i>paradisissimilis</i> Rozeboom	29 ²	12	6	46(a)
<i>luzonensis</i> Rozeboom	4	—	6	46(a)
<i>Aedes</i> (<i>Lorrainea</i>)	—	—	—	—
<i>lamelliferus</i> Bohart and Ingram	2 ²	2	5	62(a, b)
<i>Aedes</i> (<i>Mucidus</i>)	—	—	—	—
<i>ferinus</i> Knight	2	4	6	46(g)
<i>painei</i> Knight	1	—	1	73(d)
<i>Aedes</i> (<i>Neomelaniconion</i>)	—	—	—	—
<i>albothorax</i> (Theobald)	—	6	9	35
<i>circumluteolus</i> (Theobald)	—	8	9	35, 50

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Aedes (Ochlerotatus)</i>				
<i>aboriginis</i> Dyar	—	9	11	7
<i>angustivittatus</i> Dyar and Knab	24 ²	39	12	9, 10, 22, 41, 59
<i>atlanticus</i> Dyar and Knab	5	16	11	58
<i>atropalpus</i> (Coquillett)	18 ²	46	11	58
<i>aurifer</i> (Coquillett)	—	2	11	58
<i>bimaculatus</i> (Coquillett)	—	2	11	58
<i>campestris</i> Dyar and Knab	—	19	11	7, 58
<i>canadensis</i> (Theobald)	25 ²	100	11	58
<i>cantator</i> (Coquillett)	19 ²	15	11	58
<i>caspius</i> (Pallas)	—	8	11	58
<i>cataphylla</i> Dyar	—	7	11	7
<i>communis</i> (De Geer)	1	113	11	7, 58
<i>condolexscens</i> Dyar and Knab	1	—	12	96
<i>dorsalis</i> (Meigen)	2 ²	5	11	58
<i>dupreei</i> (Coquillett)	6	3	11, 12	9 [?], 58
<i>edgari</i> Stone and Rosen	7 ²	15	1	72(a)
<i>epactius</i> Dyar and Knab	—	16	12	10, 20
<i>eucephalaeus</i> Dyar	2 ²	—	11	9
<i>euplocamus</i> Dyar and Knab	2 ²	8	12	10, 22
<i>excrucians</i> (Walker)	1 ²	1	11	58
<i>fitchii</i> (Felt and Young)	—	10	11	58
<i>flavescens</i> (Mueller)	—	2	11	58
<i>fluviatilis</i> (Lutz)	8 ²	30	12, 13	6, 9, 22, 41, 59
<i>fulvus</i> (Wiedemann)	1 ²	15	12, 13	5, 6, 9, 41
<i>fulvus pallens</i> Ross	13 ²	9	11	58
<i>grossbecki</i> Dyar and Knab	—	4	11	58
<i>hastatus</i> Dyar	3	7	12, 13	4, 6
<i>hexodontus</i> Dyar	—	6	11	58
<i>impiger</i> (Walker)	1	25	11	58
<i>infirmatus</i> Dyar and Knab	—	3	11	58
<i>intrudens</i> Dyar	—	3	12	58
<i>jacobinae</i> Serafim and Davis	—	2	13	6
<i>mitchellae</i> (Dyar)	3 ²	5	11	58
<i>nigromaculis</i> (Ludlow)	15 ²	15	11	58
<i>nubilus</i> Theobald	—	14	12, 13	22 [?], 45 [?]
<i>oligopistus</i> Dyar	3 ²	1	12	9 [?], 41 [?]
<i>pullatus</i> (Coquillett)	7 ²	12	11	58
<i>punctor</i> (Kirby)	7	32	11	7, 58
<i>scapularis</i> (Rondani)	28 ²	29	12, 13	6, 9, 59, 87, 89(b)
<i>serratus</i> (Theobald)	34 ²	36	12, 13	6, 9, 22, 41, 45, 59, 103
<i>sollicitans</i> (Walker)	9 ²	14	11	58
<i>spencerii itahaensis</i> (Theobald)	—	8	11	58

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>sticticus</i> (Meigen)	5 ²	64	11	58
<i>stimulans</i> (Walker)	1 ²	2	11	58
<i>taeniorhynchus</i> (Wiedemann)	38 ²	62	11-13	10, 11, 22, 41, 58, 59, 84, 87, 89(b), 96
<i>thelcter</i> Dyar	6	2	11	58
<i>thibaulti</i> Dyar and Knab	2	—	11	58
<i>thorntoni</i> Dyar and Knab	7	4	11	58
<i>tortilis</i> (Theobald)	3 ²	—	12	89(b)
<i>trivittatus</i> (Coquillett)	23 ²	45	11, 12	22, 58
<i>vigilax</i> (Skuse)	5 ²	5	6	46, 56
<i>Aedes</i> (<i>Protomacleaya</i>)				
<i>terrans</i> (Walker)	21 ²	108	12	9, 41 [?], 59
<i>thorntoni</i> Dyar and Knab	3 ²	—	12	9, 41 [?], 59
<i>triseriatus</i> (Say)	37 ²	53	11	58
<i>zoosophus</i> Dyar and Knab	2	6	11	58
<i>Aedes</i> (<i>Rhinoskusea</i>)				
<i>longirostris</i> (Leicester)	2 ²	1	6	46(j), 56
<i>wardi</i> Reinert	3	3	6	46
<i>Aedes</i> (<i>Scutomyia</i>)				
<i>albolineatus</i> (Theobald)	74 ²	102	4	24(a, f), 46(g, k), 73(a, d, e, h)
<i>arboricola</i> Knight and Rozeboom	2 ²	2	6	46(k)
<i>boharti</i> Knight and Rozeboom	25	70	6	46(a, d, g, j, k)
<i>hoogstraali</i> Knight and Rozeboom	2	2	6	46(f)
<i>laffooni</i> Knight and Rozeboom	4	—	6	46(g, i)
<i>Aedes</i> (<i>Skusea</i>)				
<i>pembaensis</i> Theobald	—	5	9	35
<i>Aedes</i> (<i>Stegomyia</i>)				
<i>aegypti</i> (Linnaeus)	51 ²	124	6, 7, 9, 10, 12, 13	6, 10, 18, 30, 39, 41, 45, 50, 51, 54-56, 58, 59, 63, 69, 87, 89
<i>africanus</i> (Theobald)	5 ²	6	9	57
<i>albopictus</i> (Skuse)	41 ²	34	4, 6, 7, 10, 11	8, 24(a, f), 27(a), 46(c, e, h, j), 54, 56, 58
<i>alcastidi</i> Huang	1	—	6	46(j)
<i>annandalei</i> (Theobald)	3	—	7	54
<i>apicoargenteus</i> (Theobald)	—	6	9	30, 39, 57
<i>chemulpoensis</i> Yamada	2	—	10	8
<i>desmotoides</i> (Giles)	—	15	6	46
<i>flavopictus</i> Yamada	2	2	10	27(a)
<i>gardnerii</i> (Ludlow)	5	18	6	46(g)
<i>guamensis</i> Farner and Bohart	6 ²	27	1, 5	66(a), 71(a)
<i>hebrideus</i> Edwards	33 ²	28	1	67, 69, 73(d)
<i>heischi</i> van Someren	8 ²	—	9	28

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>hensilli</i> Farnier	—	2	7	27
<i>horrescens</i> Edwards	8	—	1	63
<i>leuteocephalus</i> (Newstead)	—	7	9	18, 40, 55
<i>malayensis</i> Colless	7 ²	—	6	56 (Colony, Bangkok strain)
<i>metallicus</i> (Edwards)	7	1	9	28, 40
<i>pandani</i> Stone	16 ²	17	5	66(a)
<i>paullusi</i> Stone and Farnier	4	—	6	46(h, k)
<i>pernotatus</i> Farnier and Bohart	11 ²	—	1	69(b, c)
<i>polynesiensis</i> Marks	100 ²	54	1	64, 67, 71(c-g), 72(a), 76(a)
<i>pseudoscutellaris</i> (Theobald)	49 ²	33	1	63 (Colony specimens)
<i>quasicutellaris</i> Farnier and Bohart	49 ²	57	1	73(d)
<i>riversi</i> Bohart and Ingram	1	—	7	27(c)
<i>rotanus</i> Bohart and Ingram	—	2	5	66
<i>scutellaris</i> (Walker)	80 ²	103	1, 4, 6	24(a, f), 46(k), 77(a) (Lab colony)
<i>scutoscriptus</i> Bohart and Ingram	1	2	5	62(a, b)
<i>simpsoni</i> (Theobald)	3	6	9	30, 55, 57
<i>tongae</i> Edwards	27 ²	31	1	97(b)
<i>upolensis</i> Marks	2 ²	3	1	71(b, g)
<i>w-albus</i> (Theobald)	2 ²	—	6, 7	56
<i>Aedes</i> (<i>Verrallina</i>)				
<i>andamanensis</i> Edwards	—	1	6	46
<i>butleri</i> Theobald	3	3	6, 7	46, 56
<i>campylostylus</i> Laffoon	2	7	6	46(g)
<i>carmentis</i> Edwards	2	15	4	24(a)
<i>cucctoi</i> Belkin	1 ²	—	1	73(d)
<i>dux</i> Dyar and Shannon	16 ²	22	6	46, 56
<i>hamistylus</i> Laffoon	1 ²	2	1	73
<i>johnsoni</i> Laffoon	3 ²	10	6	46(b)
<i>lineatus</i> (Taylor)	11 ²	9	1	69(a), 73(d)
<i>macrodixoa</i> Dyar and Shannon	1 ²	18	6	46, 46(g)
<i>margarsen</i> Dyar and Shannon	—	3	6, 7	23, 31(b)
<i>nigrotarsis</i> (Ludlow)	5	18	6	46(g)
<i>Anopheles</i> (<i>Anopheles</i>)				
<i>aithenii</i> James, Group	1 ²	3	6, 7	23, 31(b)
<i>alboaenatus</i> (Theobald)	—	2	6	31(b)
<i>algeriensis</i> Theobald	19	5	10	14, 19, 25(a)
<i>annulipalpis</i> Lynch Arribalzaga	—	1	13	2
<i>apicimacula</i> Dyar and Knab	8	19	12	4, 10, 41, 59
<i>argyropus</i> (Swellengrebel)	2	3	6	31(b)
<i>atroparvus</i> van Thiel	—	17	10	37
<i>atropus</i> Dyar and Knab	12	—	12	34, 58

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>aztecus</i> Hoffmann	54 ²	22	12	34 (Colony, Mexico City)
<i>baezai</i> Gater	12 ²	11	6	31(b), 46(a, k)
<i>barberi</i> Coquillett	10 ²	—	11	58
<i>barbrosiris</i> van der Wulp	58 ²	109	6, 7, 10	6, 23, 31(b), 36, 46(g, h, j, k), 49
<i>barianensis</i> James	1	—	7	23
<i>bengalensis</i> Puri	14	—	6	31(b)
<i>bradleyi</i> King	16 ²	17	11	58
<i>brevipalpis</i> Roper	1	—	6	31(b)
<i>claviger</i> (Meigen)	28	10	10	13, 14 [?], 16(a), 25(a)
<i>costani</i> Laveran	7	14	8, 9	28, 40, 55, 79(a)
<i>crawfordi</i> Reid	1	3	6	31(b)
<i>crucians</i> Wiedemann	21 ²	14	11	54
<i>culiciformis</i> Cogill	2	—	7	23
<i>earlei</i> Vargas	223 ²	167	11	58
<i>eiseni</i> Coquillett	20 ²	24	12, 13	4, 6, 9, 10, 22, 41, 45, 59
<i>fausti</i> Vargas	1	—	12	34
<i>franciscanus</i> McCracken	20	22	11	58
<i>franciscoi</i> Reid	1	25	6	46(g)
<i>freeborni</i> Aitken	2 ²	15	11	58
<i>georgianus</i> King	14	—	11	58
<i>grabhamii</i> Theobald	25 ²	11	12	69(a), 90, 96
<i>hectoris</i> Giaquinto-Mira	1	—	12	20
<i>hyrcanus</i> (Pallas)	1	—	10	19
<i>hyrcanus</i> (Pallas), Group	12 ²	—	6, 7, 10	8, 14, 31(b), 36, 46, 56
<i>intermedius</i> (Peryassu)	10 ²	13	12, 13	6, 21
<i>labranchiae</i> Falleroni	8	—	10	16(a)
<i>lesteri</i> Baisas and Hu	80	80	6	46(f, g, k)
<i>letifer</i> Sandosham	2	5	6	31(b)
<i>lindesayi japonicus</i> Yamada	1	—	10	24(a)
<i>lindesayi lindesayi</i> Giles	2	—	7	23
<i>lindesayi pleicau</i> Koidzumi	7	—	7	54
<i>lindesayi benguetensis</i> King	14	1	6	46(g)
<i>lindesayi benguetensis</i> King	19 ²	31	10, 11	13, 17, 19, 35(a), 58
<i>maculipennis</i> Meigen, Group	—	25	13	6
<i>maculipes</i> (Theobald)	—	21	12	41
<i>malefactor</i> Dyar and Knab	—	10	6	46(g)
<i>manalangi</i> Mendoza	2	—	10	25(a)
<i>marteri</i> Senevet and Prunelle	1	—	10	6, 59
<i>mattogrossensis</i> Lutz and Neiva	2	11	12, 13	6, 9, 21, 59, 103
<i>mediopunctatus</i> (Lutz), Group	61 ²	46	12, 13	37
<i>messeae</i> Falleroni	—	18	10	6
<i>minor</i> da Costa Lima	7	11	13	6
<i>neomaculipalpus</i> Curry	60 ²	55	12	10, 41, 59, 103

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>nigerrimus</i> Giles	3 ²	9	7, 10	8, 23, 31(b)
<i>nitidus</i> Harrison, Scanlon, and Reid	2	—	6	31(b)
<i>obscurus</i> (Grunberg)	5	6	9	30
<i>occidentalis</i> Dyar and Knab	4 ²	—	11	58
<i>pallidus</i> Theobald	4	5	9	30
<i>paraliae</i> Sandosham	2	—	9	31(b)
<i>parapunctipennis</i> Martini	2	5	12	22, 34
<i>peditaeniatus</i> (Leicester)	28	10	6	31(b)
<i>peryssus</i> Dyar and Knab	17 ²	37	12, 13	6, 9
<i>pilinoctum</i> Harrison and Scanlon	7	—	6	46(g, h)
<i>plumbeus</i> Stephens	12	—	10	16(a)
<i>pseudobarbivrosus</i> Ludlow	31 ²	93	4	24(a), 46(g, k)
<i>pseudomaculipes</i> (Peryassu)	2	—	13	6
<i>pseudopunctipennis</i> Theobald	68 ²	124	11-13	2, 5, 6, 9-11, 34, 41, 45, 57, 59, 85, 87
<i>pseudosinensis</i> Baisas	2	—	6	46(g)
<i>punctimacula</i> Dyar and Knab	45 ²	18	12, 13	9, 10, 41, 45, 59
<i>punctipennis</i> (Say)	30 ²	61	11	58
<i>quadrimaculatus</i> Say, Complex	21 ²	65	11	58
<i>roperi</i> Reid	1	3	6	31(b)
<i>sacharovi</i> Favre	7	3	10	14, 19
<i>samarensis</i> Rozeboom	31 ²	80	6	46(k)
<i>separatus</i> (Leicester)	2	7	6	31(b)
<i>shannoni</i> Davis	—	12	12, 13	6, 21
<i>sinensis</i> Wiedemann	23 ²	89	6, 7, 10	8, 23, 27(a), 31(b), 36, 54
<i>sineroides</i> Yamada	—	6	10	27
<i>umbrosus</i> (Theobald)	4	9	6, 7	23, 31(b)
<i>vanus</i> Walker	4	—	6	46(h)
<i>vestitipennis</i> Dyar and Knab	6 ²	50	12	4, 10, 21, 59, 90, 96
<i>walkeri</i> Theobald	21	42	11	58
<i>Anopheles (Cellia)</i>				
<i>aconitus</i> Doenitz	9 ²	5	6, 7	23, 31(b), 56
<i>annularis</i> van der Wulp	12 ²	31	6, 7, 10	23, 31(b), 36, 46(g), 54, 56
<i>annulipes</i> Walker	1 ²	3	2, 3	3
<i>baisasi</i> Colless	—	1	6	46(g)
<i>balabacensis</i> Baisas	11	10	6	46(a, j), 56
<i>christyi</i> (Newstead and Carter)	1	4	9	18, 55
<i>cinereus</i> Theobald	1	2	9	55
<i>cinereus hispaniola</i> (Theobald)	5	2	10	16, 25(a)
<i>cristatus</i> King and Baisas	1	—	6	46(h)
<i>culicifacies</i> Giles, Complex	3 ²	19	7, 10	23, 36, 56

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>demeilloni</i> Evans	1	3	9	35
<i>dthali</i> Patton	1	—	7	23
<i>farauti</i> Laveran	32	265	1	69
<i>flitipinae</i> Manalang	2 ²	2	6	46(g)
<i>flavivirostris</i> (Ludlow)	20 ²	88	6	46(g, h, j, k)
<i>fluvialitilis</i> James	—	4	7, 10	8, 23, 36
<i>funestus</i> Giles	3	15	8, 9	18, 40, 50, 55, 79(a)
<i>gambiae</i> Giles, Complex	5 ²	34	6, 8, 9, 13	6, 18, 30, 40, 47, 50, 55, 61, 79(a)
<i>garnhami</i> Edwards	1	2	9	55
<i>hackeri</i> Edwards	2	1	6	46(a, j)
<i>indefinitus</i> (Ludlow)	18	35	6	46(j)
<i>jamesii</i> Theobald	2 ²	4	7	23
<i>jeyporiensis</i> James	4 ²	1	7, 10	23, 36
<i>karwari</i> (James)	6 ²	36	4, 7	24(a), 25
<i>keniensis</i> Evans	—	1	9	Locality unknown
<i>kochi</i> Doenitz	61 ²	107	4, 6, 7	23, 24, 46(j, k), 31(a, b), 56
<i>kolambuganensis</i> Baisas	1	—	6	46(h)
<i>koliensis</i> Owen	2	167	1, 4	24(a), 73(d)
<i>leucosphyrus</i> Doenitz, Group	2	—	6	31
<i>limosus</i> King	115	176	6	46(f, g, j, k)
<i>litoralis</i> King	29	36	6	46(g, j)
<i>longirostris</i> Brug	—	10	1	44
<i>ludlowae</i> (Theobald)	5	29	6, 7	46(g), 54
<i>lungae</i> Belkin and Schlosser	38	30	7	56
<i>macarthuri</i> Colless	—	1	6	55
<i>machardyi</i> Edwards	1	2	9	8(a), 24(b), 31(b), 36, 46(g, h, j, k), 54, 56
<i>maculatus</i> Theobald, Group	35 ²	134	6, 7, 10	79(a)
<i>macutipalpis</i> Giles	6 ²	6	8	23
<i>majidi</i> Young and Majid	2	1	7	46(g, k)
<i>mangyanus</i> (Banks)	10	56	6, 7	32
<i>marshallii</i> (Theobald)	—	1	9	30
<i>melas</i> Theobald	—	9	9	8(a), 23, 54
<i>minimus</i> Theobald, Group	2	2	7	14 [?]
<i>multicolor</i> Chambouliu	4	—	10	73(d)
<i>nataliae</i> Belkin	1	3	1	Locality unknown
<i>nili</i> (Theobald)	—	2	9	8
<i>pallidus</i> Theobald	1 ²	1	7	46
<i>parangensis</i> (Ludlow)	—	2	6	55
<i>pharoensis</i> Theobald	1	2	9	31(b), 46(g, k), 56
<i>philippinensis</i> Ludlow	17	22	6	55
<i>pretoriensis</i> (Theobald)	1	2	9	

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>pujutensis</i> Colless	2	—	6	31(b)
<i>pulcherrimus</i> Theobald	—	18	10	1, 23
<i>punctulatus</i> Doenitz	33	210	1	24(a), 44, 73(d)
<i>rhodesiensis</i> Theobald	—	4	9	48
<i>riparis</i> King and Baisas	3	3	6	46(h, k)
<i>rufipes</i> (Gough)	—	1	9	50
<i>sergentii</i> (Theobald)	5	—	10	14 [?]
<i>solomonis</i> Belkin, Knight, and Rozeboom	—	14	1	73(d)
<i>splendidus</i> Koidzumi	2 ²	5	7, 10	23, 36, 54
<i>squamosus</i> Theobald	—	1	9	Locality unknown
<i>stephensi</i> Liston	3	8	7, 10	23, 29, 42
<i>subpictus</i> Grassi	23	13	6, 7, 10	23, 24, 31(b), 36, 54, 56
<i>sundaicus</i> (Rodenwaldt)	3	15	6, 7	23, 31(b), 56
<i>superpictus</i> Grassi	3	3	10	19, 53
<i>takasagoensis</i> Morishita	5	6	7	54
<i>tessellatus</i> Theobald	10	124	6, 7	8, 23, 31(b), 46(g, h), 54, 56
<i>turkhuudi</i> Liston	1 ²	—	7	23
<i>vagus</i> Doenitz	2 ²	60	6, 7	8, 24(d), 31(b), 46, 49, 56
<i>wilsoni</i> Evans	3 ²	3	9	53
<i>Anopheles (Kerteszia)</i>				
<i>bambusicolus</i> Komp	23 ²	21	12, 13	9
<i>bellator</i> Dyar and Knab	9	23	12	103
<i>boliviensis</i> (Theobald)	4	11	12, 13	5, 9
<i>cruzii</i> Dyar and Knab	—	16	13	6
<i>homunculus</i> Komp	2	16	12	103
<i>neivai</i> Howard, Dyar, and Knab	21 ²	6	12	10, 41
<i>Anopheles (Lophopodomyia)</i>				
<i>gilesi</i> (Peryassu)	2 ²	—	13	6
<i>pseudoitibiamaculatus</i> Galvao and Barretto	—	2	13	6
<i>Anopheles (Nyssorhynchus)</i>				
<i>albimanus</i> Wiedemann	141 ²	85	11–13	6, 10, 11, 34, 39, 41, 59, 82, 89(b)
<i>albitalaris</i> Lynch Arribalzaga	125 ²	95	12–14	2, 6, 9, 10, 41, 43, 59, 103
<i>anomalophyllus</i> Komp	5	—	12	10
<i>aquasalis</i> Curry	115 ²	131	12, 13	6, 10, 21, 41, 59, 87, 89(b), 102, 103

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>argyritarsis</i> Robineau-Desvoidy	82 ²	43	12, 13	5, 9, 10, 41, 59, 87
<i>benarrochi</i> Gabaldon, Cova Garcia, and Lopez	11 ²	28	12, 13	6, 9
<i>braziliensis</i> (Chagas)	16 ²	61	12, 13	6, 9
<i>darlingi</i> Root	68 ²	53	10, 12, 13	4, 6, 9, 21, 52, 59
<i>galvaoi</i> Causey, Deane, and Deane	—	5	13	6
<i>lanei</i> Galvao and Amaral	—	2	13	6
<i>lutzi</i> Cruz	—	4	12	6
<i>nuneztovari</i> Gabaldon	18 ²	3	12, 13	6, 9, 59
<i>oswaldoi</i> (Peryassu)	194 ²	337	12, 13	2, 6, 9, 41, 59, 103
<i>parvus</i> (Chagas)	6 ²	30	13	6
<i>rangeli</i> Gabaldon, Cova Garcia, and Lopez	18 ²	242	12, 13	9, 59
<i>rondoni</i> (Neiva and Pinto)	11	16	13	2
<i>sawyeri</i> Causey, Deane, and Sampaio	—	4	13	Locality unknown
<i>strodei</i> Root	113 ²	35	12, 13	6, 9, 10, 21, 41, 59
<i>triannulatus</i> (Neiva and Pinto)	137 ²	110	12, 13	2, 6, 9, 41, 43, 59
<i>Anopheles</i> (<i>Stethomyia</i>)				
<i>komp</i> Edwards	8 ²	14	12, 13	6, 9, 41
<i>nimbus</i> (Theobald)	81 ²	67	12, 13	6, 9, 21
<i>thomasi</i> Shannon	1	1	12	9
<i>Armigeres</i> (<i>Armigeres</i>)				
<i>aureolineatus</i> (Leicester)	—	5	6, 7	23, 46
<i>baisasi</i> Stone and Thurman	1 ²	9	6	46(g)
<i>breinli</i> (Taylor)	1 ²	—	1	73(d)
<i>joloensis</i> (Ludlow)	1 ²	14	6	46(g)
<i>kuchingensis</i> Edwards	4	—	6	56
<i>malayi</i> (Theobald)	1	6	1	73(d)
<i>subalbatus</i> (Coquillett)	6 ²	17	7, 10	8, 23, 29, 54
<i>Armigeres</i> (<i>Leicesteria</i>)				
<i>digitatus</i> (Edwards)	2	2	6, 7	46(g), 54
<i>flavus</i> (Leicester)	—	2	6	46
<i>magnus</i> (Theobald)	—	15	6	46
<i>Bironella</i> (<i>Bironella</i>)				
<i>gracilis</i> Theobald	—	6	1	69
<i>Bironella</i> (<i>Brugella</i>)				
<i>hollandi</i> Taylor	43	41	1	73(d)
<i>Bironella</i> (<i>Neobironella</i>)				
<i>papuae</i> (Swellengrebel and Swellengrebel de Graaf)	—	34	1	69

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Chagasia</i>				
<i>bathana</i> (Dyar)	13	6	12	10, 41
<i>bonneae</i> Root	15	14	12	9
<i>fajardi</i> (Lutz)	32	29	12, 13	6, 9
<i>Coquillettidia</i> (<i>Coquillettidia</i>)				
<i>annettii</i> (Theobald)	—	2	9	30
<i>aurites</i> (Theobald)	—	1	9	51
<i>crassipes</i> (van der Wulp)	—	4	6	46
<i>fuscopennata</i> (Theobald)	—	4	9	57
<i>giblini</i> (Taylor)	—	5	4	44
<i>maculipennis</i> (Theobald)	—	1	9	30
<i>metallica</i> (Theobald)	2 ²	2	9	30
<i>perturbans</i> (Walker)	5	36	11, 12	58
<i>pseudoconopas</i> (Theobald)	—	2	9	57
<i>xanthogaster</i> (Edwards)	5 ²	22	1	69(a)
<i>Coquillettidia</i> (<i>Rhynchoaenia</i>)				
<i>arribalzaga</i> (Theobald)	1 ²	5	12, 13	6, 10
<i>chrysonotum</i> (Peryassu)	—	3	12	41
<i>fasciolata</i> (Lynch Arribalzaga)	2 ²	7	12, 13	6, 10, 41
<i>juxtamansonia</i> (Chagas)	11	2	12, 13	6, 41
<i>nigricans</i> (Coquillett)	—	3	12	41
<i>Culex</i> (<i>Acalytntrum</i>)				
<i>binigrolineatus</i> Knight and Rozeboom	4 ²	2	4	24(a)
<i>perkinsi</i> Stone and Penn	2	8	1	73(d)
<i>Culex</i> (<i>Anoediopora</i>)				
<i>bamborum</i> Rozeboom and Komp	22 ²	27	12, 13	9
<i>conservator</i> Dyar and Knab	25 ²	20	12, 13	6, 9, 41, 87
<i>corriganii</i> Dyar and Knab	1 ²	1	12	41
<i>Culex</i> (<i>Carrallia</i>)				
<i>anduzei</i> Cerqueira and Lane	—	2	12	9
<i>bihaicolus</i> Dyar and Nunez Tovar	2 ²	7	12, 13	9, 59
<i>bonnei</i> Dyar	3 ²	50	12	9
<i>infoliatatus</i> Bonne-Wepster and Bonne	4 ²	6	12	9
<i>iridescens</i> (Lutz)	36 ²	—	12, 13	9
<i>secundus</i> Bonne-Wepster and Bonne	74 ²	87	12, 13	9
<i>urichii</i> (Coquillett)	74 ²	56	12, 13	6, 9
<i>wilsoni</i> Lane and Whitman	1 ²	156	12, 13	9

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Culex (Culex)</i>				
<i>abnormalis</i> Lane	7 ²	4	12	9
<i>acharistus</i> Root	8	—	13	6
<i>alis</i> Theobald	18	—	6	46(a, k)
<i>annulioris</i> Theobald	—	9	1	73(i)
<i>annulirostris</i> Skuse	107 ²	80	1, 4	24(a), 44, 69(c), 76(a)
<i>annulus</i> Theobald	4	2	7	23
<i>atriceps</i> Edwards	4	—	1	72(a)
<i>bahamensis</i> Dyar and Knab	9 ²	—	12	89(b)
<i>bidens</i> Dyar	—	6	90(b)	
<i>bitaeniorhynchus</i> Giles	101 ²	8	6, 7, 10	8, 46(c, g-k), 54
<i>chidesterei</i> Dyar	32 ²	1	12, 13	6, 59
<i>coronator</i> Dyar and Knab	129 ²	123	12, 13	2, 6, 10, 41, 59
<i>decens</i> Theobald	—	2	12	9
<i>declarator</i> Dyar and Knab	79 ²	54	11-13	6, 9, 22, 41, 58, 59, 103
<i>duplicator</i> Dyar and Knab	7 ²	—	12	89(b)
<i>fasyi</i> Baisas	—	2	6	46
<i>fuscocephala</i> Theobald	185 ²	57	6, 7	33, 46(c, g, h, j, k)
<i>gellidus</i> Theobald	44 ²	6	6, 7	46(e, f, j), 56
<i>guiarti</i> Blanchard	—	1	9	30
<i>habilitator</i> Dyar and Knab	5 ²	—	12, 13	45, 92, 93
<i>incognitus</i> Baisas	14 ²	7	1, 4, 6	24(a, c), 46(g), 73(d)
<i>infictus</i> Theobald	13 ²	7	12	22, 41, 59, 87, 101
<i>interrogator</i> Dyar and Knab	7 ²	1	11, 12	41, 58
<i>infula</i> Theobald	16	—	6	46(g, h, k)
<i>litoralis</i> Bohart	—	2	5	66(b)
<i>lygrus</i> Root	—	3	13	6
<i>maracayensis</i> Evans	9 ²	6	12, 13	9, 59
<i>marquesensis</i> Stone and Rosen	—	51	1	63
<i>mimeticus</i> Noe	2	3	7, 10	8, 25, 29, 46(k)
<i>mimulus</i> Edwards	30 ²	113	4, 6	24(a), 46(b, k), 56
<i>mollis</i> Dyar and Knab	39 ²	88	12, 13	6, 9, 10, 41
<i>nigripalpus</i> Theobald	111 ²	34	11-13	6, 9, 10, 22, 39, 41, 45, 58, 59, 87, 89, 90, 96
<i>omani</i> Belkin	10	7	1	34, 73(b)
<i>orientalis</i> Edwards	—	1	7, 10	27
<i>pacificus</i> Edwards	45 ²	79	1	69(c)
<i>perfuscus</i> Edwards	—	6	9	35, 50
<i>philippinensis</i> Sirivanakarn	21	19	6	46(h)
<i>piptiens</i> Linnaeus	38 ²	64	10, 11, 13	9, 13, 27, 58
<i>poitcilipes</i> (Theobald)	—	2	9	40
<i>pseudovishnui</i> Colless	2 ²	6	6, 7	23, 46(c)

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>quinquefasciatus</i> Say	178 ²	129	1, 6, 7, 10-13	2, 6, 8, 9, 23, 39, 41, 43 [?], 46(f, g, j, k), 54, 56, 58, 59, 63, 69, 71, 87, 89, 92, 96, 97
<i>restuans</i> Theobald	22 ²	30	11, 12	58
<i>roseni</i> Belkin	5 ²	8	4	72(a), 76(a)
<i>salinarius</i> Coquillett	26 ²	27	11	58
<i>secutor</i> Theobald	12 ²	4	12, 13	41, 45, 89, 96
<i>sitiens</i> Wiedemann	75 ²	19	1, 6, 8	46(f, g, h, j), 69, 79(a)
<i>squamosus</i> (Taylor)	15 ²	44	1, 4	24(a), 73(d)
<i>starckeae</i> Stone and Knight	4 ²	26	1	69
<i>stigmatosoma</i> Dyar	—	14	11	58
<i>tarsalis</i> Coquillett	37 ²	54	11	58
<i>thalassius</i> Theobald	—	6	9	35
<i>tritaeniorhynchus</i> Giles	18 ²	11	7, 10	8, 27
<i>vagens</i> Wiedemann	7	14	7	23
<i>vicinus</i> (Taylor)	—	12	4	44
<i>vishnui</i> Theobald	69 ²	11	6, 7	8, 23, 46(f, g, h, k), 56
<i>whimorei</i> (Giles)	12 ²	44	4, 6, 7	44, 46(g), 56
<i>zombaensis</i> Theobald	—	6	9	Locality unknown
<i>Culex (Culicomyia)</i>				
<i>fragilis</i> Ludlow	3 ²	2	6, 7	46, 56
<i>macfiei</i> Edwards	4 ²	8	9	30
<i>nebulosus</i> Theobald	3 ²	33	9	30, 40
<i>nigropunctatus</i> Edwards	99 ²	7	6, 7	46(c, e-g, j, k), 56
<i>pallidithorax</i> Theobald	15 ²	—	6, 7, 10	8, 46(g, k), 54
<i>papuensis</i> (Taylor)	22 ²	12	4, 6	24(a), 46(h, j, k)
<i>pullus</i> Theobald	10 ²	90	1, 4	24(a), 73(d)
<i>ryukyensis</i> Bohart	—	2	10	27(b)
<i>scanloni</i> Bram	69 ²	—	6	46(f-h, j, k)
<i>spathifurca</i> (Edwards)	34 ²	—	6, 7	46(k), 56
<i>Culex (Eumelanomyia)</i>				
<i>brevipalpis</i> (Giles)	—	23	4	24(a), 46
<i>femineus</i> Edwards	2	20	1	69
<i>foliatus</i> Brug	12 ²	—	4	24(g), 46(g, k)
<i>hayashii</i> Yamada	3	1	10	27
<i>horridus</i> Edwards	—	1	9	Locality unknown
<i>inconspicuus</i> (Theobald)	—	4	9	30
<i>insignis</i> (Carter)	—	2	9	40
<i>laureli</i> Baisas	31 ²	—	6	46(g, k)

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>malayi</i> (Leicester)	10 ²	—	6, 7	8, 56
<i>wigglesworthi</i> Edwards	2	2	9	30
<i>yeageri</i> Baisas	4	—	6	46(j)
<i>Culex</i> (<i>Lophoceraomyia</i>)				
<i>atracus</i> Colless	—	1	1	73(d)
<i>bergi</i> Belkin	1	—	1	73(d)
<i>buxtoni</i> Edwards	8 ²	29	1	69
<i>carolinensis</i> Bohart and Ingram	1	2	5	62(a)
<i>fraudatrix</i> (Theobald)	10 ²	54	1	24(a), 73(d)
<i>hurlbuti</i> Belkin	—	1	1	73
<i>infantulus</i> Edwards	1 ²	—	6	46(h)
<i>josephineae</i> Baisas	6 ²	—	6	46(k)
<i>kuhnsi</i> King and Hoogstraal	1	—	6	46
<i>lairdi</i> Belkin	1	—	1	73(d)
<i>lavatae</i> Stone and Bohart	2 ²	—	6	46(k)
<i>mammilifer</i> (Leicester)	4	1	6	46(j, k)
<i>minor</i> (Leicester)	—	6	6	46(k)
<i>ornatus</i> (Theobald)	2 ²	58	4	24(a)
<i>reidi</i> Colless	1 ²	—	7	56
<i>solomonis</i> Edwards	4 ²	3	1	73(d)
<i>wakasi</i> Belkin	1 ²	4	1	73(d)
<i>Culex</i> (<i>Lutzia</i>)				
<i>allostigma</i> (Howard, Dyar, and Knab)	5	13	12	9
<i>bigoti</i> Bellardi	1	—	13	6
<i>fuscanus</i> Wiedemann	46 ²	—	6, 7	8, 46(c, e, g, h, j, k)
<i>haitifaxii</i> Theobald	—	34	6, 7	8, 46(a, c, g, h, j, k)
<i>tigrisipes</i> de Grandpre & de Charmoy	2	46	1	73(g)
<i>Culex</i> (<i>Melanoconion</i>)				
<i>abominator</i> Dyar and Knab	4 ²	—	11	58
<i>albinensis</i> Bonne-Wepster and Bonne	8 ²	—	12, 13	9, 59
<i>alogistus</i> Dyar	3 ²	1	12	9
<i>atratus</i> Theobald	12 ²	1	12	41, 86, 89(b), 95, 96
<i>bastagarius</i> Dyar and Knab	73 ²	54	12, 13	6, 9, 10, 41, 59, 103
<i>batesi</i> Rozeboom and Komp	9 ²	6	12	9, 59
<i>caudelli</i> (Dyar and Knab)	4 ²	10	12	9
<i>clarki</i> Evans	—	11	13	6
<i>commevynensis</i> Bonne-Wepster and Bonne	4 ²	2	12	9
<i>conspirator</i> Dyar and Knab	33 ²	26	12	9, 41, 59, 103

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>corethynensis</i> Dyar	2 ²	2	12	9, 59
<i>distinguendus</i> Dyar	3 ²	1	12	9, 41, 103
<i>dunni</i> Dyar	48 ²	24	12, 13	6, 9, 41, 59
<i>eastor</i> Dyar	7 ²	—	12	22, 41
<i>educator</i> Dyar and Knab	58 ²	42	12, 13	6, 9, 22, 41, 59
<i>egcyon</i> Dyar	13 ²	—	12	9, 41
<i>elevator</i> Dyar and Knab	25 ²	4	12, 13	8, 9, 22, 41, 92, 103
<i>erraticus</i> (Dyar and Knab)	75 ²	51	11-13	6, 22, 41, 58, 59
<i>evansae</i> Root	2 ²	4	12, 13	6, 100
<i>idotus</i> Dyar	7 ²	3	12	9, 87
<i>inadmirabilis</i> Dyar	34 ²	18	12	9, 59
<i>inhibitor</i> Dyar and Knab	26 ²	—	11-13	6, 9, 41, 58, 59, 89(b), 103
<i>innovator</i> Evans	2	5	12, 13	6, 59
<i>intrincatus</i> Brethes	4 ²	6	12, 13	10, 41
<i>iolambis</i> Dyar	13 ²	2	12	87
<i>jocasta</i> Komp and Rozeboom	2	—	12	103
<i>jubilifer</i> Komp and Brown	12 ²	25	12	9, 41, 59
<i>lucifugus</i> Komp	30 ²	26	12	41, 59
<i>ocossa</i> Dyar and Knab	14 ²	—	12, 13	6
<i>oedipus</i> Root	1 ²	—	12	41
<i>panocosa</i> Dyar	5	—	12	41
<i>paracrybda</i> Komp	1 ²	—	12	41
<i>peccator</i> Dyar and Knab	8 ²	6	11, 12	58
<i>phlogistus</i> Dyar	14 ²	2	12, 13	6, 9, 59
<i>pilosus</i> (Dyar and Knab)	61 ²	18	11, 12	9, 41, 58, 59, 103
<i>plectoporce</i> Root	7 ²	5	12, 13	6, 9
<i>psatharus</i> Dyar	1 ²	—	12	41
<i>rooti</i> Rozeboom	6 ²	3	12	9, 41
<i>serratimarge</i> Root	2 ²	3	12	9, 59
<i>spissipes</i> (Theobald)	5 ²	—	12	41
<i>sursuptor</i> Dyar	4 ²	6	12	9
<i>taeniopus</i> Dyar and Knab	11 ²	—	12	41
<i>tecmarsis</i> Dyar	4 ²	—	12	41
<i>theobaldi</i> (Lutz)	39 ²	38	12, 13	6, 9, 41, 59
<i>unicornis</i> Root	3 ²	5	12	41, 59
<i>zeteki</i> Dyar	1 ²	—	13	6
<i>Culex</i> (<i>Micraeodes</i>)	—	6	12	Locality unknown
<i>antillummagnorum</i> Dyar	—	—	—	—
<i>Culex</i> (<i>Microculex</i>)	—	—	—	—
<i>albipes</i> Lutz	14 ²	—	13	6
<i>aphylactus</i> Root	—	2	12	9
<i>chryselatus</i> Dyar and Knab	—	2	12	9
<i>consolator</i> Dyar and Knab	2	12	12, 13	6, 9
<i>davisi</i> Kumm	34 ²	66	12, 13	6, 103

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>elongatus</i> Rozeboom and Komp	9 ²	25	12	9
<i>gairus</i> Root	14 ²	15	13	6
<i>hedys</i> Root	6 ²	23	12	9
<i>imitator</i> Theobald	12 ²	14	12, 13	6, 41, 59, 103
<i>inimitabilis</i> Dyar and Knab	9 ²	9	12, 13	6, 9
<i>jenningsi</i> Dyar and Knab	2	2	12	10
<i>pleuristriatus</i> Theobald	39 ²	24	12, 13	6, 59, 103
<i>stonei</i> Lane and Whitman	4 ²	10	12	103
<i>Culex</i> (<i>Neoculex</i>)				
<i>apicalis</i> Adams	31 ²	64	11	58
<i>boharti</i> Brookman and Reeves	8 ²	—	11	58
<i>leonardi</i> Belkin	—	1	1	73
<i>martinii</i> Medschiid	5	—	10	Locality unknown
<i>territans</i> Walker	31 ²	11	11, 12	58
<i>Culex</i> (<i>Phenacomyia</i>)				
<i>corniger</i> Theobald/ <i>lactator</i> Dyar and Knab	29 ²	64	12	6, 9, 10, 41, 89(b), 103
<i>Culex</i> (subgenus uncertain)				
<i>ocellatus</i> Theobald	3 ²	9	12	9
<i>Culiseta</i> (<i>Allotheobaldia</i>)				
<i>longiareolata</i> (Marquart)	—	6	10	25, 53
<i>Culiseta</i> (<i>Climacura</i>)				
<i>melanura</i> (Coquillett)	4	3	11	58
<i>Culiseta</i> (<i>Culicella</i>)				
<i>morsitans</i> (Theobald)	—	8	10	Locality unknown
<i>Culiseta</i> (<i>Culiseta</i>)				
<i>incidens</i> (Thomson)	9 ²	—	11	58
<i>inornata</i> (Williston)	15 ²	70	11	58
<i>Deinocerites</i>				
<i>cancer</i> Theobald	24 ²	28	12	59, 87, 89(b), 95, 96, 103
<i>epitedeus</i> (Knab)	5 ²	—	12	41
<i>pseudes</i> Dyar and Knab	12 ²	3	12	10, 41
<i>spanius</i> (Dyar and Knab)	1 ²	—	12	59
<i>Eretmapodites</i>				
<i>chrysoaster</i> Graham	3 ²	6	9	30, 40
<i>Ficalbia</i>				
<i>minima</i> (Theobald)	2	—	7	56

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Haemagogus (Conopostegus)</i>				
<i>leucocelaenus</i> (Dyar and Shannon)	—	6	12, 13	6, 9
<i>Haemagogus (Haemagogus)</i>				
<i>anastasionis</i> Dyar	2	5	12	9, 10
<i>andinus</i> Osorno-Mesa	2	12	12	9
<i>argyromeris</i> Dyar and Ludlow	13 ²	34	12	41
<i>boshelli</i> Osorno-Mesa	—	9	12	9
<i>celestes</i> Dyar and Nunez Tovar	8 ²	41	12	59
<i>chalcospilans</i> Dyar	—	3	12	9, 10
<i>equinus</i> Theobald	2 ²	8	12, 13	6, 9, 59, 90
<i>iridicolor</i> Dyar	—	5	12	10
<i>janthinomys</i> Dyar	35 ²	5	12, 13	2, 6, 9
<i>lucifer</i> (Howard, Dyar, and Knab)	5 ²	25	12	9, 41
<i>panarchys</i> Dyar	2 ²	—	13	11
<i>spgazzinii</i> Brethes	24 ²	13	12, 13	6, 9
<i>splendens</i> Williston	5 ²	—	12	2, 94, 105
<i>Hodgesia</i>				
<i>cairmsensis</i> Taylor	6	—	1	73(d)
<i>malayi</i> Leicester	21	4	12	46
<i>solomonis</i> Belkin	55	10	1	73(d)
<i>spoliata</i> Edwards	14	20	4	24(a)
<i>Isostomyia</i>				
<i>paranensis</i> (Brethes)	1	—	13	6
<i>lunata</i> (Theobald)	6	11	13	6
<i>Johnbelkinia</i>				
<i>longipes</i> (Fabricius)	5 ²	3	12, 13	9, 41, 59
<i>Limatus</i>				
<i>asulleptus</i> (Theobald)	3	1	12	22
<i>durhamii</i> Theobald	31 ²	50	12, 13	6, 9, 22, 41, 59
<i>flavisetosus</i> de Oliveira-Castro	7 ²	23	12	9
<i>Malaya</i>				
<i>genurostris</i> Leicester	18	15	6	46
<i>leet</i> (Wharton)	36	45	4	24(a)
<i>Mansonia (Mansonia)</i>				
<i>humeralis</i> Dyar and Knab	11 ²	9	12, 13	5, 6, 43
<i>indubitans</i> Dyar and Shannon	3	1	11-13	6, 9, 58
<i>titillans</i> (Walker)	26 ²	—	12, 13	9, 13, 41, 59

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Mansonia (Mansonioides)</i>				
<i>africana</i> (Theobald)	1	6	9	18
<i>annulifera</i> (Theobald)	1	6	9	18
<i>bonneae</i> Edwards	9 ²	28	6	46
<i>dives</i> (Schiner)	1	2	4, 6	24(a), 46
<i>papuanis</i> (Taylor)	—	7	4	24(a)
<i>uniformis</i> (Theobald)	6	23	4, 6, 7, 9, 10	18, 25, 24(a), 46, 54, 57
<i>Mimomyia (Etorleptomyia)</i>				
<i>elegans</i> (Taylor)	7	—	1	73(d)
<i>luzonensis</i> (Ludlow)	4 ²	—	6, 7	46(g, h), 54
<i>solomonis</i> Belkin	19	7	1	73(d)
<i>Mimomyia (Mimomyia)</i>				
<i>chamberlaini</i> Ludlow	3	4	6	46
<i>chamberlaini metallica</i> (Leicester)	6	1	6, 7	46, 54
<i>Mimomyia (Ingramia)</i>				
<i>deguzmanae</i> (Mattingly)	17	—	6	46(g, j, k)
<i>Opifex</i>				
<i>fuscus</i> Hutton	5 ²	3	1	38
<i>Orthopodomyia</i>				
<i>albicosta</i> (Lutz)	1 ²	—	12, 13	Locality unknown
<i>anopheloides</i> (Giles)	4	1	6, 7	46, 54
<i>fascipes</i> (Coquillett)	23 ²	22	12	4, 9, 41, 59, 103
<i>kummi</i> Edwards	—	3	12	10
<i>phylozoa</i> (Dyar and Knab)	2	1	12	9, 41
<i>signifera</i> (Coquillett)	15 ²	16	11, 12	41, 58
<i>Phonimomyia</i>				
<i>incaudata</i> (Root)	1 ²	—	13	6
<i>longirostris</i> (Theobald)	3	—	13	6
<i>pallidiventer</i> Theobald	1 ²	—	13	6
<i>piticauda</i> (Root)	3 ²	3	13	6
<i>trinidadensis</i> (Theobald)	10 ²	4	12	103
<i>Psorophora (Grabhamia)</i>				
<i>cingulata</i> (Fabricius)	27 ²	35	12, 13	6, 41, 45, 59, 87, 89, 103
<i>columbica</i> (Dyar and Knab)	33 ²	—	11, 12	58
<i>confinis</i> (Lynch Arribalzaga)	10 ²	135	11, 12	9, 10, 34, 58, 59
<i>discolor</i> (Coquillett)	30 ²	56	11, 12	58
<i>jamaicensis</i> (Theobald)	7 ²	2	12	89(b), 96
<i>pygmaea</i> Theobald	8 ²	—	12	89
<i>signipennis</i> (Coquillett)	28 ²	54	11, 12	58
<i>varinervis</i> Edwards	—	3	13	43

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Psorophora (Janthinosoma)</i>				
<i>albipes</i> (Theobald)	—	8	13	6
<i>cyaneus</i> (Coquillett)	30 ²	72	11–13	58, 59
<i>ferox</i> (von Humboldt)	52 ²	44	11–13	6, 9, 10, 22, 41, 45, 58, 59, 103
<i>horrida</i> (Dyar and Knab)	9	36	11, 12	58
<i>longipalpus</i> Randolph and O'Neill	17 ²	—	11, 12	58
<i>lutzi</i> (Theobald)	4 ²	6	12	9
<i>varipes</i> (Coquillett)	10 ²	15	11, 12	5, 159
<i>Psorophora (Psorophora)</i>				
<i>ciliata</i> (Fabricius)	34 ²	21	11–13	2, 6, 58, 59
<i>cillipes</i> (Fabricius)	2 ²	18	12, 13	6, 9, 22, 43
<i>howardii</i> Coquillett	3 ²	4	11, 12	10, 58
<i>lineata</i> (von Humboldt)	6 ²	13	12, 13	6, 59, 103
<i>pallescens</i> Edwards	—	1	13	43
<i>Runchomyia (Ctenogoeeldia)</i>				
<i>magna</i> (Theobald)	3	6	12	10, 59
<i>Sabethes (Peytonulus)</i>				
<i>xenismus</i>	—	2	12	9
<i>Sabethes (Sabethes)</i>				
<i>albiprivus</i> Theobald	—	13	12	9
<i>belisarioi</i> Neiva	—	10	12	9
<i>bipartipes</i> Dyar and Knab	—	3	12	89(b)
<i>cyaneus</i> (Fabricius)	—	3	12	9, 41
<i>tarsopus</i> Dyar and Knab	—	1	12	9, 41
<i>Sabethes (Sabethinus)</i>				
<i>intermedius</i> (Lutz)	—	12	12	9
<i>melanonymphe</i> Dyar	1	1	12	9
<i>Sabethes (Sabethoides)</i>				
<i>chloropterus</i> (von Humboldt)	2	19	12	9, 10
<i>Topomyia (Topomyia)</i>				
<i>pseudobarbus</i> Baisas	2	7	6	46
<i>Toxorhynchites (Afrorhynchus)</i>				
<i>aeneus</i> (Evans)	—	1	9	40
<i>Toxorhynchites (Ankylorhynchus)</i>				
<i>purpureus</i> (Theobald)	8 ²	5	13	6
<i>trichopygus</i> (Wiedemann)	1	2	13	6

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Toxorhynchites (Lynchiella)</i>				
<i>bambusicola</i> (Lutz and Neiva)	—	8	12	9
<i>gadeloupensis</i> Dyar and Knab	13	2	12	9, 59
<i>haemorrhoidalis</i> (Fabricius)	12	14	12	9
<i>haemorrhoidalis superbus</i> (Dyar and Knab)	—	1	13	6
<i>moctezuma</i> Dyar and Knab	4	—	12	41
<i>portoricensis</i> (Roeder)	2	2	12	89(b)
<i>rutilus septentrionalis</i> (Dyar and Knab)	8 ²	15	11, 12	58
<i>theobaldi</i> (Dyar and Knab)	16	16	12	9, 10, 41, 59
<i>violaceus</i> (Wiedemann)	—	5	13	6
<i>Toxorhynchites (Toxorhynchites)</i>				
<i>amboinensis</i> (Doleschall)	—	2	6	46
<i>brevipalpis</i> Theobald	—	3	9	15, 40, 55
<i>nepenthis</i> Dyar and Shannon	2 ²	1	6	46
<i>splendens</i> (Wiedemann)	1 ²	1	6, 7	31(b), 56
<i>Trichoprosopon</i>				
<i>compressum</i> Lutz	2	1	13	6
<i>digitatum</i> (Rondani)	17	99	12	9, 10, 41, 59, 103
<i>evansae</i> Antunes	—	1	12	59
<i>pallidiventer</i> (Lutz)	4	3	12	9
<i>Tripteroides (Rachionotomyia)</i>				
<i>argenteiventris</i> (Theobald)	2	—	4	24(a)
<i>foliicola</i> Belkin	6	—	1	69(c)
<i>melanesiensis</i> Belkin	24 ²	—	1	69(b)
<i>microlepis</i> (Edwards)	4	—	4	24(a)
<i>solomonis</i> (Edwards)	12	—	1	73(d), 44
<i>standfasti</i> Peters	2	—	4	44
<i>Tripteroides (Rachisoura)</i>				
<i>bisquamatus</i> Lee	25	—	4	24(a)
<i>brevirhynchus</i> Brug	4 ²	—	4	24(a)
<i>filipes</i> (Walker)	8	61	4	24(a)
<i>fuscipleura</i> Lee	2	—	4	24(a)
<i>latisquama</i> (Edwards)	25	24	4	24(a)
<i>leei</i> Peters	3	—	4	24(a)
<i>longipalpatus</i> Lee	1	—	4	24(a)
<i>mathesoni</i> Belkin	1	2	1	73(d)
<i>plumiger</i> Bonne-Wepster	1	—	1	73(d)
<i>stonei</i> Belkin	1	21	1	73(d)
<i>vanleeuweni</i> (Edwards)	2 ²	—	4	24(a)

Table 2. Continued.

Taxon	World faunal areas			Localities ¹
	Slides	Adults	World faunal areas	
<i>Tripteroides (Tripteroides)</i>				
<i>alboscuteellatus</i> Lee	2	—	4	44
<i>antennalis</i> Bohart and Farmer	—	2	6	46
<i>bambusa</i> (Yamada)	1	1	7	54
<i>bimaculipes</i> (Theobald)	12 ²	—	4	24(a), 44
<i>binotatus</i> Belkin	1	—	4	44
<i>elegans</i> Brug	6 ²	—	4	44
<i>lipovskyi</i> Belkin	1	2	1	73(d)
<i>mabini</i> Baisas and Ubaldo-Pagayon	1	—	6	46(g)
<i>monetifer</i> (Dyar)	16	—	6	46(g)
<i>nissanensis</i> Lee	1	—	4	44
<i>quasiornatus</i> (Taylor)	1	—	4	44
<i>riverai</i> Miyagi, Toma, and Tsukamoto	15	—	6	46(g)
<i>Uranotaenia (Pseudoficalbia)</i>				
<i>anhydor syntheta</i> Dyar and Shannon	3 ²	1	11	58
<i>annulata</i> Theobald	—	3	9	40
<i>bicolor</i> Leicester	—	2	6	46
<i>jacksoni</i> Edwards	1	—	7	27
<i>nigrripes</i> (Theobald)	—	3	9	18, 48
<i>quadrinaculata</i> Edwards	12 ²	12	1	73(b)
<i>Uranotaenia (Uranotaenia)</i>				
<i>apicalis</i> Theobald	1	—	13	6
<i>argyrotarsis</i> Leicester	—	1	6	46
<i>barnesi</i> Belkin	4	2	1	69, 73(d)
<i>briseis</i> Dyar	1 ²	—	12	59
<i>catosomata</i> Dyar and Knab	4 ²	7	13	6, 9, 41
<i>civinskii</i> Belkin	1	5	1	73(d)
<i>coazacoalcos</i> Dyar and Knab	1 ²	—	12	9, 10, 41, 59
<i>cooki</i> Root	1	—	12	89
<i>geometrica</i> Theobald	28 ²	57	12, 13	6, 9, 48, 59
<i>leucoptera</i> (Theobald)	—	2	13	6
<i>lowii</i> Theobald	20 ²	13	12, 13	6, 41, 58, 59, 89(b)
<i>macfarlanei</i> Edwards	1	—	7	Locality unknown
<i>nataliae</i> Lynch Arribalzaga	—	1	13	6
<i>pallidiventer</i> Theobald	—	1	12	Locality unknown
<i>pulcherrima</i> Lynch Arribalzaga	40 ²	55	12	6, 9, 41, 59
<i>sapphirina</i> (Osten Sacken)	20 ²	22	11, 12	58
<i>sexaueri</i> Belkin	23	—	1	73(d)
<i>socialis</i> Theobald	1	—	12	96
<i>solomonis</i> Belkin	4	2	1	73(d)

Table 2. Continued.

Taxon	Slides	Adults	World faunal areas	Localities ¹
<i>Wyeomyia (Decamyia)</i>				
<i>pseudopecten</i> Dyar and Knab	8 ²	24	12	9, 59
<i>ulocoma</i> (Theobald)	19 ²	21	12	9, 59
<i>Wyeomyia (Dendromyia)</i>				
<i>complosa</i> (Dyar)	1 ²	3	12	59
<i>felicita</i> (Dyar and Nunez Tovar)	3 ²	20	12	59
<i>luteoventralis</i> Theobald	11	10	12, 13	6, 9, 12, 59
<i>undulata</i> del Ponte and Cerqueira	3	—	13	6
<i>ypsipola</i> Dyar	3 ²	—	12	9
<i>Wyeomyia (Nunezia)</i>				
<i>bicornis</i> (Root)	—	1	12	59
<i>Wyeomyia (Wyeomyia)</i>				
<i>arthrostigma</i> (Lutz)	21 ²	24	12, 13	6, 9, 41, 59
<i>caracula</i> Dyar and Nunez Tovar	—	1	12	59
<i>celanocephala</i> Dyar and Knab	10 ²	19	12	59
<i>codiocampa</i> Dyar and Knab	11	12	12	9
<i>gaudians</i> Dyar and Nunez Tovar	—	3	12	59
<i>guasapata</i> Dyar and Nunez Tovar	3 ²	11	12	41, 59
<i>guatemala</i> Dyar and Knab	—	3	12	10
<i>haynei</i> Dodge	1	—	11	58
<i>medioalbipes</i> Lutz	13 ²	7	13	6, 9, 59
<i>mitchelli</i> (Theobald)	7	—	12	89(b)
<i>oblita</i> (Lutz)	12 ²	20	12, 13	6, 9
<i>pertinans</i> (Williston)	10 ²	—	12	87
<i>scotinomus</i> (Dyar and Knab)	2 ²	—	12	41
<i>smithii</i> (Coquillett)	1	—	11, 12	58
<i>vanduzeei</i> Dyar and Knab	1	—	11, 12	58
<i>Wyeomyia</i> (no subgenus)				
<i>aporonoma</i> Dyar and Knab	—	1	12	10
<i>bourrouli</i> (Lutz)	5	12	13	6
<i>chalcoccephala</i> Dyar and Knab	3	7	12	22
<i>circumcincta</i> Dyar and Knab	—	2	12	9
<i>intonca</i> Dyar and Knab	—	2	12	41
<i>mystes</i> Dyar	4 ²	15	12	59
<i>occulta</i> Bonne-Wepster and Bonne	15	17	12	9

¹ See Table 1 for locality numbers and letters.² At least 1 male genitalia slide is present.

mon Islands combined have more than 15,000 islands, some of which are up to 1,600 km apart and extend into different faunal regions.

The collection contains 763 species from 32 genera and 78 subgenera. Included are 12,015 pinned adults and 11,337 slide mounted specimens, distributed on 7,179 slides as follow: 2,668 larval and pupal exuviae, 1,673 larval exuviae, 290 pupal exuviae, 3,166 larvae, and 3,540 genitalia (97% male). Collection records from the Rozeboom, Knight, and Laffoon South Pacific collections and an inventory of all slides and their label data are on file in the NMNH. Specimens without recognizable localities were not included in the above figures, except for a few cases in which they represented the only example for the species.

Where known, we changed synonymous label names to current usage. The listings of genera and subgenera often were left as determined by Rozeboom or others, even though we may not necessarily agree; for example, the tribe Sabethini is under revision and current usage is not given here. How-

ever, we did confirm or change the identification of many specimens when needed. This was particularly the case for obviously erroneous distributions.

ACKNOWLEDGMENTS

We are grateful to Ronald A. Ward and Daniel A. Strickman for reviewing the manuscript.

REFERENCES CITED

- Belkin, J. N. 1962. The mosquitoes of the South Pacific (Diptera, Culicidae), Volumes I and II. Univ. Calif. Press, Berkeley and Los Angeles, CA.
- Harbach, R. E., T. V. Gaffigan and J. E. Pecor. 1990. The J. Pedro Duret mosquito collection (Diptera: Culicidae). *Mosq. Syst.* 22:192-195.
- Kitzmiller, J. B. 1982. Anopheline names: their derivations and histories, Volume VIII. Thomas Say Foundation, Entomological Society of America, College Park, MD.
- Knight, K. L. and Stone, A. 1977. A catalog of the mosquitoes of the world (Diptera: Culicidae). Volume VI. The Thomas Say Foundation, Entomological Society of America, College Park, MD.