

A REARING METHOD FOR *Aedes abserratus*  
(F. AND Y.)

R. A. BRUST AND K. S. KALPAGE

Department of Entomology, University of  
Manitoba, Winnipeg, Canada

Univoltine *Aedes* species of Manitoba are difficult to rear in the laboratory and *A. abserratus* is no exception. A method of rearing the larvae is described here and the same technique has been adapted to other species of Northern Manitoba and the North West Territories.

Eggs of *A. abserratus* required at least 2 months of cold or winter temperatures (2–4° C.) before they would hatch. Maximum hatch was obtained when eggs were kept for 2 months at 20° C. following oviposition, and 4 months at 4° C. A hatch of 90 percent of the viable eggs was obtained when eggs were taken directly from 4° C. and placed in dilute nutrient broth (1:200) at 12–15° C. Larvae hatched after 36–48 hours. They were reared at constant temperatures of 10°, 20°, 23°, 24°, 25°, 26° C., and also at a diurnal temperature cycle of 10° and 23°.

At constant temperatures the writers found that 30–50 percent of the larvae reached the adult stage. Larvae completed their development to pupae in 21–25 days at 10° C.; in 10–12 days at 20° C., and in 5 days at 26° C. Mortality was greatest at the highest temperature, and intersexes resulted at temperatures of 24°–26° C. (Brust, 1966).

By using a daily cycle of low and high temperatures, the writers were able to get 80–85 percent of the larvae through to adults. Temperatures of 10° C. for 12 hours, and 23° C. for 12 hours daily were found to be most suitable. The larval medium consisted of 100–300 ml. of distilled water, 300 mg. of Fleischman's "live" dry yeast, 300 mg. of fine ground dog food (Gaines pellets) and 5 ml. of "wetted" sphagnum peat moss (purchased at garden supplies stores) per 50 larvae. The yeast and dog food (weighed dry) were suspended in distilled water and the floating particles decanted. Air was removed from the peat by vacuum, and the peat wetted down with distilled water. Only the suspended particles of peat were used in the medium; the floating particles were decanted. A stock supply of the prepared peat and food were kept frozen until needed.

Larvae were reared in a closed, unbreakable plastic pan 21 x 30 x 8 cm. The water level in the pan was kept to 3 mm. for first and second instar larvae. Fresh medium was prepared after the third or fourth day and the water level raised to 6 mm. The medium was changed again during the fourth instar, and this time the water level raised to 10 mm.

Pupae were removed to a cage and kept in distilled water at a constant temperature of 20° C. Adult males began to emerge after the third day and females after the fourth day. Adults lived up to 2 months in cages where the humidity was kept at or above 80 percent R.H.

The adults have been mated artificially using the method described by McDaniel and Horsfall, 1957, but the authors have not attempted to maintain a continuous colony.

References

BRUST, R. A. 1966. Gynandromorphs and intersexes in mosquitoes (Diptera: Culicidae). *Canad. J. Zool.* 44:911–921.

MCDANIEL, I. N., and HORSFALL, W. R. 1957. Induced copulation of aedine mosquitoes. *Science* 125:745.

NEW RECORDS FOR *Aedes* SPECIES IN MANITOBA

R. A. BRUST AND K. S. KALPAGE

Department of Entomology, University of  
Manitoba, Winnipeg, Canada

Four species not previously reported from Manitoba include *Aedes abserratus* (Felt and Young), *A. triseriatus* (Say), *A. detritus* (Howard, Dyar and Knab), and *A. barri* (Rueger). The first two have been reported from several provinces in Canada, but *A. detritus* only from Labrador and Ontario (Smith 1952, Steward and McWade 1961) and *A. barri* only from Ontario in Canada (Steward and McWade 1961).

*A. barri* larvae occur in temporary snow-melt pools in Manitoba, together with *A. excrucians* (Walker), *A. fitchii* (F. and Y.), *A. punctator* (Kirby) and *A. communis* (De Geer). Larvae have been collected from the Sandilands Forest Reserve, and from Oak Bluff, Manitoba. Adults, which laid eggs in the laboratory, were collected from Churchill, Flin Flon, Porcupine Forest Reserve, Turtle Mountain Forest Reserve, Riding Mountain National Park, and Sandilands Forest Reserve. The larvae are easily identified if one is using Barr's key (1958) to Minnesota mosquitoes, but may be confused with *A. excrucians* if one is using a key which does not include *A. barri*.

Larvae from Manitoba are similar to those described from Minnesota. The number of comb scales ranged from 10 to 21 with an average of 15. The comb scales were arranged in an irregular double row or in a triangular patch. The upper and lower head hairs were usually double, but occasionally triple. The siphon was 4 times as long as broad at the middle (range 3½–4½); the pecten extended beyond the middle to the outer third with 3 or 4 detached teeth. The ventral tuft on the siphon consisted of 2 or 3 branches and was closely associated with the last pecten tooth. Usually the distal pecten tooth, and often the penultimate tooth, was not straight or gently curved as in *A. excrucians*, but had a slight kink near its base. The most reliable character for separating *A. barri* larvae from *A. excrucians* was found to be the spine on each ventral valve of the siphon. The valves are triangular shaped and are located on the most distal part of