

usual in 1968 in the vicinity of London. Their occurrence in and around houses is in accord with the report of Horsfall (1955) that *P. ciliata* enters buildings and of Headlee (1945) that it is not infrequently sent with house captures.

### References

- CARPENTER, S. J. and LACASSE, W. J. 1955. Mosquitoes of North America (north of Mexico). Univ. Calif. Press, Berkeley and Los Angeles. 360 p.
- HEADLEE, T. J. 1945. The mosquitoes of New Jersey and their control. Rutgers University Press, New Brunswick, N. J. 326 p.
- HORSFALL, W. R. 1955. Mosquitoes, their bionomics and relation to disease. Ronald Press, New York. 723 p.
- JUDD, W. W. 1962. The mosquito, *Psorophora ciliata* (Fabr.), at London, Ontario. Mosq. News 22:304.
- STEWART, C. C. and MCWADE, J. W. 1961. The mosquitoes of Ontario (Diptera: Culicidae) with keys to the species and notes on distribution. Proc. Entomol. Soc. Ontario (1960), 91:121-188.

### *Culex pipiens* L. FEEDING ON THE OLIGOCHAETE *Aeolosom hemprichi* EHRENBERG

B. R. BERRY AND M. G. MAW

Research Institute, Research Branch, Canada  
Agriculture, Belleville, Ontario

Most mosquito larvae feed indiscriminately upon algae and small organisms such as rotifers, protozoa, bacteria and fungal spores (Clements, 1963) and a few species, including apparently some filter feeders, are predators (Petersen *et al.*, 1969). Since *Culex pipiens* L. has not hitherto been regarded as being carnivorous on larger, more complex organisms, it is noteworthy that we have observed fourth instar larvae of this species feeding upon the oligochaete worm, *Aeolosoma hemprichi* Ehrenberg.

We suspected that *C. pipiens* might be ingesting the oligochaetes while we were observing the fate of mosquito larvae hatching from egg rafts deposited upon floating patches of decaying algae. Two fourth instar larvae had been collected inadvertently with the algal sample and one appeared to have devoured one of the many oligochaetes that were browsing among the strands of algae.

Ten worms were placed in a small evaporating dish with 5 fourth instar *C. pipiens* larvae. The mosquitoes did not actively pursue the worms but ingested them when they encountered them during normal feeding activities. When mature worms were drawn toward the larval oral cavity tail-first they were sometimes able to escape. On the other hand, worms carried into the oral cavity head-first were always ingested, apparently with ease.

Mature *Aeolosoma* are approximately 1 to 2 mm long and thrive in decaying algae and hay infusions (Ward and Whipple, 1945). They could well be an important item in the mosquito larval diet under certain conditions, especially where organic material is abundant in the water.

### Literature Cited

- CLEMENTS, A. N. 1963. The physiology of mosquitoes. Macmillan Co., N. Y.
- PETERSEN, J. J., CHAPMAN, H. C., and WILLIS, O. R. 1969. Predation of *Anopheles barberi* Coquillett on first instar mosquito larvae. Mosq. News 29:134-135.
- WARD, H. B., and WHIPPLE, G. C. 1945. Fresh-water biology. John Wiley and Sons Inc., N. Y.
- COMPARISON OF THE EFFECT OF SIX PYRETHROIDS AGAINST A BACKSWIMMER, *Notonecta undulata* SAY

G. D. MILLS, JR., J. H. FALES AND  
C. G. DUREIN, JR.

Entomology Research Division, Agr. Res. Serv.,  
USDA

Beltsville, Maryland 20705

Since 0.0025-0.02 p.p.m. Abate® *o,o*-(dimethyl phosphorothioate *o,o*-diester with 4,4'-thiodiphenol) killed 10-93 percent of the backswimmers, *Notonecta undulata* Say, in laboratory tests after 2 days of exposure (Fales *et al.*, 1968), it was used as a standard to evaluate the effectiveness of six pyrethroids against the same species in laboratory tests.

MATERIALS AND METHODS. The six pyrethroids tested were allethrin, *d-trans*-allethrin, dimethrin, Neopynamin® (2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylic acid ester with *N*-(hydroxymethyl)-1-cyclohexene-1,2-dicarboximide), S. B. Penick 1382 ((5-benzyl-3-furylmethyl (±)-*cis-trans*-2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate), and S. B. Penick 1390 ((5-benzyl-3-furylmethyl *d-trans*-2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate). As in the tests with Abate in 1966 and 1967, the WHO beaker method of determining susceptibilities of mosquito larvae (World Health Organization 1960) was used. However, in 1966, the stock solutions (prepared with acetone) were diluted with acetone and then added to distilled water containing the test insects; in 1967, the dilutions from the same stock solution were made with distilled water and then added to lake water containing the test insects. Since the results were the same for both methods, the tests reported here followed the procedure used in 1967 except that the test insects were placed in distilled water instead of in lake water.