

Table 3. Numbers of female mosquitoes collected in dwellings at Irosin and Putiao and infected with the *Wuchereria bancrofti* filaria.

	Irosin		Putiao	
	No. collected & examined	No. infected	No. collected & examined	No. infected
<i>C. quinquefasciatus</i>	56	0	115	2*
<i>Ae. poicilius</i>	2	0	0	-
<i>Ae. ananae</i>	2	0	0	-
<i>Ae. albopictus</i>	2	0	3	0
<i>Armigeres baisasi</i>	3	0	3	0
<i>Mansonia uniformis</i>	1	0	0	-
<i>Anopheles</i> sp.	2	0	0	-

* See the text.

ananae were collected through the dwelling survey, although *Cx. quinquefasciatus* was predominant in collections at both places. Out of 171 *Cx. quinquefasciatus* collected, 82 were dead up to the 15th day. Larvae of *Cx. quinquefasciatus* were easily found at both places. The main breeding places were drains, cesspools and ground pools. These breeding places of this species will increase year by year in the area.

Considering total population contact with man, *Cx. quinquefasciatus* would appear to be responsible for transmission (Baisas 1957) and thus the epidemiological value of this species in Bicol Region should be clarified. It would also be interesting to study the behavior of the abundant *Ae. ananae*, regarding the human blood index and susceptibility to filarial worms.

We wish to thank Dr. Francisco Valeza, Chief, Filariasis Division Bureau of Disease Control, Department of Health, and Prof. B. D. Cabrera, Dean, Institute of Public Health, University of the Philippines, for their assistance and cooperation in undertaking the survey.

This study was supported by an overseas re-

search grant of the Ministry of Education, Science and Culture of Japan, and the Sasakawa Memorial Health Foundation.

Literature cited

- Baisas, F. B. 1957. Notes on Philippine mosquitoes, 19. The mosquito problem in the control of filariasis in Sorsogon Province. *Phil. J. Sci.* 86: 71-120.
- Baisas, F. B., L. F. L. Banez & N. Leuenberger. 1960. Notes on Philippine mosquitoes, 22. The axil-breeding species. *Phil. J. Sci.* 89: 183-199.
- Cabrera, B. D. & M. Tubangui. 1951. Studies on filariasis in the Philippines. 3. *Aedes (Finlaya) poicilius* (Theobald), the mosquito intermediate host of *Wuchereria bancrofti* in the Bicol Region. *Acta Med. Phil.* 7: 221-229.
- Cabrera, B. D. & P. V. Arambulo III. 1973. Human filariasis in the Philippines. *Acta Med. Phil.* 9: 160-173.
- Camacho, V. M. 1969. Control of filariasis in the province of Sorsogon imperative to economic growth. *Phil. J. Pub. Health.* 8: 19-44.

CULISETA MELANURA LARVAE IN TIRES—A RECURRING PHENOMENON?

HENRY R. RUPP

Somerset County Mosquito Extermination Commission, Somerville, NJ 08876

While making a larval survey on 9 July 1977 in Bridgewater Township, New Jersey, of an area littered with discarded automobile tires, I collected several specimens of *Culiseta melanura*

(Coquillett) from water in the tires. Although Wallis and Whitman (1967) reported the occurrence of *Cs. melanura* larvae in both shaded and unshaded areas, these specimens were found in

shaded areas like those described by Eads (1976). Water in the tires was clear, ranging from almost colorless to amber in hue, depending upon the amount of litter in the tires. Associated species were *Aedes triseriatus*, *Anopheles punctipennis* and *Culex restuans*.

A second location of *Cs. melanura* larvae in tires occurred on 24 July in Warren Township, New Jersey, in a densely shaded site. The water in these tires was the color of coffee, and larvae could not be seen unless the liquid was placed in a shallow white container. Associated species in this instance were *Ae. triseriatus* and *An. barberi*.

Siverly (1972) indicated that "receptacles such as tires and tin cans are seldom used" for *Cs. melanura* breeding; however, Judy A. Hansen, Superintendent of the Cape May County Mosquito Extermination Commission, has informed me that *Cs. melanura* larvae have been found in tires in that southernmost county of New Jersey. Thus, it seems possible that the use of tires as an oviposition site by *Cs. melanura* may

not be so rare as once thought but only not so well noted or reported. With tires in an urban or suburban area one expects to find *Culex* and/or *Ae. triseriatus* breeding, and while one is not likely to mistake *Ae. triseriatus* for *Cs. melanura*, it is possible that *Cs. melanura* larvae may have been casually identified as *Culex* on the basis of previous experience with *Culex* in the tire habitat. In any event, this further finding of *Cs. melanura* in this particular habitat may reinforce Wallis and Whitman's comments concerning the need for additional survey activity in light of this species' known vector potential.

References Cited

- Eads, R. B. 1976. An unusual larval habitat for *Culiseta melanura*. Mosquito News 36:544.
 Siverly, R. E. 1972 Mosquitoes of Indiana. Indiana State Board of Health. 126 pp.
 Wallis, R. C. and Whitman, L. 1967. *Culiseta melanura* (Coquillett) breeding in artificial containers. J. Med. Entomol. 4:273-274.

A RELIABLE CHILLED WATER STREAM FOR RHEOPHILIC INSECTS

D. A. CRAIG

Department of Entomology
University of Alberta
Edmonton, Alberta
T6G 2E3

While attempting to rear larval Tanyderidae (Diptera: Nematocera) which may have a larval period in excess of 2 years (Exner and Craig 1976), the need for a highly reliable artificial stream became evident. Motor driven stirrers have been used effectively by Fredeen (1959) to rear simuliid larvae, and others have used similar techniques (Craig 1966). Now, however, magnetic stirrers offer the best approach as they are reliable and readily available in most laboratories (Colbo 1970). I required a temperature controlled system, so have used a Stir-kool® magnetic stirrer (produced by Thermoelectrics Unlimited, Inc., 1202 Harrison Avenue, Wilmington, Delaware, 19809) which has the upper plate cooled thermoelectrically (Peltier Effect).

The stream (Fig. 1) consists of a 4½" high piece of 5½" inside diameter plexiglass tubing,

glued to a piece of flat plexiglass to make a base. A piece of 1¼" high, 2¼" inside diameter plexiglass tubing, is glued to the center of the base. This provides a chamber for the magnetic stirring bar.

Fine sand and gravel were placed in the groove between the stirring chamber and the outer wall. By varying the speed of the stirring bar, water velocities can be varied from almost stationary to approximately 14 cm/sec, quite sufficient to erode the substrate. A subsidiary benefit of the stream design is that any material swept into the stirring chamber is immediately flicked clear by the bar.

The cooling plate allows the temperature of the stream to be controlled closely, for my purposes, 12°C. Such a simple device lends itself to modification for other purposes.

This stream has operated continuously with-