

- Catalogue of Arboviruses Including Certain Other Viruses of Vertebrates. USDHEW Publ. No. (CDC) 75-8301. 789 pp.
- Christensen, B. M., W. A. Rowley, Y. W. Wong, D. C. Dorsey and W. J. Hausler. 1978. Laboratory studies of transovarial transmission of trivittatus virus by *Aedes trivittatus*. *Am. J. Trop. Med. Hyg.* 27:184-186.
- Coz, J., M. Valade, M. Cornet and Y. Robin. 1976. Transmission transovarienne d'un flavivirus, le virus Koutango, chez *Aedes aegypti* L. *C. R. Acad. Sci. Paris* 283:109-110.
- Kokernot, R. H., J. Hayes, R. L. Will, B. Radivojevic, K. R. Boyd and D. H. M. Chan. 1969. Arbovirus studies in the Ohio-Mississippi Basin, 1964-1967. III. Flanders virus. *Am. J. Trop. Med. Hyg.* 18:762-767.
- LeDuc, J. W., W. Suyemoto, B. F. Eldridge, P. K. Russell and A. R. Barr. 1975. Ecology of California encephalitis viruses on the Del Mar Va Peninsula. II. demonstration of transovarial transmission. *Am. J. Trop. Med. Hyg.* 24:124-126.
- Main, A. J., S. W. Hildreth, R. C. Wallis and J. Elston, 1979a. Arbovirus surveillance in Connecticut. III. Flanders virus. *Mosq. News.* 39:560-565.
- Main, A. J., A. L. Smith, R. C. Wallis and J. Elston. 1979b. Arbovirus surveillance in Connecticut. I. Group A viruses. *Mosq. News.* 39:544-551.
- Rosen, L., R. B. Tesh, J. C. Lien and J. H. Cross. 1978. Transovarial transmission of Japanese encephalitis virus by mosquitoes. *Science* 99:909-911.
- Watts, D.M., S. Pantuwatana, G. R. DeFoliart, T. M. Yuill and W. H. Thompson. 1973. Transovarial transmission of La Crosse virus (California encephalitis group) in the mosquito, *Aedes triseriatus*. *Science* 182:1140-1141.
- Whitney, E. 1964. Flanders strain, an arbovirus newly isolated from mosquitoes and birds of New York State. *Am. J. Trop. Med. Hyg.* 13:123-131.

A REPORT OF MERMITHID PARASITES
IN LARVAE OF *Aedes hexodontus*
DYAR IN NORWAY

LEWIS T. NIELSEN,¹ JEAN R. FINNEY² AND
CHRISTINE J. DAHL³

Mermithid nematodes have been reported from at least 63 species of mosquitoes from Africa, Alaska, Australia, Canada, England, France, India, Mexico, Russia, Sumatra and the United States. To our knowledge, these are the first records of a mermithid parasite occurring in mosquitoes in Scandinavia and naturally occurring in *Aedes hexodontus* (Dyar) (Poinar 1975).

On June 17, 1978, a collection of *Ae. hexodontus* larvae was taken along Highway E76, about 6 km west of Vagslid, Norway, and

1 km north of the lake, Kjelavatnet. The area is in southern Norway, about 59° 49' north latitude. The larvae, all in the fourth instar, were collected from a grassy, tundra pool at an elevation of 1,000 m. The site at this latitude is well above timberline. Upon examination most of the larvae were found to be infected with mermithid nematodes.

Intensive collecting at the site produced a total of about 100 living larvae and numerous bodies of dead larvae, some of which were floating on the surface. No pupae or pupal skins were collected and there were no adults in the vicinity of the pool. There were many other snow pools within 1 km of the infected site. Several of these contained fourth instar larvae. *Aedes hexodontus* was the only species present and no infected larvae were found in any of these other pools.

The collected material was brought to the Department of Systematics entomology labo-

¹ Department of Biology, University of Utah, Salt Lake City, Utah 84112.

² Research Unit on Vector Pathology, (RUV), Memorial University of Newfoundland, St. John's, Newfoundland, Canada A1C 5S7.

³ Department of Entomology, University of Uppsala, Sc 75122, Uppsala, Sweden.

ratory at Lunds University, Sweden. Upon examination only 2 larvae from the infected site were found to be free of parasites. These were reared to adults and produced one male and one female of *Ae. hexodontus*. Ten larvae which were dissected had multiple infections, 2-5 worms occurred in each. All of the remaining larvae died upon emergence of the worms.

As the worms emerged they were transferred to jars containing clean, damp, baltic sand. Some of the worms which emerged quickly became covered with a fungus mycelium; an estimated 20% of the worms died from these infections. The *Ae. hexodontus* larvae were kept at 20-25°C until all worms had emerged. The baltic sand containing the mermithid worms was then transferred to a cold room and stored at 8-10° C. On July 25, 1978, in preparation of the senior author's return to the United States, the majority of living nematodes remaining in the baltic sand were heat killed in water and fixed in F.A. 4:1 (Seinhorst 1962).

The preserved nematodes and baltic sand were examined at R.U.V.P. in October 1978. The nematodes were processed (Baker 1953) and mounted in glycerine for identification. Unfortunately, no adult nematodes were found among the preserved specimens; 25% of those fixed had just started to molt to the adult stage. Only 4% of those fixed were immature females, the rest were immature males. This predominance of males is commonly observed when multiple infections occur in the larval stage of mosquitoes. There was evidence of fungal infections on the cuticles of 31% of the nematodes fixed. This may have been partially responsible for the failure of maturation of any worms remaining in the baltic sand, examination of which for additional worms or eggs proved negative.

As complete identification of mermithid nematodes is only feasible when adult specimens are available, it was not possible to accurately identify the specimens. However, all possessed a rounded tail and a long, sharply pointed, conic tail appendage similar to that found in species of the genus *Gastromermis*.

It is hoped that additional collections can be made from this Norwegian locality and a complete description will be published when the adult nematodes are obtained. Not only is there a strong likelihood that it may be a mermithid new to science, but its establishment in an austere arctic-type tundra environment merits its study as a potential biological control agent against the hordes of *Aedes* mosquitoes

which inhabit the arctic regions in areas where man has settled.

Literature Cited

- Baker, A. D. 1953. Rapid method for mounting nematodes in glycerine. *Canad. Entomol.* 85:77-78.
- Poinar, G. O. Jr. 1975. *Entomogenous nematodes*. E. J. Brill, Publ. Leiden, Netherlands. 317p.
- Seinhorst, J. W. 1962. On the killing, fixation and transferring to glycerine of nematodes. *Nematologica* 8:29-32.

COLONIZATION OF *CULICOIDES WISCONSINENSIS* JONES (DIPTERA: CERATOPOGONIDAE)

BRADLEY A. MULLENS AND EDWARD T. SCHMIDTMANN

Department of Entomology, Cornell University, Ithaca, New York 14853

Culicoides wisconsinensis was described by Jones (1956), who collected immatures from a peat-muck substrate in Wisconsin. Jamnback (1965) recovered larvae from soft mud in a brackish marsh and a freshwater lagoon in New York State. Rowley (1967) and Wirth (personal communication) have reared this species from the margins of saline and alkaline lakes in Washington State and the Midwest, respectively. Very little else is known of its biology.

On 5 May 1980, *Culicoides* pupae were collected from a cattail bog polluted with milk-center effluent on Hutchinson Farm, St. Lawrence Co., NY. Reared females held without access to blood oviposited on wet filter paper, and the species was subsequently colonized. Specimens were held at 22°C at a 14:10 (L:D) photoperiod to gather life history data. The colony is currently in the 7th generation.

Follicle maturation in females dissected within 30 min. of eclosion was at late stage III. The pre-oviposition period was 2-3 days, during which time the females mated on contact with males in 237 ml screened holding containers or in the confines of an aspirator