Feeding interactions between native freshwater mussels (Bivalvia: Unionidae) and zebra mussels (*Dreissena polymorpha*) in the Ohio River

Bruce C. Parker¹, Matthew A. Patterson¹, and Richard J. Neves²

¹ Department of Biology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, U. S. A.

² Virginia Cooperative Fish and Wildlife Research Unit, Department of Fisheries and Wildlife Sciences, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, U.S.A.

Abstract: The effects of zebra mussel infestation on the feeding of native unionids in the Ohio River were evaluated through gut contents and available food in the water column. In 1996, heavily infested *Amblema plicata* (Say, 1817) and *Quadrula pustulosa* (I. Lea, 1831) had significantly less (p < 0.01) organic matter in their guts (1.4 and 0.6 mg ash-free dry weight [AFDW], respectively) than lightly infested specimens (4.6 and 1.8 mg AFDW, respectively), and heavily infested *Q. pustulosa* had a significantly lower (p < 0.05) mean algal cell number (1.8×10^4) in the guts than lightly infested specimens (3.9 x 10⁵). However, mean algal cell numbers in the guts of heavily infested and lightly infested *A. plicata* (5.7×10^5 versus 9.1 x 10⁵, respectively) were not significantly different (p = 0.17). In 1997, significant reductions (p < 0.05) in total algal cells and organic matter in gut samples again occurred for heavily versus lightly infested individuals of both species. In addition, gut contents of individual *A. plicata* from one of two sites contained significantly less (p < 0.05) organic matter (0.92 versus 4.55 mg AFDW) and fewer algal cells (9.4×10^4 versus 2.3×10^5) than the combined gut contents of all zebra mussels (18-33 mm in length) attached to their shells. Gut analyses also revealed significant diet overlap between native unionids and infesting zebra mussels. Water samples collected from just above the mussel beds in 1997 showed that algal densities and total suspended solids at the heavily infested site (> 360 zebra mussels/m²) were reduced by more than 50%, when compared to samples collected from the surface. Thus, competitive interactions or interference by zebra mussels likely reduced the availability of algal and detrital food resources for consumption by unionids.

Key words: algae, zebra mussels, unionids, Ohio River, competition

Since its introduction into Lake St. Clair, the zebra mussel, Dreissena polymorpha (Pallas, 1771), has greatly reduced phytoplankton and bacterioplankton levels in the Great Lakes (Wu and Culver, 1991; MacIsaac et al., 1992; Cotner et al., 1995; Fanslow et al., 1995; Heath et al., 1995). Phytoplankton levels in Lake Erie, for example, dropped 62-92% (Leach, 1993), and planktonic diatoms decreased 85% despite sufficient nutrients for growth (Holland, 1993). Consequently, Secchi disk transparencies in Lake Erie have increased 85% (Leach, 1993). Phytoplankton grazing by zebra mussels also can alter the composition of the phytoplankton community. In Lake Huron, for example, zebra mussel feeding has shifted dominance from diatoms to filamentous green algae (Lowe and Pillsbury, 1995), and recent studies show selective rejection of the nuisance bluegreen alga Microcystis by zebra mussels, such that Microcystis becomes dominant in the plankton (H. Vanderploeg, NOAA, pers. comm.)

Zebra mussel colonization of the Great Lakes also has caused dramatic declines in the survival and fitness of native freshwater mussel populations (Hebert *et al.*, 1991; Hunter and Bailey, 1992; Haag *et al.*, 1993; Gillis and Mackie, 1994; Nalepa, 1994; Schloesser and Nalepa, 1994). By attaching to the shells of unionids, zebra mussels can directly affect unionid survival by interfering with feeding, respiration, balance, burrowing, and locomotion (Mackie, 1991; reviewed by Schloesser et al., 1996). Large densities of zebra mussels, however, also can affect unionid survival indirectly by reducing or removing food resources from the water column (Lewandowski, 1976; Hebert et al., 1991; Mackie, 1991; Haag et al., 1993). A large gill-area to body-dry-weight ratio, and a large number of gill cirri in individual zebra mussels, allow for increased filtration efficiency and filtration rate relative to those of native unionids (Silverman et al., 1995). Filtration rates of the freshwater mussel, Lampsilis siliquoidea (Barnes, 1823), for example, were found to be only one-tenth the filtration rate of individual zebra mussels (Heath et al., 1995). In laboratory experiments, Baker and Hornbach (1997) reported that Amblema plicata (Say, 1817) filtered 74 ml/hr, while the 28 infesting zebra mussels filtered 130 ml/hr as a group. Thus, relatively small populations of zebra mussels can affect the feeding of unionids.

Zebra mussel populations in the lower Ohio River

have achieved densities comparable to those in the Great Lakes (350,000/m²; A. Miller, USACOE, pers. comm.). Because of documented impacts to the phytoplankton communities and native mussel populations of the Great Lakes, large populations of zebra mussels in the Ohio River could have similar consequences for native mussel populations. Strayer and Smith (1996) found that low zebra mussel infestation rates in the Hudson River were associated with high unionid mortality and hypothesized that reduced food resources might be the cause. No studies, however, have directly confirmed whether zebra mussels affect the feeding of unionids in a riverine environment where organic materials are continually supplied from upstream. Thus, the objective of this study was to determine whether zebra mussels reduce unionid ingestion of phytoplankton and organic matter by (1) ingesting similar food resources, and (2) reducing food resources at the sediment-water interface.

METHODOLOGY

On 23 July 1996, ten specimens each of the threeridge, Amblema plicata, and the pimpleback, Quadrula pustulosa (I. Lea, 1831), were collected from a lightly infested site on the Ohio River near Parkersburg, West Virginia, which had a mean density of 0.3 zebra mussels/m², and a maximum of one zebra mussel/unionid (P. Morrison, USFWS, pers. comm.). On 16 August 1996, ten specimens of A. plicata were collected from a heavily infested site near Paducah, Kentucky, which had 3,600 zebra mussels/m² (A. Miller, USACOE, pers. comm.). Specimens of Q. pustulosa were difficult to find at this site, so on 21 August 1996, ten specimens were collected from another heavily infested site near Maysville, Kentucky, which had 360 zebra mussels/m² and a maximum of 92 zebra mussels/unionid (P. Morrison, USFWS, pers. comm.). In the field, mussel bodies were removed from shells, weighed, preserved in 95% ethanol, and transported to the laboratory for analysis.

In 1997, ten specimens each of *A. plicata* and *Q. pustulosa* were collected from a highly infested (370 zebra mussels/m²) and a lightly infested (< 1 zebra mussel/m²) site on the Ohio River for gut content analysis. In addition, all zebra mussels, 18-33 mm in length, attached to the shells of *A. plicata* were removed and preserved in 95% ethanol for gut content analysis. Zebra mussels 18-33 mm in length were chosen, because it is difficult to remove the entire gut contents of smaller individuals. At each collection site, water samples with algae were collected from the the surface and overlying the mussel bed, fixed with acid Lugol's solution (Saraceni and Ruggiu, 1969), and placed in settling chambers to compare the density and relative abundance of algal genera using inverted microscopes. Aliquots of 100 ml were then filtered through pre-ashed

Whatman GF/F filters, dried (100°C), and ashed (500°C) to determine the ash-free dry weight (AFDW) of seston.

Gut contents of unionids and of zebra mussels attached to Amblema plicata were individually removed from each specimen, pooled, then suspended in 3 ml of water, and fixed with 50 µl of acid Lugol's solution for analysis. A 50 µl aliquot of the gut contents was placed on a microscope slide. Ocular grids divided the field of view into 59 transects, and algal cells were counted and identified to genus from two transects using an Ausjena/Nomarsky microscope at 400X. The variability of this semi-quantitative method ($\pm 20\%$, $\alpha = 0.05$) was determined using ten counts from the same sample. The remaining gut contents were collected on pre-ashed Whatman GF/F filters, dried (100°C), and ashed in a muffle furnace (500°C) overnight to determine AFDW. Mean algal cell numbers and mean AFDW values in the gut samples of each species were compared by ANOVA.

RESULTS

In 1996, significant differences in total algal cells and organic matter were observed in guts of lightly and heavily infested unionids (Table 1). While mean algal cell numbers in guts of lightly and heavily infested *Amblema plicata* (5.7 x 10⁵ versus 9.1 x 10⁵ cells) were not significantly different (p = 0.17), the gut contents of lightly infested *A. plicata* had significantly more (p < 0.01) organic matter (4.6 mg AFDW) than heavily infested specimens (1.4 mg AFDW). Heavily infested *Quadrula pustulosa* showed significantly lower (p < 0.05) organic matter and mean algal cell number (0.6 mg AFDW and 1.8 x 10⁴ cells, respectively) than lightly infested specimens (1.8 mg AFDW and 3.9 x 10⁵ cells, respectively).

In 1997, significant reductions in organic matter content and total algal cells also were observed in guts of heavily infested unionids (Table 2). Organic matter content and total algal cells were significantly lower (p < 0.05) in the guts of heavily infested *Amblema plicata* (0.9 mg

Table 1. Mean algal cell number and ash-free dry weight (AFDW; \pm SD) in guts of *Amblema plicata* and *Quadrula pustulosa* heavily infested (H) and lightly infested (L) with zebra mussels, July-August 1996.

Species	N	Algal Cell Number	AFDW (mg)	
A. plicata (L)	11	9.1 x $10^5 \pm 6.0$ x 10^5	4.6 ± 0.9	
A. plicata (H)	10	$5.7 \times 10^5 \pm 4.9 \times 10^5$	1.4 ± 0.7	
Q. pustulosa (L)	10	$3.9 \times 10^5 \pm 2.8 \times 10^5$	1.8 ± 1.0	
Q. pustulosa (H)	10	$1.8 \times 10^4 \pm 9.2 \times 10^3$	0.6 ± 0.3	

Table 2. Mean algal cell number and ash-free dry weight (AFDW; \pm SD) in guts of *Dreissena polymorpha* and of *Amblema plicata* and *Quadrula pustulosa* heavily infested (H) and lightly infested (L) with zebra mussels, July-August 1997.

Species	Ν	Algal Cell Number	AFDW (mg)	
A. plicata (L)	10	$5.5 \times 10^6 \pm 2.4 \times 10^6$	5.1 ± 1.7	
A. plicata (H)	10	9.4 x 104 ± 7.6 x 104	0.9 ± 0.8	
Q. pustulosa (L)	10	1.9 x 10 ⁶ ± 1.3 x 10 ⁶	2.0 ± 1.2	
Q. pustulosa (H)	9	6.9 x 10 ⁴ ± 7.3 x 10 ⁴	0.3 ± 0.2	
D. polymorpha	9	2.3 x 10 ⁵ ± 1.3 x 10 ⁵	4.6 ± 3.6	

AFDW and 9.4 x 10⁴ cells, respectively) than lightly infested specimens (5.1 mg AFDW and 5.5 x 10⁶ cells, respectively). Significant reductions in both organic matter content and total algal cells also were observed for heavily infested (0.3 mg AFDW and 6.9 x 10⁴ cells, respectively) versus lightly infested (2.0 mg AFDW and 1.9 x 10⁶ cells, respectively) *Quadrula pustulosa*.

Examination of unionid guts in 1996 and 1997 indicated that Amblema plicata and Quadrula pustulosa readily ingested a significant amount of detritus (ca. 90%) along with algal cells between 4 and 80 µm in length. Diatoms (Bacillariophyta) and green algae (Chlorophyta) dominated gut samples, and the dominant algal genera in 1996 (Chlorella, Cyclotella, Navicula, Melosira, and Scenedesmus; Table 3) and 1997 (Chlorella, Cyclotella, Mougeotia, Melosira, and Scenedesmus; Table 4) were similar. Usually, the relative abundance of algal genera within unionid guts was very similar to relative abundances of algae in water samples collected from the river bottom (Table 5). Interestingly, the pennate diatom Synedra dominated water samples at the lightly infested site (43%), but few if any of these > 100 μ m-long cells were found in the unionid guts. In 1997, algal cell densities and seston AFDW at the water surface (8.37 x 104 cells/ml and 9.0 mg/l, respectively) and above the mussel bed (8.42 x 104 cells/ml and 9.0 mg/l, respectively) were nearly identical at the lightly infested site (Table 5). Seston AFDW (5.0 mg/l) and algal cell densities (2.2 x 104 cells/ml) overlying the mussel bed at the heavily infested site, however, were greatly reduced compared to samples taken from the surface (7.2 mg/l and 8.3 x 104 cells/ml, respectively).

In 1997, individual Amblema plicata collected from the heavily infested site were infested with > 100 zebra mussels/unionid, averaging 50 zebra mussels between 18 and 33 mm in length. Gut contents of zebra mussels contained large amounts of detritus. Pooled samples of zebra mussel guts contained five times more (p < 0.05) organic matter (4.55 mg versus 0.92 mg, respectively) and twice as many algal cells (2.3 x 10^5 cells versus 9.4 x 10^4 cells, respectively) than the specimen of *A. plicata* to which they were attached (Table 3). In addition, the dominant algal genera – *Chlorella, Cyclotella, Mougeotia, Melosira*, and *Scenedesmus* – and range of cell sizes (4-70 µm) in zebra mussel guts were nearly identical to those in infested *A. plicata* (Table 4). Relative abundances of algal genera within zebra mussel guts were similar to relative abundances in water samples collected from the river bottom (Table 5).

DISCUSSION

Reduced food resources at the sediment-water interface can cause decreased growth rates in bivalves despite adequate food resources at the water surface (Frechette and Bourget, 1985). This phenomenon could be particularly important for unionids in the lower Ohio River because zebra mussels occur in large densities at the sediment-water interface and often attach directly to the shells of unionids. In fact, water samples collected from just above the mussel beds in 1997 showed that algal densities and AFDW at the heavily infested site (> 360 zebra mussels/m²) were reduced by more than 50% when compared to samples collected from the surface. Thus, it appears that zebra mussel densities in heavily infested sections of the Ohio River contribute to reduced food resources at the sediment-water interface.

Regardless of reductions in total algal cell densities, the effects of zebra mussel infestation on unionid ingestion can be reduced if zebra mussels and unionids selectively feed on different food types. Currently, studies on selective feeding in bivalves appear to be inconclusive. Some authors have concluded that bivalves select food particles of high quality (Allen, 1914; Loosanoff and Engel, 1947; Shumway et al., 1985), while others have concluded that feeding is non-selective (Churchill and Lewis, 1924; Gale and Lowe, 1971; Bayne et al., 1976). Thus, selective feeding could be species dependent. Zebra mussels have been reported to ingest a wide range of food particles between 0.7 and 450 µm (Mikheyev, 1967; Jorgensen et al., 1984). However, Sprung and Rose (1988) indicated that retention efficiency in zebra mussels is maximized for food particles between 5-35 µm; Ten Winkle and Davids (1982) also concluded from gut content analysis that zebra mussels select particles between 15-50 µm. In our study, zebra mussel gut samples contained food particles between 4-70 µm in maximum dimension. In comparison, Miura and Yamashiro (1990) indicated that the unionid Anodonta calipygos (Kobelt, 1879) ingests food particles between 0.5 and 100 µm. Like for the zebra mussel, maximum retention efficiencies were

Table 3. Percent relative abundances (SD) of algae in guts of lightly (L) and heavily (H) infeste	d
Amblema plicata and Quadrula pustulosa collected from the Ohio River, July-August 1996. (+	۰,
presence [$\leq 2\%$]; –, absence).	

Algae	A. plicata (L)	A. plicata (H)	Q. pustulosa (L)	Q. pustulosa (H)
Chlorophyta				
Ankistrodesmus	-	-	+	_
Chlamydomonas	+	-	+	-
Chlorella	15.3 (8.7)	15.0 (11.2)	42.3 (18.8)	63.0 (29.4)
Chlorococcum	+	+	+	_
Closterium	-	-	+	-
Coelastrum	-	-	+	_
Cosmarium	-	-	+	
Gonium	-	+	_	_
Mougeotia	+	+	+	_
Oedogonium	_	+	_	_
Oocystis	+	_	_	-
Pediastrum	_	+	_	-
Scenedesmus	6.2 (5.9)	5.2 (4.7)	6.6 (5.8)	_
Schroederia	+	-	-	_
Selenastrum	_	_	+	-
Spirogyra	+	_	_	
Staurastrum	+	-	-	-
Bacillariophyta				
Achnanthes	+	-	+	_
Cocconeis	+	-	+	
Coscinodiscus	+	+	+	-
Cvclotella	11.8 (2.7)	45.1 (11.1)	7.0 (3.8)	31.9 (30.8)
Cymbella	+	+	+	-
Diatoma	+	_	+	_
Fragilaria	-	-	+	-
Gomphonema	+	-	+	-
Melosira	5.7 (4.0)	16.0 (11.0)	8.8 (8.2)	2.8 (6.4)
Navicula	26.5 (7.4)	2.2 (0.6)	10.6 (6.2)	
Nitzschia	_	_	+	
Pinnularia	+	+	+	-
Pleurosigma	_	_	+	-
Stephanodiscus	-	+	-	-
Surirella	+	_	-	_
Synedra	+	+	+	-
Tabellaria	+	-	-	-
Cyanoprokaryota				
Chroococcus	+	+	+	-
Merismopedia	-	+	-	-
Oscillatoria	+	-	+	-
Spirulina	-	-	+	-
Other				
Chromulina	-	+	-	-
Dinobryon	+	-	-	-
Peridinium	-	+	+	-
Chroomonas	+	-	-	-

found for intermediate-sized particles (5-30 μ m). Maximum filtration rates for the unionid *Elliptio complanata* (Lightfoot, 1786) also were found for particles between 4 and 5 μ m (Paterson, 1984). In our study, unionid gut analyses indicate that unionids ingest food particles between 4 and 80 μ m in maximum dimension. Thus, zebra mussels and unionids in the Ohio River ingest food resources of similar size.

Relative abundances of algal genera ingested by zebra mussels and the two unionid species were very similar to those in water samples collected immediately above the mussel bed, giving no evidence of selective feeding. The only exception was the pennate diatom *Synedra* which was not readily ingested by unionids or zebra mussels,

Q. pustulosa	Q. pustulosa (H)	D. polymorpha
+	-	-
+	+	+
9.6 (8.8)	30.8 (24.7)	37.5 (16.8)
+	+	+
+	-	-
+	_	_

Table 4. Percent relative abundances (SD) of algae in guts of *Dreissena polymorpha* and lightly (L) and heavily (H) infested *Amblema plicata* and *Quadrula pustulosa* collected from the Ohio River, July-August 1997. (+, presence [$\leq 2\%$]; -, absence).

A. plicata

(H)

A. plicata

(L)

Chlorophyta					
Ankistrodesmus	-	-	+	-	-
Chlamydomonas	-	+	+	+	+
Chlorella	1.7 (1.4)	13.8 (10.1)	9.6 (8.8)	30.8 (24.7)	37.5 (16.8)
Chlorococcum	+	+	+	+	+
Cosmarium	-	-	+	-	-
Crucigenia	-	_	+	-	-
Gonium	-	-	+	-	-
Mougeotia	40.6 (7.5)	5.1 (7.0)	21.2 (11.7)	5.3 (9.3)	14.2 (12.2)
Oedogonium	+	_	_	-	_
Pandorina	+	-	+	-	-
Pediastrum	+	+	+	+	+
Scenedesmus	9.5 (3.6)	5.7 (2.0)	8.2 (5.9)	10.9 (9.6)	13.9 (10.7)
Trebouxia	+	_	-	-	-
Bacillariophyta					
Coscinodiscus	-	-	+	-	-
Cocconeis	-	-	+	-	-
Cyclotella	21.3 (11.9)	38.5 (18.6)	25.9 (6.9)	16.2 (12.7)	5.4 (4.5)
Diatoma	+	_	+	-	-
Gomphonema	+	-	-	-	-
Melosira	13.7 (9.2)	25.9 (16.7)	17.6 (9.4)	28.5 (18.3)	13.7 (10.4)
Navicula	+	+	+	-	+
Stephanodiscus	+	-	+	-	-
Synedra	-	-	+	-	-
Tabellaria	-	-	+	-	+
Cyanoprokaryota					
Aphanocapsa	-	-	+	-	_
Chroococcus	+	-	+	-	-
Oscillatoria	_	-	+	-	-
Other					
Peridinium	-	-	+	-	-

probably because of its cell length of > 100 μ m. In unionid gut samples, *Mougeotia*, *Cyclotella*, and *Melosira* did appear in slightly greater abundance than in water samples. Regardless of selective feeding, unionid and zebra mussel gut contents contained a nearly identical assemblage of dominant algal genera. Thus, by ingesting food particles of similar size and type, zebra mussels in the Ohio River can compete directly with native unionids for food resources. However, competition can only be confirmed if the food items measured to assess diet overlap constitute a significant portion of the total diet of one of the potential competitors (Buss and Jackson, 1981). Unfortunately, there are no studies on the dietary or nutritional requirements of freshwater mussels.

Algae

Significant reductions in the mean AFDW and total algal cell number from gut samples of heavily infested versus lightly infested native freshwater mussels indicate that interference competition for food resources is occurring in the lower Ohio River. Since the arrival of the zebra mussel, unionid mortality thus far at various sites in the lower Ohio River has been estimated at 20-40% (P. Morrison, USFWS, pers. comm.). Interference with unionid feeding by zebra mussels, however, also could have long term consequences for the overall fitness of native mussels. Recent studies have shown that native unionids from the heavily infested Ohio River have significantly reduced glycogen levels relative to unionids from lightly infested areas (Patterson et al., 1997). Glycogen is an important energy reserve for animals, especially bivalves (de Zwann and Zandee, 1972; Barber and Blake, 1981; Bayne and Newell, 1983; Haag et al., 1993), and significant reductions can lead to chronic mortality or declines in reproductive success. Gonad development in marine bivalves, for example, has been shown to continue despite reduced energy reserves (Gabbott and Bayne, 1973; Bayne, 1975), but subsequent growth rates and energy reserves of the developing larvae decreased (Bayne, 1972; Helm et al., 1973; Bayne et al., 1975). Thus, by reducing food resources and interfering

Table 5. Percent relative abundances of algae at the surface and just above the mussel bed in the heavily (H) and lightly infested (L) Ohio River, July-August 1997. (+, presence [$\leq 2\%$]; -, absence).

Algae	Surface (L)	Bottom (L)	Surface (H)	Bottom (H)
Chlorophyta				
Ankistrodesmus	-	+	-	-
Chlamydomonas	6	5	16	11
Chlorella	10	8	25	23
Chlorococcum	+	2	+	7
Gonium	-	+	-	-
Mougeotia	8	2	6	5
Oedogonium	+	+	-	+
Pediastrum	+	+	-	+
Scenedesmus	12	16	13	12
Staurastrum	+	+	+	-
Bacilariophyta				
Coscinodiscus	-	+	+	+
Cyclotella	4	5	3	13
Diatoma	+	+	-	+
Melosira	+	2	4	5
Navicula	-	+	-	+
Stephanodiscus	+	+	-	+
Synedra	12	43	11	10
Cyanoprokaryota				
Chroococcus	-	2	+	7
Merismopedia	8	7	-	-
Oscillatoria	+	6	3	3
Other				
Mallomonas	-	-	-	+

with normal feeding, zebra mussels could have their greatest effect on the long-term persistence of unionid beds in the lower Ohio River through reduction in reproductive success and recruitment.

ACKNOWLEDGMENTS

We thank Patty Morrison, Mitch Ellis, and others at the Ohio River Islands National Wildlife Refuge, as well as Dr. Andrew Miller and the United States Army Corps of Engineers for help in collecting mussels from the Ohio River. Thanks also go to Catherine Gatenby and the many volunteers who assisted with collection and processing of mussels in the field. Additionally, we thank Ashleigh Funk, Lana Shurts, and Doug Smith for laboratory assistance. This study was funded by Quick Response Funds of the Biological Resources Division, U. S. Geological Survey.

LITERATURE CITED

- Allen, W. R. 1914. The food and feeding habits of freshwater mussels. Biological Bulletin 27:127-147.
- Baker, S. M. and D. J. Hornbach. 1997. Acute physiological effects of

zebra mussel (Dreissena polymorpha) infestation on two unionid mussels, Actinonaias ligamentina and Amblema plicata. Canadian Journal of Fisheries and Aquatic Sciences 54:512-519.

- Barber, B. J. and N. B. Blake. 1981. Energy storage and utilization in relation to gametogenesis in Argopecten irridians concentricus (Say). Journal of Experimental Marine Biology and Ecology 52:121-134.
- Bayne, B. L. 1972. Some effects of stress in the adult on the larval development of *Mytilus edulis*. *Nature* 237:459.
- Bayne, B. L. 1975. Reproduction in bivalve mollusks under stress. In: The Physiological Ecology of Estuarine Organisms, J. Vernberg, ed. pp. 259-277. University of South Carolina Press, Columbia, South Carolina.
- Bayne, B. L. and R. C. Newell. 1983. Physiological energetics of marine molluscs. In: The Mollusca, Vol. 4, Physiology, Part 1, A. S. M. Saleuddin and K. W. Wilbur, eds. pp. 407-515. Academic Press, New York.
- Bayne, B. L., P. A. Gabbott, and J. Widdows. 1975. Some effects of stress in the adult on the eggs and larvae of *Mytilus edulis*. Journal of the Marine Biological Association of the United Kingdom 55:675-689.
- Bayne, B. L., R. J. Thompson, and J. Widdows. 1976. Physiology I. In: Marine Mussels, Their Ecology and Physiology, B. L. Bayne, ed. pp. 121-206. Cambridge University Press, Cambridge.
- Buss, L. W. and J. B. C. Jackson. 1981. Planktonic food availability and suspension-feeder abundance: evidence of *in situ* depletion. *Journal of Experimental Marine Biology and Ecology* 49:151-161.
- Churchill, E. P. and S. I. Lewis. 1924. Food and feeding in freshwater mussels. Bulletin of the United States Bureau of Fisheries 39:439-471.
- Cotner, J. B., W. S. Gardner, J. R. Johnson, R. H. Sada, J. F. Cavaletto, and R. T. Heath. 1995. Effects of zebra mussels (*Dreissena polymorpha*) on bacterioplankton: evidence for both size-selective consumption and growth stimulation. *Journal of Great Lakes Research* 21(4):517-528.
- de Zwann, A. and D. I. Zandee. 1972. Body distribution and seasonal changes in the glycogen content of the common sea mussel Mytilus edulis. Comparative Biochemistry and Physiology 43A:53-58.
- Fanslow, D. L., T. F. Nalepa, and G. A. Lang. 1995. Filtration rates of the zebra mussel (*Dreissena polymorpha*) on natural seston from Saginaw Bay, Lake Huron. Journal of Great Lakes Research 21(4):489-500.
- Frechette, M. and E. Bourget. 1985. Food limited growth of Mytilus edulis
 L. in relation to the benthic boundary layer. Canadian Journal of Fisheries and Aquatic Sciences 42:1166-1170.
- Gabbott, P. A. and B. L. Bayne. 1973. Biochemical effects of temperature and nutritive stress on *Mytilus edulis*. Journal of the Marine Biological Association of the United Kingdom 53:269-286.
- Gale, W. F. and R. L. Lowe. 1971. Phytoplankton ingestion by the fingernail clam, Sphaerium transversum (Say), in pool 19, Mississippi River. Ecology 52(3):507-513.
- Gillis, P. L. and G. L. Mackie. 1994. Impact of the zebra mussel, Dreissena polymorpha, on populations of Unionidae (Bivalvia) in Lake St. Clair. Canadian Journal of Zoology 72:1260-1271.
- Haag, W. R., D. L. Berg, D. W. Garton, and J. L. Farris. 1993. Reduced survival and fitness in native bivalves in response to fouling by the introduced zebra mussel (*Dreissena polymorpha*) in western Lake Erie. Canadian Journal of Fisheries and Aquatic Sciences 50:13-19.
- Heath, R. T., G. L. Fahnenstiel, W. S. Gardner, J. F. Cavaletto, and S. Hwang. 1995. Ecosystem-level effects of zebra mussels

(Dreissena polymorpha): an enclosure experiment in Saginaw Bay, Lake Huron. Journal of Great Lakes Research 21(4):501-516.

- Hebert, P. D., C. C. Wilson, M. H. Murdoch, and R. Lazar. 1991. Demography and ecological impacts of the invading mollusc Dreissena polymorpha. Canadian Journal of Zoology 69:405-409.
- Helm, M. M., D. L. Holland, and R. R. Stephenson. 1973. The effect of supplementary algal feeding of a hatchery breeding stock of Ostrea edulis on larval vigour. Journal of the Marine Biological Association of the United Kingdom 53:673-684.
- Holland, R. E. 1993. Changes in planktonic diatoms and water transparency in Hatchery Bay, Bass Island area, western Lake Erie, since the establishment of the zebra mussel. *Journal of Great Lakes Research* 19(3):617-624.
- Hunter, R. D. and J. F. Bailey. 1992. Dreissena polymorpha (zebra mussel): colonization of soft substrata and some effects on unionid bivalves. The Nautilus 106(2):60-67.
- Jorgensen, C. B., T. Kiorboe, F. Mohlenberg, and H. U. Riisgard. 1984. Ciliary and mucus-net filter feeding, with special reference to fluid mechanical characteristics. *Marine Ecology Progress Series* 15:283-292.
- Leach, J. H. 1993. Impacts of the zebra mussel (Dreissena polymorpha) on water quality and fish spawning reefs in western Lake Erie. In: Zebra mussels: Biology, Impacts, and Control, T. F. Nalepa and D. W. Schloesser, eds. pp. 381-397. Lewis Publishers, Boca Raton, Florida.
- Lewandowski, K. 1976. Unionidae as a substratum for Dreissena polymorpha Pall. Polskie Archiwum Hydrobiologie 23(3):409-420.
- Loosanoff, V. L. and J. B. Engel. 1947. Effect of different concentrations of micro-organisms on feeding of oysters (O. virginica). United States Fish and Wildlife Service Fisheries Bulletin 51:31-57.
- Lowe, R. L. and R. W. Pillsbury. 1995. Shifts in the benthic algal community structure and function following the appearance of zebra mussels (*Dreissena polymorpha*) in Saginaw Bay, Lake Huron. *Journal of Great Lakes Research* 21(4):558-566.
- MacIsaac, H. J., W. G. Sprules, O. E. Johannsson, and J. H. Leach. 1992. Filtering impacts of larval and sessile zebra mussels (*Dreissena polymorpha*) in western Lake Erie. *Oecologia* 92:30-39.
- Mackie, G. L. 1991. Biology of the exotic zebra mussel, *Dreissena polymorpha*, in relation to native bivalves and its potential impact in Lake St. Clair. *Hydrobiologia* 219:251-268.
- Mikheyev, V. P. 1967. [Filtration nutrition of the Dreissena.] (In Russian). Trudy vsesoyuzogo Nauchno-issledovatelnogo Instituta Prudovogo Rybnogo Khoziaistva 15:117-129. [not seen]
- Miura, T. and T. Yamashiro. 1990. Size selective feeding of *Anodonta* calipygos, a phytoplanktivorous freshwater bivalve, and viability of egested algae. Japanese Journal of Limnology 51(2):73-78.

- Nalepa, T. F. 1994. Decline of native unionid bivalves in Lake St. Clair after infestation by the zebra mussel, *Dreissena polymorpha*. *Canadian Journal of Fisheries and Aquatic Sciences* 51:2227-2233.
- Paterson, C. G. 1984. A technique for determining apparent selective filtration in the fresh-water bivalve *Elliptio complanata* (Lightfoot). *The Veliger* 27(2):238-241.
- Patterson, M. A., B. C. Parker, and R. J. Neves. 1997. Effects of quarantine times on glycogen levels of native freshwater mussels (Bivalvia: Unionidae) previously infested with zebra mussels. *American Malacological Bulletin* 14(1):75-79.
- Saraceni, C. and D. Ruggiu. 1969. Techniques for sampling water and phytoplankton. In: A Manual on Methods for Measuring Primary Production in Aquatic Environments, R. A. Vollenweider, ed. pp. 5-7. Blackwell Scientific Publications, Oxford.
- Schloesser, D. W. and T. F. Nalepa. 1994. Dramatic decline of unionid bivalves in offshore waters of western Lake Erie after infestation by the zebra mussel, *Dreissena polymorpha. Canadian Journal of Fisheries and Aquatic Sciences* 51:2234-2242.
- Schloesser, D. W., T. F. Nalepa, and G. L. Mackie. 1996. Zebra mussel infestation of unionid bivalves (Unionidae) in North America. *American Zoologist* 36:300-310.
- Shumway, S. E., T. L. Cucci, R. C. Newell, and C. M. Yentsch. 1985. Particle selection, ingestion, and absorption in filter-feeding bivalves. *Journal of Experimental Marine Biology and Ecology* 91:77-92.
- Silverman, H., E. C. Achberger, J. W. Lynn, and T. H. Dietz. 1995. Filtration and utilization of laboratory-cultured bacteria by Dreissena polymorpha, Corbicula fluminea, and Carunculina texasensis. Biological Bulletin 189:308-319.
- Sprung, M. and U. Rose. 1988. Influence of food size and food quantity on the feeding of the mussel *Dreissena polymorpha*. *Oecologia* 77:526-532.
- Strayer, D. L. and L. C. Smith. 1996. Relationships between zebra mussels (Dreissena polymorpha) and unionid clams during early stages of the zebra mussel invasion of the Hudson River. Freshwater Biology 36(3):771-779.
- Ten Winkle, E. H. and C. Davids. 1982. Food selection by Dreissena polymorpha Pallas (Mollusca: Bivalvia). Freshwater Biology 12:553-558.
- Wu, L. and D. A. Culver. 1991. Zooplankton grazing and phytoplankton abundance: an assessment before and after invasion of *Dreissena* polymorpha. Journal of Great Lakes Research 17 (4):425-436.

Date of manuscript acceptance: 6 March 1998



Parker, B C, Patterson, M A, and Neves, R J. 1998. "Feeding Interactions Between Native Freshwater Mussels (Bivalvia : Unionidae) and Zebra Mussels (Dreissena Polymorpha) in the Ohio River." *American malacological bulletin* 14, 173–179.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/172622</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/143309</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse Copyright Status: In Copyright. Digitized with the permission of the rights holder Rights Holder: American Malacological Society License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.