

LIST OF COLEOPTERA COLLECTED AT MASSETT, QUEEN  
CHARLOTTE ISLANDS, B. C.

BY REV. J. H. KEEN.

*(Continued from page 172.)*

2931. *Ptenidium pullum*, Makl.—Not rare. In grass tufts in winter.
2951. *Trichopteryx xanthocera*, Matth.—Rare. Under log in February.
2952. *Trichopteryx parallelipeda*, Matth.—Rare. In grass tufts in February.
2953. *Trichopteryx diffinis*, Matth.—Common in November in seed pods of foxglove.
- Trichopteryx* (?).—Numerous.
3058. *Coccinella 9-notata*, Hbst.—Abundant in summer. Emerge from pupal state during July.
3262. *Coxelus pacificus*, Horn.—Rare under bark.
3315. *Pediacus depressus*, Hbst.—Ground ; 7th May, 1891.
3316. *Cucujus puniceus*, Mann.—Occurs sparingly under bark. Flies in May.
3348. *Dendrophagus glaber*, Lec.—Not rare ; under loose bark of fences.
3363. *Henoticus serratus*, Gyll.—Abundant in decaying leaves and on skins.
3366. *Cryptophagus*, sp.—One only taken on dry skin.
3382. *Atomaria planulata*, Makl.—Four only, taken in rotten grass in July.
3384. *Atomaria Kamtschatica*, Mots.—In moss in February.
- Atomaria*, n. sp.—Window ; one specimen taken.
3486. *Hister foedatus*, Lec.—One only taken, under rotten crab in garden in June. [Four taken subsequently.]
- Pyromalus mancus*, Casey.—Two taken *in cop.*, March 30th, in crevice of bark of spruce stump.
3662. *Brachypterus troglodytes*, Murray.—Numerous on nettles in May and June.
3699. *Epuræa ambigua*, Mann.—Abundant in April under loose bark of felled spruce.
3707. *Epuræa fulvescens*, Horn.—Not numerous. Berry bushes in June.
3709. *Epuræa truncatella*, Mann.—Common in July, under rotten turnips.
3727. *Omosita colon*, L.—Common during summer on bones.
3765. *Rhizophagus sculpturatus*, Mann.—Not rare, under loose bark.
3767. *Rhizophagus dimidiatus*, Mann.—Under loose bark.
3768. *Rhizophagus brunneus*, Horn.—One only in June on dry log.



3779. *Stephostethus liratus*, Lec.—Common on dry skins and leaves.  
*Lathridius costicollis*, teste Casey.
3821. *Corticaria herbivagans*, Lec.—Rare. In moss in February; in rotten grass in April.
3848. *Peltis pippingskoeldi*, Mann.—One only in July.
3875. *Peltastica tuberculata*, Mann.—Common in larder, on bread, etc.
3880. *Amphicyrta simplicipes*, Mann.—Occasionally in moss at tree roots in winter.
3882. *Simplocaria nitida*, Mots.—Numerous in moss at tree roots all winter.
3884. *Pedilophorus acuminatus*, Mann.—Not rare. In moss during winter.
3960. *Heterocerus tristis*, Mann.—Common in summer on green patches covered by tide at high water.
3988. *Eucinetus infumatus*, Lec.—Occasionally in water-bits. Once occurred in large numbers under loose bark of decaying spruce (in January), where larvæ, pupæ and adults were all represented.
3990. *Eucinetus testaceus*, Lec. (pale specimens)—Occurred with above-mentioned *Eucinetus infumatus* colony.
4065. *Epiphanis cornutus*, Esch.—Rare. On skins in July.
4152. *Cryptohypnus musculus*, Esch.—Rare. Under driftwood.
4218. *Elater nigrinus*, Payk.—Occasionally beaten from bushes under loose bark, 17 May, 1891.
4266. *Megapenthes stigmosus*, Lec.—Common during August.
4391. *Athous ferruginosus*, Esch.—Abundant in July and August.
4433. *Corymbites resplendens*, Esch.—Not rare. On wing during May.
4441. *Corymbites furtivus*, Lec.
4451. *Corymbites caricinus*, Germ.—Frequent in June on spruce.
4451. *Corymbites caricinus*, Germ., var. *umbricola*, Esch.—Common on wing from May onwards. Seen *in cop.*, May 19.  
*Corymbites spectabilis*, Mann.—One only taken in July on wing.
4546. *Throscus validus*, Lec.—Common in windows in June. Never taken out of doors.
4785. *Eros simplicipes*, Mann.—Occasional. Flying during May.
4901. *Podabrus piniphilus*, Esch.—Common.
4919. *Silis pallida*, Mann.—Commonly swept from grass in June.
4956. *Telephorus divisus*, Lec.—Frequent in May and June on green patches covered by tide at high water.



5274. *Hadrobregmus gibbicollis*, Lec.—Rare. Taken on wing in July.  
*Platycerus Keeni*, Casey.—Only one colony found, under a drift-log on sandy beach in April. A few probably immature specimens still retained a slightly reddish tinge. About a dozen were taken.
5467. *Ægialia cylindrica*, Esch.—Numerous under driftwood in May.
5471. *Ægialia crassa*, Lec.—Under log on sandy beach, April 20th. Numerous in summer, crawling on sand hills.
5476. *Psammodius cælatus*, Lec.—Frequent under driftwood on beach.
5523. *Aphodius ruricola*, Mels.—Common in excrement.
5644. *Hoplia Sackenii*, Lec., var.—One specimen only, found on ground.
5970. *Asemum atrum*, Esch.—Not common. Under bark and on wing in May.
5983. *Opsimus quadrilineatus*, Mann.—Found under bark in January.  
*Opsimus quadrilineatus* (black var.).—Not uncommon; under bark of spruce fences all winter.
6232. *Rhagium lineatum*, Oliv.
6299. *Leptura oblitterata*, Hald.
6350. *Leptura Behrensii*, Lec.—Wing-cases only found. February 18; under loose spruce bark.
6367. *Plectrura spinicauda*, Mann.—Common on spruce during summer. Varies much in size.
7226. *Phellopsis porcata*, Lec.—Rare. Two specimens only taken; one under bark, the other in a window.
7501. *Phaleria globosa*, Lec.—Numerous in April, crawling on sand.
7584. *Ægialites debilis*, Mann.—Several taken together with the larvæ.  
*Marolia Holmburghii*.—Common from March onwards on spruce.
7723. *Rhinosimus viridiæneus*, Rand.—Frequently beaten from spruce and other trees in April.
7728. *Ditylus quadricollis*, Lec.—Common under logs on sandy beach in June and July.
7770. *Anaspis rufa*, Say.—Common during summer on Umbelliferae.
7997. *Dendroides ephemeroïdes*, Mann.—Under alder bark. Not rare. Emerges from pupal form in July.
8288. *Sciopithes obscurus*, Horn.—Invariably from trees; never on the ground. (See 8297.) Common during summer. Beaten from berry bushes.



8297. *Geoderces melanothrix*, Kirby.—Under chips on the ground in great numbers and seldom on trees. (See 8288.)
8353. *Trichalophus didymus*, Lec.—Not common. Under logs and in rotton grass.
8470. *Emphyastes fucicola*, Mann.—Occurs but seldom, but then in great numbers. Under logs half buried in sand, during spring.
8471. *Plinthus carinatus*, Boh.—Our commonest weevil. Under logs.
8473. *Pissodes costatus*, Mann.—Not common. Beaten from spruce in April.
8598. *Phycocoetes testaceus*, Lec.—Not rare. Under drift-logs on beach in spring.
8599. *Trachodes ptinoides*, Germ.—Common. Under drift-logs on beach in spring. Seen *in cop.*, June 2nd.
8600. *Trachodes 4-tuberculatus*, Mots.—Not common. Found in July. Occasionally in moss at tree roots in winter.
8601. *Trachodes horridus*, Mann.—Common, under logs.
9042. *Elassoptes marinus*, Horn.—Found in extensive colonies under drift-logs in May.
9044. *Rhyncolus brunneus*, Mann.—Not rare ; in rotten wood.  
*Rhyncolus*, n. sp.—Rare ; under driftwood on sandy beach in May and June.
9074. *Pityophthorus nitidulus*, Mann.—Not rare.
9093. *Xyloterus bivittatus*, Kirby.—Occurs sparingly on newly-cut logs and in moss at tree roots in winter.
9116. *Dryocoetes autographus*, Ratz.—Under spruce bark.
9119. *Xylocleptes concinnus*, Mann.—Very numerous in newly-fallen timber.
9142. *Micracis hirtella*, Lec.—Not rare.
9165. *Hylesinus sericeus*, Mann.—Not uncommon in dead bark.
9181. *Dendroctonus rufipennis*, Kirby, var. *obesus* (black form).—Fairly numerous in bark of newly-cut timber.
9187. *Dolurgus pumilus*, Mann.—Common in spruce bark.  
*Hylurgops glabratus*.—Common under spruce.
9373. *Euscaphurus saltator*, Casey.—Common on under side of logs in damp places. [3987. *Dascyllidæ*.]



Keen, John Henry. 1895. "List of Coleoptera collected at Massett, Queen Charlotte Islands, B. C." *The Canadian entomologist* 27, 217-220.

<https://doi.org/10.4039/Ent27217-8>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/22096>

**DOI:** <https://doi.org/10.4039/Ent27217-8>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/14798>

**Holding Institution**

MBLWHOI Library

**Sponsored by**

MBLWHOI Library

**Copyright & Reuse**

Copyright Status: NOT\_IN\_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.