

(detritophagous and mud suckers) are nose feeders (Table 1.)

Based on the observation of the major gut contents and food preference under normal and, abnormal situations, the various food items may be described as:

1. Basic food - major part of gut contents throughout the year.
2. Secondary food - frequent in gut contents, but lesser than basic foods.
3. Obligatory food - forced to take under stress and food scarcity.
4. Incidental food - of rare occurrence.

Reduction in availability of 'preferred' prey resources

Degradation of favourable feeding sites leads to adverse qualitative and quantitative impacts on the growth of planktonic and benthic communities. This causes in turn serious disruption of the food chain and the energy cycle in the early phases of the life cycle of omnivorous, herbi-omnivorous, carni-omnivorous and carnivorous fish species. Food availability, the nature of feeding grounds and stimuli-feeding responses are less compatible with the adaptations/specialisations for torrential rapids in the hillstreams, particularly in case of bottom dwellers and feeders; the water current

has played a significant role in their evolution.

Alterations in water quality are also brought about by the addition of silt, explosives, large rocks (a result of dam/barrage construction) as well as irrational fishing methods.

ACKNOWLEDGEMENTS

We thank Prof. Asha Chandola-Saklani, Head, Department of Zoology, HNB Garhwal University, for valuable discussions, Prof. M.K. Chandrashekar and Prof. T.J. Pandian, School of Biological Sciences, Madurai Kamaraj University, kindly spared their time to give valuable suggestions. During first author's visit to the Department of Animal Behaviour, School of Biological Sciences, under the DST's 'SERC Visiting Fellowship 1994-95' (No. SR/VS/033), valuable help was granted.

January 19, 1999

N. SINGH

Zoology Department,
HNB Garhwal University,
Srinagar, Garhwal 246 174.

R. SUBBARAJ

Department of Animal Behaviour,
School for biological Sciences,
Madurai Kamaraj University,
Madurai 625 021, Tamil Nadu, India.

REFERENCES

- ALEEV Y. G. (1969): Functional and gross morphology in fishes (Israeli Programme for Scientific Translation, Jerusalem).
- BADOLA, S. P. (1993): Ecological studies on the ichthyofauna of some freshwater resources of Garhwal region. Ph. D. thesis, HNB Garhwal University, Srinagar, Garhwal.
- DAS, S. M. & S.K. MOITRA (1963): *Ichthyologica* 2: 107.
- DAS, S. M. & S. K. MOITRA (1965): *Ichthyologica* 4: 107.
- LARKIN, P. A. (1979): In: Fisheries management, edited by H Clepper (Sport Fishing Institute, Washington).
- NIKOLSKY, G. V. (1963): The ecology of fishes (Academic Press, London).

22. A SUPPLEMENTARY LIST OF THE HOST-PLANTS OF INDIAN LEPIDOPTERA

Indian Lepidoptera are comparatively well known. The early stages and biology of all species of economic importance are known, but little emphasis has been placed on the remaining

species. These constitute the vast majority and are of significance in bio-diversity studies.

The opportunistic rearing of eggs from gravid females and larvae discovered in the field

over a period of several years resulted in the discovery of the following, hitherto unreported, hostplants. This work was carried out in the Kumaon Himalaya in northern India, at an elevation of 1500 m above msl.

The taxonomy of several groups of moths is in need of review. In cases such as the *Spilarctia* Butler species of the *sagittifera* Moore group (Arctiidae), and what was previously the *Dasychira* Steph. genus (Lymantridae), we have not assigned specific status.

Rosa sp., mentioned as the hostplant of *Eterusia leptalina* Kollar, *Dasychira inclusa* Walker and *Spilarctia multiguttata* Walker, are hybrid tea roses rather than good species. In some cases, hostplants accepted in one part of the insect's range are refused in other parts. Thus, freshly emerged larvae of *Ambulyx liturata* Butler (Sphingidae) did not accept *Quercus leucotrichophora* A. Camus and *Q. floribunda* Lindley ex A. Camus (Fagaceae) in Kumaon, although it has been bred on *Quercus* Linn. in China by Mell (Bell and Scott 1937).

The preference of most local Arctiinae for *Pouzolzia zeylanica* (Linn.) Bennet & Brown, *Setaria megaphylla* (Steud.) Dur. & Schinz, and *Plantago major* Linn. is of interest. Many well known European Arctiinae are extremely polyphagous, the larvae having accepted, in addition to the known hostplants, such diverse items as potatoes, apples and even bread! The same cannot be said of the Himalayan species, except perhaps members of the *Spilarctia casigneta* group. We did not get the opportunity to try *Setaria megaphylla* on *Estigmene imbuta* Walker, but there seems a likelihood that it will accept it as readily as *Pouzolzia* Gaud. It, however, did not accept *Plantago major*.

Gardner (quoted by Sevastopulo 1949) notes that the larvae of *Polytela gloriosae* Fabr. (Noctuidae) feed on Liliaceae and Amaryllidaceae, species of *Zephyranthes* Herb. being specially favoured. In our experience, they much prefer *Gloriosa superba* Linn. to *Zephyranthes carinata* Herb., for they will not touch the latter so long as

even a stem of the former is available. We have also bred them on *Zephyranthes* Herb.

All the following bred specimens are in our collection. We have followed Barlow (1982) in the arrangement of moth families.

Lepidoptera Species	Host Plant Species
Family: Zygaenidae	
<i>Tripanophora</i>	<i>Camellia sinensis</i>
<i>semihyalina</i> Kollar	(Linn.) Kuntze (Theaceae)
	<i>Wisteria sinensis</i> (Sims.)
	DC (Leguminosae)
	<i>Pelargonium</i> L'Herit
	(Geraniaceae)
<i>Eterusia leptalina</i> Kollar	<i>Pyrus communis</i> Linn.
	(Rosaceae)
	<i>Rosa</i> sp. (Rosaceae)
<i>Agalope bifasciata</i> Hope	<i>Crataegus crenulata</i>
	G. Koch (Rosaceae)
Family: Limacodidae	
<i>Darna ?cotesi</i> Swinhoe	<i>Cyperus paniceus</i>
	(Rottb.) Boeck.
	(Cyperaceae)
Family: Bombycidae	
<i>Bombyx huttoni</i>	<i>Morus nigra</i> Linn.
Westwood	(Moraceae)
Family: Sphingidae	
<i>Dolbina inexacta</i> Walker	<i>Olea glandulifera</i> Wall.
	ex DC (Oleaceae)
Family: Notodontidae	
<i>Chadisra bipars</i> Walker	<i>Grewia optiva</i>
	J.R. Drummond
	ex Burret (Tiliaceae)
Family: Arctiidae	
<i>Spilarctia</i> sp. of the	<i>Dioscorea bulbifera</i>
<i>sagittifera</i> group	Linn. (Dioscoreaceae)
	<i>Cuscuta reflexa</i> Roxb.
	(Convolvulaceae)
	<i>Strobilanthes</i>
	<i>dalhousianus</i> (Nees)
	C.B. Clarke (Acanthaceae)
	<i>Plantago major</i> Linn.
	(Plantaginaceae)

MISCELLANEOUS NOTES

Lepidoptera Species	Host Plant Species	Lepidoptera Species	Host Plant Species
	<i>Pouzolzia zeylanica</i> (Linn.) Bennet & Brown (Urticaceae)	<i>Euproctis anguligera</i> Butler	<i>Glochidion velutinum</i> Wight. (Euphorbiaceae)
	<i>Setaria megaphylla</i> (Steud.) Dur. & Schinz (Graminae)	<i>Dasychira inclusa</i> Walker	<i>Quisqualis indica</i> Linn. (Combretaceae)
<i>Spilarctia multiguttata</i> Walker	<i>Rosa</i> sp. (Rosaceae)	<i>Dasychira</i> sp.	<i>Rosa</i> sp. (Rosaceae)
	<i>Dioscorea bulbifera</i> Linn. (Dioscoreaceae)	<i>Ilema nigrigula</i> Walker	<i>Bauhinia vareigata</i> Linn. (Leguminosae)
<i>Estigmene imbuta</i> Walker	<i>Pouzolzia zeylanica</i> (Linn.) Bennet & Brown (Urticaceae)	Family: Agaristidae	
		<i>Episteme aduatrix</i> Kollar	<i>Dioscorea bulbifera</i> Linn. (Dioscoreaceae)
<i>Estigmene quadriramosa</i> Kollar	<i>Plantago major</i> Linn. (Plantaginaceae)	Family: Noctuidae	
	<i>Taraxacum</i> sp. (Compositae)	<i>Cocytodes coerulea</i>	<i>Bohemeria platyphylla</i> D. Don (Urticaceae)
	<i>Pouzolzia zeylanica</i> (Linn.) Bennet & Brown (Urticaceae)	<i>Thysanoplusia orichalcea</i> Fabricius	<i>Lepidium virginicum</i> Linn. (Cruciferae)
<i>Pericallia galactina</i> von. d. Hoev	<i>Pouzolzia zeylanica</i> (Linn.) Bennet & Brown (Urticaceae)	<i>Polytela gloriosae</i> Fabricius	<i>Gloriosa superba</i> Linn. (Liliaceae) preferred over <i>Zephyranthes</i> <i>carinata</i> Herbet (Amaryllidaceae)
	<i>Setaria megaphylla</i> (Steud.) Dur. & Schinz (Graminae)	Family: Epiplemidae	
<i>Pericallia imperialis</i> Kollar	<i>Plantago major</i> Linn. (Plantaginaceae)	<i>Epiplema reticulata</i> Moore	<i>Jasminum dispernum</i> Wallich (Oleaceae)
	<i>Pouzolzia zeylanica</i> (Linn.) Bennet & Brown (Urticaceae)	Family: Pyralidae	
	<i>Setaria megaphylla</i> (Steud.) Dur. & Schinz (Graminae)	<i>Agathodes ostentalis</i> Huebner	<i>Erythrina suberosa</i> Roxb. (Leguminosae)
<i>Callimorpha plagiata</i> Walker	<i>Pouzolzia zeylanica</i> (Linn.) Bennet & Brown (Urticaceae)	Family: Pieridae	
		<i>Pontia daplidice</i> Linne	<i>Lepidium virginicum</i> Linn. (Cruciferae)
<i>Macrobrochus gigas</i> Walker	Lichens	<i>Artogeia canidia</i> Sparrman	<i>Lepidium virginicum</i> Linn. (Cruciferae)
Family: Lymantridae		Family: Nymphalidae	
<i>Euproctis latifascia</i> Walker	<i>Quercus</i> <i>leucotrichophora</i> A. Camus (Fagaceae)	<i>Symbrenthia lilaea</i> Hewitson	<i>Bohemeria platyphylla</i> D. Don (Urticaceae)
	<i>Glochidion velutinum</i> Wight. (Euphorbiaceae)	<i>Precis iphita</i> Cramer	<i>Aechmanthera tomentosa</i> Nees (Acanthaceae)
<i>Euproctis plagiata</i> Walker		<i>Pareba issoria</i> Huebner	<i>Debregeasia longifolia</i> (Burm. f.) Wedd. (Urticaceae)

ACKNOWLEDGEMENTS

We thank Professor Y.P.S. Pangtey and Dr. B.S. Kalakoti, Department of Botany, Th. D.S.B. College, Nainital, who kindly identified some of the plants, and Dr. Poonam Mehrotra of the Department of Ecology of the same institution for her kind help with the bibliography.

October 27, 1999

PETER SMETACEK
RAJANI SMETACEK
Jones Estate,
Bhimtal,
Nainital, Pin 263 136,
Uttar Pradesh,
India.

REFERENCES

- BARLOW, H.S. (1982): An Introduction to the Moths of South East Asia. Malayan Nature Soc., Kuala Lumpur.
- BELL, T.R.D. & F.B. SCOTT (1937): The Fauna of British India including Ceylon and Burma, Moths, Vol. V, Spingidae, Taylor & Francis, London.
- SEVASTOPULO, D.G. (1949): A supplementary list of the foodplants of the Indian Bombycidae, Agaristidae and Noctuidae. *J. Bombay nat. Hist. Soc.* 48: 265-276.

23. ON THE PREDATION OF THE GIANT REDEYE *GANGARA THYRSIS* (FABRICIUS) (FAMILY : HESPERIIDAE; ORDER : LEPIDOPTERA)

The Giant Redeye *Gangara thyrsis* Family HesperIIDae is not an uncommon butterfly in Bangalore. It is often seen in gardens around its food plants — *Areca lutea*, *Cocos nucifera* and other palms.

Observations on the predators of the Giant Redeye were made on nine *Areca lutea* plants ranging in height from 1-4 m, and frequented by these insects. It was observed that the bonnet macaque (*Macaca radiata*) and the house crow (*Corvus splendens*) fed on the larvae and pupae of the Giant Redeye.

One individual of a troop of bonnet macaques which visited the premises where observations were made, systematically searched all the palms for larvae and pupae. The macaque searched the leaves rolled up by the larvae, opened them, and ate the larvae (which have long, loosely attached, white thread-like outgrowths amidst which are red spots).

Similarly, the macaque opened the tubes made of palm fronds which conceal the pupae and ate the pupae.

A house crow which visited the premises seemed to have noticed a pupa of the Giant Redeye. It gave up its efforts to procure the pupa as it was unable to balance itself on the slender palm fronds. A good half hour had elapsed before the bird returned and perched on the neighbouring *Colocasia* sp.(?) growing amidst the palms. From the new perch, it successfully ripped open the tube and swallowed the pupa whole.

These are probably new records of predators of the Giant Redeye.

May 25, 1999

S. KARTHIKEYAN
24, Opp. Banashankari Temple,
Shakambarinagar,
8th Block Jayanagar P.O.,
Bangalore 560 082, Karnataka, India.

24. MATING BEHAVIOUR OF THE COMMON MORMON *PAPILIO POLYTES* (FAMILY: PAPILIONIDAE)

During February 1998, I was studying the metamorphosis of different species of Papilionidae and Nymphalidae in my home laboratory. The Common Mormon (*Papilio*

polytes) was one of the species reared successfully. After a pupal period of ten days, a female Common Mormon emerged from its chrysalis at about 0900 hrs. The Common



Smetacek, Peter and Smetacek, Rajani. 2000. "A Supplementary List of the Host plants of Indian Lepidoptera." *The journal of the Bombay Natural History Society* 97, 157–160.

View This Item Online: <https://www.biodiversitylibrary.org/item/189503>

Permalink: <https://www.biodiversitylibrary.org/partpdf/155288>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.