MISCELLANEOUS NOTES

ACKNOWLEDGEMENTS

We are thankful to the Indian Council of Agricultural Research, New Delhi, for financial support, and to Dr. D.N. Yadav, Officer-incharge, for encouragement.

November 9, 1998 AESHITA MUKHERJEE B.M. PARASHARYA

AINP on Agricultural Ornithology, Gujarat Agricultural University, Anand Campus, Anand 388 110, Gujarat, India.

REFERENCES

ALI, S. & S.D. RIPLEY (1983): Handbook of the Birds of India and Pakistan. Compact edition. Oxford University Press, New Delhi.

Hancock, J. & J. Kushlan (1984): The Herons Handbook. Croom Helm, London.

8. PALLAS'S FISHING EAGLE *HALIAEETUS LEUCORYPHUS* (PALLAS) PIRATES FISH FROM AN OTTER *LUTRA LUTRA* (LINN.)

I observed four common otters (Lutra lutra) on boulders in the middle of the River Manas. The otters were fishing; they would come out of the water with the fish and feed while sitting on the boulder. A Pallas's fishing eagle was noticed circling above the otters, repeatedly trying to snatch the fish. It made two or three futile attempts by diving at the otter. After a while, the bird made another attempt. This time, the otter was frightened and dropped the fish, which was neatly picked up by the eagle in one swoop. The Pallas's fishing eagle is regularly known to pirate fish from other fish eating birds, or smaller eagles, but very few have been recorded robbing otters or other mammals

(Prakash 1989).

ACKNOWLEDGEMENTS

I am thankful to Dr. Vibhu Prakash, Principal Scientist, BNHS, Bharatpur for his guidance and to the Pigmy Hog Conservation Programme personnel for their help during field trips.

October 8, 1998 BIBHUTI PRASAD LAHKAR
c/o Niramoy Pharmacy
Near Ganesh Mandir, Ganeshguri Tiniali,
Dispur, Guwahati 781 006,
Assam, India.

REFERENCE

PRAKASH, V. (1989): General Ecology of Raptors in Keoladeo National Park. Ph. D. thesis submitted to the University of Bombay.

9. GROUP SIZE AND VIGILANCE IN INDIAN PEAFOWL *PAVO CRISTATUS* (LINN.), FAMILY: PHASIANIDAE

Flocking in birds is considered to be an adaptive social strategy against danger, and the evolution of gregariousness is an effective solution to the problem of the individual's vigilance load (Dimond and Lazarus 1974). Birds

feeding in large flocks are safer (in terms of number of birds alert at any moment) and have more time to feed (in terms of the proportion of its time that an individual spends alert). It is known that the time budget of a species is related



Lahkar, Bibhuti Prasad. 2000. "8. Pallas's Fishing Eagle Haliaeetus Leucoryphus (Pallas) Pirates Fish from An Otter Lutra Lutra (Linn.)." *The journal of the Bombay Natural History Society* 97, 425–425.

View This Item Online: https://www.biodiversitylibrary.org/item/189503

Permalink: https://www.biodiversitylibrary.org/partpdf/155350

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: http://creativecommons.org/licenses/by-nc/3.0/ Rights: https://www.biodiversitylibrary.org/permissions/

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.