NEW DESCRIPTIONS

A NEW SPECIES OF *OLIGODON* FROM THE PALNI HILLS, SOUTH INDIA (SERPENTES: COLUBRIDAE)¹

ROMULUS WHITAKER AND SHEKAR DATTATRI² (With a plate)

INTRODUCTION

While on a herpetological survey of the Palni Hills, South India, we collected a hitherto unreported species of kukri snake of *Oligodon* along with several other interesting reptiles and amphibians. It is named here as:

Oligodon nikhili sp. nov.

Description:

Seven supralabials, the third and the fourth touching the eye; the sixth rising slightly above the labial border; one anterior temporal; internasals present, loreal absent; nostril in elongated nasal completely divided by a vertical suture. Scales around mid body in fifteen rows. Ventrals: 144; Caudals: 33, paired; anal divided.

Hemipenis extending to the 13th caudal plate; proximal quarter of the organ smooth, the next quarter with equal sized spines and the distal half flounced, the flounces edged with numerous small spines; three prominent longitudinal folds present.

Colour:

Ground colour light brown, each scale with a minute black speck. Two chocolate brown dorso-lateral stripes 1.5 scales wide extending from the posterior end of the characteristic

¹ Accepted September 1982.

² Madras Snake Park Trust, Madras 600 022.

chevron head pattern to the tip of the tail. Two indistinct lateral lines on scale row three, extending from above the fifth ventral to the anal plate; belly whitish with prominent ventero-lateral speckling and two broken lines of black spots.

Holotype:

Museum of the Madras Snake Park Trust, an adult male from Tiger shola, 1500 mts., Palni Hills, Tamilnadu, South India, 20.9.81, collected by Romulus and Zahida Whitaker and Shekar Dattatri. No other specimens reported.

Measurements:


Snout-vent: 350 mm Vent-tail: 73 mm Total length: 423 mm Head length: 11 mm Head width: 9 mm Neck width: 8 mm Diameter of eye: 2 mm

Scalation:

Ventrals: 144 Caudals: 33 Scale rows: 17-15-15

DIAGNOSIS AND DISCUSSION

The cylindrical body with smooth scales, presence of the nostril in an elongated nasal, one pre- and two post-oculars, circular pupi! of eye, typical head shape and pattern estaJ. BOMBAY NAT. HIST. Soc. 79 Whitaker & Dattatri: Oligodon nikhili sp. nov.



Oligodon nikhili sp. nov. from Palni Hills, Tamil Nadu. (Photos: Rom Whitaker)

blish beyond doubt that this snake belongs to the Genus Oligodon.

Of the 33 species of this Genus recorded from the oriental region, eleven other species have 15 scale rows at mid-body. Tabulated below are the differences exhibited by the holotype compared to the most similar forms. collected five species uropeltids (Uropeltis, Platyplectrurus and Teretrurus spp.), one species of Xylophis, several forest frogs and lizards. The new kukri snake was found by the three year old son of one of us (R.W.), Nikhil, who was left at the roadside to play at "snake hunting". The snake was hiding behind a piece

Character		O. taeniolatus	brevicaudata	-1940	erythrorachis	sublineatus	new species
1.	Supra labials:	7	7	- statt	7	7	7
2.	Scale rows:	15	15		15	15	15
3.	Ventrals:	158-218	158-173		154	134-161	144
4.	Caudals:	29-56	25-29		46	23-37	33
5.	Loreals:	present	absent		absent	present	absent
6.	Inter nasals:	present	absent		present	present	present
7.	Distribution	India	Western		Assam	Sri Lanka	Western
		Sri Lanka	Ghats				Ghats

From the characters enumerated above, it is evident that the new species closely resembles *O. erythrorachis* described by Wall (1910) based on a single specimen collected in Assam. However, it differs from the latter in coloration and in the ventral and subcaudal counts.

The next most closely resembling species is O. sublineatus described from Sri Lanka. However, O. sublineatus differs in having complete head shields and in the hemipenis being spinose throughout, the spines being almost uniform in size and regularly arranged. This is not the case with the new species.

Habitat:

Tiger shola, where our specimen was collected, is a medium altitude rain forest (1500 mts. above MSL) a few kilometres below Shembaganur on the Kodaikanal ghat road. It remains one of the few intact pockets of 'shola' forest in the area and on the same trip we of cement on the roadside retaining wall and fell at the child's feet when he removed the loose slab of cement.

CONCLUSION

When we started out on this trip, a new species of snake was the last thing we expected. Our discovery shows only too clearly that a great deal lies hidden in the complex and unique forest hill ranges of the Western ghats. It is appalling to see the destruction man has wrought in many of these areas. Hill slopes that were lush with vegetation in the senior author's younger days are deforested, barren, hot and eroded today. What we are losing in terms of our fauna and flora is anybody's guess.

ACKNOWLEDGEMENT

We are grateful to the Madras Snake Park Trust for funding the survey.

REFERENCE

WALL, F. (1910): A new snake from Assam., Oligodon erythrorhachis. J. Bombay nat. Hist. Soc. 19: 923-924.



Whitaker, Romulus and Dattatri, Shekar. 1982. "A new species of Oligodon from the Palni Hills, south India (Serpentes: Colubridae)." *The journal of the Bombay Natural History Society* 79, 630–631.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/191543</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/156930</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse Copyright Status: In Copyright. Digitized with the permission of the rights holder License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.