# WILDLIFE IN BANGLADESH MANGROVE ECOSYSTEM<sup>1</sup>

Mohammad Ali Reza Khan<sup>2</sup>

(With a text-figure)

The wildlife of the mangrove ecosystems of the Sunderbans, Chakoria Sunderbans and other mangrove formations in Bangladesh comprises of about 400 species, including eight amphibians, 50 reptiles, 261 birds and 49 mammals. Several of these are either endangered or vulnerable particularly species restricted to these ecosystems.

## INTRODUCTION

A great variety of wildlife, defined here as all organisms from Amphibia to Mammalia, has enriched the mangrove ecosystems of the Sunderbans, Chakoria Sunderbans and tidal forests of Bangladesh. It is quite evident from the century-old as well as current literature, *viz.*, Baker 1887, O'Malley 1908, Law 1945, 1948a & b, 1954, Mitra 1957, Mukherjee 1959, Mandal 1964, Acharji & Mukherjee 1964, Mukherjee & Gupta 1965, Mountfort 1969, Biswas 1973, Hendrichs 1975, Mukherjee 1975, Green 1978, Seidensticker & Hai 1978, Gittins 1981, Khan 1981, 1982a & b, Khan & Ahsan 1981, and Khan & Rahman 1982.

Most of the mangrove vegetation of Bangladesh lies within the Sunderbans of Khulna district. This covers 62% of the total Sunderbans of Bangladesh and West Bengal of India, the

<sup>1</sup> Accepted March 1984. The paper was presented in the 'Regional symposium on the mangrove, coral and island ecosystems' jointly organised by SACEP/ESCAP and the Environment Pollution Control Department of Bangladesh at Dhaka in August 1983.

<sup>2</sup> Department of Zoology, University of Dhaka, Dhaka-2, Bangladesh. *Present Address*: Curator, Al Ain Zoo & Aquarium, P. O. Box 1204, Al Ain, Abu Dhabi, U.A.E. latter comprising 38% (Hendrichs 1975). The total area of Bangladesh Sunderbans is about 5800 km<sup>2</sup>, of which 4100 km<sup>2</sup> are land and 1700 km<sup>2</sup> water. Bangladesh Sunderbans have been divided into four forest ranges, 14 blocks and 55 compartments varying in size from 40 to 160 km<sup>2</sup> (Fig. 1). There are some isolated, small patches of both planted and naturally growing mangrove vegetation along the southern parts of the districts of Patuakhali, Barisal, Naokhali and Chittagong, mostly on the inshore and offshore islands and coast. This type of forest also occurs in the Chakoria Sunderbans, 21°45'N and 92°E; Whykeong, 21°05'N and 92°12'E, and Teknaf 21°N and 92°15'E along the River Naaf, bordering Burma; and on the lone coral island of the country - St. Martin's, 23.35°N and 92.22°E (Fig. 1).

From the wildlife point of view, the forests of the Sunderbans of Bangladesh and India were studied by Hendrichs (1975) and Mukherjee (1975), respectively. The Bangladesh Sunderbans lie between 21°31'N to 22°30'N and 89°E to 90°E. Gittins (1981) working with the Rhesus Macaque of the Sunderbans divided the forest into three zones according to the salinity of the surrounding water, but without providing the range of salinity.

(i) the fresh-water zone — to the North and East of a line drawn from Cobadak





Forest station (22°15'N, 89°20'E) in the North to the mouth of Katka Khal (21°50'N, 89°45'E) on the sea face, consisting of 1920 km<sup>2</sup> of *Heritiera fomes* dominated forest:

- (ii) moderately salt-water zone West of the above line to the Malancha River (21°40'N, 89°18'E), consisting of 1324 km<sup>2</sup> of *Excoecaria agallocha* dominated forest: and
- (iii) the salt-water zone West of the Malancha River to the international boundary with India, consisting of 781 km<sup>2</sup> of sparse *E. agallocha* and dense patches of the palm *Phoenix paludosa*.

The plants of the Sunderbans show marked adaptation to life under saline condition and to frequent inundation by the tides. Thus, they have developed succulent leaves, stilt roots, pneumatophores and vivipary. *H. fomes* and *E. agallocha* cover the most part of the Sunderbans. Oryza coarctata, Nipa fruticans and Imperata cylinderica are prevalent on the mudflats.

The Chakoria Sunderbans, 84 km<sup>2</sup>, is dominated by Dalbergia spinosa and Aegialitis rotundifolia. The planted areas and the mangrove formations of the islands, including those along the River Naaf, consist mainly of Sonneratia spp., Avicennia spp. and Acanthus ilicifolius, whereas that of the St. Martin's has stunted Lumnitzera racemosa, Aegialitis rotundifolia and Caesalpinia crista.

The Sunderbans ecosystem presents a unique opportunity of studying wildlife which no other ecosystems viz., the dry and moist deciduous sal forests, semi-evergreen and evergreen forests of the central and eastern Bangladesh, could provide. As far as wildlife is concerned, the Bangladesh Sunderbans is one of the most open forests of all, in spite of the compactness of the trees. This is mainly because innumerable khals (channels) and rivers cut the whole of the Sunderbans into thousands of fragments. The peripheries of all fragmented pieces of forest make it convenient for the wildlife to utilize these open and well-lighted areas, which are to some extent similar to the ecotonal zones of other forest ecosystems.

## METHOD

During 1980 and 1982 I and one of my research fellows paid a total of five visits to the Khulna Sunderbans, where we covered 608 km, including 64 km on foot, 187 km in a dinghy and 357 km in powered boats. About 200 hours spread over 30 days, were devoted to the field observations. A little over 15 days were spent on the St. Martin's Island and 15 days on various other islands, and many visits were made to the mangrove formations along the River Naaf. A research fellow monitored the activities of the Crab-eating Macaque and other wildlife of the River Naaf mangrove formation at Whykeong. I paid one visit to the Bay of Bengal on board the fishing vessel 'Anushandhani' in January 1981 and spent four days in the bay, south of the Khulna Sunderbans.

Most of the wild animals, given in appendix I, were observed by myself and a few by the research fellows. Records of the extinct and some uncommon wildlife have been taken from Mitra (1957), Hendrichs (1975) and Mukherjee (1975). The wildlife were noted either from direct visual observation, from foot-prints, pugmarks, scats on the muddy shores, faecal materials or calls. Nocturnal observations were made with the help of a 4-cell spotlight and headlight of the powered boats. Animals which occurred in the human habitations or forest villages bordering the Sunderbans have also been included in the appendix and in all calculations. The number of such wildlife species probably does not exceed five per cent of the total number of species listed.

The stratification patterns or vertical layering of the wildlife of the various mangrove ecosystems are based on the mode of food gathering and not on the habitat preferences. They have been categorized as aquatic, terrestrial, amphibious, arboreal, arbo-terrestrial and shore-dwellers, depending upon collection of food from water, land, and both, from foliage, foliage and land, and from mud-flats and/or sand-flats. The arboreals included those animals which fed on both plant and animal matter from the trees or shrubs/herbs, and winged insects/animals through aerial pursuits, e.g., flycatchers, bee-eaters, bats, etc.

The wildlife has been classified as carnivores, omnivores and herbivores basing on the natural feeding behaviour. Those found by us or known to be feeding on live or dead animals, from invertebrates to mammals, have been categorized as carnivores. These include both insectivores, piscivores, carrion-feeders and other flesh-eaters. Omnivores include these wildlife which fed almost equally on animal and plant matter. The herbivores either fed on ground vegetation through grazing; on climbers, lianas and shrubs through browzing; or on leaves, fruits, buds and bark of mangrove trees. Wildlife which normally fed on animal matter and casually collected nectar and other such plant matter have been included under carnivores, e.g., drongos, magpie-robins, babblers, etc.

Attempts have been made to provide a population estimate of chital (spotted deer) and rhesus macaque.

### **RESULTS AND DISCUSSION**

The amphibia included eight species belonging to four genera of four families (appendix I). It is dominated by *Rana hexadactyla* — a brackish water species, the toad *Bufo melanostictus* and the tree frog *Rhacophorus maculatus*. During the monsoon, between June and September, the last two species occur almost all over the Sunderbans, except the southern part, facing the sea, The remaining species were noted in the peripheral areas of the Sunderbans.

The reptilian fauna included 10 species each of chelonians and lizards, 29 species of snakes and one species of crocodile. Of these, the common batagur Batagur baska, does not occur outside the Sarankhola Range of the Sunderbans Forest Division. Four species of marine turtles, Chelonia mydas, Caretta caretta, Lepidochelys olivacea and Eretmochelys imbricata, visit the sandy southern islands such as Katka, Supati, Nilkamal, Dubla and Putney. The other turtles occur in the north-eastern rivers and in the villages bordering the forests. The lizards belonged to five genera and four families. The house lizards, Hemidactylus frenatus and H. brooki, wall lizard Gekko gecko, and ring lizard Varanus salvator were found almost everywhere. The snakes belonged to 22 genera and nine families. The keelbacks, water snakes, cobra, sea snakes and pit vipers were commonly found in the Sunderbans, either inside the forest, or in the khals and rivers. Two rare species the rock python Python molurus and king cobra, Opiophagus hannah are represented by fairly good populations in the Sunderbans. The sand boa Eryx conicus, and the wart snake Acrochordus granulatus have been reported by Mukherjee (1975) from the West Bengal part of the Sunderbans. I presumed these to be present in our part too. The whitebellied mangrove snake Fordonia leucobalia, and glossy marsh snake Gerardia prevostiana and the Malacca sea snake. Hydrophis caerulescens have been seen so far in the Sunderbans only.

The estuarine or salt water crocodile Crocodylus porosus, occurs only in the Sunderbans. Although there is a single report of its occurrence in the coast of Cox's Bazar, eastern Bangladesh (Fr. R. W. Timm, pers. comm.). Hendrichs's (1975) report of the gharial Gavialis gangeticus, from the Sunderbans is possibly erroneus as it appears to be a purely freshwater species and no one else has seen it in the Sunderbans. The marsh crocodile, Crocodylus palustris, is possibly extinct from the Sunderbans and it occurs nowhere else in the country, excepting two small captive populations: 4-5 in a tank at Bagerhat, under Khulna district, and three in the only zoo of the country at Dhaka (Khan 1982c).

It is most likely that the reptilian list will expand once an extensive collection of lizards and snakes is made in the Sunderbans.

Of the 261 species of birds, recorded from the mangrove ecosystems, 14 have been taken from Hendrichs (1975) and Mukherjee (1975) and one from a Yale University collection reported by Ripley (1982). Altogether, 180 species of 105 genera and 32 families were non-passerine; 81 species of 50 genera and 19 families passerine. One hundred and sixty three species were resident and 98 migratory. All locally migratory and summer visitors have been considered under the resident category, as they breed either in the Sunderbans or elsewhere in the country. The non-passerines included 107 resident and 73 migratory species

So far 11 species of kingfishers have been reported from Bangladesh (Khan 1982a), of which eight are present in the Sunderbans. Out of these the brownwinged kingfisher Pelargopsis amauroptera and the ruddy kingfisher Halcyon coromandra were found only in the Sunderbans. Two other species, blackcapped and whitecollared kingfishers, Halcyon pileata and H. chloris, do not occur outside the mangrove ecosystems of the Sunderbans, Chakoria Sunderbans, coastal and the St. Martin's islands. The whitebellied and goliath herons, Ardea insignis & A. goliath, white stork Ciconia ciconia, whitebellied sea eagle Haliaeetus leucogaster, oriental hobby Falco severus & Indian skimmer Rhynchops albicollis (reported by Mukherjee 1975), swamp partridge Francolinus gularis, masked finfoot Heliopais personata, parasitic skua Stercorarias parasiticus (found in the Swatch-of-no-ground, at the Bay of Bengal), lesser & large crested terns Sterna bengalensis & S. bergii (reported by Mukherjee 1975), oystercatcher Haematopus ostralegus, avocet Recurvirostra avosetta, European starling Sturnus vulgaris (seen on St. Martin's Island), mangrove whistler Pachycephala grisola and orangebellied flowerpecker Dicaeum trigonostigma have so far been sighted in the mangrove formations.

There are many species of birds which are opportunistic in the sense that they roost inside the Sunderbans and other mangrove areas but gather food from the neighbouring areas, mostly cultivated fields, e.g. most of the mynas, parakeets, doves, pigeons, egrets, etc. These birds breed in the mangrove vegetations also. Many smaller species of passerine birds such as warblers and non-passerine charadriids could not be identified. Netting followed by specimen collection may reveal the presence of several more species, thereby raising the total figure to 300 or so, that means half of the Bangladesh avifauna.

The living mammals of the Sunderbans are

represented by 42 species belonging to 37 genera and 22 families. In addition the crabeating macaque Macaca fascicularis, occurs in the mangrove formations of the River Naaf. The species of mammals which have disappeared in the recent past from the mangrove areas of Bangladesh include onehorned rhinoceros, Rhinoceros unicornis: smaller onehorned rhinoceros, R. sondaicus; wild buffalo Bubalus bubalis; swamp deer, Cervus duvauceli; hog deer, Axis porcinus; and the leopard, Panthera pardus. The royal Bengal tiger, Panthera tigris and chital, Axis axis have disappeared from the whole of Bangladesh, excepting the Kulna Sunderbans. Possibly the biggest populations of tiger, chital, rhesus macaque Macaca mulatta; smooth Indian otter Lutra perspicillata and wild boar Sus scrofa occur in the Sunderbans. The Irrawaddy dolphin, Orcaella brevirostris; shortfinned pilot whale, Globicephala macrorhynchus; finless porpoise Neophocaena phocaenoides; and the Malay dolphin Stenella malayana are rather restricted to the Sunderbans estuary.

### Stratification pattern

In all, 15 species of mammals were terrestrial; 12 arboreal, including 11 species of flying mammals; 7 arbo-terrestrial (crabeating macaque also come under this category); 6 aquatic and two amphibious. All extinct mammals were terrestrial.

About 83 non-passerine species of birds were aquatic, 32 terrestrial, 30 shore-dwellers, 21 arboreal, 11 amphibious and 3 arbo-terrestrial. Twenty passerine species were terrestrial, 44 arboreal and 17 arbo-terrestrial.

Among the reptiles all the chelonians were aquatic, 4 lizards arboreal, two skinks and three monitor lizards were normally terrestrial and casually arboreal, 11 species of snakes were aquatic, 10 terrestrial, 3 arboreal, 3 arboterrestrial and 2 amphibians. The crocodile is amphibious too. The amphibians comprised 6 terrestrial and 2 aquatic species.

Out of the total, 112 species of wildlife of the mangrove ecosystem of Bangladesh gathered food from the aquatic environment, 89 from land, 84 from air and foliage, 30 from mud-, sand-flats, 36 from land, air and foliage and the remaining 17 from both water and land.

## Feeding habits

Of the wildlife recorded from the mangrove ecosystem, 24 mammals were carnivorous, 11 herbivorous and 8 omnivorous; 157 non-passerine birds were carnivorous, 12 omnivorous and 11 herbivorous; 60 passerine species were carnivorous, 13 omnivorous and 8 herbivorous; 46 reptiles were carnivorous, and 4 omnivorous when 7 amphibians were carnivorous and one, *Rana hexadactyla*, almost invariably fed on dragon and damsel flies although some algal materials were found in the stomach. The presence of algal material in the stomach might be merely accidental.

A total of 294 species were carnivorous, including insectivores, piscivores and flesheaters; 38 omnivorous and 30 herbivorous, both folivorous and frugivorous.

## International status

International Union for Conservation of Nature and Natural Resources (IUCN) defines endangered species as those in danger of extinction and whose continued survial is unlikely if the casual factors continue operating. The common batagur, green turtle, olive ridley turtle, hawksbill turtle, rock python, peregrine falcon, royal Bengal tiger and leopard of the mangrove ecosystems are endangered so far as the Red Data Book (RDB) of IUCN is concerned. The RDB has included the estuarine crocodile as a vulnerable species, meaning that this species is believed likely to become extinct in the near future if the adverse casual factors continue operating.

Schedule I of the Convention on International Trade in Endangered Species of Wild Fauna & Flora (CITES) includes 7 turtles and tortoises, Bengal and yellow lizards, rock python, white stork, spotted green shank, Ganges river dolphin, finless porpoise, royal Bengal tiger and leopard cat. Another seven species — ring lizard, marsh crocodile, rhesus macaque, smooth Indian and clawless otters, fishing and jungle cats — are in Schedule II of the CITES.

## Population estimation

Hendrichs.(1975) for the first time attempted estimating the populations of tiger, otter, wild boar, chital, rhesus macaque and some other vertebrates and invertebrates of the Khulna Sunderbans. Gittins (1981), and Khan & Ahsan (1981) attempted estimating the population of the rhesus macaque, while I have made an attempt to provide a rough estimate of the spotted deer (chital).

Hendrichs (1975) estimated the total population of the tiger in the Sunderbans to be 350, otter 20000, wild boar 20000; chital 80000 and rhesus macaque 40000. His estimated density of these were tiger —  $0.1/\text{km}^2$ , otter —  $5/\text{km}^2$ , boar  $5/\text{km}^2$ , chital —  $20/\text{km}^2$  and macaque —  $10/\text{km}^2$ . Gittins (1981) noted 2.6 groups of macaques with 20 individuals in each group, that is a density of  $52/\text{km}^2$ . He has estimated the total population of macaque in  $2274\text{-km}^2$ of natural forests to be 118,248, and another 7972 in 1533-km<sup>2</sup> scrub forest of the Sunderbans. In the scrub forest the density of the macaque was  $5.2/km^2$ .

Ahsan and I covered 608 km linear distance, equivalent to 20.83 km<sup>2</sup> of transects, where we encountered 1.58 groups/km<sup>2</sup> with 17.05 macaques/km<sup>2</sup> and the total population was estimated to be 68,200 in an area of about 4000 km<sup>2</sup>. The population estimation done by the three authorities showed much variation. Our estimation of 68,200 macaques is much nearer to Hendrich's 40,000 than Gittins' 126,220.

In the 608 km of linear distance and 15.2 km<sup>2</sup> of transects which we covered we encountered 200 deer. The density of chital was 13.15/km<sup>2</sup>. The Sunderbans, therefore, supports an estimated 52,600 deer. This is 34.25% less than the estimated population given by Hendrichs (1975). But the difference is close to his 25% range of deviation. This difference is mainly because he devoted more time and covered each of his sampling areas in great detail, which we could not do because of lack of time and logistic support.

### RECOMMENDATIONS

The Government of Bangladesh has already gazetted three areas of the Sunderbans mangrove forest as wildlife sanctuaries. These are the East, West and South Wildlife Sanctuary, consisting of compartments no. 6 ( $54 \text{ km}^2$ ), part of no. 54 ( $90 \text{ km}^2$ ), and part of no. 43 and the whole of no. 44 ( $177 \text{ km}^2$ ) respectively. Unfortunately, these declarations have paid little or no attention to the earlier recommendations made by several expeditions and an enquiry committee meetings by the experts of FAO, IUCN, WWF and the Government Forest Department, *viz.*, Mountfort & Poore 1968, Seidensticker and Hai 1978, and Oliver 1979. Although these recommendations were based on sound ecological considerations.

To save the wildlife from extinction, to stop appreciable changes in the mangrove ecosystem, and to make the already declared sanctuaries meaningful so that these support viable populations of wildlife I strongly recommend that:

- Compartment no. 3 to 8, 11, 12 and 45 be declared as the Sunderbans National Park, incorporating the existing East Sanctuary;
- 2. The remaining portion of the Compartment no. 43 be added to the existing South Sanctuary;
- 3. Instead of West Sanctuary a new one be declared in the North, comprising compartments no. 30 to 33;
- 4. The sanctuaries/national parks be sufficiently manned by technical staff so that they can monitor the faunal and floral changes year round.
- 5. Cheaper tourist facilities be developed to compensate the revenue lost due to discontinuation of forestry practises.

#### ACKNOWLEDGEMENTS

Fr. R. W. Timm CSC, Director, CARITAS, Dhaka has kindly read and corrected the

manuscript. Messers Shah Ali Imam, Conservator, Coastal Circle, Ghulam Habib DFO Khulna, A. K. Fazlul Haque DFO Noakhali Coastal Afforestation, Munshi Anwarul Islam DFO Coastal Afforestation Chittagong, A. Z. M. Shamshul Huda DFO Cox's Bazar and S. Choudhury ADFO Khulna, all the range officers of the four ranges of Khulna, ranges of Noakhali, Chittagong Coastal Afforestation Divisions and Cox's Bazar Forest Division provided me necessary guides and boat and generous hospitality. Directors of SPARRSO and Environment Pollution Control Department; President, Natural History Society of Bangladesh allowed me to take part in the Expeditions to the Bay of Bengal and St. Martin's Island, and SACEP/ESCAP Symposium on the mangrove, coral and island ecosystems. Mr. Noor Mohammed Sarker, Conservator, Wildlife Circle, organised a survey of the islands of Chittagong and Noakhali Coastal Afforestation Divisions, when Mr. Mukhlesur Rahman, JRO, Wildlife Circle; Messers Md. Farid Ahsan, M. A. Wahab, SMA Rashid, all research fellows of the Department of Zoology of Dhaka University offered their assistance in the field work conducted between 1980 and 1983. I thank them all.

#### REFERENCES

ACHARJI, M. N. & MUKHERJEE, A. K. (1964): Report of a collection of snakes from Lower Bengal (Reptilia: Ophidia). J. Zool. Soc. India. 16(1&2): 76-81.

BAKER, E. B. (1887): Sport in Bengal and how, when and where to seek it. Ledger, Smith & Co., London.

BISWAS, R. N. (1973): On the domestication of the otter by fishermen in Bangladesh. J. Bombay nat. Hist. Soc. 70(2): 374.

GITTINS, S. P. (1981): A survey of the primates of Bangladesh. Report to Fauna Preservation Society of London and Conder Conservation Trust, England. 64 pp. GREEN, K. M. (1978): Primates of Bangladesh a preliminary survey of population and habitat. *Biol. Conserv.* 13: 141-160.

HENDRICHS, H. (1975): The status of the tiger (*Panthera tigris* L., 1758) in the Sunderbans Mangrove Forest (Bay of Bengal). *Saugetierk*, *Mitt.* 23 (3): 161-199.

KHAN, M. A. R. (1981): The non-human primates of Bangladesh. *Tigerpaper* 8(1): 12-15.

- a checklist. Dhaka University, Bangladesh. 173 pp. (1982b): On the distribution

of the mammalian fauna of Bangladesh. Proc. 2nd Bangladesh Nat. Conf. Forestry, 560-575. (1982c): Present status and distribution of the crocodiles and gharial of Bangladesh. Proc. 5th Working Meeting Crocodile Specialist Group of \$SC/IUCN. 229-236.

KHAN, M. A. R. & AHSAN, M. F. (1981): The population status of the non-human primates of Bangladesh. Report to Univ. Grants Commission, Dhaka, 25 pp.

KHAN, M. A. R. & RAHMAN, M. M. (1982): Avifauna of the Char Rahman and Sonadia Island of Chittagong district of Bangladesh. Proc. 2nd Bangladesh Nat. Conf. Forestry. 547-555.

LAW, S. C. (1945): Note on the occurrence of some hitherto unrecorded birds in Central and south Bengal. *Ibis* 87: 405-408.

(1948a): On the occurrence of Sitta frontalis Swainson and Sitta castanea Lesson in Khulna Sunderbans. J. Bombay nat. Hist. Soc. 47 (3): 733-734.

——— (1948b): Occurrence of the Large Himalayan Malkoha [*Rhopodytes t. tigris* (Less.)] in Khulna Sunderbans. ibid. 48(1): 183-184.

(1954): A contribution to the ornithology of the Sunderbans. J. Bengal nat. Hist. Soc. 27(1): 59-65.

(1956): A contribution to the ornitheology of the Sunderbans. ibid. 28(1): 149-152.

MANDAL, A. K. (1964): The behaviour of the Rhesus monkeys (*Macaca mulatta* Zimmerman) in the Sunderbans. ibid. 33: 153-165.

MITRA, S. N. (1957): Banglar Shikar prani (in Bengali). Bengal Govt. Press, Calcutta.

MOUNTFORT, G. (1969): The vanishing jungle. Collins, London.

MOUNTFORT, G. & POORE, D. (1968): Report on the 1967 World Wildlife Fund expedition to Pakistan. WWF, Switzerland.

MUKHERJEE, A. K. (1959): Pakhirala, Sajnakhali — an introduction to a bird sanctuary in the Sunderbans. J. Bengal nat. Hist. Soc. 30: 161-165.

(1975): The Sunderbans of India and its biota. J. Bombay nat. Hist. Soc. 72(1): 1-20.

MUKHERJEE, A. K. & GUPTA, S. (1965): Habits of the Rhesus Macaque Macaca mulatta (Zimmerman) in the Sunderbans, 24 Parganas, West Bengal. ibid. 62(1): 145-146.

OLIVIER, R. C. D. (1979): Wildlife conservation and management in Bangladesh. Report to FAO, Rome.

O'MALLEY, L. S. S. (1908): Bengal district gazetteers, Khulna. The Bengal Secretariat Book Depot, Calcutta.

RIPLEY, S. D. (1982): Synopsis of the birds of India and Pakistan. 2nd rev. edn. Bombay Natural History Society, Bombay (India).

SEIDENSTICKER, J. & HAI, M. A. (1978): The Sunderbans wildlife management plan: Conservation in the Bangladesh coastal zone. Report to Govt. of Bangladesh and WWF, Switzerland.

#### APPENDIX I

CHECKLIST OF THE WILDLIFE OF BANGLADESH MANGROVE ECOSYSTEMS

Sl. No. Scientific name

#### Common name

#### AMPHIBIA

- 1. Bufo melanostictus
- 2. Microhyla ornata
- 3. Rana hexadactyla
- 4. Rana cyanophlyctis
- 5. Rana tigerina

9.

- 6. Rana limnocharis
- 7. Rana temporalis

Batagur baska

8. Polypedates maculatus

Common Toad Ornate Microhylid Pond/Green Frog Skipper Frog Indian Bull Frog Cricket Frog

. .

Maculated Tree Frog

#### REPTILIA

Common Batagur/Tuntong Roofed Turtle Threekeeled Land Tortoise Yellow Turtle Spotted Flap Shell Bibron's Soft Shell Green Turtle Loggerhead Turtle Olive Ridley Turtle Hawksbill Turtle House Lizard Common House Lizard Common House Lizard Wall Lizard Garden Lizard Skink Common Skink Bengal/Grey Lizard **Ring/Monitor** Lizard Yellow/Common Lizard Slender Worm Snake **Rock** Python Common Sand Boa Common Wolf Snake Striped Keelback Checkered Keelback Darkbellied Marsh Snake Olive Keelback File/Wart Snake Rat Snake/Dhaman Common Bronzeback Tree Snake Common Vine Snake Common Smooth Water Snake

10. Kachuga tecta 11. Melanochelys tricarinata 12. Morenia petersi 13. Lissemys punctata 14. Pelochelys bibroni 15. Chelonia mydas 16. Caretta caretta Lepidochelys olivacea 17. 18. Eretmochelys imbricata 19. Hemidactylus brooki 20. Hemidactylus flaviviridis 21. Hemidactylus frenatus 22. Gekko gecko Calotes versicolor 23. 24. Mabuya dissimilis 25. Mabuya carinata 26. Varanus bengalensis 27. Varanus salvator 28. Varanus flavescens 29. Typhlina porrectus 30. Python molurus 31. Eryx conicus 32. Lycodon aulicus Amphiesma stolata 33. 34. Xenochrophis piscator 35. Xenochrophis cerasogaster 36. Atretium schistosum 37. Acrochordus granulatus 38. Ptyas mucosus 39. Dendrelaphis tristis 40. Ahaetulla nasutus

Sl. no	o. Scientific name	Common name
42.	Cerberus rhynchops	Dogfaced Water Snake
43.	Gerardia prevostiana	Glossy Marsh Snake
44.	Fordonia leucobalia	Whitebellied Mangrove Snake
45.	Bungarus caeruleus	Common Krait
46.	Bungarus fasciatus	Banded Krait
47.	Naja naja	Monocellate/Bengal Cobra
48.	Ophiophagus hannah	King Cobra
49.	Enhydrina schistosa	Hooknosed Sea Snake
50.	Hydrophis nigrocinctus	Blackheaded Sea Snake
51.	Hydrophis obscurus	Estuarine Sea Snake
52.	Hydrophis caerulescens	Malacca Sea Snake
53.	Microcephalophis gracilis	Common Narrowheaded Sea Snake
54.	Microcephalophis cantoris	Cantor's Narrowheaded Sea Snake
55.	Vipera russelli	Russell's Viper
56.	Trimeresurus gramineus	Bamboo Pit Viper
57.	Trimeresurus erythrurus	Spot-tailed Pit Viper
58.	Crocodylus porosus	Estuarine/Saltwater Crocodile
		Aves: Non-passerine birds
		Status
59.	Podiceps ruficollis	Little Grebe/Dabchick R (resident)
60.	Pelicanus onocrotalus	White/Rosy Pelican M(migratory)
61.	Phalacrocorax niger	Little Cormorant R
62.	Anhinga rufa	Darter/Snakebird R
63.	Ardea insignis	Great Whitebellied Heron R
64.	Ardea goliath	Goliath/Giant Heron M
65.	Ardea cinerea	Grey Heron R
66.	Ardea purpurea	Purple Heron R
67.	Ardea alba	Large Egret R
68.	Ardeola grayii	Pond Heron/Paddy Bird R
69. 70	Ardeola striatus	Little Green Heron R
70.	Bubulcus ibis	Cattle Egret R
/1.	Egretta intermedia	Median/Intermediate Egret R
12.	Egretta garzetta	Little Egret R
13.	Nyclicorax nyclicorax	Night Heron R
74.	Gorsachius melanolophus	Malay/liger Bittern R
76	Ixobrychus minutus	Little Bittern R
70.	Ixobrychus cinnamomeus	Chestnut Bittern R
78	Mysteria lausses hele	Black Bittern R
70.	Angstomus oscitzer	Painted Stork R
80	Cicopia apisoopus	Upendill Stork R
81	Ciconia episcopus	White Stork R
82	Lententiles delle	White Stork
83	Leptophilos audius	Greater Adjutant Stork* R
05.	Leptopulos Javanicus	Lesser Adjutant R

\* possibly extinct now.

# WILDLIFE IN BANGLADESH MANGROVE ECOSYSTEM

		Common name	Status
Sl. no.	Scientific name		Status
84.	Threskiornis aethiopica	White Ibis	R
85.	Anser fabalis	Bean/Pinkfooted Goose	M
86.	Anser anser	Grey Lag Goose	M
87.	Anser indicus	Barheaded Goose	M
88.	Dendrocygna javanica	Lesser Whistling Teal	R
89.	Tadorna ferruginea	Ruddy Shelduck	M
90.	Anas acuta	Pintail	Μ
91.	Anas crecca	Common Teal	Μ
92.	Anas platyrhynchos	Mallard	M
93.	Anas strepera	Gadwall	M
94.	Anas penelope	Wigeon	M
95.	Anas clypeata	Shoveller	Μ
96.	Anas querquedula	Ganganey	Μ
97.	Netta rufina	Redcrested Pochard	Μ
98.	Aythya ferina	Common Pochard	Μ
99.	Aythya nyroca	White-eyed Pochard	Μ
100.	Aythya fuligula	Tufted Duck	Μ
101.	Nettapus coromandelianus	Cotton Teal	R
102.	Elanus caeruleus	Blackwinged Kite	R
103.	Pernis ptilorhynchus	Honey Buzzard	R
104.	Milvus migrans	Pariah Kite	R
105.	Haliastur indus	Brahminy Kite	R
106.	Accipiter badius	Shikra	R
107.	Buteo rufinus	Longlegged Buzzard	Μ
108.	Buteo buteo	Buzzard	Μ
109.	Aquila rapax	Tawny Eagle	Μ
110.	Aquila pomarina	Lesser Spotted Eagle	М
111.	Haliaeetus leucogaster	Whitebellied Sea Eagle	R
112.	Haliaeetus leucoryphus	Pallas's Fishing Eagle	R
113.	Icthyophaga ichthyaetus	Greyheaded Fishing Eagle	R
114.	Gyps bengalensis	Whitebacked Vulture	R
115.	Circus macrourus	Pale Harrier	M
116.	Circus melanoleucos	Pied Harrier	M
117.	Circus aeruginosus	Marsh Harrier	M
118.	Circaetus gallicus	Short-toed Eagle	R
119.	Spilornis cheela	Crested Serpent Eagle	R
120.	Pandion haliaetus	Osprev	M
121.	Falco peregrinus	Peregrine Falcon	M
122.	Falco subbuteo	Hobby	R
123.	Falco chiquera	Redheaded Merlin	R
124.	Falco tinnunculus	Kestrel	M
125.	Francolinus gularis	Swamp Partridge	D
126.	Gallus gallus	Red Jungle Fowl	P
127	Rallus aquaticus	Water Rail	M
128	Porzana fusca	Ruddy Crake	D
129	Amaurornis phoenicurus	Whitebreasted Water Hen	R D
130	Gallicrex cinerea	Water Cock	R
			K

Sl. no.	Scientific name	Common name	Status
131.	Porphyrio porphyrio	Moorhen	R
132.	Fulica atra	Coot	Zita R
133.	Heliopais personata	Masked Finfoot	R
134.	Hydrophasianus chirurgus	Pheasant-tailed Jacana	R
135.	Metopidius indicus	Bronzewinged Jacana	R
136.	Haematopus ostralegus	Oystercatcher	R
137.	Rostratula benghalensis	Painted Snipe	R
138.	Himantopus himantopus	Blackwinged Stilt	R
139.	Recurvirostra avosetta	Avocet	Μ
140.	Glareola lactea	Small Indian Pratincole	R
141.	Vanellus cinereus	Greyheaded Lapwing	М
142.	Vanellus indicus	Redwattled Lapwing	R
143.	Vanellus spinosus	Spurwinged Lapwing	nitthan R
144.	Pluvialis dominica	Eastern Golden Plover	M
145.	Charadrius hiaticula	Ringed Plover	Μ
146.	Charadrius dubius	Little Ringed Plover	M
147.	Charadrius alexandrinus	Kentish Plover	Μ
148.	Charadrius placidus	Longbilled Ringed Plover	M
149.	Charadrius mongolus	Mongolian Plover	Μ
150.	Numenius phaeopus	Whimbrel	Μ
151.	Numenius arquata	Curlew	М
152.	Limosa limosa	Blacktailed Godwit	Μ
153.	Tringa erythropus	Spotted Red Shank	M
154.	Tringa totanus	Common Red Shank	M
155.	Tringa stagnatilis	Marsh Sandpiper	M
156.	Tringa nebularia	Green Shank	tana <b>M</b> inana (11
157.	Tringa ochropus	Green Sandpiper	M
158.	Tringa glareola	Wood Sandpiper	М
159.	Tringa terek	Terek Sandpiper	Μ
160.	Tringa hypoleucos	Common Sandpiper	Μ
161.	Arenaria interpres	Turnstone	M
162.	Limnodromus semipalmatus	Snipebilled Godwit	Mathematical Mathematical Addition
163.	Gallinago stenura	Pintail Snipe	M
164.	Gallinago gallinago	Fantail Snipe	Μ
165.	Calidris tenuirostris	Eastern Knot	M State
166.	Calidris alba	Sanderling	M
167.	Calidris minuta	Little Stint	Minaria M
168.	Calidris temminckii	Temminck's Stint	M
169.	Calidris alpina	Dunlin	M
170.	Calidris testacea	Curlew-Sandpiper	M
171.	Philomachus pugnax	Ruff and Reeve	Μ
172.	Stercorarius parasiticus	Parasitic Skua/Jaegar	Marthan and Marthan 1951
173.	Larus argentatus	Herring Gull	M
174.	Larus ichthyaetus	Great Blackheaded Gull	M
175.	Larus brunnicephalus	Brownheaded Gull	M
176.	Larus ridibundus	Blackheaded Gull	Μ
177.	Chlidonias hybrida	Whiskered Tern	М

# WILDLIFE IN BANGLADESH MANGROVE ECOSYSTEM

Sl. no.	Scientific name	Common name	Status
178	Chlidonias leucopterus	Whitewinged Black Tern	Μ
179	Gelochelidon nilotica	Gullbilled Tern	R
180	Hydroprogne caspia	Caspian Tern	Μ
181	Sterna aurantia	Indian River Tern	R
182	Sterna hirundo	Common Tern	Μ
183	Sterna acuticauda	Blackbilled Tern	R
184	Sterna fuscata	Sooty Tern	Μ
185	Sterna albitrons	Little Tern	R
186	Sterna hergij	Large Crested Tern	R
187	Sterna bengalensis	Lesser Crested Tern	R
188	Rhynchops albicollis	Indian Skimmer	R
189	Treron pompadora	Greyfronted Green Pigeon	R
190	Treron phoenicoptera	Green Pigeon	R
191	Columba livia	Blue Rock Pigeon	R
192	Streptopelia orientalis	Rufous Turtle Dove	M
193	Streptopelia decaocto	Ring Dove	R
194.	Streptopelia tranquebarica	Red Turtle Dove	R
195	Streptopelia chinensis	Spotted Dove	R
196	Psittacula krameri	Roseringed Parakeet	R
197	Psittacula finschii	Slatyheaded Parakeet	R
198.	Clamator jacobinus	Pied Crested Cuckoo	R
199.	Cuculus varius	Brainfever Bird	R
200.	Cuculus micropterus	Indian Cuckoo	R
201.	Cacomantis sonneratii	Banded Bay Cuckoo	R
202.	Cacomantis merulinus	Rufousbellied Plaintive Cuckoo	R
203.	Eudynamys scolopacea	Koel	R
204.	Rhopodytes tristis	Large Greenbilled Malkoha	R
205.	Centropus sinensis	Crow-Pheasant/Coucal	R
206.	Tyto alba	Barn Owl	R
207.	Otus scops	Scops Owl	R
208.	Otus bakkamoena	Collared Scops Owl	R
209.	Bubo bubo	Eagle-Owl/Great Horned Owl	R
210.	Bubo zevlonensis	Brown Fish Owl	R
211.	Bubo flavipes	Tawny Fish Owl	R
212.	Ninox scutulata	Brown Hawk-Owl	R
213.	Athene brama	Spotted Owlet	R
214.	Asio flammeus	Shorteared Owl	М
215.	Caprimulgus indicus	Jungle Nightjar	R
216.	Caprimulgus macrurus	Longtailed Nightjar	R
217.	Cypsiurus parvus	Palm Swift	R
218.	Cervle rudis	Lesser Pied Kingfisher	R
219.	Alcedo atthis	Common Kingfisher	R
220.	Pelargopsis amauroptera	Brownwinged Kingfisher	R
221.	Pelargopsis capensis	Storkbilled Kingfisher	R
222.	Halcyon coromandra	Ruddy Kingfisher	R
223.	Halcyon pileata	Blackcapped Kingfisher	R
224.	Halcyon smyrnensis	Whitebreasted Kingfisher	R
225.	Halcyon chloris	Whitecollared Kingfisher	R
			and a good of the opposite the state of the second state of the se

45

Sl. no.	Scientific name	Common name	Status
226	Merops orientalis	Green Bee-eater	R
227	Coracias benghalensis	Indian Roller/Blue Jay	R
228	Upupa epops	Ноорое	R
229	Megalaima lineata	Lineated Barbet	R
230.	Megalaima haemacephala	Coppersmith	R
231.	Jvnx torquilla	Wryneck	Μ
232.	Micropternus brachyurus	Rufous Woodpecker	R
233.	Picus myrmecophoneus	Little Scalybellied Green Woodpecke	r R
234.	Dinopium benghalense	Lesser Goldenbacked Woodpecker	R
235.	Picoides macei	Fulvousbreasted Pied Woodpecker	R
236.	Picoides mahrattensis	Yellowfronted Pied Woodpecker	R
237.	Picoides nanus	Pigmy Woodpecker	R
238.	Chrysocolaptes lucidus	Larger Goldenbacked Woodpecker	R
	Aves: Pas	serine birds	
239.	Mirafra assamica	Bush Lark	R
240.	Alauda gulgula	Eastern Skylark	R
241.	Hirundo rustica	Common Swallow	Μ
242.	Hirundo daurica	Redrumped/Striated Swallow	Μ
243.	Lanius schach	Blackheaded/Rufous Shrike	R
244	Lanius cristatus	Brown Shrike	М
245	Oriolus xanthornus	Blackheaded Oriole	R
246	Dicrurus adsimilis	Black Drongo	R
247.	Dicrurus leucophaeus	Grey/Ashy Drongo	М
248.	Dicrurus aeneus	Bronzed Drongo	М
249.	Dicrurus paradiseus	Greater Racket-tailed Drongo	R
250.	Artamus fuscus	Ashy-Swallow Shrike	R
251.	Sturnus malabaricus	Greyheaded Myna	R
252.	Sturnus vulgaris	Starling	М
253.	Sturnus contra	Pied Myna	R
254.	Acridotheres tristis	Common Myna	R
255.	Acridotheres ginginianus	Bank Myna	R
256.	Acridotheres fuscus	Jungle Myna	R
257.	Dendrocitta vagabunda	Tree Pie	R
258.	Corvus splendens	House Crow	R
259.	Corvus macrorhynchos	Jungle Crow	R
260.	Tephrodornis pondicerianus	Common Wood Shrike	R
261.	Coracina novaehollandiae	Large Cuckoo-Shrike	R
262.	Coracina melaschistos	Smaller Grey Cuckoo-Shrike	М
263.	Coracina melanoptera	Blackheaded Cuckoo Shrike	М
264.	Pericrocotus cinnamomeus	Small Minivet	R
265.	Aegithina tiphia	Common Iora	R
266.	Chloropsis aurifrons	Goldfronted Chloropsis	R
267.	Pycnonotus melanicterus	Blackheaded Yellow Bulbul	R
268.	Pycnonotus jocosus	Redwhiskered Bulbul	R
269.	Pycnonotus cafer	Redvented Bulbul	R
270.	Pellorneum ruficeps	Spotted Babbler	R
271.	Trichastoma abotti	Abott's Babbler	R

## WILDLIFE IN BANGLADESH MANGROVE ECOSYSTEM

Sl. n	o. Scientific name	Common name	Status
272.	Turdoides striatus	Jungle Babbler	R
273.	Alcippe poioicephala	Quaker Babbler	R
274.	Muscicapa parva	Redbreasted Flycatcher	Μ
275.	Muscicapa rubeculoides	Bluethroated Flycatcher	M doubtful
276.	Muscicapa thalassina	Verditer Flycatcher	М
277.	Culicicapa ceylonensis	Greyheaded Flycatcher	R
278.	Rhipidura albicollis	Whitethroated Fantail Flycatcher	R
279.	Terpsiphone paradisi	Paradise Flycatcher	R
280.	Hypothymis azurea	Blacknaped Flycatcher	R
281.	Pachycephala grisola	Mangrove Whistler	R
282.	Bradypterus luteoventris	Brown Bush Warbler	М
283.	Cisticola exilis	Fantail Warbler	R
284.	Cisticola juncidis	Streaked Fantail Warbler	R
285.	Prinia hodgsoni	Franklin's Wren-Warbler	R
286.	Prinia socialis	Ashy Wren-Warbler	R
287.	Orthotomus sutorius	Tailor Bird	R
288.	Acrocephalus stentorius	Great Reed Warbler	М
289.	Acrocephalus dumetorum	Blyth's Reed Warbler	M
290.	Phylloscopus affinis	Tickell's Leaf Warbler	M
291.	Erithacus svecicus	Blue Throat	M
292.	Copsychus saularis	Magpie-Robin (National Bird)	R
293.	Phoenicurus ochruros	Black Redstart	M
294.	Saxicola torquata	Collared Bush Chat	M
295.	Saxicola caprata	Pied Bush Chat	R
296.	Monticola solitarius	Blue Rock Thrush	M
297.	Zoothera citrina	Orangeheaded Ground Thrush	M
298.	Turdus ruficollis	Redthroated Thrush	M
299.	Parus major	Grev Tit	R
300.	Sitta castanea	Chestnutbellied Nuthatch	R
301.	Sitta frontalis	Velvetfronted Nuthatch	R
302.	Anthus hodgsoni	Indian Tree Pipit	M
303.	Anthus novaeseelandiae	Paddyfield Pipit	M
304.	Motacilla flava	Yellow Wagtail	M
305.	Motacilla citreola	Yellowheaded Wastail	M
306.	Motacilla cinerea	Grev Wagtail	M
307.	Motacilla alba	White Wagtail	M
308.	Dicaeum trigonostigma	Ornagebellied Flowerpecker	D
309.	Dicaeum erythrorhynchos	Tickell's Flowerpecker	R P
310.	Dicaeum cruentatum	Scarletbacked Flowerpecker	R D
311.	Nectarinia zeylonica	Purplerumped Sunbird	R D
312.	Nectarinia asiatica	Purple Sunbird	R D
313.	Zosterops palpebrosa	White-eve	R D
314.	Passer domesticus	House Sparrow	T.
315.	Ploceus philippinus	Bava/Weaver Bird	D
316.	Ploceus manyar	Streaked Weaver Bird	R D
317.	Lonchura malabarica	Common Silverbill	N
318.	Lonchura striata	Whitebacked Munia	R
319.	Lonchura punctulata	Spotted Munia	R
		spotted munia	K

Sl. no	o. Scientific name	Common name	
- des	Mammalia		
320.	Suncus murinus	Grey Musk Shrew	
321.	Pteropus giganteus	Flying Fox	
322.	Rousettus leschenaultii	Fulvous Fruit Bat	
323.	Cynopterus sphinx	Shortnosed Fruit Bat	
324.	Rhinopoma hardwickei	Lesser Rat-tailed Bat	
325.	Taphozous saccolaimus	Pouchbearing Sheathtailed Bat	
326.	Megaderma lyra	False Vampire	
327.	Coelops frithi	Tailless Leafnosed Bat	
328.	Pipistrellus mimus	Indian Pygmy Pipistrelle	
329.	Pipistrellus coromandra	Indian Pipistrelle	
330.	Hesperoptenus tickelli	Tickell's Bat	
331.	Scotophilus temmincki	Lesser Yellow Bat	
332.	Macaca mulatta	Rhesus Macaque	
333.	Macaca fascicularis	Crabeating Macaque	
334.	Canis aureus	Jackal	
335.	Lutra perspicillata	Smooth Indian Otter	
336.	Aonyx cinerea	Clawless Otter	
337.	Viverra zibetha	Large Indian Civet	
338.	Viverricula indica	Small Indian Civet	
339.	Paradoxurus hermaphrodites	Palm Civet/Toddy Cat	
340.	Herpestes auropunctatus	Small Mongoose	
341.	Herpestes edwardsi	Common Mongoose	
342.	Panthera tigris	Royal Bengal Tiger (National Animal)	
343.	Panthera pardus	Leopard/Panther (extinct)	
344.	Felis bengalensis	Leopard-Cat	
345.	Felis viverrina	Fishing Cat	
346.	Felis chaus	Jungle Cat	
347.	Rhinoceros unicornis	Onehorned Rhinoceros (extinct)	
348.	Rhinoceros sondaicus	Smaller Onehorned Rhinoceros (extinct)	
349.	Bubalus bubalis	Wild Buffalo (extinct)	
350.	Cervus duvauceli	Swamp Deer/Barasingha (extinct)	
351.	Axis porcinus	Hog Deer (extinct)	
352.	Axis axis	Chital/Spotted Deer	
353.	Muntiacus muntjac	Barking Deer/Muntiac	
354.	Sus scrofa	Wild Boar	
355.	Lepus nigricollis	Rufoustailed Hare	
356.	Funambulus pennanti	Fivestriped Palm Squirrel	
357.	Bandicota bengalensis	Lesser Bandicoot	
358.	Bandicota indica	Bandicoot Rat	
359.	Mus booduga	Indian Porcupine	
360.	Mus musculus	House Mouse	
361.	Rattus rattus	Common House Rat	
362.	Hystrix indica	Indian Poroupine	
363.	Orcaella brevirostris	Irrawaddy Dolphin	
364.	Globicephala macrorhynchus	Shortfinned Pilot Whale	
365.	Peponocephala electra	Broadbeaked/Melonheaded Dolphin	
366.	Neophocaena phocaenoides	Finless/Little Porpoise	
367.	Stenella malayana	Malay Dolphin	
368.	Platanista gangetica	Ganges Susu/River Dolphin	



Khan, Mohammad Ali Reza. 1986. "WILDLIFE IN BANGLADESH MANGROVE ECOSYSTEM." *The journal of the Bombay Natural History Society* 83, 32–48.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/191627</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/157143</u>

**Holding Institution** Smithsonian Libraries and Archives

**Sponsored by** Biodiversity Heritage Library

**Copyright & Reuse** Copyright Status: In Copyright. Digitized with the permission of the rights holder License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.