

indicates the variety and condition of the fruit; quantities of  $n/10$  NaOH required to neutralize; and the per cent of acidity in terms of citric acid.

TABLE SHOWING ACID CONTENT OF TOMATO FRUITS

Variety	Condition			Average no. of cc. of $n/10$ NaOH, to neutralize	Total per cent of acid as citric
	When picked	Interval or incubation	When titrated*		
Dwarf Stone	Ripe	0	Red	1.695	.52
Dwarf Stone	Half grown	0	Green	1.82	.56
Dwarf Stone	Half grown	Incub. 32° C. 10 days	Artif. yellow	2.135	.66
Dwarf Stone	Half grown	Lab. 24 days	Red	1.375	.42
Dwarf Stone	Half grown	Incub. 32° C. 10 days	Green	1.485	.46
Sparks' Earliana	Ripe	0	Red	1.695	.52
Sparks' Earliana	Half grown	0	Green	1.87	.58
Truckers' Favorite	Half grown	Incub. 32° C. 22 days	Artif. yellow	2.56	.79
Truckers' Favorite	Half grown	Lab. 24 days	Red	1.66	.51
Red Peach	Half grown	Incub. 32° C. 22 days	Artif. yellow	2.115	.65
Red Peach	Half grown	Lab. 24 days	Red	1.675	.52
Yellow Peach	Half grown	Incub. 32° C. 22 days	Artif. yellow	2.47	.76
Yellow Peach	Half grown	Lab. 24 days	Yellow	2.065	.64
Yellow Plum	Ripe	0	Yellow	2.12	.65
Yellow Plum	Half grown	0	Green	1.92	.59
Yellow Pear	Half grown	Incub. 32° C. 20 days	Artif. yellow	1.60	.49
Yellow Pear	Half grown	Lab. 24 days	Yellow	1.395	.43

\* All fruits designated "red," "yellow," and "artificial yellow" were, at the same time, ripe.

The results above reported may not yet be as extensive as might be desired in order to follow closely the changes in acidity under different conditions; but they consistently point out certain relations of interest which may be briefly enumerated as follows: (1) A comparison of the acid content of green and normally ripened fruits was made, using Dwarf Stone, Sparks' Earliana, and Yellow Plum, all direct from the field. There were no marked differences between the green and ripe stages within the variety; yet the acidity of the green fruits of the red varieties in these tests is somewhat higher, while the acid content of the green fruits of the one yellow variety tested is somewhat lower. (2) Fruits of Dwarf Stone, Truckers' Favorite, Red Peach, Yellow Peach, and Yellow Pear which



were picked green and ripened in the incubator at 32–33°C. (10–22 days) exhibit a higher acid content than either those ripened on the vines or those ripened at the temperature of the laboratory. (3) There are considerable differences in the acidity of varieties, but judging from the results of these tests the normally ripened fruits of yellow varieties commonly contain as much acid as those of red varieties.

The several facts brought out by these tests render it obvious that there is now no sufficient evidence to justify relating pigmentation to total acidity. The acidity changes are, however, interesting in themselves, in these as well as in other fruits. No attempt was made to follow progressively any changes in acidity induced by conditions; but in titrating on one occasion, after an interval of two days, new samples of both red and yellow fruits which had been ripened in the laboratory, it was found that the acidity had noticeably declined since the previous titrations from the same lots of fruits.

We have reckoned the acidity of the tomato in terms of citric acid, as is customary. It should be noted, however, that while Bowman (3) and others report citric as the chief acid of the tomato, Albahary (1), on the contrary, gives .48 per cent as the malic acid content and .09 per cent as that of citric acid in the fresh fruits. The author last mentioned gives no indications respecting the variety or condition of the fruit employed. In a later contribution (2) he reports the results of analyzing tomato fruits in different stages of maturation, as follows: "1° le fruit vert avant l'apparition de la graine dans la pulpe; 2° le fruit vert au moment où la graine est complètement formée; 3° le fruit rouge arrivé à sa pleine maturation." In the second stage, corresponding to practically full grown, green, he finds .58, and in the ripe fruits .42 per cent of organic acids. This is in complete agreement with our findings. In the earliest stage of fruit development Albahary finds an acid content of only .116 per cent. Wehmer (5), after quoting Albahary (1) as to the percentage of the various acids in the fruit, remarks, "Die Acidität wechselt stark je nach dem Reifestadium (von 0,06–0,697% des Saftes auf Citronensäure berechnet)." He does not indicate the



source of these data, and certainly the smaller percentage given can refer only to the youngest stages of fruit development.

*Graduate Laboratory, Missouri Botanical Garden.*

#### LITERATURE CONSULTED

1. Albahary, J. M. Analyse complète du fruit du *Lycopersicum esculentum* ou Tomate. Compt. rend. acad. Paris **145**: 131-33. 1907.
2. ———, Étude chimique de la maturation du *Lycopersicum esculentum* (Tomate). Compt. rend. acad. Paris **147**: 146-47. 1908.
3. Bowman, W. Tomatoes: chemical examination of fruits. Va. Agr. Exp. Sta. Bul. 9: 16-18. 1891.
4. Duggar, B. M. Lycopersicin, the red pigment of the tomato, and the effects of conditions upon its development. Washington Univ. Studies **1**: 22-45. 1913.
5. Wehmer, C. Die Pflanzenstoffe 685-86. 1911.



Duggar, Benjamin M. and Merrill, M C. 1914. "The Effect of Certain Conditions Upon the Acidity of Tomato Fruits." *Annals of the Missouri Botanical Garden* 1, 237-240. <https://doi.org/10.2307/2989994>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/54249>

**DOI:** <https://doi.org/10.2307/2989994>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/15919>

**Holding Institution**

Missouri Botanical Garden, Peter H. Raven Library

**Sponsored by**

Missouri Botanical Garden

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.