

A NEW VARIETY AND FOUR NEW COMBINATIONS IN
PITTOCAULON AND *TELANTHOPHORA*
(ASTERACEAE: SENECEONEAE) FROM MEXICO

Bonnie L. Clark

8201 Hauser Drive
Lenexa, KS 66212-2542, U.S.A.

ABSTRACT

A new variety, *Pittocaulon hintonii* var. *cerrograndensis* (Asteraceae: Senecioneae) is described, and four new combinations are proposed: *Pittocaulon velatum* var. *tzimolensis*, *Telanthophora cobanensis* var. *molinae*, *Telanthophora grandifolia* var. *serraquitchensis*, and *Telanthophora sublaciniatus*.

KEY WORDS: *Senecio*, *Telanthophora*, *Pittocaulon*, Mexico, Central America

RESUMEN

Se describe una variedad nueva, *Pittocaulon hintonii* var. *cerrograndensis*, y se propon cuatro combinaciones nuevas, *Pittocaulon velatum* var. *tzimolensis*, *Telanthophora cobanensis* var. *molinae*, *Telanthophora grandifolia* var. *serraquitchensis*, y *Telanthophora sublaciniatus*.

Recent studies in certain Mexican and Central American segregates of *Senecio*, s.l. (Clark 1996) have led to the recognition a new variety and the need for four new nomenclatural combinations. These are recorded here to make the names available for curatorial purposes. More complete explanations will be presented when the revisionary studies are published.

Pittocaulon hintonii H. Rob. & Brettell var. ***cerrograndensis*** B.L. Clark, var. nov. TYPE: MEXICO: JALISCO: Sierra de Manantlán, on very steep E-facing slopes of Cerro Grande on road from Lagunillas to Campo Cuatro & Juluapan, on limestone cliffs above camp 4, dry, deciduous woodland, together with the gigantic creeping stems of *Philodendron warczewiczii*, *Selenicereus* spp., and *Pedilanthus* in full sun, lat. 19° 23'–25' N and long. 103° 51' 30"–52° 30' W, elev. 1480 m, 15 Mar 1987, H.H. Ilitis et al. 29713 (HOLOTYPE: WIS!; ISOTYPES: IBUG, KSCI, TEX, ZEA).

A var. *hintonii* cujusque capituli flosculis radiatis 14 (nec 7–9) perspicue diversa.

Pittocaulon hintonii var. *cerrograndensis* differs notably from var. *hintonii* in having capitula with 14 ray florets instead of 7–9. It is apparently restricted to the Sierra de Manantlán, a region known for a number of endemics (Vázquez et al. 1995). The nomenclaturally typical var. *hintonii* occurs farther to the north, in the region of Sierra Naranjillo, Michoacán.

Pittocaulon velatum (Greenm.) H. Rob. & Brettell var. ***tzimolensis*** (T.M. Barkley) B.L. Clark, comb. nov. *Senecio praecox* (Cav.) DC. var. *tzimolensis* T.M. Barkley, Phytologia 69:142. 1990. TYPE: MEXICO: CHIAPAS: Tzimol, tropical deciduous forest 15 km S of Comitán on road to Tzimol and Tuxtal Gutierrez, elev. 1200 m, 20 Mar 1981, D.E. Breedlove 50266 (HOLOTYPE: CAS!).

Telanthophora cobanensis (J.M. Coult.) H. Rob. & Brettell var. **molinae** (H. Rob. & Brettell) B.L. Clark, comb. et stat. nov. *Telanthophora molinae* H. Rob. & Brettell, Phytologia 27:428. 1974. TYPE: GUATEMALA: DEPARTMENT OF SAN MARCOS: near Alta Fraternidad, between San Rafael Pie de la Cuesta and Palo Gordo, W facing slope of the Sierra Madre Mountains, wet mountain forest, elev. 1800–2400 m, 10–18 Dec 1963, L.O. Williams, A. Molina & T.P. Williams 26271 (HOLOTYPE: US!; ISOTYPES: FI, GHI, MO!, NY!)

Telanthophora grandifolia (Less.) H. Rob. & Brettell var. **serraquitchensis** (Greenm.) B.L. Clark, comb. et stat. nov. *Senecio serraquitchensis* Greenm., Field Mus. Nat. Hist., Bot. Ser. 2(6):286. 1907. *Telanthophora serraquitchensis* (Greenm.) H. Rob. & Brettell, Phytologia 27:428. 1974. TYPE: GUATEMALA: DEPARTMENT OF ALTA VERAPAZ: Serraquitché, elev. 760 m, Apr 1889, J. Donnell Smith 1598 (HOLOTYPE: GHI; ISOTYPE: CHII).

Telanthophora sublaciniatus (Greenm.) B.L. Clark, comb. et stat. nov. *Senecio cobanensis* J.M. Coult. var. *sublaciniatus* Greenm., Ceiba 1(2):120. 1950. TYPE: GUATEMALA: DEPARTMENT OF SACATEPEQUEZ: hills of finca Carmona southeast of Antigua, in damp forest, elev. 1590–1800 m, 27 Jan 1939, P.C. Standley 63680 (HOLOTYPE: MO!; ISOTYPE: FI!).

ACKNOWLEDGMENTS

This study derives from a doctoral dissertation prepared at Kansas State University, Manhattan, Kansas, under the direction of T.M. Barkley. Rupert Barneby of the New York Botanical Garden kindly rendered the Latin diagnosis.

REFERENCES

- CLARK, B.L. 1996. A study of variation in *Senecio* Sect. *Terminales* Greenm. (Compositae; Senecioneae). Ph.D. dissertation, Kansas State University, Manhattan, Kansas.
- VÁZQUEZ G., J.A., R. CUEVAS G., T.S. COCHRANE, H.H. ILTIS, F.J. SANTANA M., and L. GUZMÁN H. 1995. Flora de Manantlán. Sida, Bot. Misc. 13, Botanical Research Institute of Texas, Fort Worth.



Clark, Bonnie L . 2000. "A NEW VARIETY AND FOUR NEW COMBINATIONS IN PITTOCAULON AND TELANTHOPHORA (ASTERACEAE: SENECTIONEAE) FROM MEXICO." *SIDA, contributions to botany* 19, 235–236.

View This Item Online: <https://www.biodiversitylibrary.org/item/36567>

Permalink: <https://www.biodiversitylibrary.org/partpdf/163302>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.