# MACROCYSTELLA CALLAWAY, THE EARLIEST GLYPTOCYSTITID CYSTOID

# by C. R. C. PAUL

ABSTRACT. Macrocystella mariae Callaway 1877, type species of Macrocystella, has a stem which is divisible into proximal and distal portions; a theca composed of 4 basal, 5 infra-lateral, 5 lateral, 6 radial, and some oral plates; a large periproct surrounded by 5 thecal plates; biserial unbranched brachioles grouped into 5 ambulacra and arising from the margins of the flattened oral surface. In all these respects it agrees with Mimocystites bohemicus Barrande 1887, type species of Mimocystites which becomes a subjective junior synonym of Macrocystella. Macrocystella azaisi (Thoral) has 7 orals and thus Macrocystella differs from the rhombiferan Cheirocrinus Eichwald only in the absence of pectinirhombs. The Macrocystellidae are therefore transferred to the rhombiferan superfamily Glyptocystitida.

*Macrocystella* evolved into *Cheirocrinus* by the acquisition of pectinirhombs. In *Macrocystella* respiration probably took place through all the thecal plates which are very thin. In *Cheirocrinus* respiration was restricted to the pectinirhombs thus allowing much thicker and stronger thecal plates to develop.

*Macrocystella* led a freely vagrant existence and may have had internal buoyancy devices. The stem did not provide permanent fixture and may have been used as a organ of locomotion in conjunction with the brachioles.

THE cystoidea, as currently defined (Kesling 1963) is probably an artificial group. The main character which is used to unite the cystoids as a class is the possession of porestructures (rhombs and dipores) developed in the thecal plates. However similar porestructures are found in at least some representatives of other Palaeozoic echinoderm classes (blastoids, crinoids, paracrinoids, eocrinoids, for example) and one genus of rhombiferan cystoids entirely lacks pore-structures. This paper deals with another genus, Macrocystella Callaway 1877, which lacks true pore-structures but which is thought to be the oldest known representative of the Glyptocystitida, one of the three major rhombiferan superfamilies. Macrocystella has a complex taxonomic history (see below) and has been variously regarded as an eocrinoid, a rhombiferan cystoid or as a link between these classes. Close comparison indicates that Macrocystella is identical to the rhombiferan genus Cheirocrinus Eichwald in all details except the possession of pectinirhombs. Hence Macrocystella is regarded as a rhombiferan. It is believed that the absence of pectinirhombs in Macrocystella is a primitive character. Many Ordovician pelmatozoans independently developed thecal or calycinal pore structures, apparently in response to respiratory needs. To group all such echinoderms together obscures their true relationships. It is essential to consider other characters in addition to the possession of pore-structures, especially when the latter are so variable.

Acknowledgements. The author is indebted to Dr. R. P. S. Jefferies and Mr. H. G. Owen, British Museum, Natural History (BMNH), Professor B. Kummel and Mr. J. Sprinkle, Museum of Comparative Zoology, Harvard (MCZ), Mr A. G. Brighton, Sedgwick Museum, Cambridge (SM), and Dr. R. A. Robison, United States National Museum, Washington (USNM) for the loan of, or access to, specimens. Dr. I. Strachan, Birmingham University (BU) supplied latex impressions of Callaway's original specimens of *M. mariae* and Professor K. E. Caster, University of Cincinnati (UC) impressions of *Macrocystella* and *Cheirocrinus* from France and Bohemia. Part of this work was undertaken during the tenure of a National Environmental Research Council research studentship at the Sedgwick [Palaeontology, Vol. 11, Part 4, 1968, pp. 580-600, pls. 111-13.]

Museum, Cambridge and part during the tenure of a Post-doctoral Fellowship at the University of Michigan (NSF grant GB-3366). Both are gratefully acknowledged.

# PREVIOUS RESEARCH

*Macrocystella* (type species *M. mariae* Callaway 1877) was first described from the Lower Ordovician (Tremadoc) Shineton Shales of Shropshire. Barrande (1887, p. 163) described a closely similar genus, *Mimocystites*, for a single species, *M. bohemicus* Barrande. Jaekel (1899, p. 171) suggested that these two genera were synonymous but used the name *Mimocystites*. Jaekel regarded *Mimocystites* as the progenitor of the cystoids and most closely related to the rhombiferan *Cheirocrinus* Eichwald. Bather (1899, p. 920) proposed the family Macrocystellidae which he assigned to the Rhombifera. He did not elucidate the composition of the Macrocystellidae but later (1900, p. 56) included *Macrocystella*, *Mimocystites* hardly differed he used both names. Bather's reconstruction of *Macrocystella mariae* (1900, p. 95, fig. 18) was inaccurate in depicting branched arms and considerably influenced subsequent opinions on the affinities of *Macrocystella*.

Jaekel (1918, p. 27) retained both Macrocystella and Mimocystites (possibly because of Bather's reconstruction of the former) and placed them, with his new genus Polyptychella, in the Macrocystellidae. A separate family was proposed for Lichenoides. The Macrocystellidae and Lichenoidae were assigned respectively to the orders Plicata and Reducta of the Eocrinoidea. Thoral (1935, p. 113) considered Macrocystella and Mimocystites to be distinct but based his opinion of the former on the original description. Bassler and Moodey (1943) reverted to Bather's classification, included Lichenoides in the Macrocystellidae and that family in the Rhombifera. Cuénot (1948) also assigned Macrocystella to the Rhombifera and later (1953) regarded Mimocystites as a junior synonym. Moore (1954, p. 127, fig. 2a) published an inaccurate plate diagram of Macrocystella which depicts 5 basals, 5 radials, 5 orals, and a minute periproct. Moore regarded Macrocystella as an eocrinoid. Sdzuy (1955) accepted Macrocystella and *Mimocystites* as separate genera only if published reconstructions were accurate. He based his opinion of Macrocystella on Bather's (1900) and Moore's (1954) work, the accuracy of which he doubted. More recently Prokop (1966) and Ubaghs (1968) have suggested that *Macrocystella* and *Mimocystites* are synonymous. Ubaghs regards Macrocystella as most closely related to Cheirocrinus.

Quite apart from the synonymy of *Macrocystella* and *Mimocystites* another problem arises as to the status of *Cystidea* Barrande 1868. Barrande (1867, p. 179) published two nomina nuda, *Cystidea sedgwicki* and *C. bohemicus*. Later he introduced *C. bavarica* (Barrande 1868, p. 106) this time accompanied by a description and figures. Barrande made it quite clear in both publications that he intended *Cystidea* as a collective group name not a formal generic name and he so used it again (1887), erecting several more species. *Cystidea bavarica* Barrande is a valid binomen and could be construed to be type species of the genus *Cystidea* Barrande by monotypy. Pompeckj (1896, p. 90) and Sdzuy (1955, p. 170) have attributed *Cystidea bavarica* to *Macrocystella*. There is no doubt they are correct in this action and therefore *Cystidea* Barrande 1868, if accepted as an available generic name, should take precedence over both *Macrocystella* and *Mimocystites*. Such action is not in the interests of nomenclatorial stability. No author

has accepted *Cystidea* as a valid generic name whereas *Macrocystella* has been widely used and is figured and described in standard text-books in English, French, and German. Application has therefore been made to the International Commission on Zoological Nomenclature for the suppression of *Cystidea* Barrande 1868 under the Plenary Powers (Paul 1967b). In anticipation of a favourable decision, *Macrocystella* is used throughout this work.

The systematics and composition of the Macrocystellidae and the suggested synonymy between *Macrocystella* and *Mimocystites* can only be settled after a revised account of the morphology of *Macrocystella mariae* has been given. Latex impressions of the original specimens of *M. mariae* have been re-examined and additional material studied. This has been compared with Barrande's (1887) and Jaekel's (1899) descriptions and figures of *Mimocystites bohemicus*, with latex impressions of some of Barrande's original material and additional material of *M. bohemicus* in the Schary Collection, Museum of Comparative Zoology, Harvard. Latex impressions of *M. azaisi* (Thoral) have added further information.

*Macrocystella* and *Mimocystites* are identical and quite distinct from *Lichenoides* to judge from Ubagh's (1953) account of *L. priscus* Barrande. *Polyptychella* Jaekel was founded on isolated plates and its systematic position cannot be settled without further information. The Macrocystellidae thus contains the single genus *Macrocystella*. As previously stated the Macrocystellidae is assigned to the Rhombifera (Glyptocystitida).

# SYSTEMATIC PALAEONTOLOGY

## Superfamily GLYPTOCYSTITIDA Bather 1899

*Diagnosis*. A superfamily of Rhombifera with well-developed stem divided into proximal and distal portions; theca composed of 4 basals, 5 infra-laterals, 5 laterals, 4–6 radials, and 7 orals; with pectinirhombs (when pore structures are developed).

All Glyptocystitida are characterized by a theca composed of 25–7 thecal plates arranged in five circlets termed basal, infra-lateral, lateral, radial, and oral. All but two genera—*Macrocystella* and *Amecystis* Ulrich and Kirk—have pectinirhombs. These characters distinguish glyptocystitids from members of the other two major rhombi-feran superfamilies, the Hemicosmitida and Caryocystitida. The former have thecal plates arranged in three or four circlets and slightly different rhombs. The latter have a large variable number of plates, some of which may be added during growth, and a completely different type of rhomb (Paul 1968).

# Family MACROCYSTELLIDAE Bather 1899 emend. Jaekel 1918

*Diagnosis.* A family of Glyptocystitida without pectinirhombs; with cylindrical theca having 6 radials; large periproct surrounded by 5 thecal plates and covered with a flexible plated integument; brachioles confined to oral surface, grouped into 5 ambulacra.

### Genus MACROCYSTELLA Callaway 1877

1868 Cystidea Barrande, p. 106.

1877 Macrocystella Callaway, p. 669, pl. 24, fig. 13.

- 1880 Macrocystella Callaway; Zittel, p. 420.
- 1887 Mimocystites Barrande, p. 163, pl. 28 (1), figs. 1-20.
- 1891 Macrocystella Callaway; Carpenter, p. 13.
- 1891 Mimocystis [sic] Barrande; Carpenter, p. 13.
- 1896 Mimocystis [sic] Barrande; Haeckel, p. 149.
- 1899 Macrocystella Callaway; Jaekel, p. 171.
- 1899 Mimocystites Barrande; Jaekel, p. 172, fig. 33.
- 1900 Macrocystella Callaway; Bather, p. 56, fig. 18.
- 1900 Mimocystis [sic] Barrande; Bather, p. 56.
- 1913 Macrocystella Callaway; Springer, p. 157.
- 1918 Macrocystella Callaway; Jaekel, p. 27.
- 1918 Mimocystites Barrande; Jaekel, p. 27.
- 1935 Mimocystites Barrande; Thoral, p. 110, 113.
- 1943 Macrocystella Callaway; Bassler and Moodey, p. 6.
- 1943 Mimocystites Barrande; Bassler and Moodey, p. 6.
- 1948 Macrocystella Callaway; Regnéll, p. 11.
- 1948 Macrocystella Callaway; Cuénot, p. 18, fig. 17.
- 1953 Macrocystella Callaway; Cuénot, p. 619.
- 1953 Mimocystites Barrande; Choubert, Termier, and Termier, p. 137.
- 1954 Macrocystella Callaway; Moore, p. 127, fig. 2a.
- 1954 Mimocystites Barrande; Termier and Termier, p. 92, figs. a-e.
- 1955 Macrocystella Callaway; Sdzuy, p. 269.
- 1966 Macrocystella Callaway; Prokop, p. 820.
- non *Cystidea* Barrande 1867 (nomen nudum) nec Barrande 1887 nec Haeckel 1896 (indeterminate echinoderm fragments).

#### *Diagnosis*. As for family.

*Regional distribution and stratigraphic range. Macrocystella* is recorded from the Tremadoc of England and Wales (*M. mariae* Callaway), Bavaria (*M. bavarica* (Barrande) 1868), Bohemia (*M. bohemicus* Barrande 1887), and France (*M. azaisi* (Thoral) 1935). *Macrocystella* is also recorded from Greenland, the South American Cordillera and Korea (for detailed references see Regnéll, 1948, pp. 11–12). Choubert, Termier, and Termier record *Macrocystella* from the Llandeilo of Morocco. Available specimens confirm the genus from the Llandeilo of Pu-piao, Northern Shan States, Burma, (SM), from the Tremadoc of Mexico (USNM) and possibly from the Caradoc of Corwen, Wales, and Girvan, Scotland (BMNH). *M. pachecoi* Meléndez (1944) from the Ashgill of Aragon, Spain is probably a *Heliocrinites. Macrocystella* thus ranges from the Tremadoc to the Llandeilo and possibly Caradoc (Lower–Middle Ordovician).

## Macrocystella mariae Callaway 1877

Plate 111, figs. 1, 3-6; Plate 112, figs. 1-3, 5-10; Plate 113, fig. 2

- 1877 Macrocystella mariae Callaway, p. 670, pl. 24, fig. 13.
- 1896 Macrocystella mariae Callaway; Haeckel, p. 149, pl. 4, fig. 30.
- 1900 Macrocystella mariae Callaway; Bather, p. 56, fig. 18.
- 1905 Macrocystella mariae Callaway; Fearnsides, p. 617.
- 1911 Macrocystella mariae Callaway; Kirk p. 16, pl. 2, fig. 17.
- 1913 Macrocystella mariae Callaway; Springer, p. 157, fig. 249.
- 1927 Macrocystella mariae Callaway; Stubblefield and Bulman, pp. 111, 118.
- 1943 Macrocystella mariae Callaway; Bassler and Moodey, pp. 27, 175.
- 1952 Macrocystella mariae Callaway; Termier and Termier, p. 363, fig. 9.
- 1953 Macrocystella mariae Callaway; Cuénot, p. 618, fig. 15.
- 1955 Macrocystella mariae Callaway; Sdzuy, p. 270, pl. 1, fig. 14.
- 1964 Macrocystella mariae Callaway; Castell, p. 58, pl. 3, fig. 6.

*Diagnosis*. A species of *Macrocystella* of small size; with 10–15 brachioles, triangular in section; circular outer proximal columnals with thin, blade-like external flanges.

*Type.* BU 409 (Pl. 113, fig. 2) is selected as lectotype. It is possibly the original of Callaway 1877, pl. 24, fig. 13 and is from the Shineton Shales of Shineton, Shropshire. Parts of other specimens on this slab are accepted as paralectotypes.

*Horizon and locality*. Stubblefield and Bulman (1927) record *M. mariae* from the *Clonograptus tenellus* and *Shumardia pusilla* zones (Middle and Upper Tremadoc respectively) of the Wrekin. *M. mariae* is also recorded from the *Shumardia pusilla* zone of Arenig (Fearnsides, 1905, p. 617) and *Macrocystella* sp. from the same zone near Portmadoc, Caernarvonshire (Fearnsides 1910, pp. 161–2).

*Material*. Crushed remains of four more or less complete thecae, one complete stem and many isolated thecal plates and fragments.

# Description. A. Stem

The stem has a proximal and a distal portion. One complete proximal portion (Pl. 112, fig. 8) has 20 outer proximals each with a blade-like unornamented external flange. This crushed portion tapers from 5 mm. adorally to 2 mm. in approximately 15 mm. At the junction with the distal stem small distal columnals appear between the flanged columnals. Throughout the preserved portion of the distal stem, flanged and unflanged columnals alternate but this alternation becomes less obvious distally.

The distal portion of the stem (Pl. 111, fig. 1) tapers gradually from 1 mm. proximally to 0.5 mm. at the tip. It is about 35 mm. long. The topmost distal columnal is a thin annulus; the terminal distals are cylindrical and about three times as high as wide. There is no alternation of flanged and unflanged distals in this stem. Preserved proximal stems are straight or quite strongly curved. Curved distal stems are also preserved but there is no evidence to support Bather's (1900) interpretation of the distal stem with a distinct distal coil.

Counterparts of isolated columnals indicate the mode of construction and articulation of the stem. Both proximal and distal portions are composed of two types of columnals. Outer proximals (text-figs. 1a-b) are annular. Each has a smooth, sharpedged outer flange and a narrow inner flange (text-fig. 2, Pl. 112, fig. 7). Outer proximals alternate with inner proximals and the latter abut against the inner flanges of the former (text-fig. 2). The inner wall of each outer proximal has two sockets set opposite each other on both the upper and lower surfaces. The inner flanges are thickened adjacent to these sockets to form fulcra (text-fig. 2). If the sockets on one surface are orientated N.-S., those on the opposite surface of the same columnal lie NW.-SE. (text-figs. 1a-b). Both upper and lower surfaces of the inner proximals are flattened to form facets,

#### EXPLANATION OF PLATE 111

Stereophotos of Macrocystella mariae Callaway and M. azaisi (Thoral)

Fig. 2. *M. azaisi* (Thoral). Proximal and part of distal stem to show ornament of flanges on outer proximals. BMNH E23697.

All figures of latex impressions whitened with ammonium chloride sublimate. All  $\times 2$ .

584

Figs. 1, 3–6. *M. mariae* Callaway. 1. Complete stem showing proximal and distal portions. BMNH E29113. 3. Left lateral view of crushed theca. BMNH E29110a. 4. Right lateral view of same theca. BMNH E29109a. 5. Anterior lateral view of another crushed theca. BMNH E29109b. 6. Posterior view of crushed theca to show outline of periproct and small periproctal plates. BMNH E29113.

# Palaeontology, Vol. 11

# PLATE 111



PAUL, Macrocystella



at the same two opposite points (text-figs. 1c, 2). The outer margin of an inner proximal protrudes at these points and the protrusions key into the sockets in the inner wall of the outer proximals above and below (text-fig. 2). The fulcra on the outer proximals and the facets on the inner proximals articulate and the axis of articulation changes by approximately  $45^{\circ}$  with each outer proximal. This results in a right-handed spiral arrangement of articulation facets in *M. azaisi* and presumably in other *Macrocystella*.



TEXT-FIG. 1. Proximal stem columnals of *Macrocystella mariae* Callaway. *a*, *b*. Opposite sides of one outer proximal columnal (OP) in the same orientation to show different orientations of fulcra (Fu). *c*. Inner proximal columnal (IP). Ef, external flange; Fa, facet; If, internal flange.



TEXT-FIG. 2. Diagrammatic reconstruction of part of proximal stem of *Macrocystella mariae* Callaway to show arrangement of outer (OP) and inner (IP) columnals and spiral arrangement of facets (Fa) and fulcra (Fu). Ef, external flange; If, internal flange; IW, inner wall of outer proximal columnal. This has been drawn as a left-handed spiral although *M. azaisi* is known to show a right-handed spiral.

Each inner proximal is keyed into the sockets of the outer proximals above and below. Thus although highly flexible, the stem was quite resistant to rotation about its axis. Both inner and outer proximals are annular and the proximal stem has a wide lumen. Flexing of the proximal stem was probably achieved by muscles housed in this lumen. The mechanical keying of the columnals prevented rotation which would have sheared such muscles.

Larger (sometimes flanged) and smaller (unflanged) distals alternate in the distal stem but this becomes less apparent distally. The most proximal distals, which appear to be

newly formed, are annular but most distals are cylindrical. All distals have narrow lumina and the articulating surfaces are smooth (Pl. 112, fig. 3).

The nature of the outer proximals may prove to be a useful specific character. In M. mariae the outer proximals are approximately circular with thin, blade-like outer flanges (Pl. 111, fig. 1, Pl. 112, fig. 7). In M. bavarica the outer flanges are also thin and sharp-edged but are produced into ten angles or incipient spines (Sdzuy, 1955, pl. 1, figs. 8–10, text-fig. 1h). In M. azaisi the outline is circular but the flanges are thicker and have fine irregular granules or spines encircling them (Pl. 111, fig. 2, Pl. 113, figs. 1, 8). The outer flanges of M. sp. nov. from Mexico have four rounded lobes arranged in two pairs.

# в. Theca

All known thecae are crushed and it is impossible to describe all thecal plates from one specimen. BMNH E29109–10 are counterparts which show two crushed thecae in different orientations. There is another theca which shows details of the periproct.



TEXT-FIG. 3. Diagrammatic reconstruction of plate arrangement of *Macrocystella mariae* Callaway. Based on BMNH E29109a-b, E29110a-b, and E29113. B1-B4 basals, IL1-IL5 infra-laterals, L1-L5 laterals, R1-R6 radials.

A composite plate arrangement and a reconstruction based on these specimens are depicted in text-figs. 3 and 15. The theca was composed of five circlets of plates, four of which can be seen in M. mariae. The subvective system was confined to the oral surface from the margins of which the brachioles arose in five groups.

Some details of text-fig. 3 are restored but all plates shown existed. The four basals (BB) unite aborally to form an invagination around the stem. One basal (presumably B4) was hexagonal (Pl. 112, fig. 2). The infra-laterals (ILL) form a closed circlet; IL4 and IL5 contribute to the periproct border. IL1, IL2, and IL3 are roughly hexagonal. The five laterals (LL) apparently form a closed circlet; L1, L4, and L5 contribute to the periproct border. L5 is distinctly smaller than the other laterals and has a radial (R5)

586

directly adoral to it. One pair of counterparts (Pl. 111, figs. 3, 4) seem to have three hexagonal laterals which means either there were six laterals in this specimen or only L1 and L5 contributed to the periproct border. Unfortunately the relevant portion of this theca is crushed and this appearance may be misleading. The six radials (RR) form a closed circlet; one is directly adoral to L5. There are seven orals (OO) in M. *azaisi.* 



TEXT-FIGS. 4–5. Camera lucida drawings of the periproct (Pe) of *Macrocystella mariae* Callaway. 4. BMNH E29110b cf. Pl. 2, fig. 9. 5. BMNH E29113 cf. Plate 1, fig. 6. IL4–IL5 infra-laterals, L1, L4, L5 laterals.

The mouth, gonopore, and hydropore have not been detected in M. mariae but were all on the oral surface in M. azaisi (text-figs. 11a, b). Critical details of the periproct show in BMNH E29110b, and more clearly, in BMNH E29113 (text-figs. 4, 5). There are five plates around the periproct which is large. The periproct was covered by a thin, flexible, plated integument in life. Some small periproctal plates are preserved (Pl. 111, fig. 6) but the position of the anal pyramid in unknown.

The outlines of the individual thecal plates vary with their positions in the theca. Periproct border plates can be recognized easily, as can basals and radials. The remaining five laterals and infra-laterals are difficult to distinguish from each other. All plates have raised umbones from which ridges radiate to the middles of the sides, connecting centres of adjacent plates (text-fig. 6). Auxiliary ridges are developed parallel to the primary ridges to form 'rhombs'. These 'rhombs' are not true rhombs as they are composed of folds in the thecal plates. There is no development of thin-walled thecal canals. The external ridges of Macrocystella are formed by the folds in the plates and are not solid strengthening struts such as occur in *Cheirocrinus*. The thecal plates of M. mariae are approximately 0.1 mm. thick.

C 5934



TEXT-FIG. 6. Isolated thecal plate of *Macrocystella mariae* Callaway to show folds. BMNH E29119.

# c. Subvective system

The exothecal portion of the subvective system consists entirely of brachioles. These are long, slender, biserial, unbranched structures. The most complete brachioles (Pl. 112, fig. 10) have approximately 120 brachiolar plates and are slightly longer than the thecal height (up to 16 mm.). The brachioles have a triangular cross-section with sides twice the width of the adoral surface (text-fig. 7). The food groove ran down the centre of the adoral surface and was covered by lappets in life. The lappets are between one and a half times and twice as numerous as the brachiolars and apparently were flexible. Some have been preserved covering the food groove; others in an 'open' position (Pl. 112, fig. 10). The lappets alternate and each one imbricates over its more distal neighbour (text-fig. 8).

### EXPLANATION OF PLATE 112

Figs. 1–3, 5–10. *M. mariae* Callaway. 1. Internal and external views of two isolated thecal plates. The internal view (above) shows the folds in the plates. The other plate was one of five bordering the periproct (probably L1). BMNH E29112,  $\times$  5. 2. External view of B4 and an isolated distal columnal. BMNH E29112,  $\times$  5. 3. Three isolated thecal plates. All originally bordered the periproct. BMNH E29112,  $\times$  5. 5. Internal mould of isolated thecal plate BMNH E29119,  $\times$  3. 6. Inner proximal columnal within outer proximal columnal. BMNH E29112,  $\times$  2. 7. Outer proximal columnal showing inner flange and fulcra. BMNH E7574,  $\times$  1·5. 8. Crushed stem and theca. Note the larger distals are flanged. BMNH E7574,  $\times$  1·5. 9. Stereophotos of posterior lateral view of crushed theca showing periproct. Counterpart to Plate 1, fig. 5. BMNH E29110b,  $\times$  2. 10. Detail of brachioles to show lateral, aband adoral views. BMNH E29109,  $\times$  3.

Fig. 4. *M. azaisi* (Thoral). Lateral view of theca and stem. BMNH E23697,  $\times 2$ .

All except Fig. 5 latex impressions, all whitened with ammonium chloride sublimate.

# Palaeontology, Vol. 11



PAUL, Macrocystella



They coalesce towards the margins of the brachiole to form a continuous narrow band (text-fig. 9). The free portions were able to curl in on themselves. This curling in is not preservational and the lappets may have been only partially calcified in life.



TEXT-FIGS. 7–9. Brachioles of *Macrocystella mariae* Callaway. 7. Diagrammatic section through brachiole without lappets. 8. Camera lucida drawing of portion of brachiole with lappets (La) closed over food groove. Note that the lappets alternate and imbricate. 9. Camera lucida drawing of aboral view of brachiole with lappets in 'open' position. Br brachiolar plate. Text-figs. 8–9 based on BMNH E29113.

No more than three brachioles arise in any one ambulacrum and there were presumably 10–15 brachioles in all. There is no evidence to support Bather's (1900, fig. 18) reconstruction of branched brachioles. BMNH E29110a shows three brachioles in one radius (Pl. 111, fig. 3): all three are separate entities from their origin at the margin of the theca.

*Macrocystella mariae* Callaway is characterized by the following: 1. A stem which is divisible into two portions: a short, rapidly tapering, highly flexible, proximal portion composed of two types of annular columnals with a wide lumen; and a long distal portion composed of cylindrical columnals with a narrow lumen.

2. A theca with plate formula 4BB, 5ILL, 5LL, 6RR, ?OO.

3. A large periproct surrounded by five thecal plates and covered with a flexible plated integument.

4. Biserial, unbranched brachioles which arise from the margins of the flat oral surface and are grouped into five ambulacra.

# COMPARISON WITH MIMOCYSTITES BARRANDE

*Mimocystites bohemicus* Barrande has a subvective system which is confined to the flat oral surface and consists of about twenty brachioles grouped into five ambulacra.

Three brachioles were present in one radius of an available specimen (UC latex impression of original of Barrande 1887, pl. 28 (I), fig. 14). The theca is composed of 4BB, 5ILL, 5LL, 6RR, and some OO. Jaekel's analysis of the arrangement of thecal plates (1899, p. 201, fig. 36) is reproduced here in a slightly modified form (text-fig. 10). This interpretation differs slightly from that of *Macrocystella mariae*. Only two laterals



TEXT-FIG. 10. Jaekel's (1899) interpretation of the plate arrangement of *Mimocystites bohemicus* Barrande with modern notation of plates. B1–B4 basals, IL1–IL5 infra-laterals, L1–L5 laterals, R1–R6 radials. Cf. text-fig. 3.

(L1 and L5) contribute to the periproct border and L1 has a straight upper border with R6 directly adoral to it. This is an unexpected arrangement. Two specimens in the Schary collection (MCZ) apparently show the arrangement figured for *Macrocystella mariae*. However one specimen of *M. mariae* (Counterparts BMNH E29109a and 29110a) has apparently three hexagonal laterals as shown in Jaekel's figure of *Mimocystites bohemicus*. It seems possible that the plate arrangement varied slightly in different specimens. *Mimocystites azaisi* Thoral has a plate arrangement identical to that in *Macrocystella mariae*. Both the present and Jaekel's interpretations agree in most respects, particularly in the five plates around the periproct.

#### EXPLANATION OF PLATE 113

Stereophotos of *M. mariae* Callaway, *M. azaisi* (Thoral) and *M. azaisi multicristata* (Thoral). Fig. 2. *M. mariae* Callaway. Lateral view of lectotype. BU 409.

590

<sup>Figs. 1, 3, 5, 8. M. azaisi (Thoral). 1. Anterior lateral view of theca to show well-developed folds in B2 and ornament of stem flanges. CU. 3. Lateral view of another theca. CU. 5. Oral view of same.
8. Lateral view of another theca to show well-developed folds in B2 and ornament of stem flanges. CU.</sup> 

Figs. 4, 6, 7. *M. azaisi multicristata* (Thoral). 4. Oblique oro-lateral view to show ornament of orals and lateral food grooves alternating in ambulacrum IV (left). CU. 6. Interior view of oral surface of same theca. 7. Lateral view of same theca to show more strongly developed folds in radial plates. All figures of latex impressions whitened with ammonium chloride sublimate. All ×2.

# Palaeontology, Vol. 11

# PLATE 113



PAUL, Macrocystella



The thecal plates of *Mimocystites bohemicus* and *M. azaisi* are identical to those of *Macrocystella mariae* except in the number of folds, which is variable in each species.

The proximal stem of *Mimocystites bohemicus* is identical to that of *Macrocystella mariae* except that the outer proximals have thicker flanges which are less blade-like. *Mimocystites azaisi* has still thicker flanges, the peripheries of which are granulose or spinose (Pl. 111, fig. 2, Pl. 113, figs. 1, 8).



TEXT-FIG. 11. Camera lucida drawing of oral surface of *Macrocystella azaisi* (Thoral). *a*, entire surface to show oral plates and arrangement of food grooves. Cf. Plate 113, fig. 5. CU. *b*, detail of gonopore and hydropore area of same. FG, food groove; G, gonopore; H, hydropore; LFG, lateral food groove; M, position of mouth which was probably much larger than shown; O1–O7, orals. I–V, ambulacra; I<sup>1</sup>, I<sup>2</sup>, I<sup>3</sup>, facets of ambulacrum I; IV<sup>1</sup>, IV<sup>2</sup>, facets of ambulacrum IV.

The type species and one other species of *Mimocystites* therefore exhibit all the features which characterize *Macrocystella mariae*. All three are congeneric.

Specimens from the Montagne Noire, France (UC latex impressions) have yielded additional information on the morphology of *Macrocystella*. One example of *M. azaisi* (Thoral) has an almost complete oral surface (text-fig. 11*a*, Pl. 113, fig. 5). Another

example of *M. azaisi multicristata* (Thoral) shows both the internal and external surfaces of the oral area (text-fig. 12, Pl. 113, figs. 4, 6). There were seven orals, arranged as shown in text-fig. 11*a*. A slit-like hydropore and an oval gonopore are developed across the common suture of O1 and O7 (text-fig. 11*b*). There are five main ambulacral grooves each of which has lateral branches leading to brachiole facets. Apparently the branches are consistently to the left of ambulacra I and IV in the example of *M. azaisi* (Pl. 113, fig. 5) but regularly alternate in ambulacrum IV in the example of *M. azaisi multicristata* (Pl. 113, fig. 4). Details of the other ambulacra are not well preserved in either specimen.



TEXT-FIG. 12. Camera lucida drawing of internal oral surface of *Macrocystella azaisei multicristata* (Thoral).
cf. Plate 113, fig. 6. CU. G, supposed internal opening of gonopore; H, supposed internal opening of hydropore; M, mouth; O1–O7, orals; I–V ambulacra.

In internal view the mouth is large (4.6 mm.  $\times$  1.7 mm.) and is covered by ambulacral cover plates. The orals and ambulacral flooring plates are visible (Pl. 113, fig. 6). The latter are between, not on, the orals and form part of the thecal wall. Only primary ambulacral flooring plates can be detected. Close to the mouth is a deep pit which apparently connected to the hydropore. This pit is separated from the mouth and the supposed gonopore by two internal ridges, one on each side. The gonopore is apparently represented by a small circular pit some distance from the mouth (text-fig. 12). Unfortunately the critical area of the external oral surface is not preserved and it is not possible to match up external and internal openings. Ambulacra IV

and V are more deeply impressed than I, II, and III. Another pit is developed obliquely under the aboral of the two internal ridges near the hydropore (Pl. 113, fig. 6). The significance of this is unknown.

# COMPARISON WITH OTHER PELMATOZOA

Several authors have grouped *Macrocystella* with *Lichenoides* while others have placed them in separate families. In addition *Macrocystella* has been variously assigned within the Pelmatozoa.

The most recent and most complete account of the morphology of *Lichenoides* is that of Ubaghs (1953) who showed that it completely lacks a stem. The thecal plates are arranged in four circlets but the total number is variable, 5–12 basals, 5 infra-laterals, 5–7 laterals, and 5–7 radials. (The homologies with the Glyptocystitida implied by the use of the same terms for the plate circlets are unjustified.) All the infra-laterals, laterals, and radials bear epispires. These are a type of pore-structure with a single sutural pore which leads to a narrow channel in the external surface of both adjacent plates (see Ubaghs 1953, figs. 3, 11). There is no lateral periproct in *Lichenoides*. The brachioles arise from both lateral and radial plates and apparently they are not grouped into five radii.

Thus while *Lichenoides* resembles *Macrocystella* in having definite plate circlets and biserial, unbranched brachioles it differs in the absence of a stem and lateral periproct, in the presence of epispires and brachioles on lateral and radial plates, and in the total number and position of the thecal plates. These differences are considered to be important taxonomically. The Lichenoidae and Macrocystellidae are maintained as separate families as suggested by Jaekel (1918) and Ubaghs (1953).



TEXT-FIG. 13. Plate arrangement in *Cheirocrinus radiatus* Jaekel. Based on Jaekel, 1899, p. 213, fig. 36. B1–B4, basals; IL1–IL5, infra-laterals; L1–L5, laterals; R1–R6, radials.

Among cystoids *Macrocystella* most closely resembles *Cheirocrinus* Eichwald. This latter genus is characterized by a stem with proximal and distal portions. The construction and articulation of the stem are identical to that of *Macrocystella* and the helical arrangement of the articulations of the proximal stem was described by Billings (1858) in *Cheirocrinus anatiformis* (Hall) (= *Glyptocystites logani* Billings).

The theca of *Cheirocrinus* is composed of 27 plates arranged in 5 circlets: 4BB, 5ILL, 5LL, 6RR, and 7OO. The periproct is large, surrounded by 5 thecal plates (IL4, IL5, L1, L4, and L5) and was covered by a flexible plated integument in life. L5 is directly adoral to the periproct and has R5 directly adoral to it (text-fig. 13). The subvective system is restricted to the flat oral surface and there are 20–5 brachioles grouped into 5 ambulacra whose flooring plates lie between the orals, not on them. *Cheirocrinus* differs from *Macrocystella* in the possession of pectinirhombs.

*Macrocystella* and *Cheirocrinus* have in common many distinctive features of which perhaps the most important is the detailed structure of the stem. This type of stem is characteristic of and confined to the superfamily Glyptocystitida. It is most unlikely that such a complex organ developed independently in two groups which share other common characteristics. *Macrocystella* probably gave rise to *Cheirocrinus* and through it to the other Glyptocystitida as originally suggested by Jaekel (1899).

Thoral (1935) thought it possible to recognize in the Montagne Noire a lower horizon with *Macrocystella azaisi* (upper Tremadoc) and a higher horizon with *Cheirocrinus languedocianus* (basal Arenig). In Britain *M. mariae* occurs below the oldest *Cheirocrinus. Macrocystella* seems to be a characteristic fossil of the Tremadoc whereas the oldest known *Cheirocrinus* are all Arenig. Stratigraphic evidence agrees with the idea that *Macrocystella* evolved into *Cheirocrinus*.

In the past the main objections to the inclusion of *Macrocystella* in the Rhombifera were its branched arms and lack of rhombs. The former was an error but the latter is more important. Regnéll (1945) has stressed that the main character which unites the cystoids as a class is the presence of pore-structures. Detailed study of cystoid pore-structures (Paul, 1968) suggests the Rhombifera should be regarded as a distinct class. The rhomb-less *Macrocystella* is included in the Rhombifera on the same grounds that led Kesling (1963) to include *Amecystis* in the Rhombifera. *Amecystis* is effectively a *Pleurocystites* without pectinirhombs, just as *Macrocystella* is a *Cheirocrinus* without pectinirhombs. The many similarities outweigh this single distinction.

# THE EVOLUTION OF PECTINIRHOMBS

The rhombs of *Cheirocrinus* are fully developed pectinirhombs (which Bather regarded as a highly specialized type of rhomb) even in the earliest species known. There is no evidence of a gradual evolution of pectinirhombs. If *Macrocystella* evolved into *Cheirocrinus*, pectinirhombs either appeared 'suddenly' or pre-existing structures broke through the thecal plates to appear as pectinirhombs. The internal surfaces of isolated plates in both *Macrocystella* and *Cheirocrinus* show no features which could be incipient pectinirhombs. Pectinirhombs were functional throughout their growth; they are present as external features from the earliest stages. They did not develop internally and become external features later in growth. Rather sudden appearance therefore seems more likely.

All rhombs have generally been accepted as respiratory organs. In the simplest case respiration would have taken place through the thecal wall. The amount of oxygen required would have been proportional to the volume, and the amount of respiratory exchange to the surface area, of the theca. The oxygen requirements would thus increase with growth faster than the amount of exchange. This difficulty can be overcome, without materially altering the over-all thecal shape, by the production of evaginations or invaginations of the thecal wall. *Macrocystella* has the former in the folds of the thecal plates. The dichopores of pectinirhombs are invaginations and produce a slightly better volume to surface area ratio.

Exchange is facilitated by a large surface area and a thin exchange surface. Either invaginations or evaginations are almost equally effective in increasing the surface area but the latter are exposed and liable to mechanical damage. The ridges in the thecal plates of *Macrocystella* probably facilitated exchange by increasing the surface area. The thecal plates were extremely thin (0·1 mm.) and although strengthened by the ridges they were still very fragile. The dichopores of pectinirhombs are within the theca and therefore protected. Dichopore walls are much thinner than thecal plates (usually 0·01 mm.). In *Macrocystella* the entire thecal wall probably took part in exchange. In *Cheirocrinus* a differentiation of function is seen. Without decreasing the amount of exchange it



Paul, C. R. C. 1968. "Macrocystella Callaway, the earliest glyptocystitid cystoid." *Palaeontology* 11, 580–600.

**View This Item Online:** <u>https://www.biodiversitylibrary.org/item/196372</u> **Permalink:** <u>https://www.biodiversitylibrary.org/partpdf/172969</u>

**Holding Institution** Smithsonian Libraries and Archives

**Sponsored by** Biodiversity Heritage Library

**Copyright & Reuse** Copyright Status: In Copyright. Digitized with the permission of the rights holder License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.