

LOWER KIMERIDGIAN AMMONITES FROM THE DRIFT OF LINCOLNSHIRE

by the late W. J. ARKELL and J. H. CALLOMON¹

ABSTRACT. A collection of ammonites is described from the drift of Lincolnshire; they are of Lower Kimeridgian age, Mutabilis Zone. Almost all of the specimens belong to the genera *Rasenia* and *Aulacostephanus*, and include several new species which are grouped into two new subgenera of *Aulacostephanus*: *Xenostephanus* and *Xenostephanoides*. These new forms have only recently been found *in situ* in England, and are known from the Kimeridge Clay of both east and west Scotland.

THIS paper describes a collection of ammonites made mostly by Dr. R. G. Thurrell while mapping in the Lower Cretaceous of north Lincolnshire, and now deposited in the Sedgwick Museum, Cambridge. The material is all from the drift, and hence does not provide primary stratigraphic evidence; its publication seems of interest, however, for a number of reasons. The various forms were found in such intimate admixture, often in the same block of stone, that there can be little doubt that they give a fairly representative picture of a closely contemporaneous population. Their age is Lower Kimeridgian, Mutabilis Zone. The material is in rock-facies and only slightly crushed, whereas the known exposures in the outcrop of the shales of the Mutabilis Zone in this country yield, mostly, only wholly flattened shells, which have not attracted collectors in the past. Our knowledge of the fauna of the Mutabilis Zone is therefore scanty.

The fauna consists overwhelmingly of forms of the perisphinctid subfamily Aulacostephaninae at the stage where *Rasenia* changes to *Aulacostephanus*, and includes several new and previously little-known forms of great interest. The classification of the genus *Rasenia* is at present still in a state of confusion, but a revision of *Aulacostephanus* has been published by Ziegler (1962). New names will therefore be mostly confined here to the forms being described.

In referring to specimens the following abbreviations will be used:

R, MS, SW: Thurrell collection, material from Ranby, Market Stainton, and South Willingham (Lincolnshire) respectively.

BM: British Museum (Natural History).

SM: Sedgwick Museum, Cambridge.

We wish to thank Dr. R. G. Thurrell for making his material available to us for description; Dr. B. Ziegler for discussions and criticisms, and for very kindly allowing us to see the manuscript of his monograph on *Aulacostephanus*; and Mr. A. G. Brighton (Sedgwick Museum), Dr. M. K. Howarth (British Museum), and Mr. R. B. Wilson (Geological Survey, Edinburgh) for help with material in collections in their custody.

¹ This paper is based in part on a provisional manuscript prepared by Dr. Arkell shortly before his illness in 1956. I am grateful to Mr. A. G. Brighton for passing the manuscript on to me for completion after Dr. Arkell's death in 1958. J. H. C.

SYSTEMATIC DESCRIPTIONS

Family PERISPHINCTIDAE Steinmann 1890

Subfamily AULACOSTEPHANINAE Spath 1924 (= RASENIINAE Schindewolf 1925)

This subfamily encompasses a well-defined offshoot of the main perisphinctid tree. It leads back to such typically perisphinctid forms as *Pictonia* in the basal Lower Kimeridgian, with evolute coiling, round whorl-section, and smooth, regular ribbing which passes over the venter without interruption. It terminates in end-forms of *Aulacostephanus* which are of quite different appearance, with compressed, angular whorl-section, smooth tabulate or grooved venter, and strong straight ribbing reminiscent in style of the Cretaceous genus *Hoplites*. Between these extremes, further wide variations in coiling and novel forms of ornamentation occurred, so that the whole subfamily, although in itself closely delimited, covers an unusually wide range of forms. Attempts at classification have usually started from a broad division of the subfamily into two parts: an earlier part grouped around the genus *Rasenia*, leading to the later forms grouped around *Aulacostephanus*. Besides these, there has, however, sprung into existence a large number of other generic names created often on the basis of insufficient material and stratigraphy, inadequate type specimens, and the requirements of phylogenetic theory. The resulting confusion cannot be fully analysed here, but, in order to justify the classification which will be adopted in this paper, a number of points have to be discussed.

(a) *Generic classification.* It seems sufficient to divide the subfamily into only two genera: *Rasenia* and *Aulacostephanus*. Further grouping according to sculpture can be conveniently incorporated at subgeneric level.

(b) *Dimorphism.* In common with apparently most other perisphinctids, the Aulacostephaninae seem to have been dimorphic, and it is possible to classify most of the shells as macroconchs or microconchs (for full discussion, see Callomon 1957, 1963). As elsewhere, it seems helpful to incorporate this distinction at subgeneric level.

(c) *The Rasenia–Aulacostephanus boundary.* The transition between these genera is so gradual that, in drawing a dividing-line, many difficulties arise. Development occurred in more than one character of the sculpture, and the breaks in the development of different characters occurred at different times. If the development of a single character were made the sole criterion, an unnatural classification would result. Thus, for example, the presence or absence of a ventral smooth band alone might seem a tempting distinction to use as criterion; but, as is in fact seen in the material being here described, among a collection of shells which in most other respects are clearly conspecific, some have a ventral smooth band, others not. Moreover, many have it at one stage of their ontogeny, but not throughout. The best approach seems to be to group specimens of a contemporaneous population according to species, taking all their characters into account. A particular arrangement of species into genera is then of lesser importance. The arrangement of borderline species used here has been chosen to accord fairly closely with the one used by Dr. Ziegler in his monograph of the genus *Aulacostephanus*, but no claims are made for its uniqueness. It serves also to bring out resemblances between the Lincolnshire material and the fauna of the Mutabilis Zone farther south.

Genus RASENIA Salfeld 1913

Type species. *Rasenia involuta* (Salfeld MS) Spath 1935, designated by ICZN Opinion 426.

Microconchs. The following subgenera may be recognized, arranged in order of decreasing coarseness of ribbing.

1. PRORASENIA Schindewolf 1925. Type species: *P. quenstedti* Schindewolf, figured 1926, pl. xix, fig. 1. (Synonym *Desmosphinctes* Schindewolf 1925, type species: *Perisphinctes mniovnikensis* Nikitin 1884, p. 30, pl. ii, figs. 11, 12.) Small, strongly ribbed species with lappets. Its special character is the presence of strong triplicate ribbing on the inner whorls, reverting to biplicate perisphinctoid on the final body-chamber. Spath (1935, pl. 14, 15) has figured and named three species from England, and Waterston (1951, pl. 2) records them from Scotland.

2. RASENIA sensu stricto. Salfeld attached the manuscript name *involuta* to a number of specimens in the British Museum, of which Spath selected and figured only one. As, however, Spath is formally the author of the species, this specimen is holotype of *involuta* Spath rather than lectotype of *involuta* Salfeld, and both species and genus have to be interpreted by Spath's figured specimen. This is broken and slightly incomplete, and, in the absence of the last few sutures and peristome, it is impossible to tell conclusively whether the specimen is adult or not. All of the preserved last three-quarters whorl is, however, body-chamber, and it gives the impression near the end of slight contraction indicating maturity. There exist in the collections other specimens (e.g. Geol. Survey Coll. 25550-1) which agree with the holotype closely in all respects including size, and which, although also without peristomes, carry similar amounts of body-chamber. A complete adult from Scotland has been figured by Waterston (1951, pl. ii, fig. 8a, b). Its final diameter is 40 mm., and the body-chamber is terminated by a final constriction. The peristome shows the stump of a lappet. It seems certain, therefore, that *Rasenia* s.s. was fully grown at c. 40 mm. diameter with strong unmodified ribbing maintained to the end, and hence microconch; and it seems probable that it carried at least short lappets. Its special character is the differentiation of strong, sinuous primary and fine, fasciculate secondary ribbing. Inner whorls are always well covered, involute and inflated.

In a recent revision of the genus Geyer (1961, p. 86), rather than re-examine the type specimen of *involuta* Spath himself, quotes Spath's statement describing it as 'small, inner whorls . . .', which, whatever significance one may attach to size in taxonomy, it is not; and interprets *Rasenia* sensu stricto as covering large, typically macroconch forms. *Rasenia* s.s. is now capable of precise interpretation, and it is hoped that the confusion of the past has found an end.

Geyer does point out that, if *Rasenia* sensu lato is taken to include *Involuticeras* Salfeld as a subgenus (see below), *involuta* Spath 1935 is junior homonym of *involuta* (Quenstedt 1849), and proposes *Rasenia anglica* nom. nov. = *involuta* Spath non Quenstedt.

3. RASENIOIDES Schindewolf 1925, p. 335. Type species by original designation: *Nautilus striolaris* Reinecke 1818, which Schindewolf states should be interpreted by *Am. striolaris* Quenstedt pars 1888, pl. 124, fig. 8, non Reinecke. Reinecke's specimen seems to be lost, but his figure represents recognizable characters which seem to differ considerably from

those shown by Quenstedt's figure, notably in whorl-section and coiling. There is nothing in Reinecke's description to indicate a smooth venter, which, in this case, puts Quenstedt's specimen in *Aulacostephanus*; and, as small 'striolarian' Rasenids of the *thermarum* group are common in the Mutabilis Zone (see below), *Rasenioides* fills a useful role, interpreted here as microconch subgenus characterized by very fine, dense fasciculate ribbing. Lappets proper are not known, although some specimens carry a slightly raised sinuous collar on the mouth-border.

(PRORASENIOIDES Schindewolf 1925, p. 338; type species by original designation: *P. transitorius* Schindewolf, which is nom. nov. for *Am. striolaris* Quenstedt pars 1888, pl. 107, figs. 12, 13, non Reinecke, with additional specimens figured by Schindewolf 1926, p. 507, pl. xix, figs. 3-5, resembles *R. striolaris* Reinecke more than *Am. striolaris* Quenstedt 1888, pl. 124, fig. 8. However, as the latter is not strictly the type species of *Rasenioides* whereas, according to the Rules, the former is, *Prorasioides* and *Rasenioides* are here regarded as synonymous.)

Macroconchs

4. ZONOVIA Sasonov 1960. Type species by original designation: *Amm. uralensis* d'Orbigny 1845. Large, evolute, coarse-ribbed forms with smooth apertures.

5. EURASENIA Geyer 1961, p. 87, 90. Type species: *Amm. Rolandi* Oppel 1863. Large, coarsely ribbed forms intermediate between *Zonovia* and *Involuticeras*.

6. INVOLUTICERAS Salfeld 1913. Type species by monotypy (Salfeld 1914, p. 175) or subsequent designation (Spath 1931): *Amm. involutus* Quenstedt 1849. Large, involute, compressed forms, finely ribbed on inner whorls, becoming smooth.

7. ?SEMIRASENIA Geyer 1961, p. 87, 92. Type species: *Amm. Mörschi* Oppel 1863. The holotype of *mörschi* is 45 mm. in diameter, and if Oppel's statement in the legend to his figure, that the suture-line shown is the last, is to be believed, it carries three-quarters whorl body-chamber and is hence nearly complete. Geyer states the final size to be 'probably about 70 mm.' without giving reasons based either on Oppel's specimen, which he refigures but does not redescribe, or additional material, of which he cites only one additional, even smaller specimen. The peristome remains unknown, and maximum size and ontogeny of mature examples in doubt.

The only other species included by Geyer in *Semirasenia* is *Amm. thermarum* Oppel. The holotype of this species seems to us to fit equally well into *Rasenioides*, however.

Subgenus RASENIA s.s. Salfeld

1. *Rasenia* (*Rasenia*) sp. ind.

cf. *Rasenia evoluta* (Salfeld MS), Spath 1935, pp. 48-49, pl. 14, fig. 6a, b.

Material. One (MS 128).

Description. Diameter 17 mm., sutures invisible, but at least half the outer whorl appears to be body-chamber. Coiling moderately evolute; ribbing straight, coarse and strong, modifying on the last quarter-whorl to typically rasenid fasciculate style, passing uninterruptedly over the venter.

Spath assumed that *R. evoluta* was a large species (1935, p. 49, footnote). Like *R. involuta*, however, he figured only a single specimen (BM C.8046 from Market Rasen), which is therefore holotype. It is a wholly septate nucleus, maximum diameter only 24 mm., and on comparing the actual specimen with the type of *R. involuta* (which is broken and shows the inner whorls well) at similar diameters, it is hard to detect even the minutest differences between them. Whatever else Spath may have had in mind, *R. involuta* and *R. evoluta* as defined by the monotypes are the same, and the latter name may be dropped with little regret.

Subgenus RASENIOIDES Schindewolf (= PRORASENIOIDES Schindewolf)

1. *Rasenia* (*Rasenioides*) cf. *transitoria* (Schindewolf)

Plate 33, fig. 2

Ammonites striolaris Quenstedt 1888 pars, pl. 107, figs. 12, 13.

Prorasioides transitorius Schindewolf 1925, p. 338.

Material. One (MS 80a).

Description. The specimen is crushed, but agrees well with Quenstedt's figures. The ribbing passes uninterruptedly over the venter where visible. Sutures and peristome are missing, but at the maximum diameter of 32 mm. at least some of the outer whorl appears to be body-chamber. The umbilical seam uncoils slightly near the end, indicating maturity.

2. *Rasenia* (*Rasenioides*) cf. and aff. *lepidula* (Oppel)

Plate 32, figs. 19–21

cf. *Ammonites lepidulus* Oppel 1863, p. 242, pl. 67, fig. 4a, b.

Material. Five, more or less fragmentary (MS 1, 33–35, 53).

Description and comparisons. Oppel's figured type specimen was wholly inadequate to form the basis of a new species, only the final quarter-whorl being recognizable. Fortunately, this shows the specimen to have been a complete adult, and the name is used here for forms of the *R. thermarum* group which are rather larger and more evolute than *R. thermarum*, and in which the secondary ribs cross the round venter with perhaps some weakening but without actual interruption.

Agreement is best between Oppel's figure and the terminal fragment shown in Plate 32, fig. 21. This shows the peristome: the aperture is sinuous, preceded by a broad, shallow constriction and provided with a raised collar, as in *R. thermarum*, with no lappets; whereas Oppel's specimen was armed with lappets.

Oppel's species is very close to *Rasenia eulepida* Schneid (see below). This, however, has a prominent smooth band on the venter (*teste* Dr. Ziegler), but whether this serves to distinguish it from *R. lepidula* it is impossible to say in the absence of more information on the inner whorls of the latter. Schneid also stated that the distinguishing feature of *R. lepidula* was the persistence of the characteristic sheaves of multiple secondaries arising from each primary rib right to the end, whereas in *R. eulepida* near the aperture the ribbing reverts to simple bi- or triplication.

3. *Rasenia (Rasenioides) thermarum* (Oppel)

Plate 32, figs. 13–18

Ammonites thermarum Oppel 1863, p. 243, pl. 65, fig. 5a, b.*Rasenia (Semirasenia) thermarum* Geyer 1961, p. 106, pl. 8, fig. 9, (? 10).*Material.* Eight, and fragments (MS 9, 14, 31–32, 36, 64, 78, 118; R 20).

Description. All of the specimens seem to carry some body-chamber, and in one (Pl. 32, fig. 14) the peristome is preserved on one side. The impression that they are fully grown and nearly complete is supported by lack of remains of similar shells of larger ammonites. The average size of the eight specimens is 19.5 mm.: Oppel's holotype is 20 mm. in diameter, with nearly complete body-chamber, and he says: 'All the specimens found so far possess a nearly complete body-chamber. The partly visible peristome appears to be straight, without showing the formation of lappets.' The peristome observed here is also without lappet, but is sinuous and bears a slightly raised collar; it is preceded by a broad shallow constriction.

The Lincolnshire material therefore agrees well with Oppel's specimens and differs only in being slightly less inflated than the holotype. On some specimens the secondaries weaken on the venter, which is round, but they are not fully interrupted. Geyer considers *R. thermarum* to be a somewhat larger species, up to 40 mm. in diameter, effectively reinterpreting it on the basis of a specimen previously figured by de Loriol (1878, pl. xiii, fig. 5), which is variocostate like *Aulacostephanus möschi*. The total material available to him was only three specimens, however, including the holotype; and as there is no reason to believe that the holotype is in any way imperfect, incomplete, or untypical, but in fact representative of a readily recognizable group as shown by the Lincolnshire material, we prefer to continue to interpret *R. thermarum* as closely defined by Oppel's type, and retain it in *Rasenioides*.

4. *Rasenia (Rasenioides) cf. striolaris* (Reinecke)

Plate 31, figs. 6a, b

Nautilus striolaris Reinecke 1818, p. 77, figs. 52, 53.cf. *Rasenia (Rasenioides) striolaris* Geyer 1961, p. 10.*Material.* Two (MS 29, figured; MS 141).

Description. The figured specimen is only 16.5 mm. in diameter but carries half a whorl body-chamber. The umbilicus is only some 20 per cent. of the diameter, and inner whorls are almost wholly concealed. The second specimen is crushed, but similar.

Reinecke did not state the magnification of his figure, and his specimen seems to be lost. The present material agrees well with it, if we assume that his figure was about natural size. Geyer has designated and figured a neotype (1961, pl. 21, fig. 3) without, however, an adequate description. Thus, there is no mention of whether it, or supporting material, carries any body-chamber; there is no ventral view; and no indication of the form of the peristome. Moreover, the specimen differs considerably from Reinecke's figure in being much more evolute with very little inclusion of inner whorls, quite unlike both Reinecke's side-view and section.

Subgenus ZONOVIA Sasonov 1960

1. *Rasenia* (*Zonovia*) sp. ind.

Plate 33, fig. 3a, b

Material. One fragment (R 2).

Description. The specimen consists only of a quarter of two contiguous wholly septate whorls, corresponding to a diameter of about 45 mm. The whorl-section is round, inflated, depressed; the ribbing blunt, subdued, coarse. Short, straight bullate primaries trifurcate into strong secondaries, which cross the venter with slight weakening but no interruption.

Genus AULACOSTEPHANUS Sutner and Pompeckj in Tornquist 1896

Type species. *Ammonites mutabilis* d'Orbigny non Sowerby = *Ammonites pseudomutabilis* de Loriol 1874, designated by ICZN Opinion 302. A lectotype from the d'Orbigny collection has been figured by Ziegler (1962, pl. 15, fig. 8).

The boundary between *Rasenia* and *Aulacostephanus* is here being drawn such that most forms assigned to the latter have a smooth band on the venter in at least one stage of their ontogeny. Simultaneously, the whorl-section becomes more compressed and quadrate, the venter tending to flatten; and inner whorls are markedly more evolute and serpenticone than in *Rasenia* so that the secondary ribs on them are often partly exposed. The umbilici are in consequence also usually shallow, with often characteristically smooth, gently sloping or even barely perceptible walls. These features together allow a fairly sharp line to be drawn between the two genera.

The material from Lincolnshire falls into four subgenera: two pairs each consisting of a microconch and a macroconch. One of these pairs of subgenera, *Aulacostephanites* and *Aulacostephanoides*, differs but little from *Rasenia* and includes the forms well known from the Mutabilis Zone elsewhere. The other pair of subgenera, *Xenostephanoides* and *Xenostephanus*, is new. A few specimens have been known previously, but the presence of a considerable number in the material under consideration gives it a peculiar stamp of great interest.

Microconchs

1. AULACOSTEPHANITES Ziegler 1962. Type species: *Rasenia eulepida* Schneid 1939, p. 146, pl. v, figs. 13, 13a. Small, evolute forms with lappets; ornament similar to that of *Rasenioides*. Smooth band on venter, at least on inner whorls. In crushed material from the typical Mutabilis Zone shales in the south of England *Rasenioides* and *Aulacostephanites* may be indistinguishable.

2. XENOSTEPHANOIDES subgen. nov. Type species: *Aulacostephanus* (*Xenostephanoides*) *thurrelli* sp. nov. (see below). Small species (c. 40–50 mm.); coiling evolute, particularly on inner whorls; whorl-section rounded quadrate. Ribbing strong; short bullate primaries with sheaves of strong, straight secondaries often separated from the primaries by a smooth band on the whorl-side. The style of the ribbing is already much more like the 'hoplitid' of the later aulacostephanoids than the sinuous one of the rasenids. On the final body-chamber it may revert to simple biplication, as in *Prorrasenia*. The

venter carries a smooth band, sometimes accentuated by a groove. Peristomes with oblique final constrictions and short lateral projections not really long enough to call lappets.

Macroconchs

3. AULACOSTEPHANOIDES Schindewolf 1925. Type species, by original designation: *Amm. desmonotus* Oppel. The type specimen of Oppel's species (1863, p. 241, pl. 67, fig. 1a, b) seems to have been lost during the war. His figure establishes the style of ribbing, coiling, and umbilicus well, and shows clearly a smooth band on the venter. Oppel stated, however, that no sutures were visible anywhere, and that the peristome was unknown; and it is therefore impossible to say whether the specimen (the only one known to Oppel) was fully grown and complete, or merely the nucleus of a much larger form. This makes the species a highly unsuitable one on which to found a genus, and interpretations have been correspondingly varied. Dr. Ziegler has redescribed the species on the basis of additional material. Among this is one specimen (Ziegler 1962, pl. 2, fig. 14) which resembles the lost holotype extraordinarily closely, with unfortunately the same shortcomings. Others lead from this to more or less complete adults, which are large and nearly smooth with simple peristomes. We follow this interpretation of *A. desmonotus*, which is thus a macroconch.

The subgeneric characters of *Aulacostephanoides* are therefore: macroconch; rasenid ornamentation on the inner whorls which fades, becoming smooth on the final body-chamber; ventral smooth band on at least the inner whorls; simple peristomes. From general experience involuteness of coiling seems of lesser importance, and Dr. Ziegler shows that *Aulacostephanoides* can usefully accommodate a gradation of forms ranging from the discoidal *A. linealis* (Quenstedt) to the true evolute *A. mutabilis* (Sowerby). Several species of the subgenus differ but little from forms still included in *Rasenia*, the dividing-line from which is arbitrary, as discussed above.

4. XENOSTEPHANUS subgen. nov. Type species: *Aulacostephanus (Xenostephanus) ranbyensis* sp. nov. (see below). Large species (c. 120 mm.); coiling stout, depressed, and evolute—particularly on the inner whorls. Primary and secondary ribbing well differentiated on inner and middle whorls, the former tending to extreme bullae or tubercles on the umbilical margin, the latter characteristically strong, triplicate, with intercalatories, fusing into the primaries or sometimes separated from them by a narrow smooth band on the whorl sides. Venter flat on inner whorls, with smooth band, sometimes accentuated by a groove. On the body-chamber the ribbing becomes subdued and degenerates to more perisphinctid form: primary ribs biplicate smoothly or become merely single widely spaced ridges. The whorl-section becomes round, and ribbing passes over the venter without interruption. Peristomes simple.

Inner and intermediate whorls in some species are almost fully homeomorphic with the quite unrelated earlier genus *Reineckeia*; outer whorls are as in *Rasenia* (*Zonovia*), especially *R. uralensis* (d'Orbigny).

Xenostephanus and *Xenostephanoides* form a typical macroconch and microconch pair, and it would be difficult to find a clearer demonstration of the dimorphism among the perisphinctids. The inner whorls are so similar that it would be hard to sort them out in immature, incomplete, or fragmentary material.

Subgenus AULACOSTEPHANITES Ziegler 1962

1. *Aulacostephanus* (*Aulacostephanites*) cf. *eulepidus* (Schneid)

Plate 32, figs. 1–8

Rasenia eulepida Schneid 1939, p. 146, pl. v, figs. 13, 13a.*Aulacostephanus* (*Aulacostephanites*) *eulepidus* Ziegler 1962, p. 44, pl. 341, figs. 2–7, 8 *partim*, ? 11, 14.*Material.* Twenty-five (24 from MS; and R 5), and fragments.

Description and remarks. The Lincolnshire material seems to be consistently somewhat smaller than the type (33 mm.), for the average diameter of seventeen complete adults is 25 mm. (standard deviation 3.9 mm. = 15.5 per cent.). The style of ribbing, however, is the same, including the degeneration of the ribbing near the end, which Schneid cited as diagnostic compared with *Am. lepidulus* Oppel. Peristomes are preserved in several specimens. There is usually a slight terminal constriction and a raised mouth-border. Lateral extensions may be short (Pl. 32, figs. 3, 5), or long enough to qualify as lappets (figs. 2, 10). Smooth bands on the venters are generally present, but narrow and inconspicuous, and they die out on the last half-whorl of body-chamber.

The question of the generic position of the specimens here described raises in its most acute form the difficulty attending attempts to divide *Aulacostephanus* from *Rasenia*. The ventral smooth band is undoubtedly much less prominent here than in the type of *A. eulepidus* and other south German material, and the Lincolnshire forms might, on this account, rest more comfortably in *Rasenia* (*Rasenioides*), with possibly gradations to *R. thermarum* (Oppel). On the other hand, the style of coiling and ribbing resembles *A. eulepidus* much more closely than *R. thermarum* or any other *Rasenioides* so far described (with the possible exception of *R. lepidula* (Oppel), which we exclude from consideration because of the imperfection of the type); and there is gradation to the forms described as *A. ebrayoides* below, which have quite prominent smooth bands on the venter.

It seems therefore that we are here sampling the *Rasenia*–*Aulacostephanus* boundary at a slightly different stage from that observed in central Europe. There the distinction between *Rasenioides thermarum* and *Aulacostephanus eulepidus* is clear, and the latter grades into more coarsely ribbed forms which nevertheless retain the rasenid (fasciculate) style of secondary ribbing, and which Ziegler designates *A. peregrinus* nov. In contrast, these forms appear to be absent in the Lincolnshire material, which, instead, shows more or less gradation from *A. ebrayoides* (unknown in central Europe), via *A. eulepidus*, to *R. thermarum*. Whether these differences reflect slight differences in ages, or ecological factors (geographic subspeciation), it is at present hard to say.

A. eulepidus is also found in the Mutabilis Zone of Mull (cf. Pl. 33, figs. 4–6).

2. *Aulacostephanus* (*Aulacostephanites*) *ebrayoides* sp. nov.

Plate 30, figs. 14–18

Material. Nine (eight from MS, and R 16), and fragments.

Description. The holotype, MS 5 (Pl. 30, fig. 14) is 26 mm. in diameter, with half-whorl body-chamber preserved. The remaining material is of similar size. Coiling is evolute, the whorl-section quadrate. Ribbing is dense, but strong and sharp, straight, triplicate

or biplicate with intercalatories; the furcation point is very low on the whorl-side, so that the primaries (about twenty-two on the outer whorl) are reduced to short ridges on the umbilical margins. There is a distinct smooth band on the venter on inner whorls, fading somewhat on the body-chamber.

Comparisons. The species differs from *A. eulepidus* through the reduced contrast between primary and secondary ribbing, and the straightness and coarseness of the latter. Its name reflects the close homeomorphism with the Bathonian genus *Ebrayiceras* Buckman (Morphoceratidae), e.g. *E. pseudo-anceps* (Ebray) (see Arkell 1955, pl. xvii, figs. 7–11).

There are occasional specimens with more flexuous, but still coarse, secondary ribbing (Pl. 32, figs. 9, 10, with short lappets), which may link *A. ebrayoides* with *A. eulepidus*. These coarse forms of *Aulacostephanites* occur also in Mull (Pl. 33, fig. 7).

Quenstedt's *Amm.* cf. *striolaris* pars, 1888, pl. 107, fig. 18 seems to resemble *A. ebrayoides* closely in its ribbing, but differs in having a much more prominent ventral smooth band, and a deep terminal constriction (which may, however, have only varietal significance.)

3. *Aulacostephanus* (*Aulacostephanites*) sp. aff. *desmonotus* (Oppel)

Plate 32, figs. 11, 12

cf. *Ammonites desmonotus* Oppel 1863, p. 241, pl. 67, figs. 1a, b.

Material. Three (MS 76, 115; R 17).

Description and comparisons. All three specimens attain between 20 and 23 mm. and appear to carry some body-chamber. They resemble the other specimens described here under *Rasenia thermarum* in ornament, but are distinctly more compressed, have prominent ventral smooth bands and the characteristic shallow umbilicus of Oppel's *Amm. desmonotus*. The latter, however, probably attained a large size (see discussion above).

4. *Aulacostephanus* (*Aulacostephanites*) sp. aff. *mörschi* (Oppel)

Plate 32, figs. 22–24

cf. *Ammonites mörschi* Oppel 1863, p. 240, pl. 65, fig. 2a, b.


Ammonites cf. *striolaris* Quenstedt pars 1888, p. 969, pl. 107, fig. 17.

Material. Eleven (MS, ten; R 1), and fragments (MS).


Description and comparisons. These specimens are characterized by the absence of primary ribbing on the inner whorls, which carry only very dense, fine secondary ribbing. The whorl-section is compressed, umbilicus shallow, and there is a ventral smooth band. Agreement with Oppel's figure is fairly close, except that *A. mörschi* shows no trace of a ventral smooth band, is bigger, and has the primary ribs, when they appear, lower on the whorl-side. The best agreement is with Quenstedt's specimen.

EXPLANATION OF PLATE 27

Fig. 1a–c, *Aulacostephanus* (*Xenostephanus*) *ranbyensis* sp. nov. var. *elshamensis* var. nov. Adult, with half a whorl body-chamber. BM C.47908, T. B. Parks coll., from Elsham, Lincs. All natural size.



1b



1a



1c

Subgenus AULACOSTEPHANOIDES Schindewolf, emend. Ziegler

1. *Aulacostephanus (Aulacostephanoides) mutabilis* (J. de C. Sowerby)

Plate 31, fig. 3

Ammonites mutabilis J. de C. Sowerby 1823, 4, p. 145, pl. 405, fig. 1.*Rasenia mutabilis* Salfeld 1913, p. 129.*Pararasenia mutabilis* Arkell 1933, p. 612, pl. xxxix, figs. 5, 5a (holotype refigured).*Rasenia mutabilis* Arkell 1951, p. 188.*Aulacostephanus (Aulacostephanoides) mutabilis* Ziegler 1962, p. 62, figures and plates.*Material.* One (MS 80).

Description. The specimen, half an ammonite, is somewhat crushed; maximum diameter 175 mm., wholly septate, the last few septa approximated and degenerate. The complete shell must have attained some 210 mm. Inner whorls are evolute; the whorl-section is at all stages compressed with barely perceptible umbilical walls. Well-differentiated, fine, dense, secondary ribbing is visible to at least 60 mm. with occasional feeble constrictions and flared ribs; primary ribs persist to 90 mm. Thereafter the shell is wholly smooth. Venter on inner whorls not visible.

Remarks. There has been confusion in the past in interpreting this species, largely as a result of Sowerby's poor illustration. The holotype (BM 43934) has a maximum diameter of 71 mm. and is wholly septate, with none of the degeneration and approximation of the last sutures rather typical of adults of this species. Despite the sudden end of primary ribbing at a shallow constriction at 52 mm., it seems certain that the holotype is the inner part of what was a much larger shell. The secondary ribbing on the venter is interrupted by a distinct smooth band. The whorl-section is compressed and angular. All these points are exemplified in another specimen figured here (Pl. 31, fig. 3, from Cambridgeshire) which resembles the holotype very closely, and is also wholly septate to 105 mm. It also shows something of the inner whorls, invisible in the holotype, which are evolute, exposed, and densely ribbed.

As for the generic position of the species, the style of ribbing on middle whorls is that of *Rasenia*; but the compressed whorl-section, evolute inner whorls, shallow umbilical walls, and ventral smooth band place it firmly in *Aulacostephanus*. [This view is contrary to that expressed previously by Dr. Arkell (1951, p. 188); but on re-examining both the holotype and supporting material, and in the light of Dr. Ziegler's study of the whole genus *Aulacostephanus* it seems the only possible one.

J. H. C.]

2. *Aulacostephanus (Aulacostephanoides) sp. nov.?*, aff. *mutabilis* (Sowerby)

Plate 31, figs. 1, 2

Material. Two, and fragments (MS 37, 102; R 5).

Description. This species resembles *A. mutabilis* in all respects, but seems to be much smaller. Thus the specimen figured in Plate 31, fig. 1 is apparently adult, and complete with peristome at the same size (110 mm.) at which *A. mutabilis* itself (Pl. 31, fig. 3) is still septate. The same specimen is rather more evolute than *A. mutabilis*, and all ribbing has disappeared at 60 mm. The specimen shown in Plate 31, fig. 2 has inner whorls which are evolute and densely ribbed as in *A. mutabilis*; likewise the imprint of the flat

venter of the penultimate whorls in the outer (body-chamber) whorls shows the secondary ribbing to be dense, with occasional feeble constrictions and slightly flared ribs, and interrupted by a smooth band.

Similarly, small forms (for macroconchs) are known from the Kimeridge Clay of Cromarty (Scotland). They are to be described by Dr. Ziegler, who proposes the name *R. askepta* for them. They differ from those described here, however, in having oval whorl-section, involute inner whorls, and secondary ribbing uninterrupted on the venter, i.e. belong still to *Rasenia*. Their precise age is not known.

3. *Aulacostephanus* (*Aulacostephanoides*) *circumplicatus* (Quenstedt)

Plate 31, figs. 4, 5; Plate 33, figs. 9–11

Ammonites circumplicatus Quenstedt 1888, p. 978, pl. 107, fig. 19.

Ammonites cf. *trifurcatus* Quenstedt 1888, p. 971, pl. 107, fig. 21.

Tobolia pseudotrifurcata Sazonov 1960, p. 156, pl. v, fig. 4 (= nom. nov. for, and reproduction of original figure of *Amm.* cf. *trifurcatus* Quenstedt, cited above).

Material. One, and two fragments (MS 51, 120; R 19).

Description. The specimen shown in Plate 31, fig. 4 is septate to only 55 mm., and if adult, could not have exceeded 80 mm. complete. The whorl-section is slightly compressed, and in both figured specimens the secondary ribbing is weakened over the venter, but not actually interrupted.

Remarks and comparisons. The present material agrees with *A. circumplicatus* in size and style of ribbing; but the holotype has an exceptionally prominent and broad smooth ventral band. This is, however, in part due to a deformity, and, according to Dr. Ziegler, the ventral smooth band is unusually variable in the species, some specimens being practically without.

There is a considerable resemblance to *Rasenia gemmata* Schneid (1939, p. 147, pl. v (ix), figs. 9, 9a), which, however, differs in being inflated, with depressed whorl-section, and is more involute.

The species is common in the Kimeridgian exposure on Mull (Pl. 33, figs. 9–11), and some of the specimens have a clearly defined ventral smooth band.

Subgenus XENOSTEPHANOIDES nov.

1. *Aulacostephanus* (*Xenostephanoides*) *thurrelli* sp. nov.

Plate 30, figs. 1–3

Material. Eight, and fragments (R, five; MS, two; SW, eleven).


Description. The holotype (R 11; Pl. 30, fig. 1) is 44 mm. in diameter; septate to 40 mm.,

EXPLANATION OF PLATE 28

Fig. 1a–e, *Aulacostephanus* (*Xenostephanus*) *ranbyensis* sp. nov. Holotype; adult, with five-eighth whorl body-chamber preserved (1a). Suture-line at *s* shown in fig. 1e, $\times 2$. Thurrell coll. R 6, from Ranby, Lincs.

Fig. 2, *Aulacostephanus* (*Xenostephanus*) cf. *ranbyensis* sp. nov. Body-chamber fragment, showing the peristome. Thurrell coll. SW 23, from South Willingham, Lincs.

All natural size, except fig. 1e.



CALLOMON and ARKELL, Kimeridgian ammonites

with a quarter-whorl body-chamber preserved. The last three septa are approximated, with some degeneration. The coiling is extremely evolute, and the dimensions (between and through ribs or tubercles) at 45 mm. are: 24.5–26.5, 25.5–31, 49. Primary ribs are in the form of tubercles on the umbilical margin (20 per whorl at 45 mm.; 21 at 30 mm.); secondaries are trifid or bifid with frequent intercalatories, separated from the primaries by a nearly smooth lateral spiral band. On the last half-whorl, the secondaries revert to widely spaced, very coarse bifid without intercalatories indicating, together with septal approximation, that the specimen was adult. The secondary ribbing is extremely strong and coarse, resembling that of *A. undorae* (Pavlow) of the Eudoxus Zone, but is quite different on the venter. In *A. thurrelli* it is interrupted by a deep narrow smooth band, accentuated by a groove, with no tendency to form ventro-lateral tubercles.

The paratype R 25 (Pl. 30, fig. 2) resembles the holotype closely. It is septate to only 34 mm., but shows the complete body-chamber of five-eighth whorl with peristome at 50 mm. The ribbing reverts to strictly biplicate on the last quarter whorl; an oblique constriction follows an isolated triplicate rib. The lip of the aperture is flared into a collar on the venter, and has a slight lateral projection too short to be called a lappet proper.

Presumed inner whorls are shown in Plate 30, fig. 3, although no septa are visible.

The species or a close ally seems to be present in the Kimeridgian of Mull (cf. Pl. 33, fig. 8) to judge by inner whorls; but the imprint of the outer whorls is too badly distorted to be certain. There is also a specimen from the Cromarty coast (Royal Scottish Museum, Edinburgh, Hugh Miller coll.).

2. *Aulacostephanus* (*Xenostephanoides*) *lindensis* sp. nov.

Plate 30, figs. 4–13

Derivation. Lindum is the Roman name for Lincoln.

Material. Twenty-seven (SW, twenty-one; R, three; MS, four), and fragments.

Description. The material is plentiful but poorly preserved, and no single specimen is complete. It is with some hesitation, therefore, that we select the specimen MS 13 (Pl. 30, fig. 10) only formally as holotype and designate the other nine figured specimens (Pl. 30, figs. 4–9, 11–13) paratypes, although for purposes of description it would be best to regard all ten as syntypes.

Sutures are rarely visible, but from the difference in preservation between outer and inner whorls (which are usually flattened) it seems probable that most of the specimens carry some body-chamber. The holotype itself definitely does, and was presumably complete at about 30 mm. diameter. Another fragment (SW 59) shows the final peristome with oblique constriction, flared collar, and short incipient lappet (just as in *A. thurrelli*) at a final diameter of c. 40 mm. The species is thus small.

The coiling is moderately involute, the whorl-section stouter than in *A. thurrelli*. Ribbing is dense and moderately strong, with three, sometimes four, secondaries per primary, separated, as in *A. thurrelli*, by a lateral smooth band. The venter is flat, broad, with a narrow smooth band but no groove. Inner whorls are shown in Plate 30, fig. 13.

Comparison. The species most closely resembles *Aulacostephanus thurrelli* although the stouter, more involute coiling and finer ribbing appear to be consistently distinct. It stands, like *A. thurrelli*, apart from all other contemporaneous aulacostephanids; but,

in its resemblance to some later forms of the *A. eudoxus* group, it heralds the future development of the genus.

A. lindensis has also been found in the drift of Norfolk (SM J48657, one specimen, Baden-Powell coll.).

For completeness it is convenient to mention here another form which belongs to this group.

3. *Aulacostephanus* (*Xenostephanoides*) *scoticus* nom. nov. = *Aulacostephanus* cf. *autissiodorensis* (Cotteau) Spath 1935, p. 49, pl. 13, figs. 8a, b (BM C13216)

Remarks. The specimen figured by Spath from Culgower, eastern Scotland, seems unique. It carries some body-chamber on which the otherwise trifid ribbing reverts to biplication, indicating maturity, as in *A. thurrelli*; *A. scoticus* is therefore a small species and not a young specimen of giant forms like *Rasenia borealis* Spath, as Spath supposed. Its evoluteness and style of ribbing, particularly on the inner whorls, are typical of *Xenostephanoides*. It differs from *A. (X.) thurrelli* principally in having rather more inflated inner whorls, about nine more ribs per whorl, and a less-pronounced ventral smooth band, the ventral ribbing on the inner whorls being weakened without actual interruption. The exact horizon is unknown, although there are other Mutabilis Zone faunas from the same area.

Subgenus XENOSTEPHANUS nov.

1. *Aulacostephanus* (*Xenostephanus*) *ranbyensis* sp. nov.

Plate 28, figs. 1a-e, 2; Plate 29, fig. 4

Material. Two (R 6; BM C.47909), and fragments (SW).

Description. The holotype (R 6) is a splendidly preserved internal cast of 120 mm. maximum diameter, septate to 81 mm., with the last three sutures approximated and simplified. The body-chamber is broken, but, to judge from the highly degenerate last few visible ribs, represented very nearly up to the peristome and hence occupying five-eighths of a whorl.

Dimensions: at 115 mm.: 23, c. 26, 48.

80 mm.: 25-27, 30-33, 54.

55 mm.: 25-27, 29-31, 55.

Ribs: at 40-50 mm.: 19 per whorl; 70-90 mm.: 22; 115 mm.: c. 25.

EXPLANATION OF PLATE 29


Fig. 1a, b, *Aulacostephanus* (*Xenostephanus*) *anceps* sp. nov. Holotype; adult, with one-quarter whorl body-chamber preserved, and umbilical suture indicating former extent to five-eighths of a whorl. BM 50761, old collection, no history.

Fig. 2a, b, *Aulacostephanus* (*Xenostephanus*) *staintonensis* sp. nov. Syntype I; the cross marks what appears to be the last septum. Thurrell coll. MS 142, from Market Stainton, Lincs.

Fig. 3, *Aulacostephanus* (*Xenostephanus*) *staintonensis* sp. nov. Syntype II; wholly septate, last few septa approximated. SW 2/35, from South Willingham, Lincs.

Fig. 4, *Aulacostephanus* (*Xenostephanus*) cf. *ranbyensis* sp. nov. Fragment at commencement of the body-chamber. SW 34.

All natural size.



CALLOMON and ARKELL, Kimeridgian ammonites

In the umbilicus nothing is visible before a diameter of 20 mm. Thereafter the primary ribbing consists of conspicuous dorsolateral tubercles, only slightly bullate, and separated from the strong rectiradiate ribs that spring from them by a lateral smooth band. The ribs are bifid with regular intercalatories, which, on the last septate whorl, tend to join up to form trifid bundles, while fresh intercalatories come in. On the body-chamber the ribbing weakens, the primaries degenerate to bullae and triplicate, finally biplicate, ribs. The last few ribs are single, coarse and irregular, indicating the end of growth. The peristome is not preserved. The venter, where it emerges, has a groove, but no tubercles. The groove fades to a mere smooth band before septation ceases, this coming about as the ribbing loses relief, and persists to the end even when the ventral ribbing is barely perceptible. The whorl-section on inner whorls is quadrate, angular, and becomes rounded towards the end. The suture-line is a typically perisphinctid one, with slender widely spaced lobes on a straight base-line.

The peristome is preserved on another fragment (Pl. 28, fig. 2). It is straight and simple, with slight forward sweep at the umbilical seam.

There is another specimen in the British Museum (C47909, T. B. Parks coll., from Elsham, Lincs., only 72 mm. in diameter, wholly septate, somewhat crushed, which agrees with the holotype in all details, including matrix. The species has also been found in the drift of Norfolk (SM J48657, Baden-Powell coll., one specimen).

Remarks. Inner whorls are practically indistinguishable from *A. (X.) thurrelli*, and, as remarked previously, it would be hard to find a clearer demonstration of ammonoid dimorphism. Also remarkable is the close homeomorphism between *Xenostephanus* and *Reineckeia* s.s. of the Callovian, yet another instance showing the tendency of ammonites to evolve repeatedly through similar forms.

1a. *A. (X.) ranbyensis* var. *elshamensis* var. nov.

Plate 27, fig. 1a-c

Material. One (BM C.47908, T. B. Parks coll., from Elsham, N. Lincs.).

Description. Size c. 130 mm. (slight distortion), septate to 103 mm. with approximation and degeneration, half a whorl of body-chamber preserved with indications of former extent to three-quarters of a whorl, so that the maximum size of the complete shell must have been 150 mm.

Dimensions: at 100 mm.: 25, 29-34, 57.

Ribs: at 130 mm.: 25 per whorl; 120: 23; 100: 22; 80: 20; 60: 19; 40: 19; 30: 20.

The primary ribs are bullate, triplicate with low furcation point up to 80 mm.; they then revert to biplicate, perisphinctid ridges to 120 mm., and single simple ribs thereafter. The secondaries are strong, in regular triplicate sheaves, to about 80 mm., then rapidly weaken and become biplicate at high furcation points, and finally fade altogether. They show some weakening in the middle of the venter where first visible, although there is no fully smooth band, and after about 90 mm. pass over the venter with no interruption.

Comparison. Agreement with the holotype of *A. ranbyensis* is close in proportions and rib-count. The variety differs in becoming bigger, with a more inflated body-chamber,

and having less differentiation between primary and secondary ribbing so that a lateral smooth band is present only on the innermost whorls.

2. *Aulacostephanus (Xenostephanus) staintonensis* sp. nov.

Plate 29, figs. 2, 3

Material. Two (MS 142, SW 2), and fragments (SW).

Description. Neither specimen is really sufficiently complete to serve as sole basis of the species, and the two figured specimens are syntypes.

Syntype I (Pl. 29, fig. 2): maximum diameter 57 mm., apparently septate to only 35 mm. with five-eighths whorl body-chamber. Neither ribbing nor coiling, where last visible, show any signs of modification whatever, and the specimen seems to be genuinely immature.

Dimensions: at 55 mm.: 32, c. 33, 45.

Ribs: at 60 mm.: 22 per whorl; 50: 21; 40–20: 20.

Inner whorls evolute, becoming more inflated. Primary ribs moderately bullate, trifid on inner whorls, changing to regular triplicate. Strong secondaries with ventral smooth band, fading at the end.

Syntype II (Pl. 29, fig. 3): maximum diameter 115 mm., wholly septate (details of sutures invisible), flattened through crushing.

Ribs: at 115 mm.: c. 30 per whorl; 70: c. 27 (counts based on half-whorls).

Ribs on inner whorls triplicate or biplicate with intercalatories; sharp bullate primaries, no lateral smooth band; ventral smooth band. Ribbing has started to modify a 90 mm. to perisphinctid style; and at 115 mm. only straight primaries of low relief remain. The ventral smooth band persists until the entire venter is smooth.

Comparisons. The species differs from *A. ranbyensis* in being markedly more densely ribbed and involute. The primary and secondary ribbing are less well differentiated, so that there is no lateral smooth band even on inner whorls, which are less openly exposed. In this sense the species stands in the same relation to *A. ranbyensis* as *A. lindensis*


EXPLANATION OF PLATE 30

Figs. 1–3, *Aulacostephanus (Xenostephanoides) thurrelli* sp. nov. 1a, b, Holotype; adult, with one-quarter whorl body-chamber preserved. Thurrell coll. R 11, from Ranby, Lincs. 2a, b, Fragmentary but complete adult, with peristome showing slight lappet. R 19/25. 3a, b, Inner whorls, no sutures visible. SW 39.

Figs. 4–13, *Aulacostephanus (Xenostephanoides) lindensis* sp. nov. 4, SW 31, no sutures visible but probably one-half whorl body-chamber. 5, SW 29, last third whorl body-chamber. 6a, b, SW 33, no sutures visible, but probably at least one-half whorl body-chamber. 7, SW 26, ditto. 8a, b, SW 60, ditto. 9a, b, SW 9, ditto. 10a, b, Holotype, MS 13, seven-tenths whorl body-chamber. 11, SW 53, remarks as 6. 12a, b, MS 38, ditto. 13a, b, MS 18, wholly septate. All Thurrell coll.

Figs. 14–18, *Aulacostephanus (Aulacostephanites) ebrayoides* sp. nov. 14, Holotype; last half whorl body-chamber; MS 5. 15, MS 56, half whorl body-chamber. 16a–c, MS 57, ditto. 17a–c, MS 45, eight-tenths whorl body-chamber. 18a–c, MS 71, six-tenths whorl body-chamber; perhaps transitional to *A. (A.) eulepidus* (Schneid). All Thurrell coll.

All natural size.



CALLOMON and ARKELL, Kimeridgian ammonites

stands to *A. thurrelli*. An example of this species has also been found in the drift of Norfolk (SM J48658, a double mould, Baden-Powell coll.).

3. *Aulacostephanus* (*Xenostephanus*) *anceps* sp. nov.

Plate 29, fig. 1

Material. One (the holotype, BM no. 50761).

Description. Maximum size 95 mm., septate to 85 mm. with severe approximation and degeneration of the last (and only visible) sutures, and a quarter whorl body-chamber. The umbilical suture indicates a former extent of the final body-chamber to five-eighths whorl, so that the complete shell attained a maximum diameter of c. 110 mm.

Dimensions: at 90 mm.: 27, 29–33, 51 (cast).

at 75 mm.: 27, 36–43, 52 (test).

Ribs: at 100–80 mm.: 21 per whorl; 60–50: 20; 40: 19; 30–20: 18.

This is the most inflated of the species described here, with a whorl-section at 75 mm. depressed almost to the point of being coronate. In contrast, the body-chamber contracts and reverts to normal perisphinctid dimensions. Similarly, the ribbing, which is characterized by very high, sharp bullate primaries on the umbilico-lateral margin giving rise to regular sheaves of very coarse triplicate secondaries, modifies to rather widely spaced and in part biplicate ribs of low relief on the body-chamber. The secondaries show only the faintest weakening on the round venter where first visible (at 60 mm.); they smooth out altogether on the body-chamber.

Affinities and remarks. *A. anceps* bears undeniably close resemblance to various species of *Rasenia* from the Cymodoce Zone of Market Rasen, including *R. uralensis* (d'Orb.), and, in the absence of a ventral smooth band, to place it in any but this genus might seem unnatural. Its inner whorls, however, are quite different: evolute, with flat whorl sides and the differentiation, with lateral smooth band, of primary and secondary ribbing characteristic of *Xenostephanus*. The aulacostephanid characters of angular whorl-section and ventral smooth band seem here to make their first appearance in *Xenostephanus* proterogenetically.

The specimen has the test preserved on all but the last half whorl, and it is therefore conceivable that an internal cast might show a ventral smooth band not visible on the outside; and that the difference in this respect between *A. (X.) anceps* and the other forms of *Xenostephanus* described here, mostly casts, might only be apparent and attributable to differences of preservation. This seems unlikely, however, for a specimen of *A. (X.) ranbyensis* from Norfolk (SM J48657) shows the same ventral smooth band on both internal cast and external mould of the same part of the body-chamber at 60 mm. diameter.

The holotype was presented to the British Museum by Prof. J. Morris in 1867; there is no other recorded history. Its preservation and matrix are, however, so precisely similar to the other material here described that there can be little doubt that it came from a similar source. It seems to have been one of the specimens which Salfeld included in his MS species *evoluta*; but its inner whorls are quite different from those of the specimen Spath selected as type for his species (see above). Besides oysters, the matrix carries an imprint of *Rasenia* cf. *thermarum* (Oppel).

Xenostephanus seems to occur in the Kimeridgian of Skye. A single specimen of *A.* (*X.*) cf. or aff. *anceps* (Pl. 33, fig. 1) was collected by Mr. W. G. Cordey at Kildorais, Trotternish, in the spring of 1961, and we thank him for presenting it to us. It is crushed, in clay, and encrusted with an oyster on the inner whorls. Neither septa nor venter are discernible, but the ornament is typically that of *Xenostephanus*, and enough is visible of the inner whorls to show that they are evolute just as in *A. anceps*. The Skye specimen differs in having rather fewer ribs on the middle and outer whorls: at 85 mm.: 17 per whorl; 70: 16; 50: 15; 30: c. 18. The specimen was unfortunately found without any supporting fauna.

For completeness, we include here:

4. *Aulacostephanus* (? *Xenostephanus*) sp. indet. = *Ammonites* cf. *trifurcatus* Quenstedt 1888, p. 971, pl. 107, fig. 24

Remarks: Quenstedt's figure shows an extremely evolute shell, maximum size c. 85 mm., with ornament typical of *Xenostephanus*: well-spaced bullate primary ribs leading to triplicate, strong secondaries with intercalatories. Unfortunately neither venter nor sutures are anywhere visible. At 85 mm. there are 18 ribs per whorl; at 60: 17; at 40: 15. The species differs from *A. ranbyensis* therefore in being less densely ribbed; also, in having the primaries half-way up the whorl-side rather than on the umbilico-lateral margin.

Nautilus trifurcatus Reinecke (1818, p. 75, pl. v, fig. 49) is quite different: inflated, involute, with barely bullate primary ribbing. Spath (1935, p. 49, footnote) thought Quenstedt's specimen belonged to *Rasenia evoluta* Salfeld MS, and in the sense that the specimen described above as *A.* (*X.*) *anceps* appears to have been a syntype of Salfeld's unpublished manuscript species he was probably correct. It is unfortunate that he chose to publish the name on the basis of only a small nucleus (see above).

If the above identification is correct, Quenstedt's specimen seems to be the only representative of these early aulacostephanitids so far known from more southerly regions. It was found in a volcanic erratic at Floriansberg, and ascribed to White Jura δ .

There are a number of other boreal forms intermediate between *Rasenia* and *Aulacostephanus* proper which, when more is known about them, may turn out to be closely related to *Xenostephanus*.

Aulacostephanus groenlandicus Ravn (1912, p. 492, pl. xxxvii, fig. 3a-c, from Koldewey Island, 76° N.; Frebold 1930, p. ix, fig. 4, from Spitsbergen) has the evolute coiling and indications of a ventral smooth band, but is densely ribbed and lacks the tuberculate


EXPLANATION OF PLATE 31

Figs. 1, 2, *Aulacostephanus* (*Aulacostephanoides*) sp. nov. ? aff. *mutabilis* (J. de C. Sowerby). 1, Complete adult, with seven-tenths whorl body-chamber, and peristome transposed unto the figured side (white dotted line) as it is developed on the reverse side. Thurrell coll. MS 37. 2, MS 102; the outer whorl is body-chamber.

Fig. 3, *Aulacostephanus* (*Aulacostephanoides*) *mutabilis* (J. de C. Sowerby). Wholly septate example, from a nodule-bed in Kimeridge Clay near Haddenham Station, Cambridgeshire. SM J.29157.

Figs. 4, 5, *Aulacostephanus* (*Aulacostephanoides*) *circumplicatus* (Quenstedt). 4, MS 120, with some body-chamber. 5a, b, MS 51b, outer whorl body-chamber.

Fig. 6a, b, *Rasenia* (*Rasenioides*) cf. *striolaris* (Reinecke). MS 29, with half whorl body-chamber. All natural size.



CALLOMON and ARKELL, Kimeridgian ammonites



Arkell, W J and Callomon, John H. 1963. "Lower Kimeridgian ammonites from the drift of Lincolnshire." *Palaeontology* 6, 219–245.

View This Item Online: <https://www.biodiversitylibrary.org/item/196367>

Permalink: <https://www.biodiversitylibrary.org/partpdf/174441>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.