

Post-embryonic Development of *Oppia nitens* (Acarina: Oribatei)

H. G. SENGBUSCH AND C. H. SENGBUSCH

STATE UNIVERSITY COLLEGE AT BUFFALO
BUFFALO, NEW YORK 14222

RECEIVED FOR PUBLICATION SEPTEMBER 2, 1970

Abstract: The present research on the life history of *Oppia nitens* C. L. Koch adds another species of oribatoid mite which has been successfully reared under standard laboratory conditions. For the first time culturing was carried through the F₂ generation. For experimentation employing oribatoid mites as test organisms, particularly in regard to the effects of pesticides and herbicides, a standard reference table of their life histories is necessary. A table of published life histories, given herewith, provides the most comprehensive summary of information available.

The laboratory methods followed those of Sengbusch (1954, 1963). A temperature of 20° C. and a relative humidity of 82% were maintained. Adults were isolated from moist mor humus collected from a coniferous forest floor. The average length of time in days for each of the immature stages of the F₁ generation was: egg—8.5; larva—7.4; protonymph—7.6; deutonymph—9.3; tritonymph—12.3. The duration of the combined developmental stages varied from 39 to 55 days. The quiescent period which precedes each molt lasted approximately 2 to 3 days except the one before the exuviation of the adult which continued 4 to 5 days. The average length of time for the post-embryonic development of the F₂ generation was not different from that of the F₁.

INTRODUCTION

The present research on the life history of *Oppia nitens* C. L. Koch represents another species added to the list of Oribatei which have been successfully cultured under standard conditions of temperature and humidity. For the first time a species of oribatoid mite was reared through the F₂ generation.

Oribatoid mites, although once thought to be harmless plant feeders, are now known to be of economic significance. Sellnick (1928) believed that they play an important role in the economy of nature by contributing to the fertility of the forest soil. Woodring (1963) in a discussion of the value of oribatoids to litter reduction stated, "it seems hard to believe that in view of their habits, nutrition, and vast numbers they would not be important." Jacot (1930), on the other hand, pointed out injurious effects induced by their phytophagous habits. However, it was the work of Stunkard (1937) which gave impetus to research on these relatively little-known animals. It was he who established that the oribatoids may act as the intermediate hosts of certain anoplocephaline cestodes.

MATERIALS AND METHODS

The laboratory methods followed those of Sengbusch (1954, 1958, 1963). While they are somewhat tedious, they have produced approximately one-third of the life histories of oribatoid mites which have been reported (Table 3).

Adults were collected by means of a modified Tullgren funnel (Sengbusch, 1951) from the moist mor humus layer of a coniferous forest floor. They were isolated in separate glass microcells modified from the description given by Jacot (1937) and were observed daily under a binocular dissecting microscope using a magnification of $30\times$. The microcells consisted of ordinary 3×1 inch microslides on which were glued one or two small glass rings. Filter paper was cut to fit snugly in the bottom and moistened to provide the water which is so necessary for this group of mites. A cover was provided by another microslide held in place by a small rubber band. No attempt was made to make this in air-tight seal as these culture cells were then placed in Scheibler desiccators containing a saturated solution of ZnSO_4 to provide a relative humidity of 82% (Ludwig and Landsman, 1937). The desiccators were kept in a constant temperature room set at 20°C .

OBSERVATIONS

Like most Acari, the oribatoid mites leave the egg as six-legged larvae and subsequently pass through four molts. After the first molt, they are in the first eight-legged nymphal stage (protonymph), and then pass through two further nymphal stages, the deutonymph and tritonymph.

Each molt is initiated by a quiescent period of various lengths. At its onset the mite fastens its claws (unguiculae) into the filter paper and arches the dorsum. It stays in this position without feeding for two to three days. However, it would be incorrect to refer to this time as a resting period, for internally the mite is undergoing the morphological and physiological changes which will be evidenced in the following stadium. The period is terminated by the splitting of the old exoskeleton (exuvium) and the emergence of the next stage.

All stages nourish themselves in a similar fashion and also carry on the same modes of life. *Protococcus*, a green alga found on bark, was used throughout the investigation as food for the immature and adult *Oppia nitens*.

Ontogeny occurs by a gradual development from the egg to the sexually mature adult. The female mite, attempting to hide the eggs, may insert the long ovipositor into clumps of *Protococcus* or into the crevice between the edge of the filter paper and the glass and extrude the eggs. The eggs are typically oval to ovate and have one side flattened as do many insect eggs. Eggs recently laid have a whitish, almost translucent appearance and are relatively easy to locate in the small culture cells. Within three to four days development becomes evident externally as an amber pigmentation becomes visible at both poles. The thin shell is almost transparent and slightly plastic so that major variations in form and color are noticeable with low magnification. A day or two before hatching, the outline of the larva can easily be perceived using a magnification of $60\times$. Just before emergence, the exoskeleton of the larva becomes totally pigmented with the amber color noted above.

TABLE 1. Length of time in days of developmental stages of *Oppia nitens* (F₁) at 20° C.

Emergence of larva	Emergence of proto.	Emergence of deuto.	Emergence of trito.	Emergence of adult	Total
9	7	7	8	11	42
10	7	7	9	14	47
10	7	8	5	9	39
7	9	7	13	11	47
8	8	7	18	14	55
8	8	7	11	18	52
11	7	8	7	6	39
6	7	8	9	10	40
8	10	6	6	10	40
6	7	11	7	11	42
10	5	7	9	21	52
Av. 8.5	Av. 7.4	Av. 7.6	Av. 9.3	Av.12.3	Av.45.0

From Table 1 it is evident that there is some variation in the duration of the stadia in this species. From egg to adult 39 to 55 days were required in the eleven specimens that reached maturity. A predominance of the seven and eight days required for the emergence of the protonymph and deutonymph is noticeable. It seems probable that with a large enough sampling and with the environmental conditions approaching optima, the average times would be relatively constant. The greater length of time necessary before the emergence of the adult is due to the greater morphological changes necessary before the final molt.

The F₁ females began oviposition approximately three months after emergence. Because the experiment had to be terminated, only six specimens attained adulthood. From Table 2 agreement can be seen between the total development time in the F₂ generation (46.0 days) and that of the F₁ (45.0 days). No statistical difference exists in the duration of post-embryonic development in the two generations, which supports the prediction that the method of Sengbusch (1954) can be used to culture any oribatoid mite.

We saw no spermatophores in any of the culture cells containing adults. Spermatophores have been described from several species of oribatoids (Seng-

TABLE 2. Length of time in days of developmental stages of *Oppia nitens* (F₂) at 20° C.

Emergence of larva	Emergence of proto.	Emergence of deuto.	Emergence of trito.	Emergence of adult	Total
8	6	7	10	12	43
9	5	10	13	7	44
9	6	6	11	11	43
7	6	8	4	21	46
7	7	7	8	21	50
9	5	7	11	18	50
Av. 8.2	Av. 5.8	Av. 7.5	Av. 9.5	Av.15.0	Av.46.0

TABLE 3. Length of time in days of developmental stages of oribatoid mites.

	Emergence of larva	Emergence of protonymph	Emergence of deutonymph	Emergence of trityonymph	Emergence of adult	Total
<i>Hypochthonius rufulus</i> Koch	30.0	21.8	19.1	21.5	24.5	122.0
<i>Steganacarus diaphanus</i> Jacot						150
<i>Nanherrmannia nana</i> (Nicolet)	13.2	20.4	24.1	22.5	31.4	111.3
<i>Herrmannia scabra</i> (Koch)	15	16	41	42	70	184
<i>Nothrus silvestris</i> Nicolet	15.8	32.9	37.3	28.3	40.0	152.5
<i>Camisia spinifer</i> (Koch)	6.7	18.2	26.9	29.1	50.8	125.6
<i>Camisia segnis</i> (Hermann)			36			200
<i>Platynothrus peltifer</i> (Koch)		35	27	31	24	150
<i>Platynothrus peltifer</i> (Koch)						117
<i>Platynothrus peltifer</i> (Koch)						330-360
<i>Trhypochthonius tectorum</i> (Berlese)	11-35	15-20	13-27	15-32	18-50	72-164
<i>Dameus clavipes</i> (Hermann)						75
<i>Dameus clavipes</i> (Hermann)	5.1	12.3	12.1	14.5	20.0	64.0
<i>Dameus clavipes</i> (Hermann)						330-360
<i>Dameus onustus</i> Koch	8.7	28.3	26.7	29.3	40.0	150
<i>Dameus onustus</i> Koch	17.4	58.0	40.0			130.3
<i>Dameus onustus</i> Koch						75
<i>Dameus auritus</i> Koch	20	19	22	25	30	120
<i>Dameus boreus</i> Bulanova-Zakhvatkina	14±2	15±3	9±2	17±1	9-18	63-76
<i>Belba kingi</i> Hartenstein						65
<i>Metabelba montana</i> (Kulczynski)	4	53	40	78	200	375
<i>Cepheus palmicinctum</i> (Nicolet)	15	10-15	15-18	14-16	11-15	68-75
<i>Eremobelba nervosa</i> Hartenstein						79
<i>Ceratoppia bipilis</i> (Hermann)	3-5	14-24	9-13	9-13	14-31	49-87
<i>Ceratoppia bipilis</i> (Hermann)	13	24	25	28	46	102
<i>Tectocephus velatus</i> (Michael)						

TABLE 3. (Continued)

	Emergence of larva	Emergence of protonymph	Emergence of deutonymph	Emergence of tritonymph	Emergence of adult	Total
<i>Oppia nitens</i> Koch	Michael (1888)					40(-)
<i>Oppia nitens</i> Koch	Sengbusch & Sengbusch (1970)	8.5	7.4	7.6	9.3	12.3
	Woodring & Cook (1962)					45.0
<i>Oppia nova</i> (Oudemans)	Woodring & Cook (1962)					23
<i>Minuthozetes semirufus</i> (Koch)	Sengbusch (1958)	6.1	6.8	7.0	7.1	11.2
<i>Ceratozetes gracilis</i> (Michael)	Haarløv (1960)					38.0
<i>Ceratozetes cisalpinus</i> Berlese	Hartenstein (1962c)	32	20	36	28	19
<i>Ceratozetes jewelli</i> Rockett & Woodring	Woodring & Cook (1962)					119-149
<i>Galumna elimatus</i> var. (Koch)	Rockett & Woodring (1966)	12	11	9	9	32
<i>Galumna nervosus</i> (Berlese)	Sengbusch (1954)	14.9	16.1	15.6	15.9	12
<i>Galumna nervosus</i> (Berlese)	Sengbusch (1954)	10.7	8.3	7.9	8.5	24.8
<i>Pergalumna omniophagous</i> Rockett & Woodring	Sengbusch (1958)	8.8	7.5	7.6	7.3	11.7
						43.1
<i>Galumna confusa</i> Woodring	Rockett & Woodring (1966)	11	7	7	7	10
<i>Galumna longipluma</i> (Berlese)	Woodring (1965)	7-14	5-8	7-13	11-15	12-17
<i>Acrogalumna longiplumus</i> (Berlese)	Sengbusch (1954)	10.8	10.8	11.5	12.6	15.3
<i>Galumna parva</i> Woodring	Sengbusch (1958)	11.8	13.4	12.9	13.6	17.2
<i>Rostrozetes flavus</i> Woodring	Woodring (1965)	7-8	4-10	7-9	9-11	9-13
<i>Scheloribates laevigatus</i> (Koch)	Woodring (1965)	8-9	6-9	8-10	9-13	10-14
	Cleat (1952)					33-41
						35-45
						42-115
						av. 61
<i>Scheloribates laevigatus</i> (Koch)	Woodring & Cook (1962)					64
<i>Scheloribates parabilis</i> Woodring	Woodring (1965)	8-11	3-6	3-7	3-8	3-8
<i>Scheloribates nudus</i> Woodring	Woodring (1965)	8-11	3-6	3-7	3-7	14-24
<i>Protoribates lophotrichus</i> (Berlese)	Hartenstein (1962e)	23	23	27+	32+	28
<i>Euphthiracarus</i> sp.	Rhode (1955)					150
						64

busch, 1961), but there is also evidence that parthenogenesis occurs in some species (Grandjean, 1947; Sengbusch, 1958). *Oppia nitens* probably falls into the latter category.

DISCUSSION

The present research is the first consideration of the life cycle of *Oppia nitens* C. L. Koch since Michael's original work in the late 1800s. Michael reported only the total longevity utilizing crude culturing techniques. The current study lists the duration in days of each immature stage plus the total egg to imago period. These latter figures are more realistic than Michael's because they were obtained under standard laboratory conditions.

For the first time, data are presented from the development of an F₂ generation. Previous workers have generally assumed a lack of distinction between the progeny of parents obtained from nature and the offspring of laboratory raised mites. We were able to demonstrate the similarity of development times between the two groups in this species. Our culturing conditions do not appear to change the duration of the stadia or the total elapsed time for the life-cycle of *Oppia nitens*. Although the earlier assumption was found to be correct, the specific results obtained by experimentation reduce the variables involved in the use of these mites as test organisms.

The forty-nine life histories which have been published, beginning with the initial investigations of Michael in 1884, and continuing to contemporary studies are listed in Table 3. There is wide latitude in the reported times, plus the frequent lack of data for the individual immature stages. A comparison of similar species reveals discrepancies which probably reflect differences in culturing techniques. According to Woodring (1963), ". . . oribatoids have been reared by several . . . but they have been cultured only by Sengbusch (1954), and Woodring and Cook (1962)." For experimentation employing oribatoid mites as test organisms a standard reference table of their life histories is necessary. Table 3 provides the most comprehensive summary of information available.

Literature Cited

- BHATTACHARYYA, S. K. 1962. Laboratory studies on the feeding habits and life cycles of soil-inhabiting mites. *Pedobiologia* **1**: 291-298.
- BLOCK, W. C. 1965. The life histories of *Platynothrus peltifer* (Koch 1839) and *Damaeus clavipes* (Hermann 1804) (Acarina: Cryptostigmata) in soils of Pennine moorland. *Acarologia* **7**(4): 735-743.
- CLEAT, N. D. 1952. Growth in the laboratory of economically important Oribatid mites. *Nature* (London) **169**: 280-281.
- GRANDJEAN, F. 1947. Observations sur les oribates (18e Série). *Bull. Mus. Nat. d'Hist. Nat. Paris*, sér. 2, **19**: 395-402.
- . 1950. Observations Ethologiques sur *Camisia segnis* (Herm). et *Platynothrus peltifer* (Koch) (Acariens). *Bull. Mus.* **22**(2): 224-231.

- HAARLØV, N. 1960. Microarthropods from Danish soils. Ecology, Phenology. Oikös, suppl. **3**: 1-176.
- HARTENSTEIN, R. 1962a. Soil Oribatei. II. *Belba kingi*, new species (Acarina: Belbidae), and a study of its life history. Ann. Entomol. Soc. Amer. **55**(4): 357-361.
- . 1962b. Soil Oribatei. III. Studies on the development, biology, and ecology of *Metabelba montana* (Kulcz.) (Acarina: Belbidae) and *Eremobelba nervosa* n. sp. (Acarina: Eremaeidae). Ann. Entomol. Soc. Amer. **55**(4): 361-367.
- . 1962c. Soil Oribatei. IV. Observations on *Ceratozetes gracilis* (Acarina: Ceratozetidae). Ann. Entomol. Soc. Amer. **55**(5): 583-586.
- . 1962d. Soil Oribatei. V. Investigations on *Platynothrus peltifer* (Acarina: Camisiidae). Ann. Entomol. Soc. Amer. **55**(6): 709-713.
- . 1962e. Soil Oribatei. VI. *Protoribates lophotrichus* (Acarina: Haplozetidae) and its association with microorganisms. Ann. Entomol. Soc. Amer. **55**(5): 587-591.
- . 1962f. Soil Oribatei. VII. Decomposition of conifer needles and deciduous leaf petioles by *Steganacarus diaphanum* (Acarina: Phthiracaridae). Ann. Entomol. Soc. Amer. **55**(6): 713-716.
- JACOT, A. P. 1930. Moss mites as spore-bearers. Mycologia **22**: 94-95.
- . 1937. A method for culturing non-parasitic mites. In Culture Methods for Invertebrate Animals by Galtsoff, Lutz, Welch, and Needham. Comstock Publ. Co., Ithaca, New York.
- JALIL, M. 1965. Studies on the life cycle of *Hermannia scabra* (C. L. Koch 1879) (Acarina: Oribatei). Oikös **16**: 17-19.
- LUDWIG, D., AND H. LANDSMAN. 1937. The effect of different relative humidities on survival and metamorphosis of the Japanese beetle (*Popillia japonica* Newman). Physiol. Zool. **10**(2): 171-179.
- MICHAEL, A. D. 1884 and 1888. British Oribatidae, **1** and **2**. Ray Society, London.
- MURPHY, P. W., AND M. JALIL. 1964. Some observations on the genus *Tectocephus*. Acarologia **6**: 187-197.
- PAULY, F. 1956. Zur Biologie einiger Belbiden (Oribatei, Moosmilben) und zur Funktion ihrer Pseudostigmatischen Organe. Zool. Jb. Abt. Syst. **84**: 275-328.
- RHODE, C. J. 1955. Studies on Arthropods from a moss habitat with special emphasis on the life history of three oribatid mites. Diss. Abs. **15**: 1936 Northwestern University.
- ROCKETT, C. L., AND J. P. WOODRING. 1966. Biological investigations of a new species of *Ceratozetes* and of *Pergalumna* (Acarina: Cryptostigmata). Acarologia **8**: 511-520.
- SELLNICK, M. 1928. Hornmilben, Oribatei. Tierwelt Mitteleuropas 3, Lief **4**(9): 1-42.
- SENGBUSCH, H. G. 1951. Notes on some New York oribatid mites. Zoologica **35**(3): 155-162.
- . 1954. Studies on the life history of three oribatoid mites with observations on other species (Acarina: Oribatei). Ann. Entomol. Soc. Amer. **47**(4): 646-667.
- . 1958. Zuchtversuche mit Oribatiden. (Acarina). Naturwiss. **45**(20): 498-499.
- . 1961. A study of spermatophores and sperm transfer in the Oribatei. J. Parasit. **47**(4 sect. 2): 43-44.
- . 1963. Methods recommended for the preparation and culture of Oribatei. Advances in Acarology **1**: 181-190. Cornell Univ. Press, Ithaca, N. Y.
- SITNIKOVA, L. G. 1959. Life cycles of some Oribatei and methods of their cultivation. Zool. Inst. Acad. Sci. USSR (Leningrad) **38**(11): 1663-1673.
- STUNKARD, H. W. 1937. The life cycle of *Moniezia expansa*. Science **86**: 312.
- TABERLY, G. 1953. Sur L'Ethologie et le developpement post-embryonnaire de *Trhypochthonius tectorum* (Acarien, Oribate). Bull. Soc. Zool. France **77**(5-6): 330-341.

- . 1957. Observations sur les spermatophores et leur transfert chez les Oribates (Acariens). *Bull. Soc. Zool. France* **82**: 139–145.
- WOODRING, J. P. 1963. The nutrition and biology of saprophytic sarcoptiformes. *Advances in Acarologia* **1**: 89–111. Cornell Univ. Press, Ithaca, N. Y.
- . 1965. The biology of five new species of Oribatids from Louisiana. *Acarologia* **7**(3): 565–576.
- WOODRING, J. P., AND E. F. COOK. 1962. The biology of *Ceratozetes cisalpinus* Berlese, *Scheloribates laevigatus* Koch, and *Oppia neerlandica* Oudemans (Oribatei), with a description of all stages. *Acarologia* **4**(1): 101–137.



Sengbusch, H G and Sengbusch, C H . 1970. "Post-Embryonic Development of *Oppia nitens* (Acarina: Oribatei)." *Journal of the New York Entomological Society* 78, 207–214.

View This Item Online: <https://www.biodiversitylibrary.org/item/206594>

Permalink: <https://www.biodiversitylibrary.org/partpdf/179935>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: New York Entomological Society

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.