

cm. in length; stipules oblanceolate, glandular-serrate, caducous. Flowers 1.5 cm. in diameter, in small mostly 5- or 6-flowered corymbs, furnished with broad conspicuously glandular-serrate deciduous bracts; calyx glabrous with narrow acuminate glandular-serrate lobes often persistent on the ripe fruit; stamens 7-10; anthers pale pink; styles 2-4. Fruit ripening in early October, subglobose, dull crimson, punctate, about 1.5 cm. in diameter; nutlets 3-4, rounded on the back and ends, slightly broader at the apex than at the base, 5-6 mm. in length.

Usually a shrub from 4.5-5 m. high, with stout stems covered with dark corky bark, and erect or ascending branches armed with numerous slender straight or slightly curved dark purple spines 4-5 cm. in length; or often a small tree 5-6 m. high.

Rocky limestone hills near Galena, Stone County, Missouri, *E. J. Palmer*, October 13, 1913 (No. 4645, type for fruit), September 27, 1920 (No. 19,183), April 29, 1924 (No. 24,569, type for flowers), all from plant No. 3; same locality, May 23, 1923 (No. 22,798), April 29, 1924 (No. 24,568). Rocky upland woods, near Cotter, in Marion County, Arkansas, *E. J. Palmer*, June 18, 1914 (No. 6026).

The specimens of Mr. Palmer's plant No. 3 can probably best be regarded as a variety of *C. padifolia* of the *Intricatae* group which occurs in this region, and from which it differs in its generally broader and usually slightly lobed leaves and in its larger and softer crimson fruit.

Crataegus panda has been used for two different plants, first by Beadle in 1902 (*Biltmore Bot. Studies*, 1. 89) for a shrubby species of the *Flavae* group from the neighborhood of Tallahassee, Florida, and second by Ashe in 1903 for a plant from Glendon, N. Carolina (in *Jour. Elisha Mitchell Sci. Soc.* xix. 29). The description does not give the group to which this plant belongs and there is not a specimen in the Arboretum herbarium.

Crataegus cirrata was used in 1902 by Beadle (*Biltmore Bot. Studies*, 1. 101) for a shrubby species of the *Flavae* group from Girard, Alabama. The same name was used in 1916 by Ashe (in *Bull. Charleston Mus.* xii. 42) for a plant from Georgia, no information being given of the type locality or of the group to which it belongs. Ashe's plant is not represented in the Arboretum herbarium.

SYNOPSIS OF NORTH AMERICAN CRATAEGI

ERNEST J. PALMER

INTRODUCTION

The following alphabetical list and synoptical tables of the North American species and varieties of the genus *Crataegus* was compiled several years ago, in a somewhat different form, for use in the Herbarium of the Arnold Arboretum. It was not intended for publication and was

designed merely for convenient reference and to facilitate the identification of material. The very unusual size of this genus in number of species, the bulk of material in our herbarium and the large accessions constantly accumulating, as well as specimens and collections frequently being received for identification from various parts of the country became so burdensome, even with the ample library and collection of types available here, that it was thought worth while to bring together in as compact a form as possible a synopsis of the information scattered through many publications and indices. It was hoped also that it might at some future time serve as the basis for a fuller study and more complete key to the genus. It is published with some reluctance at this time in the hope that it may prove of wider use and convenience to students of American *Crataegus*, especially to those who do not have access to much of the original literature or to the collections of the larger herbaria.

It is intended in the index to give a list as complete as possible of the specific and varietal names that have been applied to North American *Crataegi*. Possibly some names have escaped the writer's attention that should have been included, and many of those listed are given only for reference. No attempt is made to give a complete synonymy under valid names nor of combinations under other generic names. Specific names that have been variously interpreted by different botanists are included but once, exception being made in a few cases where it seems evident that duplication occurred through coincidence and that the authors were referring to different plants. Several instances of this sort will be noted, some of which are corrected in a paper by Professor C. S. Sargent appearing in this issue.

As this paper is not intended as a critical study in the sense of a revision of the genus, I have carefully refrained from making new combinations and so far as possible from going into the question of the validity of names that have been published. However, for the sake of clearness it was found necessary to indicate that certain names are recognized synonyms of others and where these have been used both as of specific and varietal rank to designate the preferred form. In such cases the practice prevailing at present in the Herbarium of the Arnold Arboretum has naturally been followed and names regarded as synonyms are printed in italics. A few obvious orthographical errors in previous publication of names have been corrected and for the sake of uniformity the practice of doubling the final i after consonants except after the ending "er," as recommended in the International Rules of Nomenclature, has been adopted for all names recognized as valid.

The question of synonymy and priority of names in this genus is much complicated by the fact that many North American species were first named from specimens cultivated in European gardens. These names were often without adequate descriptions or figures or with none at all, and in almost all cases without any definite records as to the part of the

country from which the seeds or plants were derived. North America was usually considered sufficiently definite and all of the eastern parts of the continent might be included under the general designations Carolina, Virginia and Canada. These early names were in many cases taken up and variously interpreted by later writers. Some of them, through study of the type specimens where they exist or of the plants still in cultivation have been found to have precedence over later names and more careful study will probably reveal other such cases. It is possible too that duplicate descriptions of some identical plants have been published under different names by later authors working independently.

The species have as far as possible been referred to the natural sections or groups to which they belong, those recognized being the twenty wholly or partially arborescent groups of Sargent's Manual of the Trees of North America and two additional shrubby ones, *Uniflorae* and *Pulcherrimae*. Perhaps two or three other small groups should be included in a fuller treatment of the North American species, at least in the writer's opinion. The distinctions between these groups are usually clear but in some cases there is an apparent merging of characters and it is difficult to say to which of two sections certain species belong. Where the group was indicated by the author of the species his treatment has usually been followed when the species comes within the recognized sections, although in a few instances there was some doubt in the writer's mind as to the correctness of the reference. Where the original description does not make the group clear and no specimens were available for examination I have seldom ventured to supply it and such species are necessarily excluded from the synoptical tables. These, with other debatable names that could not definitely be assigned to synonymy and all others of which I have not seen specimens, are indicated by a question mark in parenthesis.

The geographical range is given usually by states and provinces except in the cases of a few widely distributed species where it did not seem necessary to enumerate all political divisions. Where a wider range is indicated than that given in earlier publications it is in all cases based upon specimens examined, most of which are in the Arnold Arboretum Herbarium. In giving type localities, details have sometimes been supplied from the type specimens where these were available.

The synoptical tables were designed merely as a temporary substitute for a more complete key to the genus, the idea being to bring together species having common characters into small groups so that in connection with the geographical range only a few would have to be considered in identifying material. Like ordinary keys they are of course expected to be used only as guides to the fuller descriptions of the manuals.

In these tables the color of the anthers, number of stamens, glabrous or pubescent character of corymbs at flowering time and general shape of

the leaves were adopted for most of the groups, in the order named. This arrangement may not prove entirely satisfactory for a fuller treatment, but these characters were selected after much experimenting for two reasons: First, because they have been clearly stated and emphasized in most of the descriptions of recently proposed species, and second, for the sake of definiteness. It is extremely difficult to find in dealing with a genus like *Crataegus*, in which the number of species is so great and the differences between them in many cases slight, any common character that will hold good throughout a large series. The taxonomic value of characters varies in different groups, but generally there appears to be considerable variability even within the species in such particulars as the size of the flowers, the number in the corymbs and the compactness or laxity of the latter. The shape of the leaves is even less stable and dependable, many types often being found on a single branch; certain sorts of serration characterize some species, but in most there is much variability and differences are only relative; the texture and veining of the leaves is sometimes characteristic, but this is often modified by ecological conditions; the color of foliage, though not without significance, is subject to similar modification. The fruit is often one of the best guides to group distinctions, but there is quite too much variation in such particulars as shape, size and color for them to be depended upon rigidly as specific criteria. The number and shape of the nutlets is rather constant in some groups and quite variable in others; but these differences likewise, though furnishing in a few exceptional cases a distinction between closely allied species, usually serve merely as a guide to the group. Even the presence or absence of pubescence on the corymbs, often one of the best specific distinctions, cannot always be relied upon. In the color of the anthers, if not too narrowly defined, is to be found at least a distinction that is definite and not one of degree. Certain species seem to be characterized by having anthers of the yellow group of shades and others by those of shades that may generally be called red. These differences are often more or less distinctly correlated with the color of the fruit and shade of foliage, especially in its autumnal aspect.

In the number of stamens there is greater variability and they must be taken with less confidence and greater latitude as specific distinctions. As the flowers of this genus are 5-merous these organs normally appear in multiples of five. Thus in certain species, especially in the *Tenuifoliae* group, five appears to be the normal number, and these vary in individual flowers from four to seven or rarely eight. There is a double series in another large array of species in which the number may be expected to vary through suppression or redundancy between eight and twelve or fifteen, although in the great majority of flowers it is actually ten. In many species the series is again doubled and we find twenty stamens or rarely a few more or less. Considerable field observation and study

of specimens in cultivation seems to indicate that these characters, as thus liberally defined, generally breed true in seedlings. Exceptions may be expected in the case of hypothetical hybrids. It would therefore appear that the number of stamens and color of anthers furnish one of the most definite criteria for distinguishing between species, although perhaps conclusive only when combined with other important though variable morphologic characters; and they can scarcely be ignored, especially in the construction of keys, unless we are willing to adopt a much broader conception of a few variable and composite species. This of course involves a controversial question upon which there is the widest latitude for difference of opinion, and which is beyond the province or purpose of this paper. It is perhaps safe to say, however, that no competent student of the genus today would be willing to consider returning to the score or so of species recognized by the botanists of a few decades ago.

A brief synopsis of the fruit characters is given in the tables following the name of each species. These have in almost all cases been taken from the original descriptions and it should be remembered that a considerable deviation from some of these specifications may be expected in individual plants under varying ecological conditions. At least a wide acquaintance with many of the species in the field and observation of specimens under cultivation has led me to this conclusion. In a few cases where it seemed evident that errors had occurred in the original descriptions or where the examination of a series of specimens showed too wide a departure from these descriptions, corrections have been made. The question mark in parenthesis (?) following name of species and author, indicates that specimens have not been seen or that the position of the species in the tables is not definitely determinable from the description drawn upon.

ALPHABETICAL LIST OF SPECIES AND VARIETIES WITH SECTION AND RANGE

CRATAEGUS

- abbreviata Sargent in Jour. Arnold Arb. III. 187 (1922).—**Crus-galli.**—
Texas (type locality: Brazoria).
- abedita Beadle in Biltmore Bot. Studies, I. 75 (1902).—**Flavae.**—Florida
(type locality: River Junction).
- abjecta Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 636 (1905).—
Intricatae.—Pennsylvania (type locality: Sellersville).
- aboriginum Sargent in Rhodora, v. 163 (1903) = C. rotundifolia var.
aboriginum Sarg.
- abstrusa Beadle in Biltmore Bot. Studies, I. 66 (1902).—**Pulcherrimae.**—
Florida (type locality: Tallahassee).
- accincta Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 154 (1910).—
Crus-galli.—Pennsylvania (type locality: McKees Rocks).

CRATAEGUS

- acclivis Sargent in Proc. Rochester Acad. Sci. iv. 115 (1903).—**Coccineae.**—New York, Pennsylvania (type locality: Rochester, N. Y.).
- acerba Sargent in N. Y. State Mus. Bull. cxxii. 93 (1908).—**Silvicolae.**—New York (type locality: Coopers Plains).
- acerifolia* Loddiges Cat. 8, ex Moench, Bäume Weiss. 31 (1785) = *C. Phaenopyrum* Medic.
- acuminata Sargent in N. Y. State Mus. Bull. cv. Bot. ix. 56 (1906).—**Tenuifoliae.**—New York, Pennsylvania, Ohio (type locality: West Albany, N. Y.).
- acutifolia Sargent in Bot. Gaz. xxxi. 217 (1901).—**Crus-galli.**—Illinois, Missouri, Tennessee, Arkansas (type locality: Carondelet).
- acutiloba Sargent in Rhodora, iii. 23 (1901).—**Tenuifoliae.**—South-eastern Canada, New England (type locality: Bar Harbor, Me.).
- admiranda Sargent in N. Y. State Mus. Bull. cxxii. 80 (1908).—**Macrancantheae.**—Ontario, New York (type locality: Niagara Falls, N. Y.).
- adunca Beadle in Biltmore Bot. Studies, i. 87 (1902).—**Flavae.**—Florida (type locality: Tallahassee).
- adusta Beadle in Biltmore Bot. Studies, i. 110 (1902).—**Flavae.**—Florida (type locality: Gainesville).
- advena Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 175 (1910).—**Pruinosae.**—Pennsylvania (type locality: Bedford).
- aemula Beadle in Biltmore Bot. Studies, i. 53 (1902).—**Silvicolae.**—Georgia, Mississippi (type locality: Rome, Ga.).
- aestivalis (Walt.) Torrey & Gray, Fl. N. Am. i. 468 (1840).—**Aestivales.**—North Carolina, South Carolina, Georgia (type locality of *Mespilus aestivalis* Walter not given).
- aestivalis var. cerasoides Sargent in Jour. Arnold Arb. i. 250 (1920).—Florida (type locality: Seville).
- aestivalis var. cerasoides f. luculenta Sargent in Jour. Arnold Arb. i. 251 (1920).—Florida (type locality: Volusia County).
- aestivalis var. maloides Sargent in Jour. Arnold Arb. i. 250 (1920).—Florida (type locality: Seville).
- affinis Sargent in Ontario Nat. Sci. Bull. iv. 71 (1908).—**Anomalae.**—Southeastern Canada, New York (type locality: Toronto, Ont.).
- agaia Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 252 (1910).—**Macrancantheae.**—Pennsylvania (type locality: Scranton).
- agrestina Beadle in Biltmore Bot. Studies, i. 72 (1902).—**Flavae.**—Alabama (type locality: Evergreen).
- alabamensis Beadle in Bot. Gaz. xxx. 342 (1900).—**Flavae.**—Alabama (type locality: Montgomery).
- alacris Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 601 (1905).—**Pruinosae.**—Pennsylvania (type locality: Sellersville).
- albanthera Sargent in Rep. Missouri Bot. Gard. xix. 53 (1908) = *C. arduennae* Sarg.

CRATAEGUS

- albemarlensis* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 10 (1901).—Group?—North Carolina (type locality: “Hyde and Pamlico Counties”). (?)
- albicans* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 20 (1901).—
Silvicolae.—Michigan (type locality: “eastern Michigan”). (?)
- albicera* Beadle in Biltmore Bot. Studies, I. 132 (1902).—**Crus-galli.**—Louisiana (type locality: Opelousas).
- algens* Beadle in Biltmore Bot. Studies, I. 135 (1902).—**Crus-galli.**—Virginia and North Carolina to Tennessee and Arkansas (type locality: Biltmore, N. Car.).
- aliena* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 157 (1910).—**Crus-galli.**—Pennsylvania (type locality: Orbisonia).
- allecta* Sargent in Rep. Geol. Surv. Mich. 1906, 534 (1907).—**Silvicolae.**—Michigan (type locality: Grand Rapids).
- alleghaniensis* Beadle in Bot. Gaz. xxx. 337 (1900).—**Flavae.**—Alabama, Tennessee (type locality: Lookout Mountain, Ala.).
- alma* Beadle in Biltmore Bot. Studies, I. 64 (1902).—**Intricatae.**—Mississippi (type locality: Meridian).
- alnorum* Sargent in Rhodora, v. 153 (1903).—**Tenuifoliae.**—Maine (type locality: Orono).
- alpestris* Hort. ex K. Koch, Dendr. I. 144 (1869) = *C. crus-galli* var. *pyracanthifolia* Ait.
- alpista* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 238 (1910).—
Intricatae.—Pennsylvania (type locality: Orbisonia).
- altrix* Ashe in Bot. Gaz. xxxiii. 232 (1902).—Group?—Illinois (type locality: “northern Illinois”). (?)
- amabilis* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 15 (1903).—
Molles.—Iowa (type locality: Keokuk). (?)
- amara* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 22 (1902).—Group?—North Carolina (type locality: Elk Cross Roads). (?)
- ambigens* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 191 (1910).—
Silvicolae.—Pennsylvania (type locality: Kittanning).
- ambitiosa* Sargent, Trees and Shrubs, II. 240 (1913).—**Pruinosae.**—Michigan (type locality: Grand Rapids).
- ambrosia* Sargent in N. Y. State Mus. Bull. cv. Bot. ix. 69 (1906).—
Macracanthae.—Ontario, New York (type locality: Albany, N. Y.).
- amica* Beadle in Biltmore Bot. Studies, I. 97 (1902).—**Flavae.**—Florida (type locality: Ocala).
- amicalis* Sargent, Trees and Shrubs, II. 238 (1913).—**Virides.**—Arkansas (type locality: Fulton).
- amicta* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 26 (1902).—
Coccineae.—Illinois (type locality: “northern Illinois”).
- amnicola* Beadle in Biltmore Bot. Studies, I. 55 (1902).—
Punctatae.—Georgia, Alabama, Tennessee (type locality: Chattanooga, Tenn.).

CRATAEGUS

- amoena Sargent in N. Y. State Mus. Bull. cxxii. 38 (1908).—**Pruinosae.**—New York, Ohio (type locality: Niagara Falls, N. Y.).
- ampla Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 615 (September, 1905).—**Tenuifoliae.**—Pennsylvania (type locality: Kutztown).
- ampla Sargent in Rhodora, vii. 208 (November, 1905) = *C. neofaxonii* Sarg.
- ampliata Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 195 (1910).—**Silvicolae.**—Pennsylvania (type locality: Orbisonia).
- amplifica Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 172 (1910).—**Pruinosae.**—Pennsylvania (type locality: Pittsburgh).
- anamesa Sargent in Jour. Arnold Arb. iii. 189 (1922).—**Virides.**—Texas (type locality: Duke, Ft. Bend County).
- ancisa Beadle in Biltmore Bot. Studies, i. 63 (1902).—**Pulcherrimae.**—Alabama, Mississippi (type locality, Meridian, Miss.).
- angulata Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 166 (1910).—**Pruinosae.**—Pennsylvania, Ohio, Kentucky (type locality: Nine Mile Run, near Pittsburgh, Pa.).
- angustata Sargent in Rep. Missouri Bot. Gard. xix. 77 (1908).—**Punctatae.**—Missouri (type locality: Swan).
- angustifolia* (Medic.) Borkhausen in Roemer, Arch. Bot. i. no. 3, 86 (1798) = *C. crus-galli* var. *pyracanthifolia* Ait.
- angustisepala Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 209 (1910).—**Tenuifoliae.**—Pennsylvania (type locality: Throop, near Scranton).
- anisophylla Beadle in Biltmore Bot. Studies, i. 99 (1902).—**Flavae.**—Florida (type locality: Jacksonville).
- annosa Beadle in Biltmore Bot. Studies, i. 83 (1902).—**Flavae.**—Alabama (type locality: Phoenix City).
- anomala Sargent in Rhodora, iii. 74 (1901).—**Molles.**—Southeastern Canada, Vermont, New York (type locality: Caughnawaga, Que.).
- antheina Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 213 (1910).—**Tenuifoliae.**—Pennsylvania (type locality: West Lockport).
- antimima Sargent in Jour. Arnold Arb. iii. 191 (1922).—**Virides.**—Texas, Oklahoma (type locality: Duke, Tex.).
- antiplasta Sargent in Jour. Arnold Arb. iii. 190 (1922).—**Virides.**—Texas (type locality: Duke).
- aperta Sargent in Rep. Missouri Bot. Gard. xix. 96 (1908).—**Pruinosae.**—Missouri (type locality: Swan).
- apiifolia* (Marsh.) Michaux, Fl. Bor. Am. i. 287 (1803).—**Microcarpae.**—Virginia to Florida and west to eastern Texas (type region: “Carolina”).
- apiifolia* var. *flavanthera* Sargent in Rep. Missouri Bot. Gard. xx. 82 (1912) = *C. spathulata* Michx.
- apiifolia* major Loddiges, Cat. 45 (1836), nomen nudum.—Loudon, Arb. Brit. ii. 824 (1838), as synonym of *C. apiifolia* Michx.

CRATAEGUS

- apiifolia* var. *minor* Loudon, Arb. Brit. II. 825 (1838) = *C. apiifolia* Michx.
apiifolia × *brachyphylla* = *C. notha*.
apiomorpha Sargent in Bot. Gaz. XXXV. 386 (1903).—**Tenuifoliae.**—
Ohio, Illinois (type locality: Fort Sheridan, Ill.).
apposita Sargent in Bot. Gaz. XXXV. 103 (1903).—**Intricatae.**—Pennsylvania, Delaware (type locality: Wilmington, Del.).
apposita var. *Bissellii* (Sarg.) Eggleston in Rhodora, x. 76 (1908) = *C. Bissellii*.
aprica Beadle in Bot. Gaz. XXX. 335 (1900).—**Flavae.**—North Carolina, Georgia, Alabama (type locality: Biltmore, N. Car.).
aquilonaris Sargent in Rhodora, v. 185 (1903).—**Macracantheae.**—
Southeastern Canada (type locality: Nipissing, Ont.).
araioclada Sargent in Jour. Arnold Arb. III. 5 (1922).—**Crus-galli.**—
Louisiana (type locality: Natchitoches).
arborea Beadle in Biltmore Bot. Studies, I. 137 (1902).—**Crus-galli.**—
Florida, Alabama (type locality: Montgomery, Ala.).
arborescens Elliot, Sketch Bot. I. 550 (1821).—**Virides.**—Georgia, Florida (type locality: Fort Argyle, on Ogeechee River, Ga.).
arcana Beadle in Biltmore Bot. Studies, I. 122 (1902).—**Pruinosae.**—
New York to North Carolina, west to Michigan and Missouri (type locality: Biltmore, N. Car.).
arcuata Ashe in Ann. Carnegie Mus. I. 387 (1902).—**Coccineae.**—
Pennsylvania, Delaware (type locality: Kutztown, Pa.).
ardua Sargent in Ontario Nat. Sci. Bull. IV. 93 (1908).—**Macracantheae.**—
Ontario (type locality: Toronto).
arduennae Sargent in Bot. Gaz. XXXV. 377 (1903).—**Crus-galli.**—
Ontario and New York, west to Michigan, Missouri and Arkansas (type locality: Glenellyn, Ill.).
arenicola Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 5 (1900).—
Uniflorae.—North Carolina (type locality: Jessup). (?)
arguta Beadle in Biltmore Bot. Studies, I. 102 (1902).—**Flavae.**—
Georgia, Alabama (type locality: Girard, Ala.).
aridula Sargent in N. Y. State Mus. Bull. CXXII. 43 (1908).—**Pruinosae.**—
New York (type locality: Niagara Falls).
aristata Sargent in N. Y. State Mus. Bull. CL. 27 (1911).—**Pruinosae.**—
New York (type locality: Rossie).
arkansana Sargent in Bot. Gaz. XXXI. 223 (1901).—**Molles.**—Arkansas (type locality: cultivated in Arnold Arboretum from seed collected at Newport, Ark.).
armata Beadle in Biltmore Bot. Studies, I. 136 (1902).—**Crus-galli.**—
Alabama, Tennessee (type locality: Nashville, Tenn.).
armentalis Beadle in Biltmore Bot. Studies, I. 117 (1902).—**Uniflorae.**—
Alabama (type locality: Albertsville).

CRATAEGUS

- armigera* Sargent in Ontario Nat. Sci. Bull. iv. 87 (1908).—**Macracthae.**—Southeastern Canada (type locality: Toronto, Ont.).
- arnoldiana* Sargent in Bot. Gaz. xxxi. 221 (1901).—**Molles.**—Massachusetts, Connecticut, New York (type locality: Arnold Arboretum, cultivated).
- arrogans* Beadle in Biltmore Bot. Studies, i. 81 (1902).—**Flavae.**—Georgia, Alabama (type locality: Phoenix City, Ala.).
- arta* Beadle in Biltmore Bot. Studies, i. 130 (1902).—**Crus-galli.**—Tennessee (type locality: Nashville).
- ascendens* Sargent in Rhodora, v. 141 (1903).—**Tenuifoliae.**—Vermont, New York (type locality: New Haven, Vt.).
- Ashei* Beadle in Bot. Gaz. xxx. 339 (1900).—**Bracteatae.**—Alabama, Mississippi (type locality: Montgomery County, Ala.).
- aspera* Sargent, Trees and Shrubs, ii. 67 (1908).—**Pruinosae.**—Missouri (type locality: Webb City).
- asperata* Sargent in Rep. Geol. Surv. Mich. 1906, 542 (1907).—**Tenuifoliae.**—Michigan (type locality: Port Huron).
- asperifolia* Sargent in Rhodora, iii. 31 (1901).—**Anomalae.**—Southeastern Canada, New England, New York (type locality: Middlebury, Vt.).
- assimilis* Beadle in Biltmore Bot. Studies, i. 68 (1902).—**Pulcherrimae.**—Florida (type locality: Chattahoochee).
- assurgens* Sargent in Bot. Gaz. xxxv. 382 (1903).—**Coccineae.**—Ohio, Illinois, Wisconsin (type locality: Glendon Park, near Chicago, Ill.).
- ater* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 21 (1903).—**Pruinosae.**—Michigan, Ohio (type locality: Port Huron, Mich.).
- atropurpurea* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 12 (1901) = *C. velata* Ashe.
- atrорubens* Ashe in Jour. Elisha Mitchell Sci. Soc. xvi. 78 (1900).—**Virides.**—Illinois, Missouri (type locality: St. Louis County, Mo.).
- attenuata* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 30 (1903).—**Crus-galli.**—Michigan, Illinois (type locality: Port Huron, Mich.).
- attrita* Beadle in Biltmore Bot. Studies, i. 98 (1902).—**Flavae.**—Alabama (type locality: Ozark).
- audens* Beadle in Biltmore Bot. Studies, i. 114 (1902).—**Flavae.**—Florida (type locality: Chattahoochee).
- augusta* Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 597 (1905).—**Pruinosae.**—Pennsylvania (type locality: Kutztown).
- aulica* Sargent in Ontario Nat. Sci. Bull. iv. 48 (1908).—**Coccineae.**—Ontario, New York, Michigan (type locality: Toronto, Ont.).
- aurea* Hort. ex Nicholson in Hand-list Arb. Kew. pt. 1, 209 (1894) as a synonym of *C. punctata* var. *xanthocarpa* = *C. punctata* var. *aurea* Ait.

CRATAEGUS

- austera Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 586 (1905).—**Pruinosae.**—Pennsylvania (type locality: Sellersville).
- austrina Beadle in Biltmore Bot. Studies, I. 59 (1902).—**Pulcherrimae.**—Alabama (type locality: Greenville).
- austromontana Beadle in Bot. Gaz. XXVIII. 412 (1899).—**Triflorae.**—Georgia, Alabama, Tennessee (type locality: Valley Head, Ala.).
- axillaris* (Pers.) Loddiges, Cat. 39 (1826).—Loudon, Arb. Brit. II. 841 (1838) as synonym of *C. parvifolia* Ait.
- baccata Sargent in Rhodora, VII. 176 (1905).—**Macracanthae.**—Massachusetts (type locality: Lancaster).
- Balkwillii Sargent in Ontario Nat. Sci. Bull. IV. 80 (1908).—**Macracanthae.**—Ontario, New York (type locality: London, Ont.).
- barbara Sargent in N. Y. State Mus. Bull. CXXII. 33 (1908).—**Punctatae.**—New York (type locality: Brighton).
- barbata Sargent in Rep. Missouri Bot. Gard. XIX. 63 (1908).—**Crus-galli.**—Missouri (type locality: Swan).
- Baroussana Eggleston in Torreya, VII. 35 (1907).—**Molles.**—Mexico (type locality: Saltillo, Coahuila).
- barrettiana Sargent in Rep. Missouri Bot. Gard. XIX. 50 (1908).—**Crus-galli.**—Ohio, Indiana, Illinois, Missouri (type locality: Barretts Station, Mo.).
- Barryana Sargent in N. Y. State Mus. Bull. CXXII. 52 (1908).—**Silvicolae.**—New York (type locality: Rochester).
- Bartoniana Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 637 (1905).—**Intricatae.**—Pennsylvania (type locality: Gray's Ferry).
- Bartramiana Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 582 (1905).—**Crus-galli.**—Pennsylvania (type locality: Bartram's Lane, Philadelphia).
- basilica Beadle in Biltmore Bot. Studies, I. 125 (1902).—**Tenuifoliae.**—North Carolina, South Carolina (type locality: Candler, N. Car.).
- Baxteri Sargent in Proc. Rochester Acad. Sci. IV. 107 (1903) = *C. foetida* Ashe.
- Beadlei Ashe in Bull. N. Car. Exper. Sta. CLXXV. 113 (1900).—**Intricatae.**—North Carolina (type locality: Salisbury). (?)
- Bealii Sargent in Rep. Geol. Surv. Mich. 1906, 553 (1907).—**Intricatae.**—Michigan (type locality: Grand Rapids).
- beata Sargent in Proc. Rochester Acad. Sci. IV. 97 (1903).—**Pruinosae.**—New York, Pennsylvania (type locality: Rochester, N. Y.).
- beata* var. *compta* (Sarg.) Eggleston in Rhodora, X. 81 (1908) = *C. compta*.
- Beckiana Sargent in N. Y. State Mus. Bull. CV. 75 (1906).—**Macracanthae.**—New York (type locality: North Greenbush).
- Beckwithae Sargent in Proc. Rochester Acad. Sci. IV. 124 (1903) = *C. diffusa* Sarg.

CRATAEGUS

- bedfordensis* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 185 (1910).—
Pruinosae.—Pennsylvania, Ohio (type locality: Bedford Springs, Pa.).
- bella* Sargent in N. Y. State Mus. Bull. cxxii. 61 (1908).—Tenuifoliae.—Ontario, New York, Pennsylvania, Ohio (type locality: Buffalo, N. Y.).
- bellatula* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 171 (1910).—
Pruinosae.—Pennsylvania (type locality: Bedford).
- bellica* Sargent, Trees and Shrubs, II. 236 (1913).—Crus-galli.—Arkansas, Oklahoma, Texas (type locality: Fulton, Ark.).
- bellula* Sargent, Trees and Shrubs I. 111 (1903).—Pruinosae.—Massachusetts, Michigan (type locality: Grand Rapids, Mich.).
- benigna* Sargent in Proc. Rochester Acad. Sci. IV. 127 (1903).—Tenuifoliae.—New York (type locality: Rochester).
- berberifolia* Torrey & Gray, Fl. N. Am. I. 469 (1840).—Crus-galli.—Louisiana (type locality: Opelousas).
- berberifolia* var. *Engelmanni* (Sarg.) Eggleston in Rhodora, X. 76 (1908) = C. Engelmannii Sarg.
- Berlandieri* Sargent in Bot. Gaz. XXXI. 230 (1901).—Molles.—Texas (type locality: Columbia).
- betulifolia* Loddiges, Cat. 45 (1836).—Loudon, Arb. Brit. II. 841 (1838) as a synonym of *C. parvifolia* Ait.—Nomen nudum.
- Bicknellii* Eggleston in Bull. Torrey Bot. Club XXXVIII. 244 (1911).—Rotundifoliae.—Massachusetts (type locality: Nantucket Island).
- bicolor* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 12 (1901).—Group?—North Carolina (type locality: Mitchell County). (?)
- biltmoreana* Beadle in Bot. Gaz. XXVIII. 406 (1899).—Intricatae.—Pennsylvania and Virginia to South Carolina, Alabama and Tennessee (type locality: Biltmore, N. Car.).
- Bissellii* Sargent in Rhodora, V. 65 (1903).—Intricatae.—Connecticut, New York (type locality: Southington, Ct.).
- bisulcata* Ashe in Bull. N. Car. Exper. Sta. CLXXV. 112 (1900).—Uniflorae.—Georgia (type locality: Stone Mountain).
- blairensis* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 193 (1910).—Silvicolae.—Pennsylvania, Ohio (type locality: Altoona, Pa.).
- Blanchardii* Sargent in Rhodora, VII. 218 (1905).—Rotundifoliae.—Ontario, Vermont (type locality: Wilmington, Vt.).
- blanda* Sargent in Bot. Gaz. XXXIII. 121 (1902).—Virides.—Alabama, Louisiana, Arkansas, Texas (type locality: Fulton, Ark.).
- blandita* Sargent in Rhodora, V. 147 (1903).—Tenuifoliae.—Quebec (type locality: Caughnawaga).
- bona* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 587 (1905).—Pruinosae.—Pennsylvania (type locality: Berks County).

CRATAEGUS

- Boothiana* Sargent in N. Y. State Mus. Bull. cxxii. 58 (1908).—**Tenuifoliae.**—New York (type locality: Rochester).
- borealis* Ashe in Bull. N. Car. Exper. Sta. clxxv. 111 (1900).—Group?—Michigan (type locality not given). (?)
- Bosciana* Roemer, Fam. Nat. Syn. III. 119 (1847) = *C. crus-galli* L.
- Boyntonii* Beadle in Bot. Gaz. xxviii. 409 (1899).—**Intricatae.**—Virginia, North Carolina, Georgia, Alabama, Tennessee (type locality: Biltmore, N. Car.).
- Boytoni* var. *Buckleyi* (Beadle) Ashe in Jour. Elisha Mitchell Sci. Soc. xxxiv. 139 (1918) = *C. Buckleyi*.
- brachyacantha* Sargent & Engelmann apud Engelmann in Bot. Gaz. VII. 128 (1882).—**Brachyacanthae.**—Arkansas, Louisiana, Texas (type locality not given).
- brachyacantha* f. *leucocarpa* Sargent in Jour. Arnold Arb. III. 10 (1922).—Louisiana (type locality: Natchitoches).
- brachyacantha* var. *maxima* Kuntze, Rev. Gen. Pl. I. 215 (1891).—Mississippi (type locality: Cairo). (?)
- brachyloba* Sargent in N. Y. State Mus. Bull. cxxii. 75 (1908).—**Anomalae.**—New York (type locality: Buffalo).
- brachiphylla* Sargent in Jour. Arnold Arb. III. 8 (1922).—**Molles.**—Arkansas (type locality: McNab, Hempstead County).
- brachiphylla* × *apiifolia* = *C. notha*.
- brachypoda* Sargent in Rep. Missouri Bot. Gard. xix. 100 (1908).—**Pruinosae.**—Kentucky, Missouri (type locality: Grandin, Mo.).
- bracteata* Sargent in Rep. Missouri Bot. Gard. xix. 91 (1908).—**Pruinosae.**—Missouri, Arkansas, Oklahoma (type locality: Webb City, Mo.).
- Brainerdii* Sargent in Rhodora, III. 27 (1901).—**Rotundifoliae.**—South-eastern Canada, Vermont (type locality: Middlebury, Vt.).
- Brainerdi* var. *asperifolia* (Sarg.) Eggleston in Rhodora, x. 82 (1908) = *C. asperifolia*.
- Brainerdi* var. *Egglestoni* (Sarg.) Robinson in Rhodora, x. 82 (1908) = *C. Egglestonii*.
- Brainerdi* var. *scabrida* (Sarg.) Eggleston in Rhodora, x. 82 (1908) = *C. scabrida*.
- brazoria* Sargent in Bot. Gaz. xxxi. 233 (1901).—**Punctatae.**—Texas (type locality: Brazoria).
- brevipedicellata* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. 6 (1900).—Group?—North Carolina (type locality: Priest Hill). (?)
- brevipes* Peck in N. Y. State Mus. Bull. cxxxix. 20 (1910).—**Pruinosae.**—New York (type locality: Corning).
- brevispina* Douglas apud Steudel, Nomencl. Bot. ed. 2, I. 432 (1841) = *C. Douglassii* Lindl.

CRATAEGUS

- bristolensis* Sargent in *Rhodora*, vii. 175 (1905).—**Macracanthae.**—Massachusetts (type locality: Somerset).
- Brittonii* Eggleston in *Bull. Torrey Bot. Club*, xxxvi. 640 (1909).—**Uniflorae.**—North Carolina, Georgia, Tennessee (type locality: Biltmore, N. Car.).
- Brockwayae* Sargent, *Trees and Shrubs*, ii. 245 (1913).—**Anomalae.**—Michigan (type locality: Clifton).
- bronxensis* Sargent in *N. Y. State Mus. Bull.* cxxii. 115 (1908).—**Pruinosae.**—New York (type locality: New York Botanical Garden).
- Brownietta* Sargent in *N. Y. State Mus. Bull.* clxvii. 78 (1913).—**Punctatae.**—New York (type locality: Hemlock Lake, Livingston County).
- Brownii* Britton in *Bull. N. Y. Bot. Gard.* i. 447 (1900) = *C. Margaretta* var. *Brownii* Sarg.
- brumalis* Ashe in *Ann. Carnegie Mus.* i. 393 (1902).—**Tenuifoliae.**—Pennsylvania (type locality: Pittsburgh).
- Brunetiana* Sargent in *Rhodora*, v. 164 (1903).—**Rotundifoliae.**—Southeastern Canada (type locality: Montmorency Falls, Que.).
- Buckleyi* Beadle in *Biltmore Bot. Studies*, i. 25 (1901).—**Intricatae.**—Virginia, North Carolina, Tennessee (type locality: Biltmore, N. Car.).
- Buistii* Hort. ex K. Koch, *Dendr.* i. 144 (1869) = *C. crus-galli* var. *pyracanthifolia* Ait.?
- Burkeana* Sargent in *Proc. Acad. Nat. Sci. Phila.* lxii. 218 (1910).—**Coccineae.**—Pennsylvania (type locality: Pittsburgh).
- Bushii* Sargent in *Bot. Gaz.* xxxiii. 109 (1902).—**Crus-galli.**—Arkansas, Louisiana, Oklahoma, Texas (type locality: Fulton, Ark.).
- caerulescens* Sargent in *Jour. Arnold Arb.* iii. 193 (1922).—**Pruinosae.**—Massachusetts (type locality: Arnold Arboretum, transplanted from Orient Heights, Breed's Island, Boston Harbor).
- caesa* Ashe in *Jour. Elisha Mitchell Sci. Soc.* xviii. 24 (1902).—**Coccineae.**—Michigan (type locality: Port Huron).
- caesariata* Sargent in *N. Y. State Mus. Bull. cv. Bot.* ix. 64 (1906).—**Rotundifoliae.**—New York (type locality: North Albany).
- calcarigera* Salisbury, *Prodr.* 357 (1796) = *C. crus-galli* L.
- caliciglabra* Schuette in *Proc. Biol. Soc. Washington*, xvi. 96 (1903).—**Rotundifoliae.**—Wisconsin (type locality not given).
- calliantha* Sargent in *Rep. Missouri Bot. Gard.* xix. 93 (1908).—**Pruinosae.**—Missouri (type locality: Pacific).
- callicarpa* Sargent in *Rep. Missouri Bot. Gard.* xix. 96 (1908).—**Pruinosae?**—Missouri (type locality: Shrewsbury).
- callida* Beadle in *Biltmore Bot. Studies*, i. 123 (1902).—**Pruinosae.**—Alabama (type locality: Gadsden).

CRATAEGUS

- callista Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 237 (1910).—
Intricatae.—Pennsylvania (type locality: Sellersville).
- callosa Sargent in Acad. Nat. Sci. Phila. LVII. 595 (1905).—**Pruinosae.**—
 Pennsylvania (type locality: North Heidelberg Township, Berks
 County).
- calophylla Sargent in Rep. Missouri Bot. Gard. xxii. 67 (1912).—**Crus-**
galli.—Tennessee, Missouri, Arkansas (type locality: Springfield,
 Mo.).
- calpodendron (Ehrh.) Medicus, Gesch. Bot. 83 (1793).—**Macracantheae.**—
 Identity and range uncertain (type region of *Mespilus calpodendron*
 Ehrhart: “Nordamerika”).
- calpodendron × punctata Eggleston in N. Y. State Mus. Bull. ccXLIII-
 iv. 65 (1923). (?)
- calva Beadle in Biltmore Bot. Studies, I. 83 (1902).—**Flavae.**—Alabama
 (type locality: Ozark).
- calvescens Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 159 (1910).—
Punctatae.—Pennsylvania (type locality: Allegheny).
- Calvinii Sargent in N. Y. State Mus. Bull. cxxii. 81 (1908).—**Macra-**
cantheae.—New York (type locality: Canandagua).
- campestris Britton in Bull. N. Y. Bot. Gard. I. 449 (1900) = C. perto-
 mentosa Ashe.
- canadensis Sargent in Rhodora, III. 73 (1901).—**Molles.**—Quebec (type
 locality: Caughnawaga).
- Canbyi Sargent in Bot. Gaz. xxxi. 3 (1901).—**Crus-galli.**—Pennsyl-
 vania, Delaware, Maryland (type locality: Wilmington, Del.).
- candens Sargent in Rep. Missouri Bot. Gard. xix. 55 (1908) = C.
 arduennae Sarg.
- carnosa Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 14 (1901).—
 Group?—North Carolina (type locality: Yancy County). (?)
- caroliniana (Poir.) Persoon, Syn. II. 36 (1806).—**Flavae.**—(Type region:
 “Carolina”). (?)
- carrollensis Sargent in Jour. Arnold Arb. III. 204 (1922).—**Macracan-**
thae.—Arkansas (type locality: Eureka Springs).
- casta Sargent in N. Y. State Mus. Bull. cv. Bot. ix. 53 (1906).—**Prui-**
nosae.—New York (type locality: North Greenbush).
- catawbiensis Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. 6 (1900).—
 Group?—North Carolina (type locality: Round Knob). (?)
- celsa Sargent in N. Y. State Mus. Bull. cxxii. 31 (1908).—**Punctatae.**—
 New York (type locality: Niagara Falls).
- cerasifera Loddiges, Cat. 45 (1836), nomen nudum.—Loudon, Arb.
 Brit. II. 849 (1838), as a synonym of C. crus-galli var. splendens Ait.
- cerasina Sargent in N. Y. State Mus. Bull. cxxii. 29 (1908).—**Crus-**
galli.—New York (type locality: Niagara Falls).
- cerasoides Sargent, Trees and Shrubs, II. 237 (1913) = C. aestivalis

CRATAEGUS

- var. *cerasoides* Sarg.
- cernua* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 25 (1902).—**Rotundifoliae**.—Wisconsin (type locality: Milwaukee). (?)
- cerronis* Nelson in Bot. Gaz. xxxiv. 370 (1902).—**Douglasianae**.—Colorado, Wyoming (type locality: Cerro Summit, Colo.). (?)
- cestrica* Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 596.—**Pruinosae**.—Pennsylvania (type locality: Newtown).
- chadfordiana* Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 660 (1905).—**Macracanthae**.—Pennsylvania (type locality: Chadford).
- champlainensis* Sargent in Rhodora, iii. 20 (1901).—**Molles**.—South-eastern Canada, Vermont, New York, Michigan (type locality: Middlebury, Vt.).
- Chapmanii* (Beadle) Ashe in Bot. Gaz. xxviii. 271 (1899).—**Macracanthae**.—North Carolina and Georgia, west to Illinois and Missouri (type locality of *C. tomentosa* var. *Chapmani*: Silver Creek, Floyd County, Ga.).
- Chapmani* var. *Plukenetii* Eggleston in Rhodora, x. 83 (1908) = *C. tomentosa* L.
- chateaugayensis* Sargent in N. Y. State Mus. Bull. cxxii. 121 (1908).—**Rotundifoliae**.—New York (type locality: Chateaugay Lake).
- cherokeensis* Sargent in Jour. Arnold Arb. iii. 1 (1922).—**Crus-galli**.—Texas (type locality: Larissa).
- chippewaensis* Sargent in Ontario Nat. Sci. Bull. iv. 44 (1908).—**Coccineae**.—Ontario (type locality: Chippewa).
- choriophylla* Sargent in Jour. Arnold Arb. iii. 201 (1922).—**Uniflorae**.—Florida (type locality: Lake City).
- chrysocarpa* Ashe in Bull. N. Car. Exper. Sta. clxxv. 110 (1900).—**Rotundifoliae**.—Manitoba, Nebraska, North Dakota, Wyoming, Colorado (type locality: foothills of Cache la Poudre Mountains, Colo.).
- chrysocarpa* var. *Faxonii* (Sarg.) Eggleston in N. Y. State Mus. Bull. ccxliii-iv. 64 (1923) = *C. Faxonii*.
- chrysocarpa* × *macrosperma* Eggleston in N. Y. State Mus. Bull. ccxlii-iv. 64 (1923). (?)
- chrysocarpa* × *punctata* Eggleston in N. Y. State Mus. Bull. ccxliii-iv. 64 (1923). (?)
- cibaria* Beadle in Biltmore Bot. Studies, i. 120 (1902).—**Molles**.—Tennessee (type locality: Nashville).
- cibilis* Ashe in Bot. Gaz. xxxiii. 232 (1902).—**Molles**.—North Carolina, Tennessee (type locality not given). (?)
- cicur* Ashe in Ann. Carnegie Mus. i. 397 (1902).—**Pruinosae**.—Pennsylvania, North Carolina (type locality not given). (?)
- ciliata* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. 13 (1900).—Group?—North Carolina (type locality: Raleigh). (?)

CRATAEGUS

cirrata Beadle in Biltmore Bot. Studies, I. 101 (1902).—**Flavae.**—Alabama (type locality: Girard).

cirrata Ashe in Bull. Charleston Mus. XII. 42 (1916). Group?—Georgia (type locality not given). (?)

clara Beadle in Biltmore Bot. Studies, I. 95 (1902).—**Flavae.**—Florida (type locality: Bristol).

Claytoniana Sargent in N. Y. State Mus. Bull. CXXII. 120 (1908).—**Tenuifoliae.**—New York (type locality: Clayton).

Clintoniana Sargent in N. Y. State Mus. Bull. CXXII. 39 (1908).—**Pruinosae.**—New York, Ohio (type locality: Buffalo, N. Y.).

coccinata Sargent in Ontario Nat. Sci. Bull. IV. 63 (1908).—**Rotundifoliae.**—Ontario (type locality: London).

coccinea Linnaeus, Spec. 476 (1753), in part = *C. rotundifolia* Moench.

coccinea var. *corallina* (Lodd.) Loudon, Arb. Brit. II. 817 (1838) = *C. Phaenopyrum* Medic.

coccinea var. *cordata* Lavallée, Icon. Arb. Segrez. 81, t. 23 (1882). (?)

coccinea var. *Ellwangeriana* (Sarg.) Eggleston in N. Y. State Mus. Bull. CCXLIII-IV. 64 (1923) = *C. Ellwangeriana*.

coccinea a. *flabellata* (Bosc) Dippel, Handb. Laubholzk. III. 435 (1893) = *C. flabellata*.

coccinea glandulosa Hort. ex Nicholson in Hand-list Hort. Kew. pt. I. 203 (1894), as a synonym of *C. mollis* Scheele.

coccinea var. *indentata* (Lodd.) Loudon, Arb. Brit. II. 817 (1838). (?)

coccinea *Kelmanni* Hort. ex Nicholson in Hand-list Arb. Kew. pt. I. 203 (1894) as a synonym of *C. mollis* Scheele. (?)

coccinea γ. *Kihlmanni* Hort. apud Lange, Rev. Gen. Crat. 30 (1897). (?)

coccinea var. *macracantha* (Lodd.) Dudley in Bull. Cornell University II. 33 (1886) = *C. macracantha*.

coccinea maxima Loddiges, Cat. 39 (1826).—Loudon, Arb. Brit. II. 817 (1838). (?)

coccinea ε. *microphylla* Lange, Rev. Gen. Crat. 30 (1897). (?)

coccinea ε? *mollis* Torrey & Gray, Fl. N. Am. I. 465 (1840) = *C. mollis*.

coccinea δ. *oligandra* Torrey & Gray, Fl. N. Am. I. 465 (1840). (?)

coccinea γ. *populifolia* (Walt.) Torrey & Gray, Fl. N. Am. I. 465 (1840). (?)

coccinea d. *pruinosa* (Wendl.) Dippel, Handb. Laubholzk. III. 435 (1893) = *C. pruinosa*.

coccinea var. *rotundifolia* (Moench) Sargent in Bot. Gaz. XXXI. 14 (1901) = *C. rotundifolia*.

coccinea var. *spinosa* Godefroy ex Loudon, Arb. Brit. II. 817 (1838) as synonym of *C. coccinea* var. *maxima* Lodd.

coccinea * *subvillosa* (Schrad.) Lange, Rev. Gen. Crat. 31 (1897) = *C. submollis* Sarg.

CRATAEGUS

- coccinea* $\beta.$ *viridis* (L.) Torrey & Gray, Fl. N. Am. I. 465 (1840) = *C. viridis*.
coccinea \times *macroisperma* Eggleston in N. Y. State Mus. Bull. CCXLIII-IV.
 65 (1923). (?)
coccinea \times *pruinosa* Eggleston in N. Y. State Mus. Bull. CCXLIII-IV. 65
 (1923). (?)
coccinoides Ashe in Jour. Elisha Mitchell Sci. Soc. XVI. 74 (1900).—
 Dilatatae.—Illinois, Missouri, Arkansas (type region: southern Illinois and eastern Missouri).
coccinoides var. *dilatata* (Sarg.) Eggleston in Rhodora, x. 81 (1908) =
 C. dilatata.
Cocksii Sargent in Jour. Arnold Arb. I. 248 (1920).—**Crus-galli**.—
 Louisiana (type locality: Winfield).
coerulea Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 207 (1910).—
 Silvicolae.—Pennsylvania (type locality: Lincoln Heights, near Scranton).
cognata Sargent in Rhodora, v. 58 (1903).—**Pruinosae**.—Southeastern Canada, New England, New York, Pennsylvania and Ohio (type locality: Great Barrington, Mass.).
Coleae Sargent, Trees and Shrubs, I. 7 (1902).—**Anomalae**.—Michigan (type locality: Grand Rapids).
collicola Ashe in Jour. Elisha Mitchell Sci. Soc. XVI. 75 (1900).—**Crus-galli**.—North Carolina (type locality not given). (?)
collina Chapman, Fl. ed. 2, Suppl. 2, 684 (1892).—**Punctatae**.—Virginia and North Carolina to Tennessee (type locality not given).
collina var. *Lettermani* (Sarg.) Eggleston in Rhodora, x. 76 (1908) =
 C. Lettermanii.
collina var. *sordida* (Sarg.) Eggleston in Rhodora, x. 76 (1908) = *C. sordida*.
Collinsiana Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 611 (1905).—
 Tenuifoliae.—Pennsylvania (type locality: Sellersville).
colonica Beadle in Biltmore Bot. Studies, I. 104 (1902).—**Flavae**.—
 South Carolina (type locality: Bluffton).
coloradensis Nelson in Proc. Biol. Soc. Washington, XVII. 175 (1904).—
 Macracanthae.—Colorado (type locality: Gregory Canyon, near Boulder).
colorado Ashe in Bull. N. Car. Exper. Sta. CLXXV. 110 (1900).—
 Douglasianae.—North Dakota and Montana to Nebraska, Colorado and Arizona (type locality: foot-hills of Cache la Poudre Mountains, Colo.).
coloradooides Ramaley in Bot. Gaz. XLVI. 383 (1908).—
 Macracanthae.—
 Colorado (type locality: Pole Canyon, near Boulder). (?)
colorata Sargent in Proc. Rochester Acad. Sci. IV. 123 (1903).—
 Tenuifoliae.—Ontario and New York to Michigan and Indiana (type locality: Rochester, N. Y.).

CRATAEGUS

- columbiana* Howell, Fl. N. W. Am. I. 163 (1898).—**Douglasianae.**—
Washington (type locality not given).
- columbiana* Sargent in Bot. Gaz. XXXI. 229 (1901) = *C. quercina* Ashe.
- columbiana* var. *Brunetiana* (Sarg.) Eggleston in Rhodora, x. 79 (1908) =
C. Brunetiana.
- columbiana* var. *Piperi* (Britton) Eggleston in Rhodora, x. 79 (1908) =
C. Piperi.
- comans* Sargent in N. Y. State Mus. Bull. CXXII. 112 (1908).—**Macra-**
canthae.—New York (type locality: Coopers Plains).
- comata* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 592 (1905).—
Pruinosae.—Pennsylvania (type locality: Gray's Ferry).
- communis* Beadle in Biltmore Bot. Studies, I. 58 (1902).—**Intricatae.**—
West Virginia, Tennessee (type locality: Cowan, Tenn.).
- compacta* Sargent in Rep. Geol. Surv. Mich. 1906, 522 (1907).—**Puncta-**
tae.—Ontario, Michigan (type locality: London, Ont.).
- comparata* Sargent, Trees and Shrubs, II. 240 (1913).—**Pruinosae.**—
Michigan (type locality: Grand Rapids).
- compatalis* Ashe in Bull. Charleston Mus. XII. 43 (1916).—Group?—
South Carolina (type locality not given). (?)
- comitalis* Beadle in Biltmore Bot. Studies, I. 93 (1902).—**Flavae.**—
Florida (type locality: Gainesville).
- compta* Sargent in Proc. Rochester Acad. Sci. IV. 102 (1903).—**Silvi-**
colae.—Ontario, New York, Pennsylvania (type locality: Rochester,
N. Y.).
- concinna* Beadle in Biltmore Bot. Studies, I. 70 (1902).—**Pulcherrimae.**—
Florida (type locality: Bristol).
- condensa* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 617 (1905).—
Tenuifoliae.—Pennsylvania (type locality: Sellersville).
- condigna* Beadle in Biltmore Bot. Studies, I. 35 (1901).—**Flavae.**—
Florida (type locality: River Junction).
- conferta* Sargent in N. Y. State Mus. Bull. CXXII. 62 (1908).—**Tenui-**
foliae.—Ontario, New York (type locality: Buffalo, N. Y.).
- confinis* Sargent in Ontario Nat. Sci. Bull. IV. 50 (1908).—**Coccineae.**—
Ontario, New York (type locality: Chippewa, Ont.).
- confragosa* Sargent in Ontario Nat. Sci. Bull. IV. 24 (1908).—**Pruinosae.**—
Ontario, Ohio (type locality: Sarnia, Ont.).
- confusa* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 235 (1910).—
Intricatae.—Pennsylvania (type locality: Allegheny).
- congestiflora* Sargent in N. Y. State Mus. Bull. CXXII. 44 (1908).—
Silvicolae.—Quebec, New York (type locality: Buffalo, N. Y.).
- conjuncta* Sargent in Rhodora, V. 57 (1903).—**Pruinosae.**—New England
and New York to Indiana, Illinois and Arkansas (type locality:
Lancaster, Mass.).

CRATAEGUS

- conjungens Sargent in Jour. Arnold Arb. III. 200 (1922).—**Triflorae.**—
Virginia (type locality: Clifton Forge).
- consanguinea Beadle in Biltmore Bot. Studies, I. 34 (1901).—**Flavae.**—
Florida (type locality: Tallahassee).
- conspecta Sargent in Ontario Nat. Sci. Bull. IV. 28 (1908).—**Coccineae.**
—Ontario, New York (type locality: Chippewa, Ont.).
- conspicua Sargent in N. Y. State Mus. Bull. CV. Bot. IX. 74 (1906).—
Macracanthae.—Quebec, Vermont, New York. (type locality:
North Albany, N. Y.).
- constans Beadle in Biltmore Bot. Studies, I. 88 (1902).—**Flavae.**—
Mississippi (type locality: Columbus).
- consueta Sargent in Rep. Missouri Bot. Gard. XIX. 67 (1908).—**Crus-**
galli.—Missouri (type locality: Swan).
- contigua Sargent in Rhodora, V. 115 (1903).—**Coccineae.**—Vermont
(type locality: Stockbridge).
- contortifolia Sargent in N. Y. State Mus. Bull. CV. Bot. IX. 59 (1906).—
Molles.—New York (type locality: North Albany).
- contorta Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 236 (1910).—
Intricatae.—Pennsylvania (type locality: Bedford).
- contrita Beadle in Biltmore Bot. Studies, I. 61 (1902).—**Pulcherrimae.**—
Florida, Georgia (type locality: River Junction, Fla.).
- corallina Hort. ex Persoon, Syn. II. 36 (1807), as a synonym of *C. cordata*
= *C. Phaenopyrum* Medic.
- cornellii Sargent in N. Y. State Mus. Bull. CXXII. 105 (1908).—**Intri-**
catae—New York (type locality: Coopers Plains).
- cornifolia (Poir.) Booth ex Loudon, Arb. Brit. II. 819 (1838), as synonym
of *C. pyrifolia* Ait. (?)
- coronata Hort. ex Lange, Rev. Gen. Crat. 33 (1897), as synonym of *C. intricata* Lange.
- corporea Sargent in Bot. Gaz. XXXV. 403 (1903).—**Macracanthae.**—
Illinois (type locality: Lake Zurich).
- corusca Sargent in Bot. Gaz. XXXIII. 117 (1902).—**Molles.**—Illinois
(type locality: Lake Zurich).
- Coursetiana* Roemer, Fam. Nat. Syn. III. 117 (1847) = *C. crus-galli* L.
- crassa Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 18 (1901).—
Flavae.—Eastern Pennsylvania (type locality not given). (?)
- crassifolia Sargent in Ontario Nat. Sci. Bull. IV. 60 (1908).—**Rotundi-**
foliae.—Ontario (type locality: London).
- Crawfordiana Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 169 (1910).—
Pruinosae.—Pennsylvania (type locality: Linesville).
- Craytonii Beadle in Biltmore Bot. Studies, I. 121 (1902).—**Intricatae.**—
North Carolina (type locality: Marshall). (?)
- cristata Ashe in Ann. Carnegie Mus. I. 392 (1902).—**Coccineae.**—
Pennsylvania, Ohio (type locality: Pittsburgh, Pa.).

CRATAEGUS

- crocata* Ashe in Ann. Carnegie Mus. I. 389 (1902) = *C. punctata* var. *aurea* Ait.
- crocea* Beadle in Biltmore Bot. Studies, I. 113 (1902).—**Flavae**.—Florida (type locality: Citra).
- crocina* Beadle in Biltmore Bot. Studies, I. 132 (1902).—**Crus-galli**.—Louisiana (type locality: Opelousas).
- Croomiana* Sargent in Jour. Arnold Arb. III. 202 (1922).—**Uniflorae**.—Florida (type locality: Tallahassee).
- cruda* Sargent in N. Y. State Mus. Bull. cxxii. 54 (1908).—**Silvicolae**.—New York (type locality: Niagara Falls).
- crudelis* Sargent in Rhodora, v. 143 (1903).—**Tenuifoliae**.—South-eastern Canada (type locality: Montmorency Falls, Que.).
- cruenta* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 16 (1901).—Group?—North Carolina (type locality: between Sprucepine and Micaville). (?)
- crus-galli* Linnaeus, Spec. 476 (1753).—**Crus-galli**.—Southeastern Canada and New England to North Carolina, west to Alabama, Tennessee, Illinois and Missouri (type region: "Virginia").
- crus-galli angustifolia* (Ehrh.) Sudworth in Bull. U. S. Dept. Agric. Div. For. xiv. 217 (1897) = *C. crus-galli* var. *pyracanthifolia* Ait.
- crus-galli* var. *arbutifolia* Hort. ex Nicholson in Hand-list Arb. Kew, pt. 1, 197 (1894), nomen nudum. (?)
- crus-galli* var. *attenuata* (Ashe) Farwell in Rep. Michigan Acad. Sci. xx. 181 (1918) = *C. attenuata*.
- crus-galli* var. *berberifolia* (Torr. & Gray) Sargent in Gard. & For. II. 124 (1889) = *C. berberifolia*.
- crus-galli* var. *capillata* Sargent in Bot. Gaz. xxxv. 100 (1903).—Delaware (type locality: Wilmington).
- crus-galli* var. *exigua* (Sarg.) Eggleston in Rhodora, x. 75 (1908).—Connecticut (type locality of *C. exigua* Sarg.: Waterford).
- crus-galli* var. *Fontanesiana* (Spach) Wenzig in Linnaea, xxxviii. 141 (1874) = *C. Fontanesiana*.
- crus-galli* ζ. *inermis* Lange, Rev. Gen. Crat. 62 (1897). (?)
- crus-galli* γ. *lancifolia* (Wend.) Lange, Rev. Gen. Crat. 62 (1897) = *C. crus-galli* var. *pyracanthifolia* Ait.
- crus-galli* β. *laurifolia* (Medic.) Lange, Rev. Gen. Crat. 61 (1897). (?)
- crus-galli* δ. *linearis* (Pers.) De Candolle, Prodr. II. 626 (1825) = *C. crus-galli* var. *pyracanthifolia* Ait.
- crus-galli* var. *nana* (Dum.) Loudon, Arb. Brit. II. 821 (1838). = *C. crus-galli* var. *pyracanthifolia* Ait.
- crus-galli* var. *oblongata* Sargent in Bot. Gaz. xxxv. 99 (1903).—Pennsylvania, Delaware (type locality: Edgemore, Del.).
- crus-galli* var. *ovalifolia* (Hornem.) Lindley, in Bot. Reg. xxii. t. 1860, (1836) = *C. prunifolia* Pers.

CRATAEGUS

- crus-galli prunellifolia* (Poir.) Sudworth, Bull. U. S. Dept. Agr. Div. For. xiv. 216 (1897). (?)
- crus-galli e. prunifolia* (Marsh.) London, Arb. Brit. II. 821, 849 (1838). = *C. prunifolia*.
- crus-galli* var. *pyracanthifolia* Aiton, Hort. Kew. II. 170 (1789).—Pennsylvania to Florida and west to Indiana and Tennessee.
- crus-galli* var. *rubens* Sargent in N. Y. State Mus. Bull. CLXVII. 75 (1913).—New York (type locality, Chapinville).
- crus-galli* var. *salicifolia* (Medic.) Aiton, Hort. Kew II. 170 (1789) = *C. crus-galli* var. *pyracanthifolia* Ait. ?
- crus-galli* var. *splendens* Aiton, Hort. Kew. ed. 2, III. 202 (1811). (?)
- crus-galli* × *macracantha* Schneider, Ill. Handb. Laubh. II. 1007 (1912). (?)
- crus-galli* × *pentagyna* Schneider, Ill. Handb. Laubh. I. 778 (1906). (?)
- crus-galli* × *prunifolia* Koehne, Deutsche Dendr. 233 (1893). (?)
- crus-galli* × *punctata* Koehne, Deutsche Dendr. 233 (1893). (?)
- crus-galli* × *succulenta* Eggleston in N. Y. State Mus. Bull. CCXLIII-IV. 63 (1923). (?)
- crux* Ashe in Jour. Elisha Mitchell Sci. Soc. XVIII. 17 (1902).—**Macracthæ**.—North Carolina (type locality: along New River, Ashe County).
- cullasagensis* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 10 (1900).—**Flavæ**.—North Carolina (type locality: along Cullasagee River, Macon County). (?)
- culta* Sargent in Rhodora, VII. 199 (1905).—**Tenuifoliae**.—Massachusetts (type locality: Shirley).
- cuneifolia* (Ehrh.) Borkhausen in Roemer, Arch. Bot. I. no. 3, 86 (1798). Group?—Range and type region of *Mespilus cuneifolia* Ehrh.: “Nordamerika.” (?)
- cuneiformis* (Marsh.) Eggleston in Britton & Brown, Ill. Fl. ed. 2, II. 299 (1913).—**Punctatae**.—“Western New York and Pennsylvania to southwest Virginia, west to central Illinois” (type locality of *Mespilus cuneiformis* Marsh. not given). (?)
- cuprea* Sargent in Bot. Gaz. XXXV. 105 (1903).—**Intricatae**.—Delaware (type locality: Wilmington).
- cupulifera* Sargent in Proc. Rochester Acad. Sci. IV. 129 (1903).—**Rotundifoliae**.—New York (type locality: Rochester).
- curabilis* Ashe in Bull. Charleston Mus. XII. 43 (1916).—Group?—South Carolina (type locality: “lower edge of the Piedmont”). (?)
- Curtisii* Ashe in Bull. N. Car. Exper. Sta. CLXXV. 109 (1900).—Group?—North Carolina (type locality: Person County). (?)
- curva* Beadle in Biltmore Bot. Studies, I. 109 (1902).—**Flavæ**.—Florida (type locality: Jacksonville).
- Cuthbertii* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 15 (1900).—Group?—North Carolina (type locality: Bladen County). (?)

CRATAEGUS

- cyanophylla* Sargent in Bot. Gaz. xxxv. 387 (1903).—**Tenuifoliae.**—Illinois, Wisconsin (type locality: Glendon Park, near Chicago, Ill.).
- cyclophylla* Sargent in Rhodora, vii. 210 (1905).—**Anomalae.**—Vermont (type locality: New Haven).
- Cydonia* Gruber in Bull. Torrey Bot. Club, xxxii. 390 (1905).—**Punctatae.**—Pennsylvania (type locality: Kutztown). (?)
- dacrioides* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 224 (1910).—**Rotundifoliae.**—Pennsylvania (type locality: Orbisonia).
- dallasiana* Sargent, Trees and Shrubs, I. 59 (1903).—**Punctatae.**—Texas (type locality: Dallas).
- Damei* Sargent in Rhodora, vii. 200 (1905).—**Tenuifoliae.**—Massachusetts (type locality: Lowell).
- dapsilis* Beadle in Biltmore Bot. Studies, I. 89 (1902).—**Flavae.**—Florida (type locality: Lane Park, near Tavares).
- Darlingtoniana* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 653 (1905).—**Intricatae.**—Pennsylvania (type locality: Serpentine Ridge, near West Chester).
- dasiphylla* Sargent in Rep. Missouri Bot. Gard. xxii. 80 (1912).—**Molles.**—Missouri, Kansas, Oklahoma (type locality: Joplin, Mo.).
- Davisii* Sargent in Rep. Missouri Bot. Gard. xxii. 77 (1912) = *C. viridis* L.
- Dawsoniana* Sargent in Rep. Missouri Bot. Gard. xix. 88 (1908).—**Virides.**—Illinois, Missouri (type locality: Arnold Arboretum, cultivated from seed collected near St. Louis, Mo.).
- Dayana* Sargent in N. Y. State Mus. Bull. cxxii. 66 (1908).—**Coccineae.**—New York (type locality: Buffalo).
- debilis* Sargent in Ontario Nat. Sci. Bull. IV. 86 (1908).—**Macracantha.**—Ontario (type locality: London).
- decens* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 19 (1903).—**Tenuifoliae.**—Ohio (type locality: Mansfield). (?)
- declivitatis* Sargent in Rep. Missouri Bot. Gard. xix. 103 (1908).—**Molles.**—Illinois, Missouri (type locality: St. Louis, Mo.).
- decora* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 17 (1902).—Group?—North Carolina (type locality: Elk Cross Roads). (?)
- decorata* Sargent in Rep. Missouri Bot. Gard. xix. 94 (1908).—**Prunifoliae.**—Missouri (type locality: Swan).
- deducta* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 602 (1905).—**Pruinosae.**—Pennsylvania (type locality: Sellersville).
- definita* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 647 (1905).—**Intricatae.**—Pennsylvania (type locality: Newtown).
- delawareensis* Sargent in Bot. Gaz. xxxv. 102 (1903).—**Pruinosae.**—Delaware (type locality: between Newport and Newcastle).

CRATAEGUS

- delecta* Sargent, Man. Trees N. Am. 451 (1905).—**Coccineae.**—Illinois (type locality: Lockport).
- delectabilis* Sargent in Ontario Nat. Sci. Bull. IV. 90 (1908).—**Macracanthae.**—Ontario (type locality: Toronto).
- delectata* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 200 (1910).—**Silvicolae.**—Pennsylvania (type locality: Keyser Valley, near Scranton).
- delicata* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 606 (1905).—**Silvicolae.**—Pennsylvania (type locality: Sellersville).
- Delosii* Sargent in Ontario Nat. Sci. Bull. IV. 56 (1908).—**Intricatae.**—Ontario (type locality: Toronto).
- deltoides* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 19 (1901).—**Pruinosae.**—New York, Pennsylvania, Maryland (type locality not given).
- delucida* Sargent in Rhodora, v. 139 (1903).—**Tenuifoliae.**—New England, New York (type locality: East Rutland, Vt.).
- demissa* Sargent in Rhodora, v. 139 (1903).—**Tenuifoliae.**—New England to New York and Ohio (type locality: Lenox, Mass.).
- denaria* Beadle in Biltmore Bot. Studies, I. 131 (1902).—**Crus-galli.**—Mississippi (type locality: Columbus).
- densiflora* Sargent in Rhodora, III. 75 (1901).—**Coccineae.**—Quebec (type locality: Rockfield).
- dentata* Loddiges, Cat. 45 (1836), nomen nudum.—Loudon, Arb. Brit. II. 849 (1838), as a synonym of *C. Crus-galli* var. *ovalifolia* Lindl.
- denudata* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 173 (1910).—**Pruinosae.**—Pennsylvania (type locality: Schenley Park, Pittsburgh).
- depilis* Sargent, Man. Trees N. Am. 419 (1905).—**Tenuifoliae.**—Illinois (type locality: Glendon Park, near Chicago.)
- depressa* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 8 (1901).—**Pruinosae.**—Northern Missouri (type locality not given).
- desertorum* Sargent in Jour. Arnold Arb. III. 187 (1922).—**Virides.**—Texas (type locality: Uvalde).
- desueta* Sargent in N. Y. State Mus. Bull. CXXII. 84 (1908).—**Punctatae.**—New York (type locality: Coopers Plains).
- Deweyana* Sargent in Proc. Rochester Acad. Sci. IV. 133 (1903).—**Macracanthae.**—New York (type locality: Rochester).
- Dewingii* Sargent in N. Y. State Mus. Bull. CXXII. 34 (1908).—**Punctatae.**—New York (type locality: Belfast).
- diaphora* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 251 (1910).—**Macracanthae.**—Pennsylvania (type locality: Keyser Valley, near Scranton).
- diffusa* Sargent in Proc. Rochester Acad. Sci. IV. 103 (1903).—**Silvicolae.**—New England, New York (type locality: Rochester, N. Y.).

CRATAEGUS

- digna Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 628 (1905).—**Molles.**—Pennsylvania (type locality: Kutztown).
- dilatata Sargent in Bot. Gaz. XXXI. 9 (1901).—**Dilatatae.**—Quebec and New England to Ohio and Kentucky (type locality: eastern Massachusetts).
- discolor Sargent in Rep. Missouri Bot. Gard. XIX. 44 (1908).—**Crus-galli.**—Missouri, Arkansas (type locality: Grandin, Mo.).
- disjuncta Sargent, Trees and Shrubs, I. 109 (1903).—**Pruinosae.**—Missouri, Arkansas (type locality: Monteer, Mo.).
- dispar Beadle in Biltmore Bot. Studies, I. 28 (1901).—**Flavae.**—South Carolina, Georgia (type locality: Aiken, S. Car.).
- disperma Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 14 (1900).—Group?—New York and Pennsylvania, west to Illinois (type locality: Wilkesbarre Mountain, Pa.). (?)
- dispessa Ashe in Jour. Elisha Mitchell Sci. Soc. XIX. 17 (1903).—**Molles.**—Missouri (type locality of *C. pyriformis* Britton, which this name replaces: Monteer).
- dissimilis Sargent in Rhodora, v. 149 (1903).—**Tenuifoliae.**—Connecticut (type locality: East Lyme).
- dissociabilis Sargent in N. Y. State Mus. Bull. CXXII. 95 (1908).—**Silvicolae.**—New York (type locality: Coopers Plains).
- dissona Sargent in Rhodora, v. 60 (1903).—**Silvicolae.**—Ontario and New England, west to northern Illinois and Wisconsin (type locality: Great Barrington, Mass.).
- divergens (Peck) Sargent in N. Y. State Mus. Bull. CV. Bot. IX. 66 (1906).—**Rotundifoliae.**—New York (type locality: North Greenbush).
- diversa Sargent in N. Y. State Mus. Bull. CXXII. 109 (1908).—**Macracanthae.**—New York (type locality: Coopers Plains).
- diversifolia Sargent in Jour. Arnold Arb. VI. 4 (1925).—**Intricatae.**—Michigan (type locality of *C. Wheeleri* Sarg. which this name replaces: Grand Rapids).
- divida Sargent in Bot. Gaz. XXXV. 401 (1903).—**Macracanthae.**—Illinois (type locality: Barrington).
- divisifolia Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 179 (1910).—**Pruinosae.**—Pennsylvania, Ohio (type locality: Kutztown, Pa.).
- Doddsii Ramaley in Bot. Gaz. XLVI. 381 (1908).—Colorado (type locality: Pole Canyon, near Boulder). (?)
- Dodgei Ashe in Jour. Elisha Mitchell Sci. Soc. XIX. 26 (1903).—**Rotundifoliae.**—Southeastern Canada and New England, west to Michigan, Wisconsin and Indiana (type locality: Port Huron, Mich.).
- Dodgei var. lumaria (Ashe) Sargent in Rep. Geol. Surv. Mich. 1906, 556 (1907).—Michigan (type locality: Port Huron).

CRATAEGUS

- dolosa Beadle in Biltmore Bot. Studies, I. 90 (1902).—**Flavae.**—Alabama (type locality: Abbeville).
- Douglasii Lindley in Bot. Reg. XXI. t. 1810 (1835).—**Douglasianae.**—British Columbia to California, east to Idaho, Montana and Wyoming (type locality: Garden of London Horticultural Society, from seed collected along lower Colorado River).
- Douglasii f. badia Sargent in Bot. Gaz. XLIV. 65 (1907).—British Columbia, Washington (type locality: Union Flat, near Pullman, Wash.).
- Douglasii* var. *rivularis* (Nutt.) Sargent in Gard. & For. II. 401 (1889) = *C. rivularis*.
- Douglasii var. *Suksdorffii* Sargent in Bot. Gaz. XLIV. 65 (1907).—Washington (type locality: western Klickitat County).
- Downingii* Hort. ex Nicholson in Hand-list Arb. Kew, pt. I. 209 (1894), as synonym of *C. tomentosa* (?).
- drymophila* Sargent in Jour. Arnold Arb. I. (1920), as “*drymopila*”.—**Silvicolae.**—Georgia, Alabama, Mississippi (type locality of *C. silvicola* Beadle, which this name replaces: Gadsden, Ala.).
- dumetosa* Sargent in Rep. Missouri Bot. Gard. XIX. 109 (1908).—**Molles.**—Missouri (type locality: Neck City).
- dumicola* Sargent in Rhodora, v. 183 (1903).—**Macracantheae.**—Quebec, New England (type locality: Fort Fairfield, Me.).
- Dunbari* Sargent in Proc. Rochester Acad. Sci. IV. 126 (1903).—**Anomalae.**—New York (type locality: Rochester).
- dunmorensis* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 174 (1910).—**Pruinosae.**—Pennsylvania (type locality: Dunmore).
- duracina* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 186 (1910).—**Pruinosae.**—Pennsylvania (type locality: Whisky Hollow, near Kittanning).
- durifolia* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 12 (1900).—**Virides.**—Illinois, Missouri (type locality “Missouri”). (?)
- durobrivensis* Sargent, Trees and Shrubs, I. 3 (1902).—**Dilatatae.**—New York (type locality: Rochester).
- Eamesii* Sargent, Manual Trees N. Am. 454 (1905).—**Coccineae.**—Connecticut (type locality: Stratford).
- Earlei* Ashe in Bull. N. Car. Exper. Sta. CLXXV. 112 (1900).—**Uniflorae.**—Florida, Alabama (type locality: Cullman County, Ala.).
- Eastmaniana* Sargent in N. Y. State Mus. Bull. CLXVII. 77 (1913).—**Punctatae.**—New York (type locality: Durand-Eastman Park, Rochester).
- Eatoniana* Sargent in N. Y. State Mus. Bull. 105, Bot. IX. 51 (1906).—**Punctatae.**—New York (type locality: Menands golf grounds, near Albany).
- ebracteata* Ashe in Bull. N. Car. Exper. Sta. CLXXV. 109 (1900).—Group?—North Carolina (type locality: Winston). (?)

CRATAEGUS

- eburnea* Ashe in Ann. Carnegie Mus. I. 395 (1902).—**Crus-galli.**—Pennsylvania and Virginia to Ohio and Tennessee (type locality not given).
- edita* Sargent in Bot. Gaz. XXXIII. 110 (1902).—**Crus-galli.**—Louisiana, Texas (type locality: Marshall, Tex.).
- Edsoni* Sargent in Rhodora, VII. 205 (1905).—**Tenuifoliae.**—New England, New York (type locality: Burlington, Vt.).
- edulis* Loddiges, Cat. 39 (1826), nomen nudum.—Loudon, Arb. Brit. II. 818 (1838), as synonym of *C. punctata* var. *aurea* Ait.
- edura* Beadle in Biltmore Bot. Studies, I. 128 (1902).—**Crus-galli.**—Louisiana (type locality: Opelousas).
- edurescens* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 180 (1910).—**Pruinosae.**—Pennsylvania (type locality: Edgemont Township, Delaware County).
- effera* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 206 (1910).—**Silvicola.**—Pennsylvania (type locality: Scranton).
- efferata* Sargent in N. Y. State Mus. Bull. CXXII. 128 (1908).—**Macrancanthea.**—New York (type locality: Hemlock Lake).
- efferta* Sargent in Rep. Missouri Bot. Gard. XIX. 51 (1908).—**Crus-galli.**—Missouri (type locality, Pacific).
- effulgens* Sargent in Rep. Missouri Bot. Gard. XXII. 71 (1912).—**Crus-galli.**—Illinois, Tennessee, Missouri (type locality: Springfield, Mo.).
- Eganii* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 15 (1900).—**Tenuifoliae.**—Illinois (type locality “northern Illinois”).
- egens* Beadle in Biltmore Bot. Studies, I. 85 (1902).—**Flavae.**—Florida (type locality: Bristol).
- Eggertii* Britton in Bull. N. Y. Bot. Gard. I. 447 (1900) = *C. coccinoides* Ashe.
- Egglestonii* Sargent in Rhodora, III. 30 (1901).—**Anomalae.**—New Hampshire, Vermont, Massachusetts (type locality: Bald Mountain, Shrewsbury, Vt.).
- egregia* Beadle in Biltmore Bot. Studies, I. 82 (1902).—**Flavae.**—Florida (type locality: Bristol).
- elliptica* Aiton, Hort. Kew, II. 168 (1789).*—**Flavae.** (?)
- elliptica* Elliot, Sketch, I. 549 (1821) = *C. aestivalis* T. & G.
- ellipticifolia* Sargent in Jour. Arnold Arb. III. 194 (1923).—**Pruinosae.**—Ohio (type locality: Delaware).
- Ellwangeriana* Sargent in Bot. Gaz. XXXIII. 118 (1902).—**Molles.**—Ontario, New York, Pennsylvania, Michigan (type locality: Rochester, N. Y.).

* I have seen only cultivated specimens of this plant, in which the principal distinction from *C. flava* appears to be in its 8–10 stamens and pale yellow anthers. Since the origin of the type cultivated in England is not definitely known the geographical range can not be given.

CRATAEGUS

- elongata* Sargent in Bot. Gaz. xxxv. 380 (1903).—**Coccineae.**—Illinois, Wisconsin (type locality: Deerfield, Ill.).
- Emersoniana* Sargent in Rhodora, vii. 181 (1905).—**Macracantheae.**—Massachusetts (type locality: Somerset).
- Engelmannii* Sargent in Bot. Gaz. xxxi. 2 (1901).—**Crus-galli.**—Southern Illinois and Kentucky to Missouri and Arkansas (type locality: Allenton, Mo.).
- ensifera* Sargent in Rep. Missouri Bot. Gard. xix. 124 (1908).—**Macracantheae.**—Missouri (type locality: Swan).
- enucleata* Sargent, Trees and Shrubs, ii. 239 (1913).—**Virides.**—Arkansas (type locality: Fulton).
- erecta* Sargent in Bot. Gaz. xxxi. 218 (1901).—**Crus-galli.**—Indiana and Illinois to Missouri and Arkansas (type locality: East St. Louis, Ill.).
- errata* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 245 (1910).—**Anomalae.**—Pennsylvania (type locality: Keyser Valley, near Scranton).
- erubescens* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 178 (1910).—**Pruinosae.**—Pennsylvania (type locality: Kittanning).
- erythrocarpa* Ashe in Bull. N. Car. Exper. Sta. clxxv. 109 (1900).—Group?—North Carolina (type locality: Raleigh). (?)
- erythropoda* Ashe in Bull. N. Car. Exper. Sta. clxxv. 113 (1900).—**Douglasianae.**—Colorado to Washington (type locality: foot-hills of Cache la Poudre Mountains, northern Colo.).
- Evansiana* Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 633 (1905).—**Rotundifoliae.**—Pennsylvania (type locality: West Fairmount Park, Philadelphia).
- exclusa* Sargent in Rhodora, v. 108 (1903).—**Molles.**—Vermont, New York, Ohio (type locality: Bald Mountain, Clarendon and Shrewsbury, Vt.).
- exigua* Sargent in Rhodora, v. 52 (1903) = *C. crus-galli* var. *exigua*.
- exigua* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 20 (1903).—**Tenuifoliae.**—Wisconsin (type locality: Milwaukee). (?)
- exilis* Beadle in Biltmore Bot. Studies, i. 76 (1902).—**Flavae.**—Georgia (type locality: Albany).
- eximia* Beadle in Biltmore Bot. Studies, i. 62 (1902).—**Intricatae.**—Alabama (type locality: Sand Mountain, near Pisgah).
- exornata* Sargent in Ontario Nat. Sci. Bull. iv. 31 (1908).—**Pruinosae.**—Ontario, New York (type locality: Chippewa, Ont.).
- extraaria* Beadle in Biltmore Bot. Studies, i. 73 (1902).—**Flavae.**—Georgia (type locality: Marietta).
- fallax* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 11 (1903).—**Macracantheae.**—Michigan (type locality: Port Huron). (?)
- fallsiana* Sargent in N. Y. State Mus. Bull. clxvii. 113 (1913).—**Anomalae.**—New York (type locality: Little Falls).

CRATAEGUS

- Farwellii* Sargent in Rep. Geol. Surv. Mich. 1906, 519 (1907).—**Crus-galli**.—Michigan (type locality: Bell Island, Detroit).
- fastosa* Sargent, Trees and Shrubs, I. 61 (1903).—**Punctatae**.—Arkansas (type locality: Fulton).
- Faxonii* Sargent in Rhodora, v. 161 (1903).—**Rotundifoliae**.—New Hampshire (type locality: Franconia).
- fecunda* Sargent in Bot. Gaz. xxxiii. 111 (1902).—**Crus-galli**.—Southern Illinois, Missouri (type locality: Arnold Arboretum from seed collected near Allenton, Mo.).
- felix* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 589 (1905).—**Silvicolae**.—Pennsylvania (type locality: Reading).
- fera* Beadle in Biltmore Bot. Studies, I. 128 (1902).—**Crus-galli**.—Louisiana (type locality: Opelousas).
- ferentaria* Sargent in Proc. Rochester Acad. Sci. IV. 135 (1903).—**Macracanthae**.—Southeastern Canada, New England, New York (type locality: Rochester, N. Y.).
- Fernaldii* Sargent in Rhodora, v. 166 (1903).—**Rotundifoliae**.—Maine (type locality: Fort Fairfield and Fort Kent, Aroostook County).
- ferox* Sargent in Rep. Missouri Bot. Gard. xix. 52 (1908) = *C. arduennae* Sarg.
- Ferrissii* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 11 (1901).—**Pruinosae**.—Illinois (type locality: "northern Illinois").
- ferta* Sargent in Ontario Nat. Sci. Bull. IV. 77 (1908).—**Macracanthae**.—Ontario (type locality: Toronto).
- fertilis* Sargent in Rhodora, v. 182 (1903).—**Macracanthae**.—Southeastern Canada, New England (type locality: Orono, Me.).
- festiva* Sargent in Rhodora, v. 54 (1903).—**Pruinosae**.—Connecticut (type locality: East Lyme).
- filipes* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 18 (1903).—**Silvicolae**.—Michigan (type locality: Port Huron).
- finitima* Sargent in N. Y. State Mus. Bull. cxxii. 78 (1908).—**Macracanthae**.—New York (type locality: Niagara Falls).
- firma* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 612 (1905).—**Tenuifoliae**.—Pennsylvania (type locality: Sellersburg).
- flabellata* (Bosc) K. Koch in Verh. Ver. Beförd. Gartenb. Preuss. ser. 2, I. 240 (Weissd. 20) (1853).—**Tenuifoliae**.—Southeastern Canada, New England and New York, west to northern Illinois (type locality of *Mespilus flabellata* Bosc: cultivated in Jardin des Plantes, Paris, from unknown source).
- flagrans* Sargent in N. Y. State Mus. Bull. cv. Bot. IX. 71 (1906).—**Macracanthae**.—New York (type locality: North Greenbush).
- flammata* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 211 (1910).—**Tenuifoliae**.—Pennsylvania (type locality: Birdseye, near Scranton).

CRATAEGUS

- flammea* Sargent in Rep. Geol. Surv. Mich. 1906, 563 (1907).—**Macracanthae.**—Ontario, Michigan (type locality: Sarnia, Ont.).
- flava* Aiton, Hort. Kew. II. 169 (1789).—**Flavae.**—Georgia, Florida (type locality: cultivated in Royal Botanical Gardens, Kew. “Native of North America”).
- flava* var. *elliptica* (Ait.) Sargent, Silva N. Am. IV. 114 (1892) = *C. elliptica*.
- flava* var. *integra* Nash in Bull. Torrey Bot. Club. XXII. 150 (1895) = *C. integra*.
- flava* var. *lobata* (Bosc) Lindley in Bot. Reg. XXIII. t. 1932 (1837) = *C. elliptica* Ait.
- flava* var. *pubescens* Gray, Man. ed. 2, 124 (1856). (?)
- flavens* Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 849 (1838), as synonym of *C. punctata flava*.
- flavida* Sargent in Rep. Geol. Surv. Mich. 1906, 554 (1907).—**Intricatae.**—Ontario, Michigan (type locality: London, Ont.).
- flavo-carnis* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 9 (1900).—Group?—North Carolina (type locality: Salisbury). (?)
- flexispina* (Moench) Sargent in Gard. & For. II. 424 (1889) = *C. flava* Ait.
- flexispina* var. *pubescens* (Gray) Millspaugh, Fl. W. Va. 360 (1892). (?)
- flexuosa* (Poir.) DeCandolle, Prodr. II. 627 (1825).—**Uniflorae.**—(Type region: “Carolina”). (?)
- floreæ* Sargent in Rhodora, v. 145 (1903).—**Tenuifoliae.**—Nova Scotia, Maine (type locality: Orono, Me.).
- florens* Beadle in Biltmore Bot. Studies, I. 94 (1902).—**Flavae.**—Georgia, Mississippi, Louisiana (type locality: Columbus, Miss.).
- florida* Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 841 (1838), as synonym of *C. parvifolia* Ait. = *C. uniflora* Muench.
- floridana* Sargent in Bot. Gaz. XXXIII. 124 (1902).—**Flavae.**—Florida (type locality: Jacksonville).
- floridula* Sargent in N. Y. State Mus. Bull. CXXII. 126 (1908).—**Anomalaæ.**—New York (type locality: Piseco).
- florifera* Sargent in Ontario Nat. Sci. Bull. IV. 14 (1908).—**Punctatae.**—Ontario (type locality: Toronto).
- fluvialis* Sargent in Rhodora, v. 117 (1903) (as “*fluvialis*”).—**Coccineæ.**—Quebec (type locality: Adirondack Junction).
- foetida* Ashe in Ann. Carnegie Mus. I. 389 (1902).—**Intricatae.**—Ontario, New England, New York, Pennsylvania (type locality: Berks County, Pa.).
- foliata* Sargent in N. Y. State Mus. Bull. CXXII. 53 (1908).—**Silvicolæ.**—New York (type locality: Niagara Falls).

CRATAEGUS

- Fontanesiana* (Spach) Steudel, Nom. Bot. ed. 2, 1. 432 (1840).—**Crus-galli**.—Pennsylvania. (Type locality of *Mespilus Fontanesiana* Spach not given, but species based upon plants cultivated in Europe.)
- Fontanesiana* β. *angustifolia* Lange, Rev. Gen. Crat. 63 (1897) = C. crus-galli var. *pyracanthifolia* Ait.?
- Fontanesiana* α. *latifolia* Lange, Rev. Gen. Crat. 63 (1897). (?)
- Fontanesiana* α. *latifolia* f. *chlorocarpa* Lange, 63 (1897). (?)
- Forbesae Sargent in Rhodora, v. 151 (1903).—**Tenuifoliae**.—Massachusetts, Connecticut (type locality: Worcester County, Mass.).
- formosa Sargent in Proc. Rochester Acad. Sci. iv. 101 (1903).—**Pruinosae**.—Ontario, New York, Ohio (type locality: Rochester, N. Y.).
- fortis Beadle in Biltmore Bot. Studies, 1. 92 (1902).—**Flavae**.—Mississippi (type locality: Columbus).
- fortunata Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 239 (1910).—**Intricatae**.—Pennsylvania (type locality: Charleroi).
- Fortunei* Hort. ex Sargent in Jour. Arnold Arb. vi. 1 (1925), as a synonym of C. *Fontanesiana* Steud.
- franklinensis Sargent in Jour. Arnold Arb. iv. 100 (1922).—**Pruinosae**.—Ohio (type locality: Columbus).
- fretalis Sargent in Rhodora, v. 112 (1903).—**Coccineae**.—Connecticut (type locality: Groton).
- Fretzii Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 604 (1905).—**Silvicolae**.—Pennsylvania (type locality: Durham).
- fructuosa Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 232 (1910).—**Intricatae**.—Pennsylvania (type locality: West Chester).
- frugalis Beadle in Biltmore Bot. Studies, 1. 100 (1902).—**Flavae**.—Georgia (type locality: Albany).
- frugiferens Beadle in Biltmore Bot. Studies, 1. 30 (1901).—**Flavae**.—Alabama (type locality: Cullman).
- frutescens Sargent in N. Y. State Mus. Bull. cxxii. 113 (1908).—**Macracantheae**.—New York (type locality: Coopers Plains).
- fruticosa Sargent, Trees and Shrubs, 1. 13 (1902).—**Uniflorae**.—Florida (type locality: Seville).
- fucata Sargent in N. Y. State Mus. Bull. cxxii. 99 (1908).—**Tenuifoliae**.—New York (type locality: Coopers Plains).
- fucosa Sargent in Rhodora, v. 137 (1903).—**Tenuifoliae**.—New Hampshire, Massachusetts (type locality: West Boylston, Mass.).
- fulgens Sargent in Rhodora, VII. 182 (1905).—**Macracantheae**.—Massachusetts, Connecticut (type locality: Stratford, Ct.).
- fulgida Sargent in Ontario Nat. Sci. Bull. iv. 84 (1908).—**Macracantheae**.—Ontario (type locality: Toronto).
- Fulleriana Sargent in Proc. Rochester Acad. Sci. iv. 111 (1903).—**Molles**.—New York (type locality: Rochester).

CRATAEGUS

- fulva* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 649 (1905).—**Intricatae**.—Pennsylvania (type locality: Forge Hill, Berks County).
- furcata* Sargent in Rep. Missouri Bot. Gard. XIX. 86 (1908) = *C. viridis* L.
- furtiva* Beadle in Biltmore Bot. Studies, I. 81 (1902).—**Flavae**.—Georgia (type locality: Albany).
- fusca* Sargent in Rhodora, VII. 193 (1905).—**Pruinosae**.—Vermont (type locality: Burlington).
- galbans* Beadle in Biltmore Bot. Studies, I. 74 (1902).—**Flavae**.—Florida (type locality: River Junction).
- Gattingeri* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 12 (1900).—**Pruinosae**.—Ohio Indiana, West Virginia, Illinois, Kentucky, Tennessee (type locality: Nashville, Tenn.).
- gaudens* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 170 (1910).—**Pruinosae**.—Pennsylvania (type locality: Riverview Park, Allegheny City).
- Gaultii* Sargent in Bot. Gaz. XXXV. 397 (1903).—**Macracantheae**.—Illinois (type locality: Milton Township, Du Page County).
- Gaylussacia* Heller in Bull. S. California Acad. Sci. II. 69 (1903).—**Douglasianae**.—California (type locality: Sebastopol, Sonoma County).
- gemmosa* Sargent in Bot. Gaz. XXXIII. 119 (1902).—**Macracantheae**.—Ontario, and New York, west to Ohio, Wisconsin and Illinois (type locality: Grand Rapids, Mich.).
- geneseensis* Sargent in N. Y. State Mus. Bull. CXXII. 27 (1908).—**Crus-galli**.—New York (type locality: Rochester).
- genialis* Sargent in Rhodora, V. 148 (1903).—**Tenuifoliae**.—Ontario, New England, New York (type locality: Stockbridge Bowl, Mass.).
- geniculata* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 16 (1900).—**Flavae**.—North Carolina (type locality: “middle North Carolina”). (?)
- georgiana* Sargent in Bot. Gaz. XXXIII. 113 (1902).—**Pruinosae**.—Georgia (type locality: Rome).
- georgica* Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 849 (1838), as synonym of *C. spathulata* Michx.
- Gilbertiana* Sargent in N. Y. State Mus. Bull. CLXVII. 101 (1913).—**Coccineae**.—New York (type locality: Warren).
- gilva* Beadle in Biltmore Bot. Studies, I. 60 (1902).—**Intricatae**.—Alabama (type locality: Albertville).
- glabrata* Sargent in Ontario Nat. Sci. Bull. IV. 85 (1908).—**Macracantheae**.—Ontario (type locality: Toronto).
- glabrifolia* Sargent in Rep. Missouri Bot. Gard. XXII. 74 (1912).—**Punctatae?**—Missouri (type locality: Grandin).

CRATAEGUS

glabriuscula Sargent in Bot. Gaz. XXXI. 235 (1901).—**Virides.**—Texas
(type locality: Dallas).

glandulifera Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb.
Brit. II. 849 (1838), as synonym of *C. pyrifolia* Ait.

glandulosa Moench, Bäume Weiss. 31 (1785) = *C. macracantha* Lodd.?
glandulosa β . *Douglasii* (Lindl.) K. Koch, Hort. Dendr. 167 (1853) = *C.*

Douglasii.

glandulosa e. *elliptica* Lauche, Deutsche Dendr. 573 (1883). (?)

glandulosa β . *macracantha* (Lodd.) Lindley in Bot. Reg. XXII. t. 1912
(1836) = *C. macracantha*.

glandulosa c. *odorata* (Wendl.) Lauche, Deutsche Dendr. 573 (1883). (?)

glandulosa γ . *purpurea* (Bosc) K. Koch, Hort. Dendr. 167 (1853). (?)

glandulosa δ . *rotundifolia* (Moench) K. Koch, Hort. Dendr. 167 (1853) =
C. rotundifolia.

glandulosa ϵ . *succulenta* (Schrad.) K. Koch, Hort. Dendr. 167 (1853) =
C. succulenta.

glandulosa α . *typica* Regel in Act. Hort. Petrop. I. 120 (1871) = *C.*
Bicknellii Eggleston.

glareosa Ashe in Jour. Elisha Mitchell Sci. Soc. XVIII. 21 (1902).—
Pruinosae?—Michigan (type locality: Port Huron).

glaucophylla Sargent in Rhodora, v. 140 (1903).—**Tenuifoliae.**—
Ontario, New England, New York, Pennsylvania and Michigan
(type locality: Great Barrington, Mass.).

globosa Sargent in Rep. Missouri Bot. Gard. XIX. 118 (1908).—**Macra-**
canthae.—Missouri (type locality: Monteer).

gloriosa Sargent in N. Y. State Mus. Bull. CXXII. 70 (1908) = *C. pedi-*
cellata var. *gloriosa* Sarg.

gracilipes Sargent in N. Y. State Mus. Bull. CXXII. 119 (1908).—**Tenui-**
foliae.—New York (type locality: Hemlock Lake, Livingston
County).

gracilis Sargent in N. Y. State Mus. Bull. CXXII. 37 (1908).—**Prui-**
nosae.—New York, Ohio, Indiana (type locality: Niagara Falls,
N. Y.).

grandis Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 9 (1901).—
Punctatae.—Illinois (type locality not given). (?)

gratiosa Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 243 (1910).—
Intricatae.—Pennsylvania (type locality: Bedford Springs).

Gravesii Sargent in Rhodora, v. 159 (1903) = *C. Dodgei* Ashe.

gravida Beadle in Biltmore Bot. Studies, I. 119 (1902).—**Molles.**—
Tennessee (type locality: Nashville).

gravis Ashe in Jour. Elisha Mitchell Sci. Soc. XX. 49 (1904).—**Silvi-**
colae.—Ontario, Michigan, Ohio (type locality: Port Huron, Mich.).

Grayana Eggleston in Rhodora, x. 80 (1908) = *C. flabellata* K. Koch.

CRATAEGUS

- gregalis Beadle in Biltmore Bot. Studies, I. 118 (1902).—**Uniflorae.**—North Carolina, South Carolina (type locality: Biltmore, N. Car.).
- Greggiana Eggleston in Bull. Torrey Bot. Club, xxxvi. 511 (1909).—**Molles.**—Mexico (type locality: Saltillo, Coahuila).
- grossa Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 223 (1910).—**Rotundifoliae.**—Pennsylvania (type locality: Bedford Springs).
- grossiserrata Ashe in Bull. N. Car. Exper. Sta. CLXXV. 112 (1900).—**Uniflorae.**—Florida (type locality not given). (?)
- grossulariaefolia Loddiges, Cat. 39 (1826), nomen nudum.—Loudon, Arb. Brit. II. 849 (1838) as synonym of *C. crus-galli* var. *prunifolia*.
- Gruberi Ashe in Ann. Carnegie Mus. I. 388 (1902).—**Tenuifoliae.**—Pennsylvania (type locality: Berks County).
- Habereri Sargent in N. Y. State Mus. Bull. cxvi. Bot. x. 21 (1907).—**Tenuifoliae.**—New York (type locality: New Hartford).
- habilis Ashe in Bot. Gaz. XXXV. 435 (1903).—**Coccineae.**—Ohio (type locality: Mansfield). (?)
- Hadleyana Sargent in N. Y. State Mus. Bull. CLXVII. 93 (1913).—**Tenuifoliae.**—Ontario, New York (type locality: Herkimer).
- haemacarpa Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 8 (1900).—**Pruinosae.**—North Carolina (type locality: banks of Cullassagee River, Macon County).
- Halliana Sargent in N. Y. State Mus. Bull. cv. Bot. IX. 73 (1906).—**Macracanthae.**—New York (type locality: North Albany).
- hamata Sargent in Rep. Missouri Bot. Gard. xix. 46 (1908).—**Crus-galli.**—Missouri (type locality: Pacific).
- Handya Sargent in Rhodora, VII. 177 (1905).—**Macrospermae.**—Massachusetts (type locality: Somerset).
- Harbisonii Beadle in Bot. Gaz. xxviii. 413 (1899).—**Bracteatae.**—Tennessee (type locality: Nashville).
- Hargeri Sargent in Rhodora, v. 66 (1903).—**Intricatae.**—Connecticut (type locality: Oxford).
- Harryi Sargent in N. Y. State Mus. Bull. cxxii. 124 (1908).—**Rotundifoliae.**—New York (type locality: Richmond).
- Harveyana Sargent in Jour. Arnold Arb. III. 200 (1923).—**Intricatae.**—Arkansas (type locality: Eureka Springs).
- heidelbergensis Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 215 (1910).—**Tenuifoliae.**—Pennsylvania (type locality: Kutztown).
- helderbergensis Sargent in N. Y. State Mus. Bull. cv. Bot. IX. 49 (1906).—**Crus-galli.**—New York (type locality: Thompsons Lake, Albany County).
- Helena Sargent in Rhodora, VII. 162 (1905).—**Macracanthae.**—Massachusetts (type locality: Southboro).
- helvina Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 28 (1903).—**Crus-galli.**—Tennessee (type locality: Clarkton). (?)

CRATAEGUS

- Hillii Sargent in Bot. Gaz. xxxv. 384 (1903).—**Coccineae.**—Ohio, Indiana, Illinois (type locality: Thatchers Park, Cook County, Ill.).
- hirtella Sargent in Rep. Missouri Bot. Gard. xix. 62 (1908).—**Crus-galli.**—Missouri (type locality: Swan).
- hirtiflora Sargent in Rep. Missouri Bot. Gard. xix. 82 (1908), as “herti-flora.”—**Punctatae.**—Missouri (type locality: Swan).
- hispidula Sargent in Rep. Missouri Bot. Gard. xix. 116 (1908).—**Macracanthae.**—Illinois, Missouri, Arkansas, Texas (type locality: Carthage, Mo.).
- Holmesiana Ashe in Jour. Elisha Mitchell Sci. Soc. xvi. 78 (1900).—**Coccineae.**—Southeastern Canada and New England to New York and Pennsylvania (type locality: “central New York”).
- Holmesiana var. tardipes Sargent in Jour. Arnold Arb. i. 254 (1920).—Southeastern Canada, New York, Ohio (type locality of *C. tardipes* Sarg.: Toronto, Ont.).
- Holmesiana var. villipes Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 11 (1901).—Pennsylvania, New York (type locality not given).
- honeoyensis Sargent in N. Y. State Mus. Bull. cxxii. 129 (1908).—**Macracanthae.**—New York (type locality: west of Honeoye Lake, Ontario County).
- honestae Sargent in Rep. Geol. Surv. Mich. 1906, 558 (1907).—**Anomalae.**—Michigan (type locality: Grand Rapids).
- horrida Medicus, Gesch. Bot. 84 (1793).—**Crus-galli.** (?)
- horridula Sargent in Rep. Geol. Surv. Mich. 1906, 526 (1907).—**Pruinosae.**—Michigan, Ohio (type locality: Grand Rapids, Mich.).
- Howeana Sargent in N. Y. State Mus. Bull. cv. Bot. ix. 52 (1906).—**Pruinosae.**—New York, Ohio (type locality: Menands, near Albany, N. Y.).
- Howellii Heiser in Am. Midland Naturalist, v. 238 (1919) = *C. columbi-ana Howell.*
- hudsonica Sargent, Man. Trees N. Am. 457 (1905).—**Dilatatae.**—New York (type locality: Albany).
- Huntiana Sargent in N. Y. State Mus. Bull. clxvii. 105 (1913).—**Molles.**—New York (type locality: between Jordanville and Mud Lake, Herkimer County).
- hystricina Ashe in Bot. Gaz. xxxv. 433 (1903).—**Macracanthae.**—Connecticut, New York (type locality: along Housatonic River, southern Connecticut).
- hybrida Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. ii. 849 (1838), as synonym of *C. crus-galli prunifolia*. (?)
- hypolasia K. Koch, Hort. Dendr. 167 (1853) = *C. mexicana* Moç. & Sessé.

CRATAEGUS

- Ideae Sargent in *Rhodora*, vii. 211 (1905).—**Anomalae.**—Vermont (type locality: Concord).
- ignava* Beadle in *Biltmore Bot. Studies*, i. 31 (1901).—**Flavae.**—Alabama (type locality: Valley Head).
- ignava* Sargent in *Proc. Acad. Nat. Sci. Phila.* LXII. 228 (1910) = *C. neosmithii* Sarg.
- ignea* Sargent in *N. Y. State Mus. Bull.* CXXII. 96 (1908).—**Tenuifoliae.**—New York, Indiana (type locality: Coopers Plains, N. Y.).
- illecebrosa* Sargent in *Jour. Arnold Arb.* iv. 103 (1923).—**Coccineae.**—Ontario (type locality: Kingston).
- illinoiensis* Ashe in *Jour. Elisha Mitchell Sci. Soc.* XVI. 76 (1900).—**Macracanthae.**—Illinois (type locality: Wady Petra).
- illudens* Beadle in *Biltmore Bot. Studies*, i. 111 (1902).—**Flavae.**—Florida (type locality: Citra).
- illuminata* Sargent in *N. Y. State Mus. Bull.* CV. Bot. ix. 65 (1906).—**Rotundifoliae.**—New York (type locality: North Albany).
- illustris* Beadle in *Biltmore Bot. Studies*, i. 68 (1902).—**Pulcherrimae.**—Mississippi (type locality: Meridian).
- immanis* Ashe in *Jour. Elisha Mitchell Sci. Soc.* XIX. 22 (1903).—**Pruinosae.**—Ohio, Michigan (type locality: Port Huron).
- immitis* Ashe in *Jour. Elisha Mitchell Sci. Soc.* XIX. 28 (1903).—**Crusgalli.**—Pennsylvania (type locality: Delaware County). (?)
- impar* Beadle in *Biltmore Bot. Studies*, i. 72 (1902).—**Flavae.**—Georgia (type locality: Marietta).
- impervia* Sargent in *Proc. Acad. Nat. Sci. Phila.* LXII. 197 (1910).—**Silvicolae.**—Pennsylvania (type locality: Dunmore, near Scranton).
- implicata* Sargent in *N. Y. State Mus. Bull.* CXXII. 49 (1908).—**Silvicolae.**—New York (type locality: Buffalo).
- improvisa* Sargent in *Ontario Nat. Sci. Bull.* IV. 69 (1908).—**Anomalae.**—Ontario (type locality: Toronto).
- inanis* Beadle in *Biltmore Bot. Studies*, i. 62 (1902).—**Intricatae.**—Alabama (type locality: Albertville).
- inaudita* Sargent in *Ontario Nat. Sci. Bull.* IV. 64 (1908).—**Rotundifoliae.**—Ontario (type locality: Chippewa).
- incaedua* Sargent, *Trees and Shrubs*, II. 3 (1907).—**Punctatae.**—Missouri (type locality: Monteer).
- incana* Beadle in *Biltmore Bot. Studies*, i. 113 (1902).—**Flavae.**—Florida (type locality: Bristol).
- incerta* Sargent in *Rep. Geol. Surv. Mich.* 1906, 523 (1907).—**Punctatae.**—Michigan (type locality: Port Huron).
- incilis* Beadle in *Biltmore Bot. Studies*, i. 41 (1901).—**Pulcherrimae.**—Alabama (type locality: Evergreen).
- incisa* Sargent in *Rhodora*, VII. 196 (1905).—**Pruinosae.**—Connecticut (type locality: Stratford).

CRATAEGUS

- incompta Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 183 (1910).—
Pruinosae.—Pennsylvania (type locality: Scranton).
- indentata Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 817 (1838), as synonym of *C. coccinea* var. *indentata*. (?)
- indicens Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 27 (1903).—**Punctatae.**—Ohio (type locality: Mansfield). (?)
- indigens Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 13 (1903).—**Uniflorae.**—North Carolina (type locality: Moore County). (?)
- inducta Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 24 (1903).—**Intricatae.**—Pennsylvania (type locality not given).
- induta Sargent, Trees and Shrubs, I. 115 (1903).—**Molles.**—Arkansas (type locality: Fulton).
- infensa Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 201 (1910).—
Silvicolae.—Pennsylvania (type locality: Charleroi).
- infera Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 651 (1905).—**Intricatae.**—Pennsylvania (type locality: Sellersville).
- infesta Sargent in Rep. Missouri Bot. Gard. xix. 45 (1908).—**Crus-galli.**—Missouri (type locality: Pacific).
- ingens Beadle in Biltmore Bot. Studies, I. 56 (1902).—**Virides.**—Tennessee (type locality: Chattanooga).
- ingestria Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 821 (1838), as var. of *C. prunifolia*; I.c., 849, as synonym of *C. crus-galli prunifolia* = *C. prunifolia* Pers.?
- ingrata Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 54 (1904).—**Pruinosae.**—Pennsylvania (type locality: Pittsburgh). (?)
- inopina Beadle in Biltmore Bot. Studies, I. 107 (1902).—**Flavae.**—Florida (type locality: Ocala). (?)
- inopinata Sargent in N. Y. State Mus. Bull. cxxii. 108 (1908).—**Anomalae.**—New York (type locality: Coopers Plains).
- inops Beadle in Biltmore Bot. Studies, I. 96 (1902).—**Flavae.**—Alabama (type locality: Ozark). (?)
- insidiosa Beadle in Biltmore Bot. Studies, I. 94 (1902).—**Flavae.**—Georgia, Alabama (type locality: Ozark, Ala.).
- insignata Sargent in N. Y. State Mus. Bull. cxxii. 101 (1908).—**Tenuifoliae.**—New York (type locality: Coopers Plains).
- insignis Sargent, Trees and Shrubs, I. 107 (1903).—**Crus-galli.**—Illinois (type locality: Kahokia).
- insolens Sargent in Rhodora, VII. 217 (1905).—**Rotundifoliae.**—Vermont (type locality: Concord).
- insolita Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 622 (1905).—
Tenuifoliae.—Pennsylvania (type locality: Collen Brook, Upper Darby).
- insperata Sargent in Rep. Missouri Bot. Gard. xix. 123 (1908).—
Macracanthae.—Missouri (type locality: Carthage).

CRATAEGUS

- insueta Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 599 (1905).—
Pruinosae.—Pennsylvania (type locality: West Fairmount Park, Philadelphia, cultivated from unknown source).
integra (Nash) Beadle in Biltmore Bot. Studies, I. 87 (1902).—**Flavae.**—
Florida (type locality of *C. flava* var. *integra* Nash: Lake Ella).
integriloba Sargent in Rhodora, III. 78 (1901).—**Macracanthea.**—
Quebec (type locality: Caughnawaga).
interior Beadle in Biltmore Bot. Studies, I. 52 (1902).—**Virides.**—
Tennessee, Alabama, Mississippi (type locality: Chattanooga, Tenn.).
intermixta Sargent in Jour. Arnold Arb. III. 5 (1922).—**Crus-galli.**—
Missouri (type locality: Riverside Park, Hannibal).
intricata Lange in Bot. Tidsskr. XIX. 264 (1895).—**Intricatae.**—New England and New York to Pennsylvania and West Virginia (type locality: Botanical Garden of Copenhagen, cultivated).
insitula Sargent in N. Y. State Mus. Bull. CXXII. 55 (1908).—**Pruinosae.**—New York (type locality: Chapinville).
invicta Beadle in Biltmore Bot. Studies, I. 37 (1901).—**Flavae.**—Georgia (type locality: Doctortown).
invisa Sargent, Trees and Shrubs, II. 147 (1911).—**Molles.**—Arkansas, Oklahoma, Texas (type locality: Fulton, Ark.).
iracunda Beadle in Biltmore Bot. Studies, I. 124 (1902).—**Pruinosae.**—
Georgia (type locality: Rome).
irrasa Sargent in Rhodora, V. 116 (1903).—**Coccineae.**—Quebec, New York (type locality: Sault au Ricollet, Montreal Island, Que.).
irrasa var. *Blanchardi* (Sarg.) Eggleston in Rhodora, X. 79 (1908) = *C. Blanchardii*.
irrasa var. *divergens* Peck in N. Y. State Mus. Bull. LXXV. Bot. VII. 51 (1904) = *C. divergens*.
iterata Sargent in Jour. Arnold Arb. IV. 102 (1923).—**Silvicolae.**—New York (type locality of *C. seclusa* Sarg., which this name replaces: Hemlock Lake, Livingston County).
Jackii Sargent in Rhodora, V. 162 (1903).—**Rotundifoliae.**—Quebec (type locality: Caughnawaga).
jasperensis Sargent in Rep. Missouri Bot. Gard. XIX. 61 (1908).—
Crus-galli.—Missouri (type locality: Webb City).
jejuna Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 593 (1905).—**Pruinosae.**—Pennsylvania, Ohio (type locality: Sellersville, Pa.).
Jenningsii Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 241 (1910).—
Intricatae.—Pennsylvania, Ohio (type locality: Kittanning, Pa.).
Jesupii Sargent in Rhodora, V. 611 (1903).—**Pruinosae.**—Vermont (type locality: West Rutland).
Jonesae Sargent in Bot. Gaz. XXXI. 14 (1901).—**Rotundifoliae.**—South-eastern Canada, Maine (type locality: Bar Harbor, Me.).

CRATAEGUS

- Keepii Sargent in *Rhodora*, v. 165 (1903).—**Rotundifoliae.**—Maine (type locality: Fort Fairfield).
- Kellermanii Sargent, *Trees and Shrubs*, II. 239 (1913).—**Pruinosae.**—Ohio, Kentucky (type locality: Dennison, O.).
- Kelloggii Sargent, *Trees and Shrubs*, I. 117 (1903).—**Molles.**—Missouri (type locality: St. Louis).
- Kilmanii Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, *Arb. Brit.* II. 849 (1838) as synonym of *C. coccinea* var. *maxima*, Lodd.
- Kennedyi Sargent, *Trees and Shrubs*, II. 73 (1903).—**Rotundifoliae.**—Vermont (type locality: summit of Willoughby Mountain, Orleans County).
- kingstonensis Sargent in *Jour. Arnold Arb.* III. 205 (1922).—**Rotundifoliae.**—Ontario (type locality: Kingston).
- Kinzerae Sargent in *Proc. Acad. Nat. Sci. Phila.* LXII. 234 (1910).—**Intricatae.**—Pennsylvania (type locality: Schenley Park, Pittsburgh).
- Knieskerniana Sargent in *N. Y. State Mus. Bull.* CLXVII. 116 (1913).—**Anomalae.**—Quebec, New York (type locality: East Herkimer, N. Y.).
- lacera Sargent in *Bot. Gaz.* XXXIII. 123 (1902).—**Tenuifoliae?**—Arkansas (type locality: Fulton).
- lacrimate Small in *Torreya*, I. 97 (1901).—**Flavae.**—Florida (type locality: Crestview).
- laetans Sargent in *Proc. Acad. Nat. Sci. Phila.* LXII. 205 (1910).—**Silvicolae.**—Pennsylvania (type locality: Pittsburgh).
- laetifica Sargent in *Proc. Acad. Nat. Sci. Phila.* LXII. 250 (1910).—**Intricatae.**—Pennsylvania (type locality: Bedford).
- laetula Sargent in *Proc. Acad. Nat. Sci. Phila.* LXII. 233 (1910).—**Intricatae.**—Pennsylvania (type locality: Newtown).
- lanata Beadle in *Biltmore Bot. Studies*, I. 86 (1902).—**Flavae.**—North Carolina, Georgia (type locality: McGuire's Mill, Gwinnett County, Ga.).
- lanceolata Sargent, *Trees and Shrubs*, II. 65 (1908).—**Virides.**—Missouri (type locality: St. Louis).
- lancifolia Wenderoth, *Ind. Sem. Hort. Marb.* (1825) ex *Flora*, IX. pt. 1, 354 (1826) = *C. crus-galli* var. *pyracanthifolia* Ait.
- Laneyi Sargent, *Trees and Shrubs*, I. 5 (1902).—**Macracantha.**—New York (type locality: Rochester).
- lanigera Sargent in *Rep. Missouri Bot. Gard.* XIX. 107 (1908).—**Molles.**—Illinois, Kentucky, Missouri (type locality: East St. Louis, Ill.).
- lanuginosa Sargent, *Trees and Shrubs*, I. 113 (1903).—**Molles.**—Missouri, Kansas, Arkansas (type locality: Webb City, Mo.).
- larga Sargent in *Rep. Missouri Bot. Gard.* XIX. 85 (1908).—**Virides.**—Illinois (type locality: Cahokia).

CRATAEGUS

- lasiantha Sargent in Rep. Missouri Bot. Gard. xix. 105 (1908).—
Molles.—Missouri, Kansas, Oklahoma (type locality: Webb City, Mo.).
- lassa Beadle in Biltmore Bot. Studies, 1. 29 (1901).—Flavae.—Alabama (type locality: Selma).
- lata Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 610 (1905).—Tenuifoliae.—Pennsylvania (type locality: Sellersville).
- latebrosa Sargent in Rep. Missouri Bot. Gard. xxii. 73 (1912).—Punctatae.—Missouri, Arkansas (type locality: Noel, Mo.).
- latiflora Sargent in N. Y. State Mus. Bull. CLXVII. 83 (1913).—Pruinosaes.—New York (type locality: Richmond, Livingston County).
- latifolia Persoon, Syn. II. 37 (1807) = C. tomentosa L.?
- latifrons Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 181 (1910).—
Pruinosae.—Pennsylvania (type locality: Orbisonia).
- latisepala Ashe in Bull. N. Car. Exper. Sta. CLXXV. 109 (1900).—
Pruinosae.—Michigan, Ohio (type locality: "eastern Michigan").
- laurentiana Sargent in Rhodora, III. 77 (1901).—Macracantheae.—
Quebec (type locality: Caughnawaga).
- laurifolia Medicus, Gesch. Bot. 84 (1793).—Crus-galli.—(?).
- lauta Sargent in Rhodora, VII. 206 (1905).—Molles.—Known only in cultivation. (Type locality: Olmstead Park, Boston, Mass.).
- lawrencensis Sargent in Jour. Arnold Arb. III. 3 (1922).—Crus-galli.—
Missouri (type locality: Larussell).
- laxa Beadle in Biltmore Bot. Studies, 1. 103 (1902).—Flavae.—Alabama (type locality: Phoenix City).
- laxiflora Sargent in Bot. Gaz. XXXV. 400 (1903).—Macracantheae.—
Illinois (type locality: Mokena).
- lecta Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 187 (1910).—Pruinosaes.—Pennsylvania (type locality: between Carnot and Stoope's Ferry, Allegheny County).
- leimonia Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 194 (1910).—
Silvicolae.—Pennsylvania (type locality: Kittanning).
- leioclada Sargent in Rep. Missouri Bot. Gard. xix. 113 (1908).—Intricatae.—Missouri, Arkansas (type locality: Swan, Mo.).
- leiophylla Sargent in Proc. Rochester Acad. Sci. IV. 99 (1903).—Pruinosaes.—New York, Pennsylvania, Ohio, Indiana (type locality: Rochester, N. Y.).
- leiophylla var. *maineana* (Sarg.) Eggleston in Rhodora, x. 81 (1908) = C. *maineana*.
- lemingtonensis Sargent in Rhodora, VII. 216 (1905).—Rotundifoliae.—
Vermont, New Hampshire (type locality: Lemington, Vt.).
- lenis Beadle in Biltmore Bot. Studies, 1. 67 (1902).—Pulcherrimae.—
Alabama (type locality: Greenville). (?)

CRATAEGUS

- Lennoniana Sargent in Proc. Rochester Acad. Sci. iv. 98 (1903).—
Pruinosae.—New York (type locality: Rochester).
- lenta Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 18 (1902).—Coccineae.—Quebec, Michigan (type locality: Port Huron, Mich.).
- lentula Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 25 (1903).—Intricatae.—North Carolina (type locality: Catawba).
- lepida Beadle in Biltmore Bot. Studies, i. 36 (1901).—Flavae.—Georgia, Florida (type locality: Waycross, Ga.).
- leptalea Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 231 (1910).—
Intricatae.—Pennsylvania (type locality: Bedford Springs).
- leptophylla Sargent in Rep. Missouri Bot. Gard. xix. 69 (1908).—
Crus-galli.—Missouri (type locality: Osage).
- leptopoda Sargent in N. Y. State Mus. Bull. cxxii. 118 (1908).—
Tenuifoliae.—New York (type locality: Hemlock Lake, Livingston County).
- Letchworthiana Sargent in N. Y. State Mus. Bull. cxxii. 68 (1908).—
Coccineae.—New York (type locality: Portage).
- Lettermanii Sargent in Bot. Gaz. xxxi. 220 (1901).—Punctatae.—
Missouri (type locality: Allenton).
- leucophleos Moench, Bäume Weiss. 31 (1785).—Macracanthea.—(?)
- levis Sargent in Rhodora, vii. 198 (1905).—Pruinosae.—Connecticut (type locality: Litchfield).
- limaria Sargent, Trees and Shrubs, ii. 149 (1911).—Molles.—Arkansas, Oklahoma, Texas, Mexico (type locality: Fulton, Ark.).
- limata Beadle in Biltmore Bot. Studies, i. 77 (1902).—Flavae.—Georgia (type locality: Warm Springs).
- limatula Sargent in Ontario Nat. Sci. Bull. iv. 16 (1908), as "limatala".—
Punctatae.—Ontario (type locality: London).
- limnophylla Sargent in Jour. Arnold Arb. iii. 3 (1922).—Crus-galli.—
Florida (type locality: St. Marks).
- limosa Sargent in N. Y. State Mus. Bull. cxxii. 67 (1908).—Coccineae.—
New York (type locality: Rochester).
- Lindenii Stapf in Kew Bull. Misc. Inform. 1914, 326 (1914).—Group?—
Mexico (type locality: Chiapas, Ciudad Real). (?)
- linearifolia Hort. ex Nicholson in Hand-list Arb. Kew, pt. 1, 197 (1894)
as synonym of *C. crus-galli* var. *linearis* DC.
- linearis Persoon, Syn. ii. 37 (1807) = *C. crus-galli* var. *pyracanthifolia* Ait.
- littoralis Sargent in Rhodora, v. 59 (1903).—Pruinosae.—Connecticut (type locality: Waterford).
- Livingstoniana Sargent in N. Y. State Mus. Bull. cxxii. 116 (1908).—
Silvicolae.—New York (type locality: Hemlock Lake, Livingston County).

CRATAEGUS

- livoniana* Sargent, Trees and Shrubs, II. 3 (1908).—**Crus-galli.**—New York (type locality: Hemlock Lake).
- lobata* (Poir.) Bosc in DeCandolle, Prodr. II. 628 (1825).—**Macracanthae.**—(?)
- lobulata* Sargent in Rhodora, III. 22 (1901).—**Coccineae.**—New England, New York (type locality: New Haven, Vt.).
- locuples* Sargent in Rep. Missouri Bot. Gard. xix. 97 (1908).—**Pruinosae.**—Missouri (type locality: Barretts Station).
- Loddigesii* Hort. ex Nicholson in Hand-list Arb. Kew, pt. 1, 207 (1894) as synonym of *C. punctata* Jacq.
- longipedunculata* Sargent in Ontario Nat. Sci. Bull. IV. 26 (1908).—**Pruinosae.**—Ontario, New York, Ohio (type locality: Chippewa, Ont.).
- longipetiolata* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 621 (1905).—**Tenuifoliae.**—Pennsylvania (type locality: Sellersville).
- longispina* Sargent in Bot. Gaz. XXXV. 398 (1903).—**Macracanthae.**—Illinois (type locality: Lake Zurich).
- lucida* Miller, Dict. ed. 8, no. 6 (1768) = *C. crus-galli* L.
- lucida* Elliot, Sketch, I. 548 (1821) = *C. aestivalis* T. & G.?
- lucorum* Sargent in Bot. Gaz. XXXI. 227 (1901).—**Tenuifoliae.**—New York, Ohio, Illinois (type locality: Barrington, Ill.).
- lucorum* var. *insolens* (Sarg.) Eggleston in Rhodora, x. 79 (1908) = *C. insolens*.
- luculenta* Sargent, Trees and Shrubs, I. 11 (1902) = *C. aestivalis* var. *cerasoides* f. *luculenta* Sarg.
- ludoviciensis* Sargent, Trees and Shrubs, II. 5 (1907).—**Virides.**—Missouri (type locality: St. Louis).
- lumaria* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 25 (1903) = *C. Dodgei* var. *lumaria* Sarg.
- luminosa* Sargent in N. Y. State Mus. Bull. CXXXII. 63 (1908).—**Tenuifoliae.**—New York (type locality: Buffalo).
- lutea* Poiret apud Jackson, Ind. Kew. I. 636 (1893), as synonym of *C. tomentosa* L.—*Mespilus lutea* Poiret in Lamarck, Encycl. Meth. Suppl. IV. 71 (1816).
- lutensis* Sargent in Rep. Missouri Bot. Gard. XXII. 75 (1912).—**Virides.**—Missouri, Oklahoma (type locality: Reding's Mill, near Joplin, Mo.).
- luteola* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 240 (1910).—**Intricatae.**—Pennsylvania (type locality: Orbisonia).
- luxuriosa* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 198 (1910).—**Silvicolae.**—Pennsylvania (type locality: Kittanning).
- Macaulayae* Sargent in Proc. Rochester Acad. Sci. IV. 130 (1903).—**Rotundifoliae.**—New York (type locality: Genesee Valley Park, Rochester).

- macera Sargent in N. Y. State Mus. Bull. cxxii. 117 (1908).—**Silvicolae.**—Ontario, New York (type locality: Hemlock Lake, N. Y.).
- macilenta Beadle in Biltmore Bot. Studies, I. 64 (1902).—**Intricatae.**—Alabama (type locality: Albertville).
- Mackensenii* Sargent, Trees and Shrubs, II. 242 (1913) = *C. limaria* Sarg.
- Mackenzii* Sargent in Mackenzie, Man. Fl. Jackson County, Mo. 108 (1902).—**Pruinosae.**—Missouri (type locality: Jackson County).
- Macnabiana* Hort. ex Nicholson in Hand-list Arb. Kew, pt. 1, 199 (1894), as synonym of *C. Douglasii* var. *rivularis* Sarg.
- Macounii* Sargent in Ontario Nat. Sci. Bull. IV. 41 (1908).—**Coccineae.**—Ontario (type locality: Toronto).
- macra Beadle in Biltmore Bot. Studies, I. 134 (1902).—**Crus-galli.**—North Carolina, Georgia, Mississippi (type locality: Rome, Ga.).
- macracantha* Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 819 (1838).—**Macracantheae.**—Ontario, Vermont and New York to northern Illinois. (First known and described from plants cultivated in European gardens.)
- macracantha* var. *minor* Loudon, Arb. Brit. II. 819 (1838). (?)
- macracantha* var. *neofluvialis* (Ashe) Eggleston in Rhodora, x. 83 (1908) = *C. neofluvialis*.
- macracantha* var. *occidentalis* (Britton) Eggleston in Rhodora, x. 82 (1908) = *C. succulenta* Schrad.?
- macracantha* var. *rhombifolia* (Sarg.) Eggleston in Rhodora, x. 82 (1908) = *C. rhombifolia*.
- macracantha* var. *succulenta* (Schrad.) Rehder in Bailey, Cycl. Am. Hort. I. 396 (1900) = *C. succulenta*.
- macrocalyx* Sargent in N. Y. State Mus. Bull. cxxii. 89 (1908).—**Pruinosae.**—New York (type locality: Coopers Plains).
- macrophylla* Sarg. in Rep. Missouri Bot. Gard. xix. 104 (1908).—**Molles.**—Missouri (type locality: Carondelet).
- macropoda* Sargent in Rep. Missouri Bot. Gard. xix. 80 (1908).—**Punctatae.**—Missouri, Kansas, Arkansas (type locality: Webb City, Mo.).
- macrosperma* Ashe in Jour. Elisha Mitchell Sci. Soc. xvi. 73 (1900).—**Tenuifoliae.**—North Carolina and West Virginia to Georgia and Tennessee (type locality: Lookout Mountain, Tenn.).
- macrosperma* var. *acutiloba* (Sarg.) Eggleston in Rhodora, x. 80 (1908) = *C. acutiloba*.
- macrosperma* var. *demissa* (Sarg.) Eggleston in Rhodora, x. 80 (1908) = *C. demissa*.
- macrosperma* var. *matura* (Sarg.) Eggleston in Rhodora, x. 80 (1908) = *C. matura*.
- macrosperma* var. *pastorum* (Sarg.) Eggleston in Rhodora, x. 80 (1908) = *C. pastorum*.

CRATAEGUS

- macrosperma* var. *pentandra* (Sarg.) Eggleston in Rhodora, x. 80 (1908) =
C. pentandra.
- macrosperma* × *pruinosa* Eggleston in N. Y. State Mus. Bull. CCXLIII-
 iv. 64 (1923). (?)
- macrosperma* × *punctata* Eggleston in N. Y. State Mus. Bull. CCXLIII-
 iv. 64 (1923). (?)
- magniflora* Sargent in Bot. Gaz. XXXV. 383 (1903).—**Coccineae.**—
 Ontario, Northern Illinois (type locality: Glendon Park, near
 Chicago, Ill.).
- magnifolia* Sargent, Trees and Shrubs, II. 69 (1908).—**Pruinosae.**—
 Missouri (type locality: Webb City).
- Maineana* Sargent in Proc. Rochester Acad. Sci. IV. 106 (1903).—
Silvicolae.—New York (type locality: Rochester).
- maligna* Sargent in N. Y. State Mus. Bull. CLXVII. 111 (1913).—**Rotundi-**
foliae.—New York (type locality: Ogdensburg).
- maloides* Sargent, Trees and Shrubs, I. 9 (1902) = *C. aestivalis* var.
maloides Sarg.
- mansfieldensis* Sargent in Jour. Arnold Arb. IV. 103 (1923).—**Rotundi-**
foliae.—Ohio (type locality: Mansfield).
- marcida* Ashe in Jour. Elisha Mitchell Sci. Soc. XX. 53 (1904).—**Tenui-**
foliae.—Ohio (type locality not given).
- Margaretta* Ashe in Jour. Elisha Mitchell Sci. Soc. XVI. 72 (1900).—
Rotundifoliae.—Ontario and West Virginia, west to Ohio, Iowa
 and Missouri (type locality: St. Louis County, Mo.).
- Margaretta* var. *Brownii* (Britton) Sargent in Jour. Arnold Arb. III. 199
 (1922).—Pennsylvania, Virginia and West Virginia, west to Ohio,
 Indiana and Missouri (type locality of *C. Brownii*: Buchanan, Va.).
- Margaretta* f. *xanthocarpa* Sargent in Jour. Arnold Arb. III. 200 (1922).—
 Iowa (type locality: Steamboat Rock).
- Maribella* Sargent in N. Y. State Mus. Bull. CLXVII. 109 (1913).—
Rotundifoliae.—New York (type locality: Little Falls).
- mariettensis* Sargent in Jour. Arnold Arb. III. 194 (1922).—**Pruinosae.**—
 Ohio (type locality: Marietta).
- Marshallii* Eggleston in Rhodora, x. 79 (1908) = *C. apiifolia* Michx.
- matura* Sargent in Rhodora, III. 24 (1901); v. 144 (1903).—**Tenui-**
foliae.—Southeastern Canada, New England, New York, Michigan
 (type locality: Middlebury, Vt.).
- McGeeae* Ashe in Jour. Elisha Mitchell Sci. Soc. XVIII. 20 (1902).—
 Group?—Iowa (type locality: “southeastern Iowa”). (?)
- media* Sargent in Rhodora, v. 150 (1903).—**Tenuifoliae.**—Connecticut
 (type locality: Oxford).
- medioxima* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 190 (1910).—
Silvicolae.—Pennsylvania, Ohio (type locality: between Carnot
 and Stope's Ferry, Pa.).

CRATAEGUS

- meiophylla* Sargent in Jour. Arnold Arb. III. 198 (1922).—**Rotundifoliae.**—Ohio (type locality: Mt. Victory).
- mellita* Sargent in N. Y. State Mus. Bull. cv. Bot. IX. 58 (1906).—**Tenuifoliae.**—New York (type locality: Sandlake).
- membranacea* Sargent in Rhodora, v. 186 (1903).—**Macracantheae.**—Vermont (type locality: Middlebury).
- menandiana* Sargent in N. Y. State Mus. Bull. cv. Bot. IX. 68 (1906).—**Macracantheae.**—New York (type locality: golf grounds, Menands).
- mendoza* Beadle in Biltmore Bot. Studies, I. 65 (1902).—**Intricatae.**—Alabama (type locality: Albertville).
- mercerensis* Sargent in Jour. Arnold Arb. III. 193 (1922).—**Rotundifoliae.**—West Virginia (type locality: Mercer Springs).
- meridiana* Beadle in Biltmore Bot. Studies, I. 115 (1902).—**Flavae.**—Alabama (type locality: Ozark).
- meridionalis* Sargent in Jour. Arnold Arb. I. 252 (1920).—**Molles.**—Alabama, Mississippi (type locality: Gallion, Ala.).
- merita* Sargent in Rep. Geol. Surv. Mich. 1906, 540 (1907).—**Tenuifoliae.**—Michigan (type locality: Grand Rapids).
- mespilifolia* Hort. ex Nicholson in Hand-list Arb. Kew, pt. 1, 203 (1894) as synonym of *C. mollis* Scheele.
- meticulosa* Sargent, Trees and Shrubs, II. 243 (1913).—**Intricatae.**—Michigan, Indiana (type locality: Grand Rapids, Mich.).
- mexicana* Mociño & Sessé apud DeCandolle, Prodr. II. 629 (1825).—Group?—Mexico (type region: “Nova Hispana”).
- mexicana* var. *microsperma* Eggleston in Bull. Torrey Bot. Club, XXXVI. 508 (1909).—Mexico (type locality: Guadalajara, Jalisco).
- Michauxii* Persoon, Syn. II. 38 (1807).—**Flavae.**—North Carolina, Georgia (type region: “Carol. Superior”).
- michiganensis* Ashe in Bull. N. Car. Exper. Sta. CLXXV. 111 (1900).—**Macracantheae.**—Michigan (type locality not given).
- micrantha* Sargent, Trees and Shrubs, I. 69 (1902).—**Virides.**—Arkansas (type locality: Fulton).
- micrantha* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 658 (1905).—**Macracantheae.**—Pennsylvania (type locality: Sellersville).
- microcarpa* Lindley in Bot. Reg. XXII. t. 1846 (1836) = *C. spathulata* Michx.
- microsperma* Sargent in Ontario Nat. Sci. Bull. IV. 82 (1908).—**Macracantheae.**—Southeastern Canada, New York (type locality: Toronto, Ont.).
- Milleri* Sargent in Jour. Arnold Arb. IV. 101 (1923).—**Pruinosae.**—Pennsylvania (type locality: Erie).
- miniata* Ashe in Jour. Elisha Mitchell Sci. Soc. XX. 51 (1904).—**Tenuifoliae.**—Pennsylvania, Ohio (type locality: Berks County, Pa.).

CRATAEGUS

- minutiflora* Sargent in Ontario Nat. Sci. Bull. iv. 54 (1908).—**Intricatae.**—Ontario, New York (type locality: Toronto, Ont.).
- mira* Beadle in Biltmore Bot. Studies, I. 78 (1902).—**Flavae.**—Georgia (type locality: Marietta).
- miranda* Sargent in Rep. Geol. Surv. Mich. 1906, 547 (1907).—**Coccineae.**—Ontario, Michigan, Ohio (type locality: Sarnia, Ont.).
- misella* Sargent in N. Y. State Mus. Bull. CLXVII. 115 (1913).—**Anomalae.**—New York (type locality: Belfast).
- missouriensis* Ashe in Bull. N. Car. Exper. Sta. CLXXV. 110 (1900).—**Macracantha?**—Tennessee, Missouri (type locality: Monteer, Mo.).
- mitis* Sargent, Man. Trees N. Am. 407 (1905).—**Virides.**—Illinois (type locality: Cahokia).
- modesta* Sargent in Rhodora, III. 28 (1901).—**Intricatae.**—New England, New York, Pennsylvania (type locality: West Rutland, Vt.).
- modica* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 624 (1905).—**Tenuifoliae.**—Pennsylvania (type locality: Sellersville).
- Mohrii* Beadle in Bot. Gaz. XXVIII. 416 (1899).—**Crus-galli.**—Georgia and Alabama, west to Kentucky, Tennessee and Arkansas (type locality: Birmingham, Ala.).
- mollicula* Sargent, Trees and Shrubs, II. 13 (1907).—**Macracantha.**—Missouri, Arkansas (type locality: Monteer, Mo.).
- mollipes* Sargent in Rep. Geol. Surv. Mich. 1906, 545 (1907).—**Molles.**—Michigan, Indiana (type locality: Grand Rapids, Mich.).
- mollis* (Torr. & Gray) Scheele in Linnaea, XXI. 569 (1848).—**Molles.**—Ontario and Michigan west to Minnesota and Nebraska, south to Tennessee, Missouri and eastern Kansas (type region: “in civitate Illinoensi Americae borealis”).
- mollis* var. *champlainensis* (Sarg.) Cowles in Am. Nat. XLII. 269 (1908) = *C. champlainensis*.
- mollis* var. *Ellwangeriana* (Sarg.) Cowles in Am. Nat. XLII. 269 (1908) = *C. Ellwangeriana*.
- mollis* var. *sera* (Sarg.) Eggleston in Rhodora, x. 82 (1908) = *C. sera*.
- mollis* var. *tiliaeefolia* (K. Koch) Koehne, Deutsche Dendr. 232 (1893) = *C. mollis* Scheele.
- mollita* Sargent, Trees and Shrubs, II. 77 (1908).—**Macracantha.**—Missouri (type locality: Monteer).
- monantha* Sargent, Trees and Shrubs, II. 237 (1913).—**Aestivales?**—Florida (type locality: Seville).
- monosperma* Sargent in Rep. Missouri Bot. Gard. XIX. 56 (1908).—**Crus-galli.**—Missouri (type locality: Swan).
- monstrata* Sargent in Rhodora, v. 146 (1903).—**Tenuifoliae.**—Massachusetts, Connecticut (type locality: Eight Mile Brook, Southbury and Middlebury, Ct.).

CRATAEGUS

- montivaga Sargent in Jour. Arnold Arb. I. 247 (1920).—**Crus-galli**.—Texas (type locality: Spring Creek, near Boerne).
- moselemensis Gruber in Bull. Torrey Bot. Club, XXXII. 391 (1905).—**Punctatae**.—Pennsylvania (type locality: Moselem). (?)
- moselemensis var. corrugata Gruber in Bull. Torrey Bot. Club, XXXII. 391 (1905).—Pennsylvania (type locality: Charming Forge, Berks County). (?)
- Moyeriana Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 618 (1905).—**Tenuifoliae**.—Pennsylvania (type locality: Sellersville).
- multifida Ashe in Jour. Elisha Mitchell Sci. Soc. XX. 52 (1904).—**Tenuifoliae**.—Michigan (type locality: Port Huron). (?)
- multipes Ashe in Bull. N. Car. Exper. Sta. CLXXV. 111 (1900).—Group?—Ohio (type locality not given). (?)
- multispina Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 7 (1900).—**Crus-galli**.—North Carolina (type locality: Round Knob, below Swannona Gap). (?)
- mundula Beadle in Biltmore Bot. Studies, I. 38 (1901).—**Flavae**.—South Carolina (type locality: Batesburg).
- munita Sargent in Rep. Missouri Bot. Gard. XIX. 73 (1908).—**Crus-galli**.—Missouri, Kansas, Oklahoma (type locality: Webb City, Mo.).
- nana* (Dum.) Sweet, Hort. Brit. ed. 3, 209 (1839).—*Mespilus nana* Dumont de Courset, Bot. Cult. ed. 2, VII. 286 (1814) = *C. crus-galli* var. *pyracanthifolia* Ait.?
- Napaea Sargent in Rhodora, VII. 203 (1905).—**Tenuifoliae**.—Connecticut (type locality: Litchfield).
- napolitana* Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 849 (1838), as synonym of *C. coccinea* L.
- Nelsonii Eggleston in Bull. Torrey Bot. Club, XXXVI. 512 (1909).—Group?—Mexico (type locality: San Cristobal, Chiapas). (?)
- nemoralis Sargent in Bot. Gaz. XXXV. 104 (1903).—**Intricatae**.—Pennsylvania, Delaware (type locality: Rockford Park, Wilmington, Del.).
- nemorosa Sargent in N. Y. State Mus. Bull. CL. 28 (1911).—**Pruinosae**.—New York (type locality: Painted Post).
- neobaxteri Sargent in N. Y. State Mus. Bull. CXXII. 74 (1908).—**Rotundifoliae**.—New York (type locality: Tuscarora).
- neobushii Sargent, Trees and Shrubs, II. 9 (1907).—**Intricatae**.—Illinois, Missouri (type locality: Monteer, Mo.).
- neocanbyi Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 638 (1905).—**Intricatae**.—Pennsylvania (type locality: between Stroudsburg and Tannersville).
- neofaxonii Sargent in Jour. Arnold Arb. VI. 2 (1925).—**Coccineae**.—Massachusetts (type locality of *C. ampla* Sarg., which this name replaces: Lanesboro.)

CRATAEGUS

- neofluvialis Ashe in Jour. Elisha Mitchell Sci. Soc. xvi. 71 (1900).—
Macracanthalae.—North Carolina, Virginia, West Virginia, Indiana
(type locality not given).
- neolondinensis Sargent, Man. Trees N. Am. 443 (1905).—**Coccineae.**—
Connecticut (type locality: New London).
- neosmithii Sargent in Jour. Arnold Arb. vi. 2 (1925).—**Pruinosae.**—
Pennsylvania (type locality of *C. ignava* Sarg., which this name
replaces: Bedford Springs).
- nescia Sargent in N. Y. State Mus. Bull. cxxii. 100 (1908).—**Tenui-
foliae.**—New York (type locality: Coopers Plains).
- nitens Sargent in Rep. Missouri Bot. Gard. xxii. 76 (1912).—**Virides.**—
Missouri (type locality: Hannibal).
- nitida (Engelm.) Sargent in Bot. Gaz. xxxi. 231 (1901).—**Virides.**—
Illinois, Missouri, Arkansas (type locality not given).
- nitidula Sargent in Rep. Geol. Surv. Mich. 1906, 521 (1907).—**Puncta-
tae.**—Michigan (type locality: Port Huron).
- noelensis Sargent in Jour. Arnold Arb. i. 253 (1920).—**Molles.**—Missouri,
Arkansas (type locality: Noel, Mo.).
- notabilis Sargent in N. Y. State Mus. Bull. cxxii. 32 (1908).—**Punc-
tatae.**—New York (type locality: Buffalo).
- × notha Sargent in Jour. Arnold Arb. iii. 9 (1922).—(*C. apiifolia* ×
brachyphylla).—Arkansas (type locality: McNab, Hempstead
County).
- noveboracensis Sargent in N. Y. State Mus. Bull. cxvi. Bot. x. 22
(1907).—**Rotundifoliae.**—New York (type locality: North Elba).
- nubicola Beadle in Biltmore Bot. Studies, i. 126 (1902).—**Tenuifoliae.**—
North Carolina (type locality: Mt. Pisgah, Alexander County).
- nuda Sargent in Jour. Arnold Arb. iii. 10 (1922).—**Macracanthalae.**—
Missouri (type locality: Swan).
- nudiflora Nuttall apud Torrey & Gray, Fl. N. Am. i. 468 (1840), as
synonym of *C. aestivalis* T. & G.
- numerosa Sargent in N. Y. State Mus. Bull. cxxii. 90 (1908).—**Pru-
nosae.**—New York (type locality: Coopers Plains).
- nupera Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 17 (1903).—**Molles.**—
Illinois (type locality: Joliet). (?)
- nutans Sargent in Rep. Geol. Surv. Mich. 1906, 544 (1907).—**Molles.**—
Michigan (type locality: Port Huron).
- Oakesiana Eggleston in Torreya, vii. 35 (1907).—**Rotundifoliae.**—
Vermont (type locality: Bloomfield).
- obesa Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 47 (1904).—**Macra-
canthalae.**—Missouri (type locality: St. Louis County). (?)
- oblita Sargent in N. Y. State Mus. Bull. cxxii. 40 (1908).—**Pruinosae.**—
New York (type locality: Buffalo).

CRATAEGUS

- oblongifolia* Sargent in N. Y. State Mus. Bull. cv. Bot. ix. 60 (1906) = *C. urbica* Sarg.
- obovatifolia* Roemer, Fam. Nat. Syn. iii. 120 (1847).—**Punctatae**.—(?)
- obscura* Sargent in Rep. Missouri Bot. Gard. xix. 119 (1908).—**Macracantha**—Missouri, Arkansas (type locality: Webb City, Mo.).
- obstipa* Sargent in N. Y. State Mus. Bull. clxvii. 80 (1913).—**Pruinosa**.—New York (type locality: Chapinville).
- obtecta* Ashe in Bot. Gaz. xxxiii. 233 (1902).—**Molles**.—Illinois (type locality not given). (?)
- occidentalis* Britton in Bull. N. Y. Bot. Gard. i. 448 (1900) = *C. succulenta* Schrad.?
- ogdensburgensis* Sargent in N. Y. State Mus. Bull. clxvii. 123 (1913).—**Macracantha**.—New York (type locality: Ogdensburg).
- ohioensis* Sargent in Jour. Arnold Arb. iii. 183 (1922).—**Crus-galli**.—Ohio (type locality: Columbus).
- olida* Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 56 (1904).—**Coccineae**.—Pennsylvania (type locality: Pyler Falls).
- olivacea* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 153 (1910) = *C. Fontanesiana* Steud.
- onusta* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 22 (1903).—**Pruinosa**.—Ohio (type locality: Mansfield).
- opaca* Hooker & Arnott in Comp. Bot. Mag. i. 25 (1835).—**Aestivales**.—Alabama, Mississippi, Louisiana, Texas (type locality not given).
- operta* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 29 (1903).—**Crus-galli**.—North Carolina, Tennessee (type locality not given). (?)
- opica* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 10 (1903).—**Macracantha**.—Pennsylvania (type locality: Berks County).
- opima* Beadle in Biltmore Bot. Studies, i. 40 (1901).—**Pulcherrimae**.—Alabama, Florida (type locality: Greenville, Ala.).
- opulens* Sargent in Proc. Rochester Acad. Sci. iv. 104 (1903).—**Silvicolae**.—New York, Michigan (type locality: Rochester, N. Y.).
- ornata* Sargent in Proc. Rochester Acad. Sci. iv. 120 (1903).—**Tenuifoliae**.—Ontario, New York (type locality: Rochester, N. Y.).
- otiosa* Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 48 (1904).—**Tenuifoliae**.—Michigan, Ohio (type locality: Summerville, Mich.).
- ovalifolia* Hornemann, Hort. Hafn. Suppl. 52 (1819), ex Sudworth, U. S. Dept. Agric. Div. For. xiv. 217 (1897), as synonym of *C. crus-galli prunellifolia* Sudw. = *C. prunifolia* Pers.?
- ovata* Sargent, Man. Trees N. Am. 402 (1905).—**Virides**.—Illinois, Missouri (type locality: Carondelet, Mo.).
- ovatifolia* Sargent in N. Y. State Mus. Bull. cxxii. 92 (1908).—**Pruinosa**.—New York (type locality: Coopers Plains).

CRATAEGUS

- pachyphylla Sargent in Rep. Missouri Bot. Gard. xix. 54 (1908).—
Crus-galli.—Missouri (type locality: Osage).
- pactilis Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 20 (1903).—**Tenuifoliae**.—Wisconsin (type locality: Milwaukee). (?)
- Paddockae Sargent in Rhodora, vii. 202 (1905).—**Tenuifoliae**.—Vermont (type locality: Lyndonville).
- padifolia Sargent, Trees and Shrubs, ii. 75 (1908).—**Intricatae**.—Missouri (type locality: Swan).
- padifolia var. incarnata Sargent in Jour. Arnold Arb. vi. 4 (1925).—Missouri, Arkansas (type locality: Galena, Mo.).
- pagensis Sargent in Jour. Arnold Arb. iv. 104 (1923).—**Intricatae**.—Oklahoma (type locality: Page).
- Paineana Sargent in N. Y. State Mus. Bull. clxvii. 97 (1913).—**Tenuifoliae**.—New York (type locality: Herkimer).
- Painteriana Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 647 (1905).—**Intricatae**.—Pennsylvania (type locality: Newtown).
- pallens Beadle in Biltmore Bot. Studies, i. 27 (1901).—**Intricatae**.—North Carolina (type locality: Biltmore).
- pallescens Sargent in N. Y. State Mus. Bull. clxvii. 81 (1913).—**Pruinosaes**.—New York (type locality: Ogdensburg).
- palliata Sargent, Trees and Shrubs, ii. 236 (1913).—*Crus-galli*.—Missouri, Arkansas (type locality: Fulton, Ark.).
- pallidula Sargent in Ontario Nat. Sci. Bull. iv. 36 (1908).—**Tenuifoliae**.—Ontario (type locality: Toronto).
- Palmeri Sargent, Trees and Shrubs, i. 57 (1903).—*Crus-galli*.—Missouri, Kansas, Arkansas, Oklahoma (type locality: Webb City, Mo.).
- paludosa Sargent, Trees and Shrubs, i. 15 (1902).—**Virides**.—Florida (type locality: Haw Creek, Volusia County).
- palustris Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. 13 (1900).—Group?—Indiana (type locality: Freeport). (?)
- panda Beadle in Biltmore Bot. Studies, i. 89 (1902).—**Flavae**.—Florida (type locality: Tallahassee).
- panda Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 29 (1903).—Group?—North Carolina (type locality: Glendon). (?)
- paradoxa Sargent in Rep. Missouri Bot. Gard. xxii. 68 (1912).—*Crus-galli*.—Missouri, Kansas, Arkansas (type locality: Joplin, Mo.).
- parca Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 19 (1903).—**Tenuifoliae**.—Wisconsin (type locality: Milwaukee). (?)
- parciflora Sargent in Rep. Missouri Bot. Gard. xix. 60 (1908).—*Crus-galli*.—Missouri, Arkansas, Texas (type locality: Webb City, Mo.).
- Parkae Sargent in Rep. Missouri Bot. Gard. xxii. 70 (1912).—*Crus-galli*.—Missouri (type locality: Springfield).
- Parryana Eggleston in Bull. Torrey Bot. Club, xxxvi. 510 (1909).—Group?—Mexico (type locality: Alvarez, San Louis Potosi). (?)

CRATAEGUS

- parviflora* Sargent in Proc. Rochester Acad. Sci. iv. 117 (1903).—
Tenuifoliae.—New York, Pennsylvania, Indiana (type locality: Rochester, N. Y.).
- parvifolia* Aiton, Hort. Kew. II. 169 (1789) = *C. uniflora* Muench.
parvifolia var. *florida* (Lodd.) Loudon, Arb. Brit. II. 842 (1838). (?)
parviflora var. *grossulariaefolia* (Lodd.) Loudon, Arb. Brit. II. 842 (1838).—(?)
- parvula* Sargent in Rep. Geol. Surv. Mich. 1906, 527 (1907).—**Pruinosaes.**—Michigan (type locality: Port Huron).
- pascens* Ashe in Jour. Elisha Mitchell Sci. Soc. XVIII. 23 (1902).—
Rotundifoliae?—Michigan, Wisconsin (type locality not given). (?)
- pastorum* Sargent in Rhodora, III. 24 (1901).—**Tenuifoliae.**—New Hampshire, Massachusetts, Connecticut (type locality: Lancaster, Mass.).
- patrum* Sargent in Rep. Missouri Bot. Gard. XIX. 95 (1908).—**Pruinosaes.**—Iowa, Missouri, Arkansas (type locality: St. Louis, Mo.).
- paucispina* Sargent in Bot. Gaz. XXXV. 391 (1903).—**Tenuifoliae.**—Illinois (type locality: Maywood).
- pausiaca* Ashe in Ann. Carnegie Mus. I. 390 (1902).—**Punctatae.**—New York, Pennsylvania (type locality not given).
- Pearsonii* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 10 (1900).—
Flavae?—North Carolina (type locality: King's Mountain). (?)
- Pechiana* Sargent in Rep. Missouri Bot. Gard. XXII. 78 (1912).—
Virides.—Illinois, Missouri (type locality: Hannibal, Mo.).
- Peckietta* Sargent in Rhodora, VII. 174 (1905).—**Macracanthae.**—New York (type locality: Piseco).
- Peckii* Sargent in Rhodora, V. 63 (1903).—**Intricatae.**—New York (type locality: Lansingburg).
- pedicellata* Sargent in Bot. Gaz. XXXI. 226 (1901).—**Coccineae.**—Ontario, New York, Pennsylvania (type locality: Rochester, N. Y.).
- pedicellata* var. *Ellwangeriana* (Sarg.) Eggleston in Rhodora, X. 82 (1908) = *C. Ellwangeriana*.
- pedicellata* var. *gloriosa* Sargent in Jour. Arnold Arb. I. 254 (1920).—
Coccineae.—New York (type locality: Rochester).
- pelacris* Sargent in N. Y. State Mus. Bull. CLXVII. 82 (1913).—**Pruinosaes.**—New York (type locality: Olean).
- pellecta* Sargent in N. Y. State Mus. Bull. CXXII. 85 (1908).—
Pruinosaes.—New York (type locality: Coopers Plains).
- pellucidula* Sargent in Rhodora, VII. 179 (1905).—**Macracanthae.**—Connecticut (type locality: Litchfield).
- penita* Beadle in Biltmore Bot. Studies, I. 57 (1902).—
Virides.—Tennessee (type locality: Chattanooga).
- pennsylvanica* Ashe in Ann. Carnegie Mus. I. 394 (1902).—
Molles.—Pennsylvania (type locality: Berks County).

CRATAEGUS

- Pennypackeri Sargent in Bot. Gaz. xxxv. 100 (1903).—**Crus-galli.**—Delaware (type locality: Stanton Station).
- pentandra Sargent in Rhodora, III. 25 (1901).—**Tenuifoliae.**—Quebec, New England, New York (type locality: New Haven, Vt.).
- pentaneura Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 12 (1903).—**Uniflorae.**—New Jersey, North Carolina (type locality not given). (?)
- pentasperma Ashe in Bot. Gaz. xxxv. 436 (1903).—Group?—North Carolina (type locality: Franklin). (?)
- peoriensis Sargent in Bot. Gaz. xxxi. 5 (1901).—**Crus-galli.**—Illinois (type locality: Princeville, Peoria County).
- pequotorum Sargent in Rhodora, v. 55 (1903).—**Pruinosae.**—Connecticut (type locality: Mumford's Point, Groton).
- peramoena Sargent in Ontario Nat. Sci. Bull. iv. 89 (1908).—**Macracanthae.**—Ontario (type locality: London).
- perampla Sargent in Rep. Geol. Surv. Mich. 1906, 525 (1907).—**Pruinosae.**—Michigan (type locality: Port Huron).
- perjucunda Sargent in Ontario Nat. Sci. Bull. iv. 27 (1908).—**Pruinosae.**—Ontario, New York (type locality: London, Ont.).
- perlaeta Sargent in Rep. Geol. Surv. Mich. 1906, 541 (1907).—**Tenuifoliae.**—Michigan (type locality: Grand Rapids).
- perlevis Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 48 (1904).—**Tenuifoliae.**—Pennsylvania (type locality: Kutztown).
- permula Sargent in Rep. Missouri Bot. Gard. xix. 65 (1908).—**Crus-galli.**—Missouri, Arkansas (type locality: Monteer, Mo.).
- perrara Sargent in N. Y. State Mus. Bull. CLXVII. 103 (1913).—**Coccineae.**—New York (type locality: Chapinville).
- persimilis Sargent in Proc. Rochester Acad. Sci. iv. 94 (1903).—**Crus-galli.**—New York (type locality: Rochester).
- persistens Sargent, Trees and Shrubs, II. 233 (1913).—**Crus-galli.**—Known only in cultivation (type locality: Arnold Arboretum, from seed received from Museum d'Histoire Naturelle, Paris, of unknown origin).
- perspicabilis Sargent in N. Y. State Mus. Bull. CLXVII. 90 (1913).—**Silvicolae.**—New York (type locality: Salamanca).
- pertomentosa Ashe in Jour. Elisha Mitchell Sci. Soc. XVI. 70 (1900).—**Macracanthae.**—Missouri, Kansas (type locality: Johnson County, Kan.).
- pexa Beadle in Biltmore Bot. Studies, I. 116 (1902).—**Flavae.**—North Carolina (type locality: Salisbury).
- Phaenopyrum (L.f.) Medicus, Gesch. Bot. 83 (1793).—**Microcarpae.**—North Carolina, Tennessee, Kentucky, Illinois and Missouri (type locality not given).

CRATAEGUS

- phaneroneura Sargent in Jour. Arnold Arb. III. 2 (1922).—**Crus-galli.**—Missouri (type locality: Clarksville).
- philadelphica Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 588 (1905).—**Pruinosae.**—Pennsylvania (type locality: West Philadelphia).
- phlebodia Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 155 (1910).—**Crus-galli.**—Pennsylvania (type locality: Bedford Springs).
- picta Sargent in Ontario Nat. Sci. Bull. IV. 91 (1908).—**Macracanthalae.**—Ontario (type locality: Toronto).
- pilifera Sargent in Rep. Missouri Bot. Gard. XIX. 64 (1908).—**Crus-galli.**—Illinois, Missouri, Arkansas (type locality: Swan, Mo.).
- pilosa Sargent in Rhodora, v. 56 (1903).—**Pruinosae.**—Massachusetts (type locality: Lancaster).
- pinetorum Beadle in Bot. Gaz. XXX. 343 (1900).—**Pulcherrimae.**—Alabama (type locality: Albertsville).
- pinguis Sargent in Rep. Geol. Surv. Mich. 1906, 559 (1907).—**Anomalae.**—Michigan (type locality: Lowell, Kent County).
- Pinshow Hort. ex Nicholson in Hand-list Arb. Kew, pt. 1. 211 (1894), as synonym of *C. uniflora* Muench.
- Piperi Britton in Torreya, I. 55 (1901).—**Douglasianae.**—Washington, Oregon, Idaho (type locality: Pullman, Wash.).
- pisifera Sargent in Rhodora, VII. 163 (1905).—**Macracanthalae.**—Ontario, Vermont, Massachusetts (type locality: Cornwall, Vt.).
- placens Sargent, Trees and Shrubs, II. 243 (1913).—**Molles.**—Ontario, Michigan (type locality: Walpole Island, Lamberton County, Ont.).
- placiva Sargent in Ontario Nat. Sci. Bull. IV. 22 (1908).—**Pruinosae.**—Ontario, New York, Michigan (type locality: Buffalo, N. Y.).
- plana Sargent in N. Y. State Mus. Bull. CXXII. 45 (1908).—**Pruinosae.**—New York (type locality: Buffalo).
- platycarpa Sargent in Rep. Missouri Bot. Gard. XIX. 92 (1908).—**Pruinosae.**—Illinois, Missouri, Arkansas (type locality: Moark, Ark.).
- poliophylla Sargent in Jour. Arnold Arb. III. 185 (1922).—**Virides.**—Texas (type locality: Columbia).
- politae Sargent in Rhodora, v. 111 (1903).—**Coccineae.**—Massachusetts, Connecticut, New York (type locality: East Lyme, Ct.).
- polita var. *Tatnalliana* (Sarg.) Eggleston in Rhodora, x. 82 (1908) = *C. Tatnalliana*.
- polybracteata Ashe in Jour. Elisha Mitchell Sci. Soc. XVI. 79 (1900).—**Rotundifoliae.**—New York, Ohio (type locality: Sugar Grove, O.).
- polyclada Sargent in Rep. Missouri Bot. Gard. XXII. 72 (1912).—**Crus-galli.**—Missouri (type locality: Springfield).
- populifolia Walter, Fl. Car. 147 (1788) = *C. Phaenopyrum* Medic.
- populnea Ashe in Ann. Carnegie Mus. I. 395 (1902).—**Tenuifoliae.**—Pennsylvania (type locality: Berks County).

CRATAEGUS

- porracea Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 19 (1901).—
Tenuifoliae.—Pennsylvania, New York (type locality: Susquehanna, Pa.). (?)
- orrecta Ashe in Ann. Carnegie Mus. I. 391 (1902).—**Punctatae.**—Pennsylvania (type locality: Pittsburgh).
- Porteri Britton in Bull. N. Y. Bot. Gard. I. 448 (1900).—**Pruinosae.**—Pennsylvania (type locality: Tannersville).
- praecincta Sargent in Ontario Nat. Sci. Bull. IV. 97 (1908).—**Macracanthae.**—Ontario (type locality: Toronto).
- praecoqua Sargent in Rhodora, v. 167 (1903).—**Rotundifoliae.**—Quebec, northern New England and New York, west to Wisconsin and northern Illinois (type locality: Crown Point, N. Y.).
- praecox Sargent in Rhodora, III. 27 (1901) = C. praecoqua Sarg.
- praestans Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 163 (1910).—**Punctatae.**—Pennsylvania (type locality: Nine Mile Run, near Pittsburgh).
- praetermissa Sargent in Rhodora, VII. 213 (1905).—**Rotundifoliae.**—Vermont (type locality: Ferrisburg).
- pratensis Sargent in Bot. Gaz. XXXI. 6 (1901).—**Punctatae.**—Illinois (type locality: Wady Petra).
- premora Ashe in Ann. Carnegie Mus. I. 391 (1902)—**Intricatae.**—Pennsylvania (type locality: eastern part of Berks County).
- Pringlei Sargent in Rhodora, III. 21 (1901).—**Coccineae.**—Ontario, New England, New York, Wisconsin, northern Illinois and Ohio (type locality: Charlotte, Vt.).
- Pringlei var. *exclusa* (Sarg.) Eggleston in Rhodora, x. 82 (1908) = C. *exclusa*.
- Pringlei var. *lobulata* (Sarg.) Eggleston in Rhodora, x. 82 (1908) = C. *lobulata*.
- prinoides Sargent in Ontario Nat. Sci. Bull. IV. 81 (1908).—**Macracanthae.**—Ontario (type locality: Sarnia).
- prismaticæ Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 15 (1901).—Group?—North Carolina (type locality: Madison County). (?)
- priva Ashe in Bull. Charleston Mus. XII. 42 (1916).—Group?—South Carolina (type locality: "south end of Piedmont"). (?)
- procax Ashe in Bull. Charleston Mus. XII. 42 (1916).—**Crus-galli.**—South Carolina (type locality not given).
- procera Sargent, Trees and Shrubs, II. 71 (1908).—**Pruinosae.**—New York, Pennsylvania (type locality: Hemlock, N. Y.).
- Proctoriana Sargent in N. Y. State Mus. Bull. CLXVII. 110 (1913).—**Rotundifoliae.**—New York (type locality: Utica).
- prominens Sargent in Ontario Nat. Sci. Bull. IV. 23 (1908).—**Pruinosae.**—Ontario, New York (type locality: Toronto, Ont.).

CRATAEGUS

- promissa* Sargent in N. Y. State Mus. Bull. cxxii. 50 (1908).—**Silvicolae.**—Ontario, New York, Vermont, Michigan (type locality: Niagara Falls, N. Y.).
- prona* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 17 (1903).—**Silvicolae.**—Michigan, Ohio (type locality: Port Huron, Mich.).
- propensa* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 214 (1910).—**Tenuifoliae.**—Pennsylvania (type locality: Keyser Valley, Scranton).
- propinqua* Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 53 (1904).—**Tenuifoliae.**—Ohio (type locality: Milan).
- propixa* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 248 (1910).—**Macracanthalae.**—Pennsylvania (type locality: Scranton).
- propria* Sargent in Rhodora, vii. 214 (1905).—**Rotundifoliae.**—Massachusetts (type locality: South Lancaster).
- pruinosa* (Wendl.) K. Koch, Hort. Dendr. 168 (1853).—**Pruinosae.**—Ontario, New England and New York, south to Delaware, west to New York, Ohio, Indiana and northern Illinois (type locality of *Mespilus pruinosa* Wendland not given).
- pruinosa* var. *conjuncta* (Sarg.) Eggleston in Rhodora, x. 81 (1908) = C. *conjuncta*.
- pruinosa* f. *dissona* (Sarg.) Eggleston in Rhodora, x. 81 (1908) = C. *dissona*.
- pruinosa* var. *latisepala* (Ashe) Eggleston in Rhodora, x. 81 (1908) = C. *latisepala*.
- pruinosa* var. *philadelphica* (Sarg.) Eggleston in Rhodora, x. 81 (1908) = C. *philadelphica*.
- pruinosa* var. *Porteri* (Britton) Eggleston in Rhodora, x. 81 (1908) = C. *Porteri*.
- pruinosa* × *punctata* Eggleston in N. Y. State Mus. Bull. ccxlvi-iv. 64 (1923). (?)
- pruinosa* × *straminea* Eggleston in N. Y. State Mus. Bull. ccliv. 422 (1924). (?)
- prunellifolia* Bosc in DeCandolle, Prodr. II. 627 (1825) = C. *prunifolia* Pers. ?
- prunifolia* (Marsh.) Persoon, Syn. II. 37 (1807).—**Macracanthalae?**—Known only in cultivation.
- prunifolia* var. *ingestria* (Lodd.) Loudon, Arb. Brit. II. 821 (1838) = C. *prunifolia* Pers. (?)
- puberis* Sargent in N. Y. State Mus. Bull. cxxii. 73 (1908).—**Rotundifoliae.**—New York (type locality: Belfast).
- pubescens* (H.B.K.) Steudel, Nomencl. Bot. ed. 2, 433 (1841).—Mexico (type locality: Real de Moran, Hidalgo).
- pubescens* var. *Botterii* Eggleston in Bull. Torrey Bot. Club, xxxvi. 506 (1909).—Mexico (type locality: Orizaba). (?)
- pubescens* f. *Humboldtii* Stapf in Kew Bull. Misc. Inform. 1914, 8, 293 (1914).—Mexico (type region: San Luis Potosi to Federal Dist.). (?)

CRATAEGUS

- pubescens f. stipulacea (Lodd.) Stapf in Kew Bull. Misc. Inform. 1914, 8, 293 (1914).—Mexico (type region: Federal District).
- pubifolia Ashe in Bull. N. Car. Exper. Sta. CLXXV. 114 (1900).—**Macracthae.**—Michigan, Indiana, Iowa (type locality: “Middle Iowa”).
- pubipes Ashe in Bull. N. Car. Exper. Sta. CLXXV. 114 (1900).—**Macracthae.**—Minnesota (type locality: “northeastern Minnesota”).
- pudens Sargent in Rep. Missouri Bot. Gard. XIX. 122 (1908).—**Macracthae.**—Missouri (type locality: Monteer).
- pulcherrima Ashe in Jour. Elisha Mitchell Sci. Soc. XVI. 77 (1900).—**Pulcherrimae.**—Florida, Georgia (type locality not given).
- pulchra Sargent in N. Y. State Mus. Bull. CXXII. 42 (1908).—**Pruinosae.**—New York, Pennsylvania (type locality: Niagara Falls, N. Y.).
- pulla Beadle in Biltmore Bot. Studies, I. 96 (1902).—**Flavae.**—Mississippi (type locality: Columbus). (?)
- pumila Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 612 (1905).—**Tenuifoliae.**—Pennsylvania (type locality: Sellersville).
- punctata Jacquin, Hort. Vind. I. 10 (1770).—**Punctatae.**—Southeastern Canada, New England and New York, south to North Carolina and Georgia, west to Minnesota, Iowa, Kentucky and Tennessee (type locality: cultivated in Botanical Garden, Vienna).
- punctata var. aurea Aiton, Hort. Kew. II. 170 (1789).—Ontario, New York, Pennsylvania, Ohio (type locality: Inverary Castle, Scotland ? “Cultivated 1746 by Archibald, Duke of Argyle”).
- punctata* β . *brevispina* (Douglas) Hooker, Fl. Bor. Am. I. 201 (1834), nomen nudum = *C. Douglasii* Lindl.
- punctata var. canescens Britton in Bull. Torrey Bot. Club. XXI. 231 (1894).—Pennsylvania (type locality: Easton).
- punctata lobata* Hort. ex Zabel in Beissner Schelle & Zabel, Handb. Laubholz-Ben. 171 (1903), nomen nudum.
- punctata var. microphylla Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 159 (1910).—Pennsylvania (type locality: Linesville).
- punctata var. mutabilis Gruber in Bull. Torrey Bot. Club. XXXII. 390 (1905).—Pennsylvania (type locality: Kutztown). (?)
- punctata* var. *rubra* Aiton, Hort. Kew. II. 170 (1789) = *C. punctata* Jacq.
- punctata* var. *xanthocarpa* (Medic.) Lavallée, Arb. Segrez. I. 53 (1880) = *C. punctata* var. *aurea* Ait.
- punctata* \times *succulenta* Eggleston in N. Y. State Mus. Bull. CCXLIII-IV. 63 (1923). (?)
- pura Sargent in Rep. Geol. Surv. Mich. 1906, 549 (1907).—**Coccineae.**—Michigan (type locality: Grand Rapids).
- pusilla Sargent in Rep. Geol. Surv. Mich. 1906, 553 (1907).—**Intricatae.**—Michigan (type locality: Grand Rapids).

CRATAEGUS

- puta Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 192 (1910).—**Silvicolae.**—Pennsylvania (type locality: Scranton).
- putata Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 244 (1910).—**Anomalae.**—Pennsylvania (type locality: Virginia, near Scranton).
- Putnamiana Sargent in Jour. Arnold Arb. IV. 102 (1923).—**Coccineae.**—Ohio (type locality: Marietta).
- pygmaea Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 650 (1905).—**Intricatae.**—Pennsylvania (type locality: Sellersville).
- pyracanthifolia* (Ait.) Sweet, Hort. Brit. ed. 3. 209 (1839) = *C. crus-galli* var. *pyracanthifolia*.
- pyracanthoides* Beadle in Biltmore Bot. Studies, I. 136 (1902).—**Crus-galli.**—Florida (type locality: Marianna).
- pyramidata* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 196 (1910).—**Silvicolae.**—Pennsylvania (type locality: Scranton).
- pyrifolia* Aiton, Hort. Kew. II. 168 (1789).—**Macracantheæ** = *C. tomentosa* L. ?
- pyrifolia* β. *aurantiaca* Lange, Rev. Gen. Crat. 82 (1897).—**Macracantheæ.**—(?)
- pyrifolia* var. *sylvestris* Schuette in Proc. Biol. Soc. Washington. XVI. 96 (1903).—Group?—Northwestern Wisconsin. (?)
- pyriformis* Britton in Bull. N. Y. Bot. Gard. I. 449 (1900) = *C. dispessa* Ashe.
- quaesita Beadle in Biltmore Bot. Studies, I. 33 (1901).—**Flavae.**—Florida (type locality: River Junction). (?)
- quercina Ashe in Jour. Elisha Mitchell Sci. Soc. XVIII. 27 (1902).—**Molles.**—Texas (type locality of *C. columbiana* Sarg., which this name replaces: Columbia).
- quinebaugensis Sargent in Rhodora, VII. 194 (1905).—**Pruinosæ.**—Connecticut (type locality: Hopeville).
- radians Sargent in N. Y. State Mus. Bull. CXXII. 64 (1908).—**Molles.**—New York (type locality: Rochester).
- radiata* Loddiges, Cat. 39 (1826).—Loudon, Arb. Brit. 819 (1838) as a synonym of *C. pyrifolia* Ait.
- radiata* Sargent in N. Y. State Mus. Bull. CXXII. 42 (1908).—**Pruinosæ.**—Ontario, New York (type locality: Buffalo, N. Y.).
- radina Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 203 (1910).—**Silvicolae.**—Pennsylvania (type locality: Schenley Park, Pittsburgh).
- radiosa Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 657 (1905).—**Macracantheæ.**—Pennsylvania (type locality: Upper Darby, Delaware County).
- raleighensis Ashe in Bull. N. Car. Exper. Sta. CLXXV. 112 (1900).—**Unifloræ.**—North Carolina (type locality: Raleigh).

CRATAEGUS

- ramosa Sargent in N. Y. State Mus. Bull. cxxii. 86 (1908).—**Pruinosae.**—New York (type locality: Coopers Plains).
- Randiana Sargent in Rhodora, v. 142 (1903).—**Tenuifoliae.**—Nova Scotia, Maine (type locality: Bar Harbor, Me.).
- rara Sargent, Trees and Shrubs, II. 241 (1913).—**Pruinosae.**—Connecticut (type locality: Hopeville).
- rava Beadle in Biltmore Bot. Studies, I. 91 (1902).—**Flavae.**—Florida (type locality: Tallahassee).
- Ravenelii Sargent in Bot. Gaz. xxxiii. 122 (1902).—**Flavae.**—South Carolina, Georgia (type locality: Augusta, Ga.).
- ravida Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. pt. 2, 6 (1901).—Group?—Tennessee (type locality: Clarkton). (?)
- recedens Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 161 (1910).—**Punctatae.**—Pennsylvania (type locality: Kutztown).
- recordabilis Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 199 (1910).—**Silvicolae.**—Pennsylvania (type locality: Kittanning).
- recta Sargent in N. Y. State Mus. Bull. cxxii. 97 (1908).—**Tenuifoliae.**—New York (type locality: Coopers Plains).
- recurva Beadle in Biltmore Bot. Studies, I. 106 (1902).—**Flavae.**—Florida (type locality: Ocala).
- redolens Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 14 (1903).—**Molles.**—Iowa, Wisconsin, Minnesota (type locality not given). (?)
- regalis Beadle in Biltmore Bot. Studies, I. 134 (1902).—**Crus-galli.**—Georgia, Alabama, Mississippi, Arkansas (type locality: Rome, Ga.).
- relicta Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 177 (1910).—**Pruinosae.**—Pennsylvania, Ohio (type locality: Orbisonia, Pa.).
- remota Sargent, Trees and Shrubs, II. 241 (1913).—**Pruinosae.**—Michigan (type locality: Grand Rapids).
- repentina Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 226 (1910).—**Rotundifoliae.**—Pennsylvania, Ohio (type locality: Bedford, Pa.).
- repulsans Sargent in N. Y. State Mus. Bull. cxxii. 107 (1908).—**Anomalae.**—New York (type locality: Coopers Plains).
- resecta Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 23 (1903).—**Intricatae.**—North Carolina (type locality: Marion). (?)
- reses Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 55 (1904).—**Intricatae.**—Pennsylvania (type locality: Berks County).
- resima Beadle in Biltmore Bot. Studies, I. 110 (1902).—**Flavae.**—Georgia (type locality: Albany).
- retrusa Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 49 (1904).—**Tenuifoliae.**—Michigan (type locality: Summerville).
- Reverchonii Sargent, Trees and Shrubs, I. 55 (1903).—**Crus-galli.**—Texas (type locality: Dallas).

CRATAEGUS

- rhodella* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 19 (1902).—
 Uniflorae.—North Carolina, Tennessee (type locality: Franklin,
 N. Car.).
- rhombifolia* Sargent in Rhodora, v. 183 (1903).—**Macracantheae.**—
 Canada, New England, New York (type locality: Trading Cove,
 Norwich, Ct.).
- Ridgwayi* Sargent in Jour. Arnold Arb. vi. 2 (1925).—**Molles.**—
 Southern Illinois (type locality: Olney).
- rigens* Beadle in Biltmore Bot. Studies, i. 54 (1902).—**Punctatae.**—
 Georgia, Alabama, Tennessee (type locality: Gadsden, Ala.).
- rigida* Sargent in Rep. Missouri Bot. Gard. xix. 99 (1908).—**Pruinosae.**—
 Missouri (type locality: Pacific).
- rimosa* Beadle in Biltmore Bot. Studies, i. 107 (1902).—**Flavae.**—
 Florida (type locality: Citra).
- riparia* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. 11 (1900).—**Silvi-**
 colae.—North Carolina (type locality: Swain County).
- rivalis* Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 580 (1905).—**Crus-**
 galli.—Pennsylvania, Ohio (type locality: Sellersville, Pa.).
- rivularis* Nuttall apud Torrey & Gray, Fl. N. Am. i. 464 (1840).—
 Douglasianae.—Idaho, Wyoming, Colorado, Utah, Nevada (type
 locality: “Oregon, along rivulets in the Rocky Mountains”).
- roanensis* Ashe in Bull. N. Car. Exper. Sta. clxxv. 114 (1900).—**Tenui-**
 foliae.—Virginia, North Carolina, West Virginia, Tennessee (type
 locality: Roan Mountain, N. Car.).
- Robbinsiana* Sargent in Rhodora, vii. 197 (1905) = *C. diffusa* Sarg.
- Robesoniana* Sargent in Rhodora, v. 110 (1903).—**Molles.**—Ontario,
 Massachusetts, New York, Pennsylvania (type locality: Lenox,
 Mass.).
- Robinsonii* Sargent in Rhodora, v. 184 (1903).—**Macracantheae.**—Nova
 Scotia (type locality: Lock Broom, near Pictou).
- robur* Beadle in Biltmore Bot. Studies, i. 69 (1902).—**Pulcherrimae.**—
 Florida (type locality: Tallahassee).
- robusta* Sargent in N. Y. State Mus. Bull. cxxii. 28 (1908).—**Crus-**
 galli.—New York (type locality: Niagara Falls).
- roribacca* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. 4 (1900).—
 Group?—North Carolina (type locality: “Oconaluftee River, in
 the mountains of North Carolina”). (?)
- Rosei* Eggleston in Bull. Torr. Bot. Club, xxxvi. 509 (1909).—Mexico
 (type locality: Alvarez, San Luis Potosi).
- rotunda* Sargent in Rep. Missouri Bot. Gard. xix. 59 (1908).—**Crus-**
 galli.—Missouri (type locality: Webb City).
- rotundata* Sargent in Ontario Nat. Sci. Bull. iv. 61 (1908).—**Rotundi-**
 foliae.—Ontario (type locality: London).

CRATAEGUS

- rotundifolia* Moench, Bäume Weiss. 29 (1785).—**Rotundifoliae.**—Nova Scotia and New England, west to New York, Pennsylvania, Michigan and Saskatchewan (type locality not given).
- rotundifolia* var. *aboriginum* Sargent in Jour. Arnold Arb. III. 196 (1922).—Quebec (type locality: Caughnawaga).
- rotundifolia* var. *Bicknellii* Eggleston in Rhodora, x. 79 (1908) = *C. Bicknellii*.
- rotundifolia* var. *chrysocarpa* (Ashe) Eggleston in Rhodora, x. 79 (1908) = *C. chrysocarpa*.
- rotundifolia* var. *Faxonii* (Sarg.) Eggleston in Rhodora, x. 79 (1908) = *C. Faxonii*.
- rotundifolia* a. *minor* Dippel, Handb. Laubholzk. III. 440 (1893). (?)
- rotundifolia* var. *pubera* Sargent in Rhodora XI. 183 (1909).—South-eastern Canada, New England and New York, west to Wisconsin and Minnesota (type locality not given).
- rotundifolia* f. *rubescens* Sargent in Jour. Arnold Arb. III. 197 (1922).—Quebec (type locality: St. Anne, Montreal Island).
- rotundifolia* b. *succulenta* (Schrad.) Dippel, Handb. Laubholzk. III. 441 (1893) = *C. succulenta*.
- rubella* Beadle in Bot. Gaz. XXX. 344 (1900).—**Intricatae.**—Virginia, West Virginia, North Carolina, Alabama and Tennessee (type locality: Valleyhead, Ala.).
- rubescens* Ashe in Bull. N. Car. Exper. Sta. CLXXV. 111 (1900).—**Intricatae.**—Alabama (type locality: Auburn). (?)
- rubicunda* Sargent in Proc. Rochester Acad. Sci. IV. 121 (1903).—**Tenuifoliae.**—Ontario, New York, Ohio (type locality: Rochester, N. Y.).
- rubicundula* Sargent, Trees and Shrubs, II. 7 (1907).—**Pruinosae.**—Missouri (type locality: Carondelet).
- rubrifolia* Sargent in Rep. Missouri Bot. Gard. XIX. 66 (1908).—**Crus-galli.**—Missouri, Kansas (type locality: Webb City, Mo.).
- rubrisepala* Sargent in Rep. Missouri Bot. Gard. XIX. 70 (1908).—**Crus-galli.**—Missouri (type locality: Joplin).
- rubrocarnea* Sargent in N. Y. State Mus. Bull. cv. Bot. IX. 55 (1906).—**Tenuifoliae.**—New York (type locality: North Albany).
- rubro-lutea* Sargent in N. Y. State Mus. Bull. CXXII. 88 (1908).—**Pruinosae**—New York (type locality: Coopers Plains).
- rudis* Sargent in Rep. Missouri Bot. Gard. XIX. 71 (1908).—**Crus-galli.**—Missouri (type locality: Monteer).
- rufipes* Ashe in Jour. Elisha Mitchell Sci. Soc. XX. 51 (1904).—**Tenuifoliae.**—Pennsylvania (type locality: Berks County).
- rufula* Sargent in Jour. Arnold Arb. I. 251 (1920).—**Aestivales.**—North Carolina, Florida, Georgia, Alabama (type locality not given).

CRATAEGUS

- rugosa* Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. V. (1900).—Group?—
North Carolina (type locality not given). (?)
- rupestris* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 27 (1902) = *C. densiflora* Sarg.*
- rupicola* Sargent in Rep. Missouri Bot. Gard. xix. 121 (1908).—**Macracantheae.**—Missouri (type locality: Swan).
- ruricola* Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 206 (1910).—
Silvicolae.—Pennsylvania (type locality: Chadsford.)
- russata* Sargent in N. Y. State Mus. Bull. clxvii. 85 (1913).—**Pruinosae.**—
New York (type locality: Painted Post, Steuben County).
- rustica* Beadle in Biltmore Bot. Studies, I. 122 (1902).—**Pruinosae.**—
North Carolina, West Virginia (type locality: Biltmore, N. Car.).
- Ruthiana* Sargent in Acad. Nat. Sci. Phila. lvii. 591 (1905).—**Pruinosae.**—
Pennsylvania (type locality: Durham).
- rutila* Sargent in Bot. Gaz. xxxv. 399 (1903).—**Macracantheae.**—Ohio,
Indiana, Wisconsin, Illinois (type locality: Chicago Heights, Ill.).
- saeva* Sargent in Ontario Nat. Sci. Bull. iv. 94 (1908).—**Macracantheae.**—
Ontario (type locality: Toronto).
- salicifolia* Medicus, Bot. Beob. II. 345 (1782) = *C. crus-galli* var.
pyracanthifolia Ait.?
- salicifolia* var. *Buistii* (Koch) Lange, Rev. Gen. Crat. 63 (1897), as
synonym of *C. Fontanesiana* a. *latifolia*. (?)
- saligna* Greene in Pittonia, III. 99 (1896).—**Brachyacantheae.**—Colorado
(type locality: “along lower Cimarron River”).
- sanguinea* Torrey & Gray, Fl. N. Am. I. 464 (1840), not Pallas.—
Macracantheae. (?)
- sanguinea* β. *Douglasii* (Lindl.) Torrey & Gray, Fl. N. Am. I. 464 (1840) =
C. Douglasii.
- Sargentii* Beadle in Bot. Gaz. xxviii. 407 (1899).—**Intricatae.**—Georgia,
Alabama, Tennessee (type locality: Valleyhead, Ala.).
- sarniensis* Sargent in Ontario Nat. Sci. Bull. IV. 34 (1908).—**Tenuifoliae.**—Ontario (type locality: Sarnia).
- saturata* Sargent in Proc. Acad. Nat. Sci. Phila. lvii. 619 (1905).—
Tenuifoliae.—Pennsylvania (type locality: Sellersville).
- Saundersiana* Sargent in Ontario Nat. Sci. Bull. IV. 66 (1908).—
Anomalae.—Ontario, New York, Ohio (type locality: London,
Ont.).
- sauratonae* Ashe in Bot. Gaz. xxviii. 270 (1899).—**Punctatae.**—North
Carolina (type locality: “along streams in the Sauraton Mountains”). (?)

* This name was proposed by Mr. Ashe to replace *Crataegus densiflora* Sargent for the reason, as he states, that the latter had been previously used in the genus (Desf. ex K. Koch, Dendr. I. 186, as a synonym). However, Desfontaines does not appear to have made such a combination, the error apparently having been made by Koch in transcribing from Spach (Hist. Vég. II. 88). Since the name *Crataegus densiflora* Desf. is thus not a valid name and has no standing, Sargent's specific name must be maintained.

CRATAEGUS

- saxatilis Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 640 (1905).—
Intricatae.—Pennsylvania (type locality: Newtown).
- scabera Sargent, Trees and Shrubs, II. 245 (1913).—**Macracantheae.**—
Missouri (type locality: Pleasant Grove).
- scabra Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 242 (1910).—
Intricatae.—Pennsylvania (type locality: Orbisonia).
- scabrida Sargent in Rhodora, III. 29 (1901).—**Anomalae.**—Southeastern
Canada, New England, New York, Pennsylvania (type locality:
Middlebury, Vt.).
- scabriuscula Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 605 (1905).—
Pruinosae.—Pennsylvania (type locality: Durham).
- schizophylla Eggleston in Bull. Torrey Bot. Club, XXXVIII. 243 (1911).—
Crus-galli.—Massachusetts (type locality: Martha's Vineyard).
- Schneckii Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. 8 (1900).—
Virides.—Illinois (type locality: Lawrenceville). (?)
- Schuettei Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 7 (1901).—
Tenuifoliae.—Wisconsin (type locality: Green Bay).
- Schweinitziana Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 652 (1905).—
Intricatae.—Pennsylvania (type locality: Sellersville).
- scitula Sargent in N. Y. State Mus. Bull. CLXVII. 84 (1913).—
Pruinosae.—New York (type locality: Chapinville).
- scopulorum Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 227 (1910).—
Rotundifoliae.—Pennsylvania (type locality: Campbell's Ledge,
Luzerne County).
- Searsii Sargent in Rhodora, VII. 183 (1905).—**Macracantheae.**—Massa-
chusetts (type locality: Southboro).
- seclusa Sargent, Trees and Shrubs, II. 239 (1913).—
Pruinosae.—Missouri, Arkansas (type locality: Pleasant Grove).
- seclusa Sargent in N. Y. State Mus. Bull. CLXVII. 89 (1913) = C. iterata
Sarg.
- secta Sargent in Rep. Missouri Bot. Gard. XIX. 83 (1908).—
Punctatae.—Missouri, Kansas, Arkansas (type locality: Carthage, Mo.).
- sectilis Ashe in Jour. Elisha Mitchell Sci. Soc. XIX. 23 (1903).—
Silvicolae.—North Carolina (type locality: Marion). (?)
- seducta Sargent in Rep. Missouri Bot. Gard. XXII. 79 (1912).—
Pruinosae.—Missouri, Arkansas (type locality: Noel, Mo.).
- Seelyana Sargent in Rhodora, VII. 209 (1905).—**Anomalae.**—Vermont
(type locality: Middlebury).
- segnis Beadle in Biltmore Bot. Studies, I. 32 (1901).—
Flavae.—Alabama (type locality: Greenville).
- sejuncta Sargent in N. Y. State Mus. Bull. CV. Bot. IX. 62 (1906).—
Coccineae.—Ontario, New England and New York, west to Ohio
and Indiana (type locality: Albany, N. Y.).

CRATAEGUS

- senta Beadle in Bot. Gaz. xxx. 341 (1900).—**Flavae.**—North Carolina (type locality: Biltmore).
- sequax Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 50 (1904).—**Tenuifoliae.**—Pennsylvania (type locality: Berks County).
- sera Sargent in Bot. Gaz. xxxiii. 115 (1902)—**Molles.**—Ontario, Michigan, Wisconsin, Illinois (type locality: Joilet, Ill.).
- serena Sargent in Rhodora, vii. 201 (1905).—**Tenuifoliae.**—Massachusetts (type locality: Lenox).
- sertata Sargent in Bot. Gaz. xxxv. 381 (1903).—**Coccineae.**—Wisconsin, Indiana, Illinois (type locality: “northern Illinois”).
- setosa Sargent in Rep. Missouri Bot. Gard. xix. 72 (1908).—**Crus-galli.**—Missouri (type locality: Grandin).
- severa Sargent, Trees and Shrubs, ii. 59 (1908).—**Crus-galli.**—Missouri (type locality: Grandin).
- sextilis Sargent in Bot. Gaz. xxxv. 390 (1903).—**Tenuifoliae.**—Illinois (type locality: Thatcher's Park, Cook County).
- Shaferi Ashe in Ann. Carnegie Mus. i. 397 (1902).—**Pruinosae.**—Pennsylvania (type locality: Carnot).
- Shallotte Ashe in Bull. N. Car. Exper. Sta. clxxv. 113 (1900).—**Macracantheae.**—North Carolina (type locality: Scranton). (?)
- sheridana Nelson in Bot. Gaz. xxxiv. 370 (1902).—**Macracantheae.**—Wyoming (type locality: Sheridan). (?)
- shirleyensis Sargent, Trees and Shrubs, ii. 244 (1913).—**Anomalae.**—Massachusetts (type locality: Shirley).
- sicca Sargent in Rep. Missouri Bot. Gard. xix. 101 (1908).—**Pruinosae.**—Missouri (type locality: Pacific).
- siderea Sargent in Proc. Acad. Nat. Sci. Phila. lxii. 212 (1910).—**Tenuifoliae**—Pennsylvania (type locality: Schenley Park, Pittsburgh).
- signata Beadle in Biltmore Bot. Studies, i. 42 (1901).—**Crus-galli.**—Alabama, Mississippi (type locality: Mobile, Ala.).
- silvestris Sargent in Ontario Nat. Sci. Bull. iv. 15 (1908).—**Punctatae.**—Ontario (type locality: London).
- silvicola Beadle in Bot. Gaz. xxviii. 414 (1899) = C. drymophila Sarg. silvicola var. *Beckwithae* (Sarg.) Eggleston in Rhodora, x. 81 (1908) = C. *Beckwithae* Sarg.
- simulans Sargent in N. Y. State Mus. Bull. cxxii. 125 (1908).—**Anomalae.**—New York (type locality: Hemlock Lake, Livingston County).
- simulata Sargent in Rep. Missouri Bot. Gard. xxii. 82 (1912).—**Macracantheae.**—Missouri (type locality: Joplin).
- singularis Sargent in N. Y. State Mus. Bull. cxxii. 106 (1908) = C. *asperifolia* Sarg.

CRATAEGUS

- sinistra Beadle in Biltmore Bot. Studies, I. 44 (1901).—**Crus-galli.**—Tennessee (type locality: West Nashville).
- sitiens Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 54 (1904).—**Pruinosa.**—Michigan, Ohio (type locality: St. Clair County, Mich.).
- Slavini Sargent in N. Y. State Mus. Bull. cxxii. 57 (1908).—**Tenuifoliae.**—New York (type locality: Buffalo).
- Smithii Sargent, Trees and Shrubs, I. 67 (1903).—**Uniflorae.**—Pennsylvania (type locality: Lowne's Run, Springfield).
- sodalis Beadle in Biltmore Bot. Studies, I. 80 (1902).—**Flavae.**—Alabama (type locality: Girard).
- sonnenbergensis Sargent in N. Y. State Mus. Bull. CLXVII. 120 (1922).—**Macracantha.**—New York (type locality: "Sonnenberg," Canandaigua, Ontario County).
- sordida Sargent in Bot. Gaz. xxxiii. 114 (1902).—**Punctatae.**—Missouri (type locality: Pleasant Grove).
- sordida var. villosa Sargent in Jour. Arnold Arb. vi. 2 (1925).—Missouri, Arkansas (type locality: Fulton, Ark.).
- sororia Beadle in Bot. Gaz. xxx. 336 (1900).—**Flavae.**—Florida, Georgia Alabama (type locality: Rome, Ga.).
- spathulata Michaux, Fl. Bor. Am. I. 288 (1803).—**Microcarpae.**—North Carolina to Florida, west to eastern Texas, Arkansas, Oklahoma and southwestern Missouri (type locality not given).
- spathulata var. *flavanthera* Sargent apud Palmer in Ann. Missouri Bot. Gard. III. 380 (1916), as "flavantha" = C. spathulata Michx.
- spatifolia Sargent in N. Y. State Mus. Bull. cxxii. 98 (1908).—**Tenuifoliae.**—New York (type locality: Coopers Plains).
- spatiosa Sargent in Rhodora, VII. 180 (1905).—**Macracantha.**—Connecticut (type locality: Mumford's Point, Groton).
- speciosa* Sargent, Trees and Shrubs, I. 65 (1903) = C. coccinoides Ashe.
- spinea Sargent in N. Y. State Mus. Bull. CLXVII. 122 (1913).—**Macracantha.**—New York (type locality: Coopers Plains).
- spinifera Sargent in N. Y. State Mus. Bull. cxxii. 111 (1908).—**Macracantha.**—New York (type locality: Coopers Plains).
- spinosissima* Loddiges, Cat. 40 (1826), nomen nudum.—Loudon, Arb. Brit. II. 849 (1838). As synonym of *C. flava lobata* Lindl.
- spinulosa Sargent in Rep. Missouri Bot. Gard. xix. 120 (1908).—**Macracantha.**—Missouri (type locality: Webb City).
- spissa Sargent in N. Y. State Mus. Bull. cxxii. 122 (1908).—**Rotundifoliae.**—New York (type locality: North Elba).
- spissiflora* Sargent in Proc. Rochester Acad. Sci. IV. 112 (1903) = C. Robesoniana Sarg.
- splendens* Loddiges, Cat. 40 (1826), nomen nudum.—Loudon, Arb. Brit. II. 820 (1838), as synonym of *C. crus-galli* var. *splendens* Ait.

CRATAEGUS

- splendida* Sargent in Ontario Nat. Sci. Bull. iv. 45 (1908).—**Coccineae.**—
Ontario (type locality: Sarnia).
- stellata* Sargent, Trees and Shrubs, II. 244 (1913).—**Intricatae.**—
Connecticut (type locality: East Lyme).
- stenophylla* Sargent in Ontario Nat. Sci. Bull. iv. 76 (1908).—**Macracanthalae.**—Ontario (type locality: Toronto).
- stenosepala* Sargent in Jour. Arnold Arb. III. 186 (1922).—**Virides.**—
Texas (type locality: Duke, Ft. Bend County).
- steubenensis* Sargent in N. Y. State Mus. Bull. cxxii. 103 (1908).—
Coccineae.—New York (type locality: Coopers Plains).
- Stevensiana* Sargent in Jour. Arnold Arb. IV. 99 (1923).—**Crus-galli.**—
Kansas, Oklahoma (type locality: Neodesha, Kas.).
- stipulacea* Loddiges, Cat. 40 (1826), nomen nudum.—Loudon, Arb.
Brit. II. 843 (1838), as a synonym of *C. mexicana* Moç. & Sessé.
- stipulacea* var. *microphylla* Hort. ex Nicholson in Hand-list Arb. Kew,
pt. 1, 197 (1894) as synonym of *C. crus-galli* var. *linearis* DC.
- stolonifera* Sargent in Bot. Gaz. xxxv. 109 (1903).—**Silvicolae.**—
New York, Pennsylvania, Delaware (type locality: Wilmington,
Del.).
- Stonei* Sargent in Rhodora, v. 62 (1903).—**Intricatae.**—Massachusetts
(type locality: Pelham).
- straminea* Beadle in Bot. Gaz. xxx. 345 (1900).—**Intricatae.**—Alabama,
Tennessee, Oklahoma (type locality: Valleyhead, Ala.).
- straminea* var. *Bissellii* (Sarg.) Eggleston in N. Y. State Mus. Bull.
CCXLIII-IV. 63 (1923) = *C. Bissellii*.
- stratfordensis* Sargent in Rhodora, VII. 178 (1905).—**Macracanthalae.**—
Connecticut (type locality: Stratford).
- Streeterae* Sargent in Proc. Rochester Acad. Sci. IV. 119 (1903).—
Tenuifoliae.—Southeastern Canada, New York, Michigan (type
locality: Rochester, N. Y.).
- strigosa* Sargent in N. Y. State Mus. Bull. cxxii. 51 (1908).—**Silvicolae.**—
New York: (type locality: Buffalo).
- strongylophylla* Sargent in Rep. Missouri Bot. Gard. xix. 44 (1908) =
C. crus-galli L.
- structilis* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 12 (1903).—
Macracanthalae.—Ontario, New York and Pennsylvania, west to
Michigan, Illinois and Missouri (type locality: Berks County, Pa.).
- suavis* Sargent in N. Y. State Mus. Bull. cxxii. 59 (1908).—**Tenuifoliae.**—Ontario, New York, Michigan, Ohio (type locality: Buffalo,
N. Y.).
- sublobulata* Sargent in Jour. Arnold Arb. III. 4 (1922).—**Crus-galli.**—
Texas (type locality: San Augustine).
- submollis* Sargent in Bot. Gaz. xxxi. 7 (1901).—**Molles.**—Southeastern
Canada, New England (type locality: Jamaica Plain, Mass.).

CRATAEGUS

- suborbiculata* Sargent in *Rhodora*, III. 72 (1901).—**Punctatae.**—Quebec (type locality: Caughnawaga).
- subpilosa* Sargent in *Jour. Arnold Arb.* III. 6 (1922).—**Crus-galli.**—Arkansas (type locality: Eureka Springs).
- subrotundifolia* Sargent in *Bot. Gaz.* xxxv. 394 (1903).—**Rotundifoliae.**—Wisconsin, Illinois (type locality: Lake Zurich, Ill.).
- subserrata* Bentham, *Pl. Hartweg.* 10 (1839) = *C. mexicana* Moç. & Sessé.
- subvillosa* Schrader ex Torrey & Gray, *Fl. N. Am.* I. 465 (1840), as a synonym of *C. coccinea* var. *mollis*.
- subviridis* Beadle in *Biltmore Bot. Studies*, I. 51 (1902).—**Virides.**—Florida (type locality: Chattahoochee).
- succincta* Sargent in *Rep. Missouri Bot. Gard.* xix. 78 (1908).—**Punctatae.**—Missouri (type locality: Pacific).
- succulenta* Schrader, *Ind. Sem. Götting.* (1823), ex Lange, *Rev. Gen. Crat.* 82 (1897).—**Macracantheae.**—Southeastern Canada, New England, New York, Pennsylvania.
- succulenta* var. *macracantha* (Lodd.) Eggleston in *N. Y. State Mus. Bull.* CCXLIII-IV. 65 (1923), as “*macrantha*,” = *C. macracantha*.
- succulenta* var. *rhombifolia* (Sarg.) Eggleston in *N. Y. State Mus. Bull.* CCXLIII-IV. 65 (1923) = *C. rhombifolia*.
- sucida* Sargent in *Rep. Missouri Bot. Gard.* xix. 80 (1908).—**Punctatae.**—Illinois, Missouri, Arkansas (type locality: Monteer, Mo.).
- superata* Sargent, *Trees and Shrubs*, II. 241 (1922).—**Pruinosae.**—Michigan (type locality: Grand Rapids).
- sutherlandensis* Sargent in *Jour. Arnold Arb.* III. 192 (1922).—**Virides.**—Texas (type locality: Sutherland Springs).
- sutherlandensis* var. *spinescens* Sargent in *Jour. Arnold Arb.* III. 193 (1922).—Texas (type locality: Sutherland Springs).
- swanensis* Sargent in *Jour. Arnold Arb.* III. 182 (1922).—**Punctatae.**—Missouri (type locality: Arnold Arboretum, cultivated from seed collected at Swan, Mo.).
- taetrica* Sargent in *Rep. Geol. Surv. Mich.* 1906, 541 (1907).—**Tenuifoliae.**—Michigan, Illinois (type locality: Port Huron, Mich.).
- tantula* Sargent in *Rep. Missouri Bot. Gard.* xix. 49 (1908).—**Crus-galli.**—Missouri (type locality: Webb City).
- tanophylla* Sargent in *Proc. Acad. Nat. Sci. Phila.* LXII. 247 (1910).—**Macracantheae.**—Pennsylvania, Ohio (type locality: Pittsburgh, Pa.).
- tarda* Sargent in *Bot. Gaz.* xxxv. 392 (1903).—**Tenuifoliae.**—Wisconsin, Illinois (type locality: Barrington, Ill.).
- tardiflora* Sargent in *Rep. Missouri Bot. Gard.* xix. 47 (1908).—**Crus-galli.**—Missouri, Arkansas (type locality: Christian County, Mo.).
- tardifolia* Ashe in *Bull. Charleston Mus.* XII. 41 (1916).—**Intricatae.**—South Carolina (type locality not given). (?)

CRATAEGUS

- tardipes* Sargent in Ontario Nat. Sci. Bull. iv. 51 (1908) = *C. Holmesiana* var. *tardipes* Sarg.
- Tatnalliana* Sargent in Bot. Gaz. xxxv. 106 (1903).—**Molles**.—Pennsylvania, Delaware (type locality: Wilmington, Del.).
- tecta* Beadle in Biltmore Bot. Studies, I. 26 (1901).—**Intricatae**.—Alabama (type locality: Marshall County).
- tenax* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 21 (1902).—**Punctatae**.—Ontario, Michigan (type locality: Port Huron, Mich.).
- tenella* Ashe in Ann. Carnegie Mus. I. 388 (1902).—**Tenuifoliae**.—Connecticut, New York, Pennsylvania, Delaware (type locality: Delaware County, Pa.).
- tenera* Ashe in Jour. Elisha Mitchell Sci. Soc. xx. 52 (1904).—**Tenuifoliae**.—Ohio (type locality: Sandusky).
- tennowana* Nelson in Bot. Gaz. liv. 408 (1912).—**Douglasianae**.—Idaho (type locality: Falk's Store, Canyon County). (?)
- tenuifolia* Britton in Bull. N. Y. Bot. Gard. I. 448 (1900).—**Tenuifoliae**.—Virginia (type locality: "south fork of Holston River").
- tenuiloba* Sargent in Proc. Rochester Acad. Sci. iv. 122 (1903).—**Tenuifoliae**.—New York (type locality: Rochester).
- tenuirama* Ashe in Bull. Charleston Mus. xii. 41 (1916).—Group?—South Carolina (type locality: "piedmont of South Carolina"). (?)
- tenuis* Sargent in Rep. Missouri Bot. Gard. xix. 48 (1908).—**Crus-galli**.—Missouri (type locality: Pacific).
- tenuisepala* Sargent in Rep. Missouri Bot. Gard. xix. 68 (1908).—**Crus-galli**.—Missouri, Arkansas (type locality: Moark, Ark.).
- tenuispina* Sargent in Rep. Missouri Bot. Gard. xix. 58 (1908).—**Crus-galli**.—Missouri (type locality: Grandin).
- tenuissima* Sargent, Trees and Shrubs, II. 243 (1913).—**Intricatae**.—Pennsylvania (type locality: Scranton).
- teres* Beadle in Biltmore Bot. Studies, I. 43 (1901).—**Flavae**.—Alabama (type locality: Montgomery).
- tersa* Beadle in Biltmore Bot. Studies, I. 129 (1902).—**Crus-galli**.—Louisiana (type locality: Opelousas).
- tetrica* Beadle in Biltmore Bot. Studies, I. 46 (1901).—**Crus-galli**.—Tennessee (type locality: Nashville).
- texana* Buckley in Proc. Acad. Nat. Sci. Phila. 1861, 454 (1861).—**Molles**.—Texas (type region: "along Brazos and Colorado Rivers").
- Thayeri* Sargent in Rhodora, v. 113 (1903).—**Coccineae**.—Massachusetts (type locality: West Boylston).
- tiliaeefolia* K. Koch in Verh. Ver. Beförd. Gartenb. Preuss. n. ser. I. 247 (Weissd. 27) (1853) = *C. mollis* Scheele.
- tinctoria* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 27 (1902).—Group?—North Carolina, Tennessee (type locality not given). (?)

CRATAEGUS

tomentosa Linnaeus, Spec. I. 476 (1753).—**Macracantha**.—Ontario and New York, south to North Carolina and Georgia, west to Wisconsin, Iowa, Missouri, Arkansas and eastern Kansas (type region: "Virginia").

tomentosa var. *Chapmani* Beadle in Bot. Gaz. xxv. 360 (1898) = C. *Chapmanii* Ashe.

tomentosa s. *flabellata* (Bosc) Wood, Cl. Book, 330 (1861) = C. *flabellata*.

tomentosa var. *microcarpa* Chapman, Fl. ed. 3, 139 (1897). (Type locality: Rome, Georgia.) (?)

tomentosa var. *mollis* (Torr. & Gray) Gray, Man. ed. 5, 160 (1867) = C. *mollis*.

tomentosa β. *plicata* Wood, Cl. Book, 330 (1861) = C. *punctata* Jacq.?

tomentosa var. *punctata* (Jacq.) Gray, Man. ed. 2, 124 (1856) = C. *punctata*.

tomentosa var. *pyrifolia* (Ait.) Gray, Man. ed. 5, 150 (1867). (?)

tomentosa var. *Smithii* (Sarg.) Eggleston in Rhodora, x. 79 (1908) = C. *Smithii*.

torta Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 176 (1910).—**Pruinosae**.—Pennsylvania (type locality: Nine Mile Run, near Pittsburgh).

tortilis Ashe in Jour. Elisha Mitchell Sci. Soc. XVIII. 19 (1902).—Group?—Wisconsin (type locality: Milwaukee). (?)

tortuosa Sargent in N. Y. State Mus. Bull. cxxii. 47 (1908).—**Silvicolae**.—New York (type locality: Buffalo).

torva Beadle in Biltmore Bot. Studies, I. 130 (1902).—**Crus-galli**.—Georgia, Alabama, Louisiana (type locality: Birmingham, Ala.).

trachyphylla Sargent in Bot. Gaz. XXXV. 388 (1903).—**Tenuifoliae**.—Illinois (type locality: Mokena).

Tracyi Ashe apud Eggleston in Bull. Torrey Bot. Club, XXXVI. 639 (1909).—**Molles**.—Texas (type locality: "foothills of Davis Mountains").

trahax Ashe in Jour. Elisha Mitchell Sci. Soc. XIX. 27 (1903).—**Crus-galli**.—Pennsylvania, Indiana (type locality: Berks County, Pa.).

transmississippiensis Sargent in Jour. Arnold Arb. III. 7 (1922).—**Molles**.—Missouri, Arkansas (type locality: Marion County, Ark., near Cotter).

Treleasei Sargent, Trees and Shrubs, I. 63 (1903).—**Molles**.—Missouri (type locality: St. Francois County).

trianthophora Sargent, Trees and Shrubs, II. 11 (1907).—**Uniflorae**.—Missouri, Arkansas, Texas (type locality: Grandin, Mo.).

tribulosa Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 182 (1910).—**Pruinosae**.—Pennsylvania (type locality: Bedford).

triflora Chapman, Fl. ed. 2, Suppl. 2, 684 (1892).—**Triflorae**.—Georgia, Alabama, Mississippi (type locality: cliffs of the Coos River, Ga.).

CRATAEGUS

- trifoliata* Bosc ex DeCandolle, Prodr. II. 630 (1825), nomen nudum.
- trilobata* Loddiges, Cat. 46 (1836), nomen nudum.—Loudon, Arb. Brit. II. 824 (1838), "A hybrid raised from seed in the Hammersmith Nursery about 1820." (?)
- Triosteum* Gruber in Bull. Torr. Bot. Club, XXXII. 392 (1905).—**Punctatae**.—Pennsylvania (type locality: Moselem). (?)
- tripartita* Sargent in Jour. Arnold Arb. III. 188 (1922).—**Virides**.—Texas (type locality: Columbia).
- trisperma* Ashe in Jour. Elisha Mitchell Sci. Soc. XVII. pt. 2, 17 (1901).—Group?—North Carolina (type locality: between north and south forks of Toe River, Yancey County). (?)
- tristis* Beadle in Biltmore Bot. Studies, I. 84 (1902).—**Flavae**.—Georgia (type locality: Rome).
- triumphalis* Sargent, Trees and Shrubs, II. 236 (1913).—**Crus-galli**.—Arkansas (type locality: Fulton).
- truculenta* Sargent in N. Y. State Mus. Bull. CLXVII. 118 (1913).—**Macracanthae**.—New York (type locality: Belfast).
- truncata* Sargent in Rep. Missouri Bot. Gard. XIX. 57 (1908).—**Crus-galli**.—Missouri (type locality: Swan).
- tumida* Sargent, Trees and Shrubs, II. 240 (1913).—**Pruinosae**.—Missouri (type locality: Dexter).
- turbinata* Pursh, Fl. Am. Sept. II. 735 (1814) = *C. spathulata* Michx.
- Twiningii* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 230 (1910).—**Intricatae**.—Pennsylvania (type locality: Scranton).
- uber* Ashe in Jour. Elisha Mitchell Sci. Soc. XX. 47 (1904).—**Tenuifoliae**.—Michigan (type locality: Summerville).
- umbratilis* Sargent in Rhodora, VII. 192 (1905).—**Punctatae**.—Connecticut (type locality: Litchfield).
- umbrosa* Sargent in Rep. Missouri Bot. Gard. XIX. 106 (1908).—**Molles**.—Illinois, Missouri (type locality: Carondelet, Mo.).
- uncta* Sargent in N. Y. State Mus. Bull. CXXII. 91 (1908).—**Pruinosae**.—New York (type locality: Coopers Plains).
- uniflora* Muenchhausen, Hausv. v. 147 (1770).—**Uniflorae**.—New York to Florida, west to Kentucky and Mississippi (type locality not given).
- uniflora* var. *florida* (Lodd.) Zabel in Beissner, Schelle & Zabel, Handb. Laubholz-Ben. 170 (1903), nomen nudum. (?)
- uniflora* var. *grossulariifolia* (Lodd.) Zabel in Beissner, Schelle & Zabel, Handb. Laubholz-Ben. 170 (1903), nomen nudum. (?)
- unilateralis* Persoon, Syn. II. 37 (1807).—**Uniflorae**.—(?)
- uniqua* Sargent, Trees and Shrubs, II. 237 (1913).—**Crus-galli**.—Louisiana, Texas (type locality: Marshall, Tex.).
- uplandia* Sargent in Proc. Acad. Nat. Sci. Phila. LVII. 594 (1905).—**Pruinosae**.—Pennsylvania (type locality: Upper Darby).

CRATAEGUS

- urasina* Hort. ex Nicholson in Hand-list Arb. Kew, pt. 1, 203 (1894), as a synonym of *C. mollis* Scheele.
- urbana* Sargent in Rep. Geol. Surv. Mich. 1906, 556 (1907).—**Anomalae**.—Michigan (type locality: Grand Rapids).
- urbica* Sargent in N. Y. State Mus. Bull. CLXVII. 105 (1913).—**Molles**.—New York (type locality of *C. oblongifolia* Sarg., which this name replaces: Menands).
- uticaensis* Sargent in N. Y. State Mus. Bull. CLXVII. 99 (1913).—**Coccineae**.—New York (type locality: Utica).
- uvaldensis* Sargent in Jour. Arnold Arb. III. 195 (1922).—**Virides**.—Texas (type locality: Uvalde).
- vaga* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 249 (1910).—**Macracantheae**.—Pennsylvania (type locality: Bedford Springs).
- Vailiae* Britton in Bull. Torrey Bot. Club, xxiv. 53 (1897).—**Macracantheae**.—Virginia, North Carolina (type locality: Roanoke, Va.).
- valens* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 15 (1903).—**Molles**.—Illinois (type locality: "southern Illinois"). (?)
- valida* Beadle in Biltmore Bot. Studies, I. 77 (1902).—**Flavae**.—Georgia (type locality: Rockmart).
- vallicola* Sargent in Rep. Missouri Bot. Gard. xix. 74 (1908).—**Crus-galli**.—Ohio, Missouri (type locality: Barretts Station, St. Louis County, Mo.).
- varians* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 225 (1910).—**Rotundifoliae**.—Pennsylvania (type locality: Bedford Springs).
- vegeta* Sargent in Bot. Gaz. xxxv. 396 (1903).—**Macracantheae**.—Illinois (type locality: Calumet).
- vegrandis* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 202 (1910).—**Silvicolae**.—Pennsylvania (type locality: Scranton).
- velata* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 26 (1902).—**Tenuifoliae**.—North Carolina (type locality of *C. atropurpurea* Ashe, which this name replaces: Yancey County). (?)
- velutina* Sargent, Trees and Shrubs, II. 238 (1913).—**Virides**.—Louisiana, Arkansas, Texas (type locality: Fulton, Ark.).
- venosa* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 16 (1903).—**Molles**.—Illinois (type locality: "southern Illinois"). (?)
- venulosa* Sargent in Ontario Nat. Sci. Bull. IV. 96 (1908).—**Macracantheae**.—Ontario (type locality: London).
- venusta* Beadle in Bot. Gaz. xxx. 338 (1900).—**Intricatae**.—Alabama (type locality: Birmingham).
- venustula* Sargent in N. Y. State Mus. Bull. cxxii. 79 (1908).—**Macracantheae**.—Ontario, New York (type locality: Niagara Falls, N. Y.).
- verecunda* Sargent in Proc. Rochester Acad. Sci. IV. 109 (1903).—**Intricatae**.—Massachusetts, New York (type locality: Rochester, N. Y.).

CRATAEGUS

- verecunda* var. *gonocarpa* Peck in N. Y. State Mus. Bull. cxxxix. 33 (1910) = *C. Dodgei* Ashe.
- verna* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 16 (1903).—**Molles.**—Illinois, Missouri (type locality not given). (?)
- vernans* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 10 (1903).—**Macracanthalae.**—Virginia, West Virginia (type locality not given). (?)
- verrucalis* Peck in N. Y. State Mus. Bull. cxii. 123, 158 (1908).—**Rotundifoliae.**—New York (type locality: Herkimer County).
- verruculosa* Sargent, Man. Trees N. Am. 394 (1905).—**Punctatae.**—Missouri, Arkansas (type locality: Springfield, Mo.).
- versuta* Beadle in Biltmore Bot. Studies, i. 112 (1902).—**Flavae.**—Georgia (type locality: Albany).
- vesca* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 13 (1903).—**Molles.**—Iowa (type locality: Keokuk). (?)
- viaria* Beadle in Biltmore Bot. Studies, i. 101 (1902).—**Flavae.**—Florida (type locality: Jacksonville).
- viatica* Ashe in Ann. Carnegie Mus. i. 398 (1902).—**Pruinosae.**—Pennsylvania (type locality: Pittsburgh).
- viburnifolia* Sargent, Trees and Shrubs, ii. 145 (1911).—**Molles.**—Texas (type locality: Columbia).
- vicana* Beadle in Biltmore Bot. Studies, i. 104 (1902).—**Flavae.**—Florida (type locality: Tavares).
- vicina* Sargent in Rep. Missouri Bot. Gard. xix. 77 (1908).—**Punctatae.**—Missouri, Kansas (type locality: Webb City, Mo.).
- vicinalis* Beadle in Biltmore Bot. Studies, i. 124 (1902).—**Pruinosae.**—Georgia, Tennessee (type locality: Chattanooga, Tenn.).
- Victorinii* Sargent in Jour. Arnold Arb. iii. 203 (1922).—**Macracanthalae.**—Quebec (type locality: grounds of College Longueuil, opposite Montreal).
- villaris* Beadle in Biltmore Bot. Studies, i. 108 (1902).—**Flavae.**—Florida (type locality: Citra).
- villicarpa* Sargent in Rep. Missouri Bot. Gard. xix. 112 (1908).—**Intricatae.**—Missouri, Arkansas (type locality: Swan, Mo.).
- villiflora* Sargent, Trees, and Shrubs, ii. 61 (1908).—**Crus-galli.**—Missouri, Arkansas, Mississippi (type locality: Grandin, Mo.).
- viminea* Ashe in Jour. Elisha Mitchell Sci. Soc. xviii. 23 (1902).—Group?—North Carolina (type locality: on Buffalo Creek, Ashe County). (?)
- virella* Ashe in Ann. Carnegie Mus. i. 396 (1923).—**Pruinosae.**—Pennsylvania (type locality: Berks County).
- virgata* Ashe in Jour. Elisha Mitchell Sci. Soc. xix. 24 (1903).—**Intricatae.**—North Carolina (type locality: Elk Cross Roads). (?)
- virginiana* Hort. ex Loudon, Arb. Brit. ii. 842 (1838) as synonym of *C. virginica* Lodd.

CRATAEGUS

- virginica* Loddiges, Cat. 40 (1826), nomen nudum.—Loudon, Arb. Brit. II. 842 (1838).—**Flavae.**—“A native of Virginia.” (?)
- viridimontana* Sargent in Rhodora, VII. 204 (1905).—**Tenuifoliae.**—Vermont (type locality: East Middlebury).
- viridis* Linnaeus, Spec. I. 476 (1753).—**Virides.**—Virginia to Florida, west to Illinois, Missouri and eastern Kansas, Oklahoma and Texas (type region: “Virginia”).
- viridis* var. *nitida* Engelmann in Britton & Brown, Ill. Fl. II. 242 (1897) = C. *nitida*.
- virilis* Sargent in Ontario Nat. Sci. Bull. IV. 68 (1908).—**Anomalae.**—Ontario (type locality: Toronto).
- visenda* Beadle in Biltmore Bot. Studies, I. 79 (1902).—**Flavae.**—Florida (type locality: Bristol).
- vitellina* Ashe in Jour. Elisha Mitchell Sci. Soc. XVIII. 27 (1902).—Group?—North Carolina (type locality: Madison County). (?)
- vittata* Ashe in Jour. Elisha Mitchell Sci. Soc. XX. 50 (1904).—**Tenuifoliae.**—Pennsylvania (type locality: Berks County). (?)
- vivida* Sargent in Ontario Nat. Sci. Bull. IV. 47 (1908).—**Coccineae.**—Ontario, New York (type locality: Toronto, Ont.).
- vulsa* Beadle in Biltmore Bot. Studies, I. 39 (1901).—**Virides.**—Georgia, Alabama (type locality: Gadsden, Ala.).
- Warneri* Sargent in Jour. Arnold Arb. III. 184 (1922).—**Crus-galli?**—Texas (type locality: Palestine).
- Watsoniana* (Spach) Roemer, Fam. Nat. Syn. III. 117 (1847).—**Crus-galli.**—(?)
- Websteri* Sargent in Rhodora, VII. 215 (1905).—**Rotundifoliae.**—New Hampshire (type locality: Holderness).
- Wheeleri* Nelson in Bot. Gaz. XXXIV. 369 (1902).—**Douglasianae.**—Colorado (type locality: Sapinero). (?)
- Wheeleri* Sargent in Rep. Geol. Surv. Mich. 1906, 552 (1907) = C. *diversifolia* Sarg.
- Wilkinsoni* Ashe in Bot. Gaz. XXXV. 435 (1903).—**Crus-galli.**—Ohio, Kentucky (type locality: Mansfield, O.).
- Williamsii* Eggleston in Bull. Torrey Bot. Club, XXXVI. 641 (1909).—**Rotundifoliae.**—Montana (type locality: Columbia Falls).
- wilmorensis* Sargent in Proc. Acad. Nat. Sci. Phila. LXII. 168 (1910).—**Pruinosae.**—Pennsylvania (type locality: between Portage and Wilmore, Cambria County).
- Wootoniana* Eggleston in Torreya, VII. 236 (1908).—**Tenuifoliae.**—New Mexico (type localities: Socorro and Lincoln Counties). (?)
- xanthocarpos* Medicus, Gesch. Bot. 85 (1793) = C. *punctata* var. *aurea* Ait.
- xanthophylla* Sargent in N. Y. State Mus. Bull. CXXII. 48 (1908).—**Silvicolae.**—New York (type locality: Buffalo).

CRATAEGUS

yadkinensis Ashe in Jour. Elisha Mitchell Sci. Soc. xvii. 17 (1900).—

Flavae.—North Carolina (type locality: along Yadkin River, Roan County). (?)

Youngii Sargent in Jour. Arnold Arb. iv. 105 (1923).—Microcarpae.—

North Carolina, South Carolina (type locality: Greensboro, N. Car.).

NAMES EXCLUDED FROM THE GENUS

CRATAEGUS

Amelanchier Desfontaines, Hist. Arb. ii. 149 (1809) = *Amelanchier canadensis* Medic.

amoena Salisbury, Prodr. 357 (1796) = *Amelanchier canadensis* Medic.

arbutifolia Aiton, Hort. Kew. ed. 2, iii. 202 (1811) = *Heteromeles arbutifolia* Roem.

coronaria (L.) Salisbury, Prodr. 357 (1796) = *Malus coronaria* Mill.

densiflora Desfontaines apud K. Koch, Dendr. i. 186 (1869), as a synonym = *Heteromeles arbutifolia* Roem.

racemosa Lamarck, Encycl. i. 84 (1783) = *Amelanchier canadensis* Medic.

spicata Lamarck, Encyl. i. 84 (1873) = *Amelanchier spicata* K. Koch (in part).

SYNOPSIS OF THE GROUPS

Nutlets not pitted or deeply concave on ventral surfaces.

Leaves merely serrate or dentate or rarely slightly lobed on shoots.

Veins of leaves not deeply impressed; fruit hard and inedible, dotted or blotched but rarely punctate..... I. **Crus-galli**.

Veins of leaves deeply impressed; fruit becoming soft and edible.

II. **Punctatae**.

Leaves of flowering branches or some of them variously lobed or divided.

Fruit usually more than 10 mm. diameter (8–20 or rarely more or less); nutlets variously ridged or rounded on back.

Leaves of flowering branches usually conspicuously narrowed at base or attenuate into winged petioles.

Corymbs few-flowered (1–5 flowers or rarely a few more).

Calyx lobes small, not conspicuously serrate; fruit usually becoming soft and edible..... XVIII. **Aestivales**.

Calyx lobes large and foliaceous, conspicuously serrate; fruit usually hard and inedible (except in *C. austromontana*).

Corymbs 1–3-flowered; leaves relatively small, mostly spatulate or obovate..... XVII. **Uniflorae**.

Corymbs 3–5-flowered (or rarely a few more); leaves larger; corymbs villous or glandular but not conspicuously bracteate.

XVI. **Triflorae**.

Corymbs 5–10-flowered (or rarely a few more); inflorescence conspicuously glandular and bracteate..... XV. **Bracteatae**.

Corymbs many-flowered (5–20 flowers or more).

Inflorescence and petioles generally conspicuously glandular; leaves subcordiaceous.

Leaves relatively large, usually of ovate or oval type; branchlets not conspicuously zig-zag..... V. **Intricatae**.

Leaves relatively small, usually of obovate or elliptic type; branchlets often conspicuously zig-zag..... XIV. **Flavae**.

Inflorescence and petioles generally not conspicuously glandular; leaves thin or thick.

Fruit blue or black (stramineous in var. of *C. brachyacantha*); leaves relatively small, thick and glossy above, those of flowering branches crenate or finely serrate, variously cut or lobed on shoots.

XIX. *Brachyacanthae.*

Fruit yellow, orange or red; leaves not as above.

Fruit usually 10–16 mm. diameter; leaves firm or subcoriaceous, mainly of ovoid or elliptic type.....XII. *Rotundifoliae.*

Fruit usually 4–10 mm. diameter; leaves thin, mostly of oblong or oval type.

Leaves very variable, irregularly lobed or merely serrate; corymbs many (10–20)-flowered; fruit usually 5–10 mm. diameter.....III. *Virides.*

Leaves less variable, with shallow rounded lobes; corymbs 4–10-flowered; fruit usually 8–15 mm. diameter.

IV. *Pulcherrimae.*

Leaves of flowering branches usually not conspicuously narrowed, or at least not attenuate at base, mostly of broadly ovate type, petioles slender.

Fruit becoming soft and succulent, bright red or scarlet, usually less than 15 mm. diameter. Corymbs usually glabrous; leaves thin, glabrous beneath, often finely villous-scabrate above.....VI. *Tenuifoliae.*

Fruit usually hard and inedible, 10–20 mm. diameter.

Fruiting calyx with distinct neck; leaves usually glabrous above; fruit often angular and pruinose.....VIII. *Pruinosae.*

Fruiting calyx sessile; young leaves usually villous-scabrate above; fruit seldom angular or pruinose.....VII. *Silvicolae.*

Fruit firm but becoming edible, larger (15–25 mm. diameter); leaves firm, mostly of ovate or deltoid-cordate type.

Mature leaves usually glabrous; fruit usually oblong or obovoid; fruiting calyx small.....IX. *Coccineae.*

Mature leaves usually pubescent, firm, with thick petioles and veins; stamens 10–20; nutlets 3–5.....X. *Molles.*

Mature leaves usually glabrous; fruit subglobose, calyx much enlarged; stamens 20; nutlets 5.....XI. *Dilatatae.*

Fruit 4–6 mm. diameter, bright scarlet, becoming soft and succulent; flowers small, many in a corymb; nutlets very small, not rounded or ridged on back.

XIII. *Microcarpae.*

Nutlets pitted or deeply concave on ventral surfaces.

Fruit dark crimson or chestnut, becoming black when fully ripe; nutlets 5.

XX. *Douglasianae.*

Fruit orange or scarlet.

Nutlets 2–3, deeply pitted or concave; flowers in usually many-flowered clusters at ends of leafy branchlets; fruit usually less than 10 mm. diameter.

XXII. *Macracanthae.*

Nutlets 3–5, with shallow pits; flower usually fewer in cluster, on short terminal or lateral branchlets; fruit usually more than 10 mm. diameter.

XXI. *Anomalae.*

SYNOPTICAL TABLES OF SPECIES ARRANGED BY GROUPS AND FLORAL AND FOLIAGE CHARACTERS

I. *Crus-galli*

A. Anthers red, rose, pink or purple (AA and AAA see p. 82)

B. Stamens about 10 (5–15) (BB see p. 81)

C. Corymbs glabrous (CC see p. 80)

D. Leaves mostly spatulate, obovate or lance-elliptic (on vigorous shoots broader but distinctly longer than broad)

- accincta Sarg. Fruit narrow-oval, scarlet, 7–8 mm. diameter.
- armata Beadle (?) Fruit globose, dull red, 5–8 mm. diameter. Color of anthers unknown.
- attenuata Ashe. Fruit oblong, dull red or green mottled, 8–12 mm. diameter. Stamens sometimes 15 or more.
- bellica Sarg. Fruit short-oblong or subglobose, dull dark red, 8–10 mm. diameter. Corymbs sometimes slightly villous.
- calophylla Sarg. Fruit short-oblong or slightly oval, orange-red, dark dotted, 8–9 mm. diameter.
- crus-galli L. Fruit short-oblong or subglobose, dull red, 8–12 mm. diameter.
- Farwellii Sarg. Fruit oval or obovoid, crimson, 9–10 mm. diameter. Corymbs often villous.
- geneseensis Sarg. Fruit short-oblong, scarlet, dark dotted, 10–12 mm. diameter.
- hamata Sarg. Fruit short-oblong or ovoid, light orange-red or green, dark dotted, 11–12 mm. diameter.
- helvina Ashe. (?) Fruit obovoid or subglobose, light red, 10–14 mm. diameter. Color of anthers not given.
- immitis Ashe. (?) Fruit globose, dark red, green mottled, 9–12 mm. diameter. Color of anthers not given.
- infesta Sarg. Fruit oblong or oval, dull yellow or tinged with orange, 9–10 mm. diameter.
- macra Beadle. Fruit subglobose or oval, red, 5–8 mm. diameter.
- multispina Ashe. (?) Fruit oblong, dull red-brown, 7–9 mm. diameter. Color of anthers not given.
- operta Ashe. Fruit oblong, dark red, 10–12 mm. diameter. Corymbs sometimes slightly villous.
- permira Sarg. Fruit short-oblong, green, red tinged, dotted, 8–9 mm. diameter.
- pyracanthoides Beadle. Fruit globose, bright red, 5–8 mm. diameter.
- schizophylla Egglest. Fruit pyriform-ellipsoid, dark red, 7–8 mm. diameter.
- setosa Sarg. Fruit subglobose or depressed-globose, crimson, dotted, 9–11 mm. diameter. Corymbs sometimes slightly villous.
- tenuis Sarg. Fruit oval to ovoid, dull green or red tinged, pale dotted, 6–7 mm. diameter.
- trahax Ashe. Fruit globose or oblong, crimson, dark dotted or mottled, 8–10 mm. diameter.
- Wilkinsonii Ashe. Fruit oblong, dark red, 10–14 mm. diameter.
- DD. Leaves broader, obovate, oval or oblong (on vigorous shoots broadly oval to suborbicular)
- Canbyi Sarg. Fruit oblong to obovoid, dark crimson, punctate, 8–10 mm. diameter.
- cerasina Sarg. Fruit short-oblong, bright cherry-red, 8–10 mm. diameter. Stamens sometimes 15–20.

- discolor Sarg. Fruit short-oblong to subglobose, dull red, blotched with green, 8–10 mm. diameter.
- fecunda Sarg. Fruit short-oblong to subglobose, orange-red, dark dotted, 10–18 mm. diameter. Corymbs and leaves sometimes slightly villous.
- lawrencensis Sarg. Fruit subglobose, green tinged with red, about 5 mm. diameter. Stamens sometimes 15 or more.
- peoriensis Sarg. Fruit oblong to obovoid, bright scarlet, 7–10 mm. diameter.
- persimilis Sarg. Fruit oblong to subglobose, dark crimson, dotted, about 11 mm. diameter. Corymbs sometimes slightly villous.
- Reverchonii Sarg. Fruit subglobose, light scarlet, 10 mm. diameter.
- robusta Sarg. Fruit short-oblong to obovoid, crimson, pale dotted, 7–8 mm. diameter. Stamens sometimes 20.
- rubrifolia Sarg. Fruit subglobose, bright orange-red, 8–9 mm. diameter.
- tantula Sarg. Fruit subglobose or short-oblong, dark red, dark dotted, 7–8 mm. diameter.
- tenuisepala Sarg. Fruit short-oblong, orange-red, dark dotted, 8–9 mm. diameter. Stamens sometimes 15 or more.
- CC. Corymbs at flowering time villous or pubescent
D. Leaves mostly spatulate, obovate or lance-elliptic (on vigorous shoots broader but usually longer than wide)
- albicera Beadle. Fruit oblong, pale yellow, 8–9 mm. diameter.
- arta Beadle. Fruit subglobose to short-ovoid, yellow-green and orange-red, about 10 mm. diameter.
- bellica Sarg. Corymbs usually nearly or quite glabrous. Fruit characters as given above.¹
- Engelmannii Sarg. Fruit globose, bright orange-red, 8–10 mm. diameter.
- Farwellii Sarg. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.
- munita Sarg. Fruit subglobose or slightly longer than thick, scarlet, pale dotted, 8–9 mm. diameter.
- operta Ashe. Corymbs usually glabrous or nearly so. Fruit characters as given above.
- setosa Sarg. Corymbs usually nearly or quite glabrous. Fruit characters as given above.
- signata Beadle. (?) Fruit oval or oblong, red, punctate, 9–10 mm. diameter.
Color of anthers not given.
- sinistra Beadle. Fruit oval, reddish, green or yellow, 6–8 mm. diameter.
- tenuis Sarg. Corymbs glabrous or nearly so. Fruit characters as given above.
- torva Beadle. Fruit subglobose, color not given, 7–9 mm. diameter.
- DD. Leaves broader, mostly elliptic, oblong or oval (on vigorous shoots broadly oval to suborbicular)

¹ Referring in this as in the following instances to the characters given under the same species in one of the preceding subdivisions of this group.

fecunda Sarg. Corymbs usually glabrous or nearly so. Fruit characters as given above.

helderbergensis Sarg. Fruit short-oblong to obovoid, crimson, dotted, about 8 mm. diameter.

montivaga Sarg. Fruit short-oblong to ellipsoidal, orange-red, 6–7 mm. diameter.

palliatia Sarg. Fruit subglobose, dark red, pale dotted, 10 mm. diameter.

persimilis Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.

tetrica Beadle. Fruit subglobose, red or yellowish-green, about 10 mm. diameter.

triumphalis Sarg. Fruit oval or short-oblong, crimson or greenish-red, pale dotted, 8–10 mm. diameter.

vallicola Sarg. Fruit short-oblong, dull crimson blotched with green, pale dotted, 14–16 mm. diameter.

Warneri Sarg. Fruit ellipsoidal or subglobose, orange-red, 6–9 mm. diameter.

BB. Stamens about 20 (15–20)

C. Corymbs at flowering time glabrous or nearly so

D. Leaves of flowering branches mostly spatulate, obovate or lance-elliptic (on vigorous shoots broader but usually longer than wide)

Cocksii Sarg. Fruit short-oblong to slightly obovoid, crimson, 8–12 mm. diameter.

Fontanesiana Steud. Fruit short-oblong or oval, bright red, 8–10 mm. diameter.

leptophylla Sarg. Fruit short-oblong to slightly obovoid, light orange-red, 8–10 mm. diameter.

limnophylla Sarg. Fruit ellipsoidal, bright red, 6–7 mm. diameter.

phlebodia Sarg. Fruit short-oblong, dull scarlet blotched with green, pale dotted, 8–11 mm. diameter.

DD. Leaves of flowering branches broader, mostly elliptic, oblong or oval (on vigorous shoots broadly oval to suborbicular)

attenuata Ashe. Stamens usually 10; fruit characters as given above.

Bushii Sarg. Fruit short-oblong, green tinged with red, 6–8 mm. diameter.

cerasina Sarg. Fruit short-oblong, bright cherry-red, pale dotted, 8–10 mm. diameter. Stamens sometimes only 10.

intermixta Sarg. Fruit subglobose to slightly obovoid, dark orange-red, 10–12 mm. diameter.

lawrencensis Sarg. Stamens sometimes 15 or less. Fruit characters as given above.

Pennypackeri Sarg. Fruit oblong, bright crimson, pale dotted, 8–10 mm. diameter. Corymbs sometimes slightly villous.

persimilis Sarg. Fruit oblong to subglobose, dark crimson, dark dotted, about 11 mm. diameter. Corymbs sometimes slightly villous.

robusta Sarg. Fruit short-oblong to obovoid, crimson, pale dotted, 7–8 mm. diameter. Stamens sometimes only 10.

- rotunda* Sarg. Fruit subglobose to depressed-globose, light cherry-red, 13–15 mm. diameter.
- sublobulata* Sarg. Fruit short-oblong, subglobose or ovoid, orange-red, 6–7 mm. diameter.
- tenuisepala* Sarg. Stamens sometimes only 10–15. Fruit characters as given above.
- CC. Corymbs at flowering time villous or pubescent
 D. Leaves of flowering branches mostly spatulate, obovate or lance-elliptic (on vigorous shoots broader but usually longer than wide)
- edita* Sarg. Fruit short-oblong, dull green tinged with red, 6–8 mm. diameter.
- limnophylla* Sarg. Corymbs slightly villous or glabrous. Fruit characters as given above.
- ohioensis* Sarg. Fruit ellipsoidal or ovoid, reddish-green, dotted, 7–8 mm. diameter.
- tersa* Beadle. Fruit subglobose, color not given, 8–9 mm. diameter.
 DD. Leaves broader, mostly elliptic, oblong or oval
- Pennypackeri* Sarg. Corymbs usually glabrous or nearly so. Fruit characters as given above.
- persimilis* Sarg. Cormbs sometimes glabrous or nearly so. Fruit characters as given above.
- AA. Anthers salmon or yellowish flushed with pink; stamens about 10
 B. Corymbs at flowering time glabrous
- consueta* Sarg. Fruit oval, orange-red, dark dotted, 8–10 mm. diameter.
- tardiflora* Sarg. Fruit short-oblong, dull red blotched with green, 10–12 mm. diameter.
 BB. Corymbs at flowering time villous or pubescent
- barbata* Sarg. Fruit short-oblong to subglobose, dark red, pale dotted, 8–9 mm. diameter.
- hirtella* Sarg. Fruit subglobose to short-oblong, green tinged with red, about 10 mm. diameter.
- AAA. Anthers white, cream or yellow
 B. Stamens about 10 (5–12) (BB see p. 84)
 C. Corymbs at flowering time glabrous or nearly so (CC see p. 84)
 D. Leaves of flowering branches mostly obovate, spatulate, narrowly oblong or lance-elliptic (on vigorous shoots broader but usually longer than broad)
- algens* Beadle. Fruit subglobose or ovoid, dull red or green and red, 8–10 mm. diameter.
- aliena* Sarg. Fruit short-oblong to slightly ovoid, crimson, pale dotted, 9–10 mm. diameter.
- arduennae* Sarg. Fruit oblong, dull dark crimson, pale dotted, 8–10 mm. diameter.
- armata* Beadle. (?) Fruit globose, dull red, 5–8 mm. diameter. Color of anthers unknown.
- barrettiana* Sarg. Fruit subglobose to slightly ovoid, dull red or bronze, 10–12 mm. diameter.

- Bartramiana** Sarg. Fruit subglobose or slightly longer than thick, crimson, dark dotted, 8–10 mm. diameter. Corymbs sometimes villous, **collicola** Ashe. (?) Fruit depressed-globose, orange-red or dull red, 8–11 mm. diameter. Color of anthers unknown.
- eburnea** Ashe. Fruit subglobose, dull dark red, about 13 mm. diameter.
- helvina** Ashe. (?) Fruit ovoid or subglobose, light red, 10–14 mm. diameter. Color of anthers unknown.
- immitis** Ashe. (?) Fruit globose, dark red or green mottled, 9–12 mm. diameter. Color of anthers unknown.
- monosperma** Sarg. Fruit oval to ovoid, dull red, pale dotted, 9–10 mm. diameter.
- multispina** Ashe. (?) Fruit oblong, dull red-brown, 7–9 mm. diameter. Color of anthers unknown.
- pachyphylla** Sarg. Fruit short-oblong, scarlet, dark dotted, 14–15 mm. diameter.
- panda** Ashe (not Beadle). (?) Fruit oval, dull red, 12–14 mm. long. Color of anthers unknown.
- Parkae** Sarg. Fruit oblong-ovoid, red, pale dotted, 8–9 mm. diameter.
- rudis** Sarg. Fruit oval to subglobose, scarlet, dark dotted, 8–10 mm. diameter.
- severa** Sarg. Fruit subglobose or depressed-globose, dark green tinged with red, 12–14 mm. diameter. (This species was described as having pink anthers, which was probably an error, as cultivated specimens and subsequent collections show anthers white or pale yellow.)
- Stevensiana** Sarg. Fruit short-oblong to slightly ovoid, green tinged with red, 4–5 mm. diameter.
- tenuispina** Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, 9–10 mm. diameter.
- truncata** Sarg. Fruit short-oblong to slightly ovoid, crimson, pale dotted, 8–10 mm. diameter.
- DD. Leaves of flowering branches broader, mostly obovate, elliptic or oblong (on vigorous shoots often suborbicular or broad as long)
- acutifolia** Sarg. Fruit oblong, bright scarlet, dark dotted, 8–10 mm. diameter.
- efferta** Sarg. Fruit short-oblong, broader than long, dark crimson, pale dotted, 12–13 mm. diameter.
- erecta** Sarg. Fruit short-oblong or subglobose, dull crimson, dark dotted, 7–10 mm. diameter.
- Palmeri** Sarg. Fruit subglobose, bright cherry-red, pale dotted, 8–9 mm. diameter.
- phaenoneura** Sarg. Fruit subglobose, dull red, 10–12 mm. diameter. Corymbs sometimes slightly villous.
- regalis** Beadle. Fruit oblong, color not given, about 8 mm. diameter.
- rubrisepala** Sarg. Fruit subglobose to short-oblong, orange-red, 7–8 mm. diameter.

CC. Corymbs at flowering time villous or pubescent

D. Leaves of flowering branches mostly obovate, spatulate, narrowly rhombic or lance-elliptic (on vigorous shoots broader but usually longer than broad)

Bartramiana Sarg. Corymbs usually glabrous or nearly so. Fruit characters as given above.

pilifera Sarg. Fruit short-oblong, dull red, pale dotted, 9–10 mm. diameter. **signata** Beadle. (?) Fruit oval or oblong, red, punctate, 7–10 mm. diameter. Color of anthers unknown.

villiflora Sarg. Fruit short-oblong, green tinged with crimson, 8–9 mm. diameter.

DD. Leaves of flowering branches broader, mostly broadly-elliptic, oval or oblong

paradoxa Sarg. Fruit oblong or ovoid, red, dark dotted, 6–7 mm. diameter.

phaneroneura Sarg. Corymbs sometimes glabrous. Fruit characters as given above.

BB. Stamens about 20 (15–20)

C. Corymbs at flowering time glabrous or nearly so

D. Leaves of flowering branches relatively narrow, mostly obovate, spatulate, oblong or lance-elliptic (on vigorous shoots broader but usually distinctly longer than broad)

araioclada Sarg. Fruit short-oblong, dull orange-red, 8–10 mm. diameter.

arborea Beadle. Fruit globose or subglobose, red or ruddy, 6–9 mm. diameter.

jasperensis Sarg. Fruit subglobose, crimson, pale dotted, 8–9 mm. diameter.

Mohrii Beadle. Fruit globose, red or greenish-red, black blotched, 7–9 mm. diameter. Corymbs sometimes slightly pilose.

polyclada Sarg. Fruit short-oblong or slightly ovoid, orange-red, pale dotted, 8–10 mm. diameter.

uniqua Sarg. Fruit short-oblong, dull red, 7–8 mm. diameter.

DD. Leaves of flowering branches broader, mostly elliptic, oval or oblong (on vigorous shoots often suborbicular or broad as long)

effulgens Sarg. Fruit short-oblong, scarlet, pale dotted, 10–12 mm. diameter.

insignis Sarg. Fruit subglobose, dull dark red, pale dotted, 14–15 mm. diameter. Corymbs sometimes slightly villous.

CC. Corymbs at flowering time villous or pubescent

D. Leaves of flowering branches relatively narrow, mostly obovate, spatulate, oblong or lance-elliptic (on vigorous shoots broader but usually distinctly longer than broad)

berberifolia T. & G. Fruit subglobose, orange, or reddish, 12–14 mm. diameter.

crocina Beadle. Fruit oval or oblong, yellow, 8–11 mm. diameter.

edura Beadle. Fruit subglobose, orange-yellow or tinged with red, 8–9 mm. diameter.

fera Beadle. Fruit globose or subglobose, bright red or scarlet, about 10 mm. diameter.

Mohrii Beadle. Corymbs usually glabrous or nearly so. Fruit characters as given above.

persistens Sarg. Fruit subglobose or short-oblong, crimson, dark dotted, about 15 mm. diameter.

subpilosa Sarg. Fruit subglobose or short-oblong, dull orange-red, 8–10 mm. diameter.

DD. Leaves of flowering branches broader, mostly oval, oblong or elliptic (on vigorous shoots often as broad as long, very large)

insignis Sarg. Corymbs usually glabrous or nearly so. Fruit characters as given above.

II. Punctatae

A. Anthers red, rose or pink (AA see p. 87)

B. Stamens about 10 (5–15)

C. Corymbs glabrous or nearly so

calvescens Sarg. Fruit subglobose, dark red, dark dotted, about 15 mm. diameter. Stamens sometimes 20.

cuneiformis Egglest. Fruit ellipsoid-pyriform, scarlet or dark red, about 10 mm. diameter. Corymbs sometimes slightly pubescent and stamens 10–20.

glabrifolia Sarg. (?) Fruit subglobose, angular, greenish red, dark dotted, 10–12 mm. diameter. Color of anthers unknown.

incerta Sarg. Fruit subglobose or short-oblong, crimson, 8–10 mm. diameter.

praestans Sarg. Fruit short-oblong to oval, dark red, minutely dark dotted, 8–10 mm. diameter.

pratensis Sarg. Fruit globose, bright scarlet, about 10 mm. diameter. Corymbs sometimes slightly pubescent.

sauratonae Ashe. (?) Fruit red, about 12 mm. diameter, shape not described. Number of stamens and color of anthers unknown.

tenax Ashe. Fruit subglobose, scarlet or mottled with yellow or olive, 14–15 mm. diameter. Stamens sometimes 20.

CC. Corymbs distinctly pubescent or villous

Brownietta Sarg. Fruit subglobose to slightly obovoid, crimson, pale dotted, about 15 mm. diameter.

cuneiformis Egglest. Corymbs usually glabrous. Fruit characters as given above.

desueta Sarg. Fruit short-oblong, scarlet, pale dotted, 8–10 mm. diameter.

pausiaca Ashe. Fruit oblong or slightly pyriform, crimson, 10–15 mm. diameter.

porrecta Ashe. Fruit globose or slightly oblong, dull red, 11–14 mm. diameter.

praestans Sarg. Corymbs usually glabrous or nearly so. Fruit characters as given above.

BB. Stamens about 20 (12–20)

C. Corymbs glabrous or nearly so

barbara Sarg. Fruit oblong-obvoid, crimson, 8–10 mm. diameter.

- calvescens* Sarg. Stamens sometimes only 10. Fruit characters as given above.
- compacta* Sarg. Fruit short-oblong, orange-red, pale dotted, 9–10 mm. diameter.
- Dewingii* Sarg. Fruit short-oblong, crimson, pale dotted, 10–12 mm. diameter.
- Eastmaniana* Sarg. Fruit short-oblong to slightly ovoid, dark crimson, pale dotted, 10–12 mm. diameter.
- grandis* Ashe. Fruit globular, bright crimson, 11–14 mm. diameter. Corymbs sometimes minutely pubescent.
- limatula* Sarg. Fruit short-oblong to subglobose, crimson, pale dotted, about 15 mm. diameter.
- nitidula* Sarg. Fruit short-oblong, dark yellow blotched with red, dark dotted, 10–12 mm. diameter.
- notabilis* Sarg. Fruit short-oblong to slightly ovoid, orange-red, dark dotted, 7–8 mm. diameter.
- sauratonae* Ashe.? Floral characters unknown. Fruit as described above.
- silvestris* Sarg. Fruit subglobose or short-oblong, dull orange-red, about 10 mm. diameter. Corymbs usually slightly villous.
- suborbiculata* Sarg. Fruit subglobose or rather longer than broad, dull red blotched with green, 12–15 mm. diameter.
- tenax* Ashe. Stamens sometimes 10–15. Fruit characters as given above.
- umbratilis* Sarg. Fruit short-oblong to subglobose, bright apple-green (unripe?), about 10 mm. diameter.
- CC. Corymbs distinctly villous or pubescent
- angustata* Sarg. Fruit subglobose or depressed-globose, crimson blotched with green, pale dotted, 12–14 mm. diameter.
- brazoria* Sarg. Fruit subglobose or longer than broad, bright canary-yellow, dark blotched, about 10 mm. diameter.
- celsa* Sarg. Fruit subglobose, crimson, pale dotted, 8–9 mm. diameter.
- dallasiana* Sarg. Fruit subglobose, dark dull red, 9–10 mm. diameter.
- Eatoniana* Sarg. Fruit subglobose to depressed-globose, bright cherry-red, 10–12 mm. diameter.
- florifera* Sarg. Fruit short-oblong, crimson with pale lenticles, 10–12 mm. diameter.
- grandis* Ashe. Corymbs sometimes glabrous. Fruit characters as given above.
- indicens* Ashe. (?) Fruit globose, dark dull red, olive mottled, 13–15 mm. diameter. Color of anthers unknown.
- punctata* Jacq. Fruit short-oblong to subglobose, dull red (yellow in var. *aurea*), pale dotted, 13–20 mm. diameter.
- recedens* Sarg. Fruit subglobose or slightly ovoid, dark red, green blotched or russet, scarcely punctate, 11–15 mm. diameter.
- silvestris* Sarg. Corymbs sometimes nearly glabrous. Fruit characters as given above.

- sordida* Sarg. Fruit subglobose, dark dull red, 10–12 mm. diameter.
- verruculosa* Sarg. Fruit subglobose, dark red, 12–16 mm. diameter.
- AA. Anthers white, cream or yellow
- B. Stamens about 10 (5–15)
 - C. Corymbs glabrous or nearly so
- glabrifolia* Sarg. (?) Color of anthers unknown. Fruit characters as given above.
- sauratonae* Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- CC. Corymbs villous or pubescent
- hirtiflora* Sarg. Fruit subglobose or depressed-globose, crimson blotched with green or pale lenticles, 15–16 mm. diameter.
- incaedua* Sarg. Fruit short-oblong, yellowish-red, pale dotted, 8–9 mm. diameter.
- latebrosa* Sarg. Characters of mature fruit not known.
- Lettermanii* Sarg. Fruit subglobose or depressed-globose, dark red with pale lenticles, 14–16 mm. diameter.
- macropoda* Sarg. Fruit subglobose or short-oblong, crimson, pale dotted, 8–10 mm. diameter.
- secta* Sarg. Fruit subglobose, crimson, pale dotted, 10–12 mm. diameter. Stamens usually only 5–8.
- BB. Stamens about 20 (12–20)
- C. Corymbs glabrous or nearly so
- fastosa* Sarg. Fruit subglobose or short-oblong, dull orange-red, pale dotted, 9–10 mm. diameter.
- sauratonae* Ashe. (?) Color of anthers and number of stamens unknown. Fruit characters as given above.
- CC. Corymbs villous or pubescent
- amnicola* Beadle. Fruit subglobose, red, 7–10 mm. diameter.
- collina* Chapm. Fruit globose, dull red, about 10 mm. diameter.
- indicens* Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- punctata* var. *aurea* Ait. Fruit short-oblong to subglobose, yellow, 13–20 mm. diameter.
- rigens* Beadle. Fruit subglobose or slightly pyriform, red, 9–12 mm. diameter.
- succincta* Sarg. Fruit subglobose, orange-red, 7–9 mm. diameter.
- sucida* Sarg. Fruit subglobose, bright red, 8–12 mm. diameter.
- vicina* Sarg. Fruit subglobose or slightly longer than wide, 10–12 mm. diameter.

III. Virides

- A. Anthers pink, rose or purple
- B. Corymbs glabrous
- antimima* Sarg. Fruit subglobose or slightly ovoid, dark orange-red, 7–8 mm. diameter. Corymbs sometimes slightly villous.

- arborescens Ell. Fruit subglobose, orange-red, 5–7 mm. diameter.
Corymbs sometimes villous.
- durifolia Ashe. (?) Fruit globular, dull red, 8–10 mm. diameter. Color of anthers unknown.
- enucleata Sarg. Fruit subglobose to short-oblong, orange-red, 7–8 mm. diameter.
- Pechiana Sarg. Fruit subglobose or slightly broader than long, orange-red, 8–9 mm. diameter.
- penita Beadle. Fruit globose or depressed-globose, red, 8–10 mm. diameter
- Schneckii Ashe. (?) Fruit nearly globular, bright red, about 8 mm. diameter. Color of anthers unknown.
- sutherlandensis Sarg. Fruit subglobose, orange-red, 7–8 mm. diameter.
- BB. Corymbs villous or pubescent
- antimima Sarg. Corymbs sometimes glabrous. Fruit characters as given above.
- ingens Beadle. Fruit globose or subglobose, red, 7–9 mm. diameter.
- micracantha Sarg. Fruit subglobose to short-oblong, bright orange-red, about 7 mm. diameter.
- paludosa Sarg. Fruit subglobose or short-oblong, orange-red, dark dotted, 5–8 mm. diameter.
- stenosepala Sarg. Fruit ellipsoidal to slightly ovoid, orange-red, 5–6 mm. diameter.
- subviridis Beadle. (?) Fruit globose, color not described, 5–7 mm. diameter. Color of anthers unknown.
- AA. Anthers cream-white or yellow
- B. Corymbs glabrous
- abbreviata Sarg. Fruit subglobose, dark red, 6–8 mm. diameter. Corymbs usually slightly villous.
- atrorubens Ashe. Fruit globular, dark red, 9–11 mm. diameter.
- blanda Sarg. Fruit subglobose to short-oblong, bright orange-red, pale dotted, about 8 mm. diameter.
- durifolia Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- glabriuscula Sarg. Fruit oblong to ovoid, dull orange-color with dark lenticles, 5–8 mm. diameter.
- interior Beadle. Fruit globose, bright red, 7–8 mm. diameter.
- lanceolata Sarg. Fruit short-oblong to slightly ovoid, dark red, 6–7 mm. diameter.
- larga Sarg. Fruit ovoid, scarlet, pale dotted, 7–8 mm. diameter.
- ludoviciensis Sarg. Fruit short-oblong, crimson, 9–10 mm. diameter.
- lutensis Sarg. Fruit orange-red, dark dotted, 4–5 mm. diameter.
- mitis Sarg. Fruit subglobose to short-oblong, dark crimson, dark dotted, 10–12 mm. diameter.
- nitens Sarg. Fruit subglobose, orange-red, 6–8 mm. diameter.

- nitida* Sarg. Fruit oblong, dull brick-red with dark lenticles, 8–15 mm. diameter.
- ovata* Sarg. Fruit subglobose or slightly longer than broad, orange-red, 5–6 mm. diameter.
- Schneckii* Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- tripartita* Sarg. Fruit short-oblong, yellow-green, 5–8 mm. diameter.
- viridis* L. Fruit subglobose or depressed-globose, bright scarlet or orange, 4–5 mm. diameter.
- BB. Corymbs villous or pubescent
- abbreviata* Sarg. Corymbs sometimes glabrous. Fruit characters as given above.
- amicalis* Sarg. Fruit subglobose, dull red, 6–8 mm. diameter.
- antiplasta* Sarg. Fruit subglobose, scarlet, 7–8 mm. diameter.
- atrorubens* Ashe. Corymbs sometimes glabrous. Fruit characters as given above.
- Dawsoniana* Sarg. Fruit ovoid, orange-red, 8–10 mm. diameter.
- poliophylla* Sarg. Fruit globose to short-oblong or ovoid, orange-red, 6–7 mm. diameter.
- subviridis* Beadle. (?) Color of anthers unknown. Fruit characters as given above.
- velutina* Sarg. Fruit subglobose, orange-red, pale dotted, 6–7 mm. diameter.

IV. *Pulcherrimae*

A. Anthers rose or purple

- abstrusa* Beadle. Fruit oval or short pyriform, color not given, 7–9 mm. diameter.
- ancisa* Beadle. Fruit globose or subglobose, red or ruddy, 10–12 mm. diameter.
- assimilis* Beadle. Fruit globose, color not described, 7–9 mm. diameter.
- austrina* Beadle. Fruit globose, green or greenish-yellow, about 10 mm. diameter.
- concinna* Beadle. Fruit subglobose, color not described, 7–8 mm. diameter.
- contrita* Beadle. Fruit subglobose yellow-green or orange, 7–9 mm. diameter.
- illustris* Beadle. Fruit depressed globose, color not described, about 10 mm. diameter.
- incilis* Beadle. Fruit subglobose, red or red and green, 7–9 mm. diameter.
- opima* Beadle. Fruit globose, bright red, 5–8 mm. diameter.
- pinetorum* Beadle. Fruit subglobose, red or ruddy, 7–10 mm. diameter.
- pulcherrima* Ashe. Fruit oval, yellow-green, 6–8 mm. diameter.
- AA. Anthers cream or yellow
- lenis* Beadle. Fruit oval or subpyriform, red or ruddy, 6–8 mm. diameter.

V. Intricatae

A. Anthers pink, rose or purple (AA see p. 91)

B. Stamens about 10 (7-12) (BB see p. 91)

C. Leaves at flowering time glabrous or nearly so

Bealii Sarg. Fruit short-oblong to subglobose, dark orange-red, dark dotted, about 10 mm. diameter. Stamens sometimes 16 or more.

Bissellii Sarg. Fruit pyriform, dull orange-red or green tinged, 9-10 mm. diameter.

Buckleyi Beadle. Fruit subglobose angled, red or russet-red, 8-12 mm. diameter.

Darlingtoniana Sarg. (?) Fruit short-oblong to depressed-globose, bright canary yellow, 12-14 mm. diameter. Flowers unknown.

definita Sarg. Fruit subglobose to short-oblong, greenish or greenish-yellow, about 10 mm. diameter. Young leaves sometimes slightly scabrate above.

diversifolia Sarg. Fruit ovoid, bright orange, dark dotted, 8-9 mm. diameter.

fortunata Sarg. Fruit short-oblong to slightly ovoid, bright canary-yellow, dark dotted, 12-14 mm. diameter.

fulva Sarg. Fruit globose or oblong-globose, red or orange, bronze tinged. Leaves sometimes slightly scabrate.

Hargeri Sarg. Fruit ovoid, dull orange-green, 10-12 mm. diameter.

inducta Ashe. Fruit globose or ovoid, dull green or russet, 8-10 mm. diameter.

Jenningsii Sarg. Fruit depressed-ovoid to short-oblong, russet-green, dark russet-red or bronze, 13-16 mm. diameter.

leioclada Sarg. Fruit subglobose to short-oblong, green or flushed with red, 10-12 mm. diameter.

luteola Sarg. Fruit subglobose to short-ovoid, light canary-yellow, 8-10 mm. diameter.

meticulosa Sarg. Fruit short-oblong to ovoid, orange-red, dark dotted, 10-12 mm. diameter.

neobushii Sarg. Fruit short-oblong to subglobose, orange-red, dark dotted, 10-12 mm. diameter.

padifolia Sarg. Fruit short-oblong, orange-red, dark dotted, 12-15 mm. diameter. Anthers described as yellow flushed with pink.

pagensis Sarg. Fruit short-oblong to slightly ovoid, dull orange-brown, about 10 mm. diameter.

Paineriana Sarg. Fruit subglobose to short-oblong, orange or orange-red, about 12 mm. diameter.

pygmaea Sarg. Fruit oblong-ovoid, dark crimson, pale dotted, 8-11 mm. diameter.

Schweinitziana Sarg. Fruit oblong, dark red, 8-9 mm. diameter.

straminea Beadle. Fruit pyriform, yellowish-green, 9-11 mm. diameter.

CC. Leaves at flowering time scabrate or villous above.

communis Beadle. Fruit globose or subglobose, red, 9–13 mm. diameter.
definita Sarg. Leaves sometimes glabrous above. Fruit characters as given above.

fulva Sarg. Leaves sometimes glabrous above. Fruit characters as given above.

infera Sarg. Fruit ovoid, dark reddish-orange, dark dotted, 10–11 mm. diameter.

minutiflora Sarg. Fruit short-oblong or subglobose, dull orange or orange and green with pale lenticles, about 10 mm. diameter.

Peckii Sarg. Fruit subglobose to short-oblong or ovoid, light yellow-green or red tinged, dark dotted, 8–10 mm. diameter.

scabra Sarg. Fruit subglobose or short-oblong, green, about 12 mm. diameter.

Stonei Sarg. Fruit ovoid, light yellow or greenish-yellow, 12–14 mm. diameter.

BB. Stamens about 20 (15–20)

C. Leaves at flowering time glabrous or nearly so

Bealii Sarg. Stamens usually about 10. Fruit characters as given above.

Darlingtoniana Sarg. (?) Flowers unknown; fruit characters as given above.

gilva Beadle. Fruit subglobose or short-pyriform, yellow or greenish-yellow, 6–8 mm. diameter.

gratiosa Sarg. Fruit short-oblong or slightly ovoid, green blotched with red, 8–10 mm. diameter. Young leaves sometimes slightly villous above.

macilenta Beadle. Fruit globose or subglobose, red, 7–9 mm. diameter.

mendosa Beadle. Fruit globose or subglobose, red, 8–10 mm. diameter.

pallens Beadle. Fruit subglobose to oval, yellow or greenish-yellow, 8–11 mm. diameter.

Sargentii Beadle. Fruit globose or subglobose, yellow or orange-yellow, flushed with red, 10–13 mm. diameter.

tecta Beadle. Fruit globose, red, 10–13 mm. diameter.

CC. Leaves at flowering time scabrate or villous

gratiosa Sarg. Young leaves sometimes glabrous or nearly so. Fruit characters as given above.

Harveyana Sarg. Fruit subglobose, orange, about 10 mm. diameter.

AA. Anthers white, cream or yellow

B. Stamens about 10 (7–15) (BB see p. 93)

C. Leaves at flowering time glabrous or nearly so

apposita Sarg. Fruit oblong-ovoid, reddish-green or yellowish-bronze, 9–11 mm. diameter.

Bartoniana Sarg. Fruit subglobose to short-oblong, dark crimson or blotched with green, 10–12 mm. diameter.

Boyntonii Beadle. Fruit globose or subglobose, yellow-green flushed with red, 10–15 mm. diameter.

- cornellii* Sarg. Fruit ovoid, light orange-yellow, dark dotted, 10–12 mm. diameter.
- cuprea* Sarg. Fruit subglobose to short-oblong, russet or dark red, 8–9 mm. diameter.
- Darlingtoniana* Sarg. (?) Flowers unknown. Fruit characters as given above.
- flavida* Sarg. Fruit subglobose to short-oblong, dull yellow or orange-yellow tinged with red, 10–12 mm. diameter.
- foetida* Ashe. Fruit subglobose, dull olive-green, red or russet mottled, 11–14 mm. diameter.
- fructuosa* Sarg. Fruit depressed-globose, deep orange-red, about 12 mm. diameter.
- intricata* Lange. Fruit subglobose to ellipsoidal, 7–9 mm. diameter.
- Kinzerae* Sarg. Fruit oblong-obvoid, russet-orange or red tinged, 10–12 mm. diameter.
- lentula* Sarg. Fruit depressed-globose, deep dull red, about 13 mm. diameter.
- nemoralis* Sarg. Fruit subglobose or slightly obovoid, bright clear red, about 10 mm. diameter.
- neocanbyi* Sarg. Fruit subglobose to short-oblong, dull orange to red-dish-orange, dark dotted, 12–15 mm. diameter.
- padifolia* Sarg. Stamens described as yellow flushed with pink (usually pink). Fruit characters as given above.
- pusilla* Sarg. Fruit short-oblong, crimson blotched with green, dark dotted, 12–14 mm. diameter. Leaves sometimes slightly scabrous above.
- resecta* Ashe. Fruit globular, bright scarlet, 14–17 mm. diameter.
- reeses* Ashe. Fruit subglobose, dark red or mottled with russet or green, 15–16 mm. diameter. Young leaves sometimes slightly scabrous.
- sexatilis* Sarg. Fruit short-oblong, dark orange or reddish-orange, dark dotted, 9–10 mm. diameter.
- stellata* Sarg. Fruit subglobose, dull reddish-green, 11–13 mm. diameter.
- tenuissima* Sarg. Fruit ovoid, dull red, 8–10 mm. diameter.
- Twiningii* Sarg. Fruit short-oblong to subglobose, green tinged with red, 10–12 mm. diameter.
- verecunda* Sarg. Fruit oblong to oblong-obvoid, crimson, about 7 mm. diameter.
- virgata* Ashe. Fruit depressed-globose, russet-red dotted with green and orange, 12–16 mm. diameter.
- CC. Leaves at flowering time scabrate or villous above
- abjecta* Sarg. Fruit subglobose or slightly longer than wide, greenish-orange, about 15 mm. diameter.
- alpista* Sarg. Fruit ovoid, dull red blotched with green, 10–12 mm. diameter.

biltmoreana Beadle. Fruit depressed-globose, green, yellow or orange, 10–15 mm. diameter.

callista Sarg. Fruit oblong, orange-red, 8–9 mm. diameter.

confusa Sarg. Fruit short-oblong to subglobose, russet-green or tinged with red, 12–17 mm. diameter.

contortula Sarg. Fruit subglobose, green blotched with red, about 10 mm. diameter.

Delosii Sarg. Fruit short-oblong to slightly ovoid, orange tinged with red, pale dotted, 10–12 mm. diameter.

modesta Sarg. Fruit subglobose, rarely oblong or pyriform, bright yellow or orange-red, dark mottled, 8–9 mm. diameter.

premora Ashe. Fruit globose, bright green, dark mottled, 10–12 mm. diameter.

pusilla Sarg. Leaves usually glabrous or nearly so. Fruit characters as given above.

reeses Ashe. Young leaves usually glabrous or nearly so. Fruit characters as given above.

villicarpa Sarg. Fruit short-oblong to subglobose, orange yellow or tinged with red, pale dotted, 10–12 mm. diameter.

BB. Stamens about 20 (15–20)

C. Leaves at flowering time glabrous or nearly so

alma Beadle. Fruit subglobose, red, 10–12 mm. diameter.

Darlingtoniana Sarg. (?) Flowers unknown. Fruit characters as given above.

eximia Beadle. Fruit globose, red, 8–10 mm. diameter.

inanis Beadle. Fruit short-oval or slightly pyriform, red or ruddy, 6–8 mm. diameter.

venusta Beadle. Fruit oval, greenish-red, dull red or bright red, 9–13 mm. diameter.

CC. Leaves at flowering time scabrate or villous above

Craytonii Beadle. Fruit oblong to ovoid, red or ruddy, 9–12 mm. diameter.

VI. Tenuifoliae

Anthers red, rose, pink or purple in this group.

A. Stamens normally 10 or less (AA see p. 97)

B. Leaves of flowering branches narrowed or attenuate at base (on vigorous shoots rounded to truncate) (BB see p. 95)

C. Corymbs glabrous or nearly so (CC see p. 95)

acuminata Sarg. Fruit short-oblong to subglobose, crimson, 8–9 mm. diameter.

acutiloba Sarg. Fruit oblong to ovoid, bright scarlet, 8–10 mm. diameter.

angustisepala Sarg. Fruit short-oblong to slightly ovoid, scarlet, about 10 mm. diameter.

antheina Sarg. Fruit short-oblong, scarlet, pale dotted, 9–12 mm. diameter.

- apiomorpha Sarg. Fruit obovoid, rarely short-oblong, bright reddish-purple, pale dotted, 8–12 mm. diameter. Corymbs sometimes slightly villous.
- ascendens Sarg. Fruit oblong or oblong-obovoid, dark crimson, 8–9 mm. diameter.
- bella Sarg. Fruit short-oblong to slightly obovoid, scarlet, 8–9 mm. diameter.
- Boothiana Sarg. Fruit short-oblong, bright orange, pale dotted, 8–9 mm. diameter.
- Collinsiana Sarg. Fruit short-oblong, dark crimson, about 8 mm. diameter.
- colorata Sarg. Fruit obovoid, crimson, pale dotted, about 12 mm. diameter.
- crudelis Sarg. Fruit oblong, crimson, pale dotted, 9–10 mm. diameter. Corymbs sometimes slightly villous.
- culta Sarg. Fruit short-oblong, ovoid or subglobose, crimson, pale dotted, 9–10 mm. diameter.
- cyanophylla Sarg. Fruit subglobose to short-oblong, reddish-purple with pale lenticles, 9–15 mm. diameter.
- delucida Sarg. Fruit oblong, bright scarlet, 9–10 mm. diameter. Leaves of terminal shoots sometimes subcordate.
- Eganii Ashe. Fruit bright red, oblong, 8–13 mm. diameter.
- exigua Ashe. Fruit obovoid, bright scarlet, 10–12 mm. diameter.
- firma Sarg. Fruit short-oblong to obovoid, crimson, 10–12 mm. diameter.
- flammata Sarg. Fruit short-oblong, scarlet, pale dotted, 8–10 mm. diameter.
- florea Sarg. Fruit oblong, scarlet, 10–12 mm. diameter.
- glaucomphyllea Sarg. Fruit short-oblong, rarely ovoid, bright scarlet, about 10 mm. diameter.
- Gruberi Ashe. Fruit globular-pyriform, light green with russet or scarlet blotches, about 11 mm. diameter.
- Hadleyana Sarg. Fruit short-oblong, scarlet, pale dotted, 9–10 mm. diameter. Corymbs sometimes slightly villous.
- leptopoda Sarg. Fruit obovoid, bright cherry-red, pale dotted, 8–9 mm. diameter.
- longipetiolata Sarg. Fruit oblong-obovoid, dark red, 8–9 mm. diameter.
- marcida Ashe. Fruit depressed-globose, dark crimson, 13–15 mm. diameter.
- merita Sarg. Fruit short-oblong to slightly obovoid, crimson, blotched with green and pale dotted, 9–10 mm. diameter.
- miniata Ashe. Fruit oblong, scarlet or mottled with olive or russet, 9–13 mm. diameter.
- nescia Sarg. Fruit short-oblong or slightly obovoid, scarlet, pale dotted, 9–10 mm. diameter.
- pallidula Sarg. Fruit short-oblong, scarlet, 7–8 mm. diameter.

- parviflora* Sarg. Fruit oblong-obvoid, crimson, pale dotted, about 10 mm. diameter.
- pastorum* Sarg. Fruit oblong, bright scarlet with pale lenticles, about 10 mm. diameter. Stamens sometimes 20.
- paucispina* Sarg. Fruit pyriform or globose-pyriform, crimson or purplish, pale dotted, 12–15 mm. diameter.
- pentandra* Sarg. Fruit dark crimson with pale lenticles, 12–15 mm. diameter.
- perlaevis* Ashe. Fruit pyriform, obovoid or rarely oblong, dark dull red blotched with orange or russet, 8–12 mm. diameter.
- porracea* Ashe. (?) Fruit oblong, dark red, 10–13 mm. diameter.
- propinqua* Ashe. Fruit globose, green, russet and dull red, 12–13 mm. diameter.
- pumila* Sarg. Fruit oblong, dark red, 8–10 mm. diameter.
- rubrocarnea* Sarg. Fruit short-oblong to subglobose, scarlet, dark dotted, 12–14 mm. diameter. Corymbs sometimes slightly villous.
- saturata* Sarg. Fruit short-oblong to subglobose, dark red, about 10 mm. diameter.
- sequax* Ashe. Fruit globose or subglobose, dark red or crimson blotched with green or russet, 12–15 mm. diameter.
- serena* Sarg. Fruit oblong, scarlet with pale lenticles, about 10 mm. diameter.
- siderea* Sarg. Fruit obovoid, yellow-green, becoming dark red, 9–10 mm. diameter.
- suavis* Sarg. Fruit short-oblong, orange-red, pale dotted, 8–9 mm. diameter.
- tarda* Sarg. Fruit oblong to subglobose, bright scarlet with pale lenticles, becoming crimson, 11–14 mm. diameter. Stamens sometimes 15–20.
- tenuifolia* Britton. Fruit pyriform, 8–9 mm. diameter. Color not described.
- viridimontana* Sarg. Fruit short-oblong, crimson, pale dotted, about 10 mm. diameter.
- Wootoniana* Egglest. (?) Fruit ellipsoidal, red, 6–10 mm. diameter. Color of anthers not described.
- CC. Corymbs sometimes slightly villous
- apiomorpha* Sarg. Fruit characters as given above.
- crudelis* Sarg. Fruit characters as given above.
- Hadleyana* Sarg. Fruit characters as given above.
- rubrocarnea* Sarg. Fruit characters as given above.
- BB. Leaves of flowering branches mostly not conspicuously narrowed at base, rounded to truncate (on vigorous shoots truncate to cordate)
- C. Corymbs glabrous
- blandita* Sarg. Fruit oblong, scarlet with pale lenticles, 8–9 mm. diameter.
- brumalis* Ashe. Fruit subglobose, dull, rather dark red, green mottled, 10–12 mm. diameter.

- Claytoniana* Sarg. Fruit subglobose to short-oblong, scarlet, about 10 mm. diameter.
- condensa* Sarg. Fruit ovoid, yellowish-red, 8–10 mm. diameter.
- Damei* Sarg. Fruit oblong to ovoid, bright cherry-red, minutely dotted, about 8 mm. diameter.
- demissa* Sarg. Fruit short-oblong to subglobose or slightly ovoid, scarlet, 8–9 mm. diameter.
- dissimilis* Sarg. Fruit short-oblong to ovoid, dull red, becoming dark red, about 12 mm. diameter.
- fucata* Sarg. Fruit short-oblong to oval, crimson, 8–9 mm. diameter.
- fucosa* Sarg. Fruit oblong, scarlet, pale dotted, 9–12 mm. diameter.
- genialis* Sarg. Fruit oblong, dark crimson, pale dotted, 8–10 mm. diameter.
- gracilipes* Sarg. Fruit narrow-ovoid, scarlet, 6–7 mm. diameter.
- Habereri* Sarg. Fruit oval to ovoid, crimson, pale dotted, about 5 mm. diameter.
- heidelbergensis* Sarg. Fruit ovoid, rarely short-oblong or globose, scarlet, blotched with russet, pale dotted, 12–15 mm. diameter.
- ignea* Sarg. Fruit short-oblong, scarlet, pale dotted, 8–9 mm. diameter.
- insignata* Sarg. Fruit short-oblong, dull red, 8–9 mm. diameter.
- insolita* Sarg. Fruit broad-ovoid, bright cherry-red, dark dotted, 12–13 mm. diameter.
- lata* Sarg. Fruit oblong, bright scarlet, about 7 mm. diameter.
- macrosperma* Ashe. Fruit somewhat longer than thick, dark but bright red, 13–18 mm. diameter.
- matura* Sarg. Fruit oblong, bright scarlet with pale lenticles, 12–15 mm. diameter.
- media* Sarg. Fruit ovoid, scarlet, 8–10 mm. diameter.
- modica* Sarg. Fruit globose, bright crimson, 8–10 mm. diameter.
- monstrata* Sarg. Fruit oblong, scarlet, pale dotted, 9–10 mm. diameter.
- Moyeriana* Sarg. Fruit ovoid to short-oblong, cherry-red, dark dotted, 10–11 mm. diameter.
- multifida* Ashe. (?) Fruit oblong, crimson, 9–11 mm. diameter.
- Napaea* Sarg. Fruit oblong-ovoid, dull red, 8–9 mm. diameter.
- ornata* Sarg. Fruit oblong, scarlet, 9–10 mm. diameter.
- otiosa* Ashe. Fruit oblong, russet becoming scarlet, 8–10 mm. diameter.
- pactilis* Ashe. (?) Fruit ovoid, scarlet, orange or russet, 11–12 mm. diameter.
- Paddockii* Sarg. Fruit short-oblong to ovoid, scarlet, 8–10 mm. diameter.
- Paineana* Sarg. Fruit long-ovoid, scarlet, pale dotted, about 10 mm. diameter.
- parca* Ashe. (?) Fruit globose or slightly oblong, scarlet-mottled with orange, about 10 mm. diameter.
- perlaeta* Sarg. Fruit short-oblong to slightly ovoid, scarlet, pale dotted, 8–9 mm. diameter.

populnea Ashe. Fruit globular or oblong, dull red mottled with green, russet and scarlet, about 14 mm. diameter.

propensa Sarg. Fruit short-oblong, scarlet, pale dotted, 8–10 mm. diameter.

Randiana Sarg. Fruit oblong, scarlet, pale dotted, 8–9 mm. diameter.

recta Sarg. Fruit short-oblong, scarlet, pale dotted, 8–9 mm. diameter.

retrusa Ashe. Fruit ovoid or oblong, orange-scarlet, 8–10 mm. diameter.

roanensis Ashe. Fruit oblong, bright red, about 8 mm. diameter.

rubicunda Sarg. Fruit oblong, scarlet, pale dotted, about 8 mm. diameter.

Corymbs sometimes slightly villous.

rufipes Ashe. Fruit oblong, scarlet mottled with green or russet, 9–11 mm. diameter.

sarniensis Sarg. Fruit short-oblong, light orange-red, pale dotted, 7–8 mm. diameter.

sextilis Sarg. Fruit subglobose to short-oblong, scarlet, 12–14 mm. diameter.

Slavinii Sarg. Fruit short-oblong or slightly ovoid, bright orange-red, pale dotted, about 10 mm. diameter.

spatifolia Sarg. Fruit short-oblong, crimson, pale dotted, 8–9 mm. diameter.

Streeterae Sarg. Fruit oblong, scarlet, pale dotted, about 8 mm. diameter.

tenella Ashe. Fruit oblong, dark orange, scarlet or mottled, 9–10 mm. diameter. Leaves sometimes narrowed at base.

tenera Ashe. Fruit oblong, bright scarlet, 9–10 mm. diameter.

tenuiloba Sarg. Fruit oblong-ovoid, scarlet, about 10 mm. diameter.

trachyphylla Sarg. Fruit ovoid, crimson, pale dotted, 12–15 mm. diameter.

vittata Ashe. Fruit subglobose or short-cylindric, red-scarlet blotched with olive or russet, 14–18 mm. diameter.

CC. Corymbs villous at flowering time

rubicunda Sarg. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.

uber Ashe. Fruit oblong, dark, dull red mottled with orange, 12–15 mm. diameter.

AA. Stamens 15–20 (rarely less)

B. Leaves of flowering branches usually narrowed or attenuate at base (on vigorous shoots rounded or truncate)

C. Corymbs glabrous at flowering time

alnorum Sarg. Fruit oblong-ovoid, bright scarlet, about 8 mm. diameter.

asperata Sarg. Fruit ovoid to short-oblong, scarlet, pale dotted, 14–16 mm. diameter. Corymbs sometimes slightly villous.

basilica Beadle. Fruit subglobose, red, 12–15 mm. diameter.

conferta Sarg. Fruit short-oblong, orange-red, pale dotted, about 8 mm. diameter.

decens Ashe. Fruit oblong, subglobose or slightly ovoid, scarlet or orange spotted, 10–12 mm. diameter.

depilis Sarg. Fruit subglobose to broad-obvoid, dark red to reddish-purple, 12–15 mm. diameter.

Edsonii Sarg. Fruit subglobose to short-oblong or obovoid, bright cherry-red, pale dotted, 13–15 mm. diameter.

lucorum Sarg. Fruit pear-shaped, becoming oblong or obovoid, crimson with pale lenticles, 12–15 mm. diameter. Corymbs sometimes slightly *villous*.

luminosa Sarg. Fruit obovoid, crimson, pale dotted, 10–12 mm. diameter.

mellita Sarg. Fruit oblong or oblong-obvoid, bright scarlet, 8–9 mm. diameter.

pastorum Sarg. Stamens sometimes 5–10. Fruit characters as given above.

taetrica Sarg. Fruit oval, scarlet-yellow, blotched and pale dotted, 9–10 mm. diameter.

tarda Sarg. Stamens sometimes 10 or less. Fruit characters as given above.

CC. Corymbs villous at flowering time

asperata Sarg. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.

lucorum Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.

BB. Leaves of flowering branches not usually narrowed at base (on vigorous shoots truncate or cordate). Corymbs glabrous

Schuettei Ashe. Fruit subglobose or globular-pyriform, dark red, about 13 mm. diameter.

VII. Silvicolae

Anthers red, rose, pink or purple. Stamens 10 or less

A. Leaves of flowering branches mostly narrowed at base (on vigorous shoots rounded or truncate)

acerba Sarg. Fruit obovoid, crimson, pale dotted, 8–10 mm. diameter.

aemula Beadle. Fruit globose or subglobose, red, 10–13 mm. diameter.

ambigens Sarg. Fruit globose or depressed-globose, greenish-yellow, becoming dark purplish-red, pale dotted, 10–12 mm. diameter. Terminal shoot leaves sometimes subcordate.

Barryana Sarg. Fruit obovoid, crimson, pale dotted, about 15 mm. diameter. Terminal shoot leaves sometimes subcordate.

blairensis Sarg. Fruit subglobose to slightly obovoid, dull red, pale dotted, 10–12 mm. diameter.

coerulea Sarg. Fruit obovoid, bright orange-red, pale dotted, 10–12 mm. diameter.

compta Sarg. Fruit oblong-obvoid, rarely oblong, light cherry-red, pale dotted, about 12 mm. diameter. Terminal shoot leaves sometimes subcordate.

congestiflora Sarg. Fruit obovoid, crimson, pale dotted, 9–11 mm. diameter.

- cruda* Sarg. Fruit obovoid, scarlet, pale dotted, about 10 mm. diameter.
- delectata* Sarg. Fruit obovoid, crimson, pale dotted, about 10 mm. diameter.
- dissociabilis* Sarg. Fruit obovoid, light cherry-red, about 9 mm. diameter. Terminal shoot leaves sometimes subcordate.
- dissona* Sarg. Fruit depressed-globose, crimson blotched with green, pale dotted, 12–16 mm. diameter.
- effera* Sarg. Fruit obovoid, orange-red, pale dotted, 10–12 mm. diameter.
- gravis* Ashe. Fruit short-oblong, red or crimson, 8–9 mm. diameter.
- impervia* Sarg. Fruit short-oblong to slightly obovoid, salmon-red, 10–12 mm. diameter.
- implicata* Sarg. Fruit subglobose to ovoid, crimson, pale dotted, 8–11 mm. diameter.
- infensa* Sarg. Fruit short-obvoid, scarlet, pale dotted, 10–11 mm. diameter. Terminal shoot leaves often subcordate.
- iterata* Sarg. Fruit subglobose, angled, light orange-red, blotched with green, 13–15 mm. (or less) diameter.
- laetans* Sarg. Fruit obovoid, rarely short-oblong, dark orange-red, pale dotted, 11–13 mm. diameter.
- leimonia* Sarg. Fruit subglobose, angled, orange-red blotched with yellow-green, pale dotted, 10–15 mm. (or less) diameter.
- Livingstoniana* Sarg. Fruit short-oblong, red, pale dotted, 10–12 mm. diameter. Terminal shoot leaves sometimes subcordate.
- luxuriosa* Sarg. Fruit subglobose to slightly obovoid, scarlet, pale dotted, 12–18 mm. diameter.
- perspicabilis* Sarg. Fruit subglobose to short-oblong, maroon, pale dotted, 10–14 mm. diameter.
- promissa* Sarg. Fruit oblong or obovoid, crimson, 7–8 mm. diameter.
- prona* Ashe. Fruit oblong-obvoid or pyriform, dark red mottled with green and russet, 8–10 mm. diameter.
- puta* Sarg. Fruit subglobose, scarlet, pale dotted, 10–14 mm. diameter.
- pyramidata* Sarg. Fruit short-oblong or slightly obovoid, scarlet, pale dotted, 10–11 mm. diameter.
- radina* Sarg. Fruit globose or slightly obovoid, yellow-green to dark purplish-red, pale dotted, 12–14 mm. diameter.
- recordabilis* Sarg. Fruit narrow-obvoid, greenish-orange to scarlet, pale dotted, 9–10 mm. diameter.
- ruricola* Sarg. Fruit obovoid, scarlet, 10–15 mm. diameter.
- sectilis* Ashe. (?) Fruit subglobose or obovoid, dark red, 12–16 mm. diameter.
- strigosa* Sarg. Fruit short-oblong, scarlet, dark dotted, 10–11 mm. diameter.
- tortuosa* Sarg. Fruit obovoid, bright orange-red, pale dotted, 10–12 mm. diameter.

xanthophylla Sarg. Fruit short-oblong, orange-red, pale dotted, 10–12 mm. diameter.

AA. Leaves of flowering branches mostly broad or rounded at base (on vigorous shoots truncate to cordate)

albicans Ashe. (?) Fruit globular, dark red, 10–13 mm. diameter.

allecta Sarg. Fruit ovoid, bright orange-red, dark dotted, 8–10 mm. diameter.

ampliata Sarg. Fruit short-oblong, dull scarlet, 9–10 mm. diameter.

delicata Sarg. Fruit ovoid to oblong, dull red, 10–12 mm. diameter.

diffusa Sarg. Fruit subglobose to short-oblong, scarlet, dark dotted, 13–15 mm. diameter.

drymophila Sarg. Fruit globose, red or greenish, 10–11 mm. diameter.

filipes Ashe. Fruit ovoid, crimson or green, spotted, 10–12 mm. diameter.

Maineana Sarg. Fruit globose, dark scarlet with light spots, about 15 mm. diameter.

medioxima Sarg. Fruit subglobose, rarely short-oblong, dark red, pale dotted, 11–14 mm. diameter.

opulens Sarg. Fruit subglobose to short-obvoid, crimson, dark dotted, 13–15 mm. diameter.

riparia Ashe. (?) Color of anthers not given. Fruit globose, orange-yellow, 7–8 mm. diameter.

stolonifera Sarg. Fruit short-oblong to subglobose, scarlet, 10–11 mm. diameter.

VIII. Pruinosae

A. Anthers pink, rose or purple (AA see p. 105)

B. Stamens about 10 (7–15) (BB see p. 101)

C. Leaves of flowering branches mostly narrowed or cuneate at base (on vigorous shoots rounded to truncate), at flowering time glabrous

alacris Sarg. Fruit short-oblong or slightly obovoid, bright red, 8–10 mm. diameter.

aridula Sarg. Fruit obovoid, dark red, pale dotted, 8–9 mm. diameter.

bellula Sarg. Fruit subglobose or broadly obovoid, dull crimson, pale dotted, 15–18 mm. diameter.

brevipes Peck. Fruit subglobose to depressed-globose, dull red blotched with red, 10–15 mm. diameter.

deducta Sarg. Fruit short-oblong to ovoid, green, 10–12 mm. diameter.

disjuncta Sarg. Fruit subglobose, green tinged with red, 12–14 mm. diameter.

divisifolia Sarg. Fruit depressed-globose, angled, dull red or blotched with green or russet, 14–17 mm. diameter.

erubescens Sarg. Fruit subglobose, pentagonal, purple-red, pale dotted, 12–15 mm. diameter. Leaves sometimes broader.

franklinensis Sarg. Fruit slightly obovoid to subglobose, dark red, 8–10 mm. diameter.

incisa Sarg. Fruit obovoid, green, red tinged or crimson, 10–12 mm. diameter.

iracunda Beadle. Fruit subglobose, red or red and green, 8–10 mm. diameter.

Jesupii Sarg. Fruit ovoid to short-oblong, bright scarlet, dark dotted, about 10 mm. diameter.

levis Sarg. Fruit ovoid, dull purple, 7–8 mm. diameter.

Milleri Sarg. Fruit ovoid, green becoming red, 8–10 mm. diameter.

nemorosa Sarg. Fruit short-oblong to ovoid, crimson, dark dotted, about 10 mm. diameter.

numerosa Sarg. Fruit ovoid, scarlet, pale dotted, 9–10 mm. diameter.

ovatifolia Sarg. Fruit ovoid, scarlet, dark dotted, 8–10 mm. diameter.

placiva Sarg. Fruit short-oblong or ovoid, bright orange-red, pale dotted, 10–12 mm. diameter.

procera Sarg. Fruit subglobose, crimson, pale dotted, 10–12 mm. diameter.

pulchra Sarg. Fruit short-oblong, cherry-red, 10–12 mm. diameter.

radiata Sarg. Fruit short-oblong, crimson, dark dotted, 15–17 mm. diam.

remota Sarg. Fruit subglobose or slightly ovoid, scarlet, 10–15 mm. diameter.

rigida Sarg. Fruit subglobose, angled, green, rose tinged, 10–12 mm. diameter.

superata Sarg. Fruit subglobose, green becoming crimson, dark dotted, 12–15 mm. diameter.

uncta Sarg. Fruit short-oblong, red, pale dotted, 10–12 mm. diameter.

CC. Leaves of flowering branches broader, mostly rounded, truncate or cordate at base

D. Leaves at flowering time glabrous or nearly so

erubescens Sarg. Fruit subglobose, pentagonal, purple-red, pale dotted, 12–15 mm. diameter.

deltoides Ashe. Fruit globular or depressed-globular, dark red, 15–17 mm. diameter.

depressa Ashe. (?) Color of anthers unknown. Fruit depressed-globose, dark red, 12–14 mm. diameter.

macera Sarg. Fruit subglobose to short-oblong, scarlet, dark dotted, about 10 mm. diameter.

plana Sarg. Fruit short-oblong, pale green becoming crimson, pale dotted, 10–12 mm. diameter.

relicta Sarg. Fruit subglobose, green flushed with red, 9–10 mm. diameter.

DD. Leaves at flowering time pubescent or scabrate above

exornata Sarg. Fruit subglobose, scarlet, dark dotted, 10–12 mm. diameter.

BB. Stamens about 20 (15–20)

C. Leaves of flowering branches mostly narrowed or cuneate at base (on vigorous shoots rounded to truncate, rarely subcordate) (CC see p. 104)

D. Leaves at flowering time glabrous or nearly so

advena Sarg. Fruit subglobose to slightly ovoid, green becoming reddish, about 15 mm. diameter. Leaves sometimes slightly villous.

ambitiosa Sarg. Fruit ovoid, crimson, green blotched, with dark dots, 10–12 mm. diameter. Leaves sometimes slightly villous.

- amoena Sarg. Fruit subglobose to short-oblong, dark red, 10–12 mm. diameter.
- aristata Sarg. Fruit subglobose, bright red, 10–12 mm. diameter.
- ater Ashe. Fruit oblong to ovoid, dark dull red mottled with green or olive, 10–12 mm. diameter.
- austera Sarg. Fruit subglobose, dull green or reddish, 10–12 mm. diameter.
- bellatula Sarg. Fruit short-oblong, green turning red, about 10 mm. diameter.
- bona Sarg. Fruit depressed-globose, angular, dark olive-green or spotted with russet and orange, 12–14 mm. diameter.
- bronxensis Sarg. Fruit subglobose, apple-green, dark dotted, 10–12 mm. diameter. Leaves sometimes villous or scabrous.
- casta Sarg. Fruit ovoid, bright cherry-red, pale dotted, 8–10 mm. diameter.
- Clintoniana Sarg. Fruit subglobose, angled, orange-red, pale dotted, 12–14 mm. diameter. Leaves sometimes slightly villous.
- comata Sarg. Fruit depressed-globose, bright green becoming light cherry-red, about 16 mm. diameter. Leaves sometimes slightly villous above.
- Crawfordiana Sarg. Fruit oval to pyriform-oval, dark orange-red blotched with yellow-green, pale dotted, 10–12 mm. diameter.
- denudata Sarg. Fruit subglobose to ovoid-globose, russet-green to dark purplish red, 10–13 mm. diameter.
- festiva Sarg. Fruit subglobose, dull red, 10–11 mm. diameter.
- fusca Sarg. Fruit short-oblong, dull brown, 10–12 mm. diameter.
- gaudens Sarg. Fruit pyriform, angled, deep orange-red, pale dotted, 11–14 mm. diameter.
- georgiana Sarg. Fruit short-oblong, dull russet-green, 10–13 mm. diameter.
- gracilis Sarg. Fruit depressed-globose, green tinged with red or orange, about 10 mm. diameter.
- horridula Sarg. Fruit ovoid, green becoming dull red, dark dotted, about 10 mm. diameter.
- Howeana Sarg. Fruit globose to depressed-globose, reddish-green to bright scarlet, 13–15 mm. (or often less) in diameter.
- immanis Ashe. Fruit "discoid," (or subglobose) bright green, crimson spotted, 13–16 mm. diameter.
- Kellermanii Sarg. Fruit subglobose, red tinged with green, 14–18 mm. (or less) diameter.
- latiflora Sarg. Fruit short-oblong, vermillion, pale dotted, 8–9 mm. diameter.
- obstipa Sarg. Fruit obconic, crimson, pale dotted, 10–12 mm. diameter.
- pallescens Sarg. Fruit short-oblong, cardinal-red, pale dotted, 10–12 mm. diameter.
- parvula Sarg. Fruit ovoid to slightly ovoid, crimson, pale dotted, 10–12 mm. diameter.

- pellecta* Sarg. Fruit short-oblong to subglobose, dark red, 12–14 mm. diameter.
- pequotorum* Sarg. Fruit short-ovoid, dark crimson with dark lenticles, 11–13 mm. diameter.
- pilosa* Sarg. Fruit subglobose to broadly ovoid, dull dark crimson, 8–10 mm. diameter. Leaves sometimes slightly villous.
- quinebaugensis* Sarg. Fruit oblong or slightly obovoid, light red or purplish, 8–9 mm. diameter.
- Ruthiana* Sarg. Fruit depressed-globose or short-oblong, bright apple-green, turning red, about 10 mm. diameter. Leaves sometimes slightly villous.
- scitula* Sarg. Fruit subglobose, crimson, pale dotted, 9–11 mm. diameter.
- sitiens* Ashe. Fruit ovoid, green mottled with russet, 12–14 mm. (or less) diameter.
- torta* Sarg. Fruit short-oblong, angled, light yellow or russet-green, dark dotted, 8–11 mm. diameter.
- tumida* Sarg. Fruit subglobose, dull crimson, 14–15 mm. diameter.
- viatica* Ashe. Fruit globose, dull green mottled with pink and russet, about 15 mm. (or less) diameter.
- virella* Ashe. Fruit depressed-globose, yellowish-green blotched with pink, olive or russet, about 13 mm. diameter. Leaves sometimes slightly pubescent.
- Wilmorensis* Sarg. Fruit subglobose to short-oblong, green becoming reddish, 10–12 mm. diameter.
- DD. Leaves at flowering time scabrous or pubescent
- advena* Sarg. Leaves sometimes glabrous. Fruit characters as given above.
- ambitiosa* Sarg. Leaves sometimes nearly or quite glabrous. Fruit characters as given above.
- bronxensis* Sarg. Leaves sometimes nearly glabrous. Fruit characters as given above.
- Clintoniana* Sarg. Leaves sometimes glabrous or nearly so. Fruit characters as given above.
- comata* Sarg. Leaves sometimes nearly or quite glabrous. Fruit characters as given above.
- decorata* Sarg. Fruit depressed-globose or short-oblong, angled, dark red, dark dotted, 8–10 mm. diameter.
- pelacris* Sarg. Fruit subglobose to ovoid, green becoming red, 10–12 mm. diameter.
- pilosa* Sarg. Young leaves sometimes nearly or quite glabrous. Fruit characters as given above.
- platycarpa* Sarg. Fruit short-oblong, angled, dull red, dark dotted, 10–12 mm. diameter. Leaves sometimes nearly glabrous.
- Ruthiana* Sarg. Leaves sometimes nearly or quite glabrous. Fruit characters as given above.

virella Ashe. Leaves sometimes glabrous. Fruit characters as given above.

CC. Leaves of flowering branches broader, mostly rounded, truncate or cordate at base

D. Leaves at flowering time glabrous or nearly so

amplifica Sarg. Fruit depressed-globose, light greenish-yellow to dark russet, 10–12 mm. diameter.

angulata Sarg. Fruit depressed-ovoid, pentagonal, light greenish-yellow becoming dark purplish-red, dark dotted, 15–16 mm. diameter.

arcana Beadle. Fruit subglobose, obtusely angled, red or ruddy, 12–15 mm. diameter.

beata Sarg. Fruit oblong to oblong-ovoid, crimson, pale dotted, 9–10 mm. diameter.

bracteata Sarg. Fruit subglobose, crimson, pale dotted, 12–14 mm. diameter.

calliantha Sarg. Fruit short-oblong, angled, pink, about 10 mm. diameter.

comparata Sarg. Fruit ovoid to oval, crimson, 9–10 mm. diameter.

confragosa Sarg. Fruit short-oblong to ovoid, scarlet, dark dotted, about 10 mm. (or more) diameter.

dunmorensis Sarg. Fruit ovoid, green becoming dull red, 10–12 mm. (or more) diameter.

Gattingeri Ashe. Fruit globular, dark red, 8–11 mm. diameter.

leiophylla Sarg. Fruit ovoid, dark green becoming bright red, 12–14 mm. diameter. (Anthers described as pale yellow, but apparently through error.)

Lennoniana Sarg. Fruit ovoid to oblong, dark crimson, pale dotted, 8–10 mm. diameter.

Mackenzii Sarg. (?) Fruit subglobose to short-oblong, green becoming lurid red, 10–12 mm. diameter. Color of anthers unknown.

macrocalyx Sarg. Fruit short-oblong, slightly angled, red, pale dotted, 10–12 mm. diameter.

oblita Sarg. Fruit ovoid, crimson, pale dotted, 9–10 mm. diameter.

perampla Sarg. Fruit short-oblong, dull red, 10–12 mm. diameter.

philadelphica Sarg. Fruit depressed-globose, green, dark dotted, 10–15 mm. diameter.

prominens Sarg. Fruit broad-ovoid or short-oblong, green with purplish cheek, 12–14 mm. diameter.

pruinosa K. Koch. Fruit subglobose, green becoming red, 10–12 mm. diameter.

ramosa Sarg. Fruit short-oblong to oval, red, punctate, 10–11 mm. diameter.

rubicundula Sarg. Fruit subglobose or depressed-globose, green becoming red tinged, pale dotted, 13–14 mm. diameter.

seclusa Sarg. Fruit subglobose, angled, light orange-red blotched with green, 13–15 mm. (or less) diameter.

seducta Sarg. Fruit depressed-globose, green blotched with red, 10–11 mm. diameter.

DD. Leaves at flowering time villous beneath
aspera Sarg. Fruit short-oblong to subglobose, scarlet, 10–12 mm. diameter.

AA. Anthers white, cream or yellow

B. Stamens about 10

C. Leaves of flowering branches mostly cuneate or narrowed at base (on vigorous shoots rounded to truncate, rarely subcordate)

D. Leaves at flowering time glabrous

brachypoda Sarg. Fruit subglobose, green, red tinged, becoming dull red, 10–12 mm. diameter.

delawarensis Sarg. Fruit subglobose, dark red, 11–12 mm. diameter.

rara Sarg. Fruit depressed-globose, dull dark red, 12–14 mm. diameter.

DD. Leaves at flowering time scabrous or pubescent

inusitula Sarg. Fruit short-oblong, dull greenish-red, pale dotted, about 10 mm. diameter. Leaves scabrate above.

sicca Sarg. Fruit subglobose to short-oblong, becoming light orange-yellow, pale dotted, about 10 mm. diameter. Leaves more or less villous above.

C. Leaves of flowering branches broader, mostly rounded to truncate at base (on vigorous shoots cordate), at flowering time glabrous

depressa Ashe. (?) Color of anthers unknown. Fruit characters as given above.

BB. Stamens about 20 (15–20)

C. Leaves of fruiting branches mostly cuneate or narrowed at base (on vigorous shoots rounded or truncate, rarely subcordate)

D. Leaves at flowering time glabrous

ambitiosa Sarg. Fruit ovoid, crimson blotched with green, pale dotted, 10–12 mm. (or more) diameter. Anthers described as yellow faintly tinged with pink in bud.

aperta Sarg. Fruit subglobose, dull green, red tinged, about 10 mm. diameter.

bedfordensis Sarg. Fruit short-oblong to slightly ovoid, green turning pale red, dark dotted, 8–10 mm. diameter.

caerulescens Sarg. Fruit ovoid, dull red, 7–8 mm. diameter.

callida Beadle. Fruit depressed-globose, green and red or russet and red, 8–10 mm. diameter.

callosa Sarg. Fruit globose or depressed-globose, green becoming dark red or cardinal, blotched and punctate with dark olive-green or russet, 13–16 mm. diameter.

cestrica Sarg. Fruit subglobose to ovoid, dark crimson, dark dotted, 12–15 mm. diameter.

cognata Sarg. Fruit pyriform or oblong (or depressed-globose), 12–14 mm. diameter.

conjuncta Sarg. Fruit subglobose or depressed-globose, dull orange-red blotched with green, dark dotted, about 10 mm. diameter.

duracina Sarg. Fruit subglobose to slightly ovoid, red blotched with green, 8–10 mm. diameter.

- Ferrissii* Ashe. Fruit oblong or pyriform, crimson, 8–13 mm. diameter.
formosa Sarg. Fruit oblong to slightly ovoid, scarlet with pale lenticles,
about 10 mm. diameter.
- glareosa* Ashe. Fruit globose, bright scarlet, about 15 mm. diameter.
Leaves sometimes slightly villous beneath.
- insueta* Sarg. Fruit pyriform, dull green tinged with red, 9–10 mm.
diameter.
- latifrons* Sarg. Fruit subglobose to short-oblong, dark red, green blotched,
14–15 mm. diameter.
- latisepala* Ashe. Fruit ovoid, dull green becoming crimson, dark
dotted, 9–12 mm. diameter.
- littoralis* Sarg. Fruit ovoid, dark crimson, 10–11 mm. diameter.
- longipedunculata* Sarg. Fruit ovoid, dark green, red tinged, dark dotted,
8–10 mm. diameter.
- magnifolia* Sarg. Fruit subglobose, angled, dull red with dark blotches,
about 15 mm. (or less) diameter.
- onusta* Ashe. Fruit globose, dark red, 14–16 mm. (or less) diameter.
- perjucunda* Sarg. Fruit short-oblong to ovoid, orange-red becoming
crimson, dark dotted, 10–12 mm. diameter.
- Porteri* Britton. Fruit pyriform or short-oblong, crimson, dark dotted,
9–10 mm. diameter.
- rubro-lutea* Sarg. Fruit subglobose or ovoid, light orange-red, dark
dotted, 12–15 mm. diameter.
- russata* Sarg. Fruit ovoid, pale red, about 10 mm. diameter.
- rustica* Beadle. Fruit subglobose, obtusely angled and swelled below
middle, red and green, becoming dull red, 7–11 mm. diameter.
- Shaferi* Ashe. Fruit subglobose, dark red or green mottled, 14–16 mm.
(or less) diameter.
- tribulosa* Sarg. Fruit short-oblong to subglobose, dull green, red tinged,
about 10 mm. diameter.
- uplandia* Sarg. Fruit subglobose to short-oblong or slightly ovoid,
dark deep red, pale dotted, 14–15 mm. diameter.
- vicinalis* Beadle. Fruit pyriform, red or ruddy, 9–12 mm. diameter.
- DD. Leaves at flowering time more or less villous or pubescent
- ellipticifolia* Sarg. Fruit short-oblong to subglobose, dull red, about 10
mm. diameter.
- ingrata* Ashe. Fruit subglobose, dull red, 12–14 mm. diameter.
- lecta* Sarg. Fruit ovoid, light yellow-green, becoming red, 13–15 mm.
diameter.
- locuples* Sarg. Fruit subglobose, dull green, red tinged, 12–14 mm. diameter.
- CC. Leaves of flowering branches broader, mostly rounded or truncate (on
vigorous shoots cordate); glabrous at flowering time
- augusta* Sarg. Fruit pyriform or at maturity globose or depressed-
globose, angled, red or crimson or blotched with olive-green, 10–14
mm. diameter.

incompta Sarg. Fruit subglobose or depressed-globose, dull red, pale dotted, about 10 mm. diameter.

jejuna Sarg. Fruit short-oblong to ovoid, angled, dull green, 7–8 mm. diameter.

Mackenzii Sarg. (?) Color of anthers unknown. Fruit characters as given above.

patrum Sarg. Fruit short-oblong to slightly obovoid, dull green, dark dotted, 8–10 mm. diameter.

IX. Coccineae

A. Anthers red, pink, rose or purple (AA see p. 109)

B. Stamens about 10 (5–10)

 D. Corymbs at flowering time glabrous or nearly so

cristata Ashe. Fruit globose or pyriform, scarlet or darker, 15–20 mm. diameter. Corymbs sometimes slightly villous.

delecta Sarg. Fruit subglobose to slightly obovoid, scarlet becoming purple, 18–20 mm. diameter. Corymbs sometimes slightly villous.

fretalis Sarg. Fruit oblong to obovoid, crimson, about 10 mm. diameter.

Holmesiana Ashe. Fruit obovoid to ellipsoid, crimson, dark dotted, 8–11 mm. diameter. Corymbs villous in varieties.

Letchworthiana Sarg. Fruit obovoid, scarlet, pale dotted, 10–11 mm. diameter.

neofaxonii Sarg. Fruit obovate, bright cherry-red, pale dotted, 10–12 mm. diameter.

pedicellata Sarg. Fruit oblong, bright scarlet with dark lenticles, 12–18 mm. diameter.

politá Sarg. Fruit subglobose to short-oblong, rarely broadly obovoid, bright cherry-red blotched with green or yellow, dark dotted, 9–11 mm. diameter.

Thayeri Sarg. Fruit subglobose to short-oblong or obovoid, orange-red with pale lenticles, 8–10 mm. diameter.

 DD. Corymbs at flowering time villous or pubescent

acclivis Sarg. Fruit short-oblong, yellowish-red, pale dotted, about 12 mm. (or more) diameter.

arcuata Ashe. Fruit oblong-globular, scarlet or mottled with orange and russet, 11–13 mm. diameter.

assurgens Sarg. Fruit oblong or slightly obovoid, dull red or crimson, 10–15 mm. diameter. Stamens sometimes 20.

aulica Sarg. Fruit short-oblong or slightly obovoid, orange-red with pale lenticles, about 15 mm. diameter.

Burkeana Sarg. Fruit subglobose to short-oblong or slightly obovoid, crimson, pale dotted, 10–12 mm. diameter.

caesa Ashe. Fruit subglobose or oblong, bright scarlet, 12 mm. diameter.

confinis Sarg. Fruit short-oblong, scarlet, pale dotted, 8–9 mm. diameter.

cristata Ashe. Corymbs usually glabrous. Fruit characters as given above.

delecta Sarg. Corymbs sometimes glabrous. Fruit characters as given above.

Eamesii Sarg. Fruit short-oblong to slightly ovoid, bright cherry-red, dark dotted, 18–20 mm. diameter.

elongata Sarg. Fruit oblong-obovoid, crimson, pale dotted, about 8 mm. diameter.

fluviatilis Sarg. Fruit oblong, rarely obovoid, crimson, 8–10 mm. diameter.

Holmesiana var. *tardipes* Sarg. Fruit as in species.

lenta Ashe. Fruit pyriform, scarlet, 10–14 mm. diameter.

lobulata Sarg. Fruit oblong, bright crimson with pale lenticles, about 10 mm. diameter.

Pringlei Sarg. Fruit oblong, dark dull red with pale lenticles, 15–16 mm. diameter.

pura Sarg. Fruit oval to slightly obovoid, bright orange-red, dark dotted, 8–10 mm. diameter.

sejuncta Sarg. Fruit subglobose to oval, crimson, pale dotted, 13–15 mm. (or less) diameter.

sertata Sarg. Fruit subglobose to slightly obovoid, bright red becoming crimson, pale dotted, 10–14 mm. diameter.

uticaensis Sarg. Fruit short-oblong, scarlet, pale dotted, 12–13 mm. diameter.

vivida Sarg. Fruit obovoid, dull orange-red blotched with yellow, 9–10 mm. diameter.

BB. Stamens about 20 (15–20)

D. Corymbs at flowering time glabrous or nearly so

contigua Sarg. Fruit oblong to obovoid, scarlet, pale dotted, 8–10 mm. diameter.

limosa Sarg. Fruit short-oblong, crimson, pale dotted, 8–10 mm. diameter.

Corymbs often slightly villous.

magniflora Sarg. Fruit short-obvoid, scarlet, pale dotted, about 14 mm. diameter.

miranda Sarg. Fruit obovoid, scarlet, pale dotted, about 10 mm. diameter.

illicebrosa Sarg. Fruit obovoid, crimson, pale dotted, 12–13 mm. diameter.

Putnamiana Sarg. Fruit depressed-globose, green turning red, punctate, about 15 mm. diameter.

splendida Sarg. Fruit short-oblong to obovoid, crimson, pale dotted, 9–10 mm. diameter.

DD. Corymbs at flowering time villous or pubescent

assurgens Sarg. Stamens sometimes 10. Fruit characters as given above.

chippewaensis Sarg. Fruit oblong-obovoid, crimson, dark dotted, 10–16 mm. diameter.

Dayana Sarg. Fruit obovoid, crimson, pale dotted, 15–18 mm. diameter.

Gilbertiana Sarg. Fruit obovoid, crimson, pale dotted, 13–15 mm. diameter.

Hillii Sarg. Fruit obovoid, crimson, pale dotted, 10–12 mm. diameter.

limosa Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.

Macounii Sarg. Fruit oblong-obvoid, scarlet mottled with green, 10–12 mm. diameter.

neolondinensis Sarg. Fruit obovoid to short-oblong, bright red, pale dotted, 13–18 mm. diameter.

steubenensis Sarg. Fruit obovoid, scarlet, pale dotted, 9–10 mm. diameter.

AA. Anthers white, cream or yellow

B. Stamens about 10 (5–10). Corymbs at flowering time villous or pubescent

perrara Sarg. Fruit short-oblong to slightly obovoid, crimson, pale dotted, 10–12 mm. diameter.

BB. Stamens about 20. Corymbs at flowering time villous or pubescent

conspicua Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, about 18 mm. diameter.

irrassa Sarg. Fruit oblong, dark red, pale dotted, about 10 mm. diameter.

X. Molles

A. Anthers red, pink or rose

B. Stamens about 10

C. Leaves of flowering branches mostly cuneate or rounded at base (on vigorous shoots truncate or subcordate)

anomala Sarg. Fruit obovoid to oblong, crimson with pale lenticles, 15–20 mm. diameter.

digna Sarg. Fruit globose to slightly oblong, scarlet dotted with yellow or orange, 12–14 mm. diameter.

Greggiana Egglest. Fruit globose, brick-red, 10–12 mm. diameter.

noelensis Sarg. Fruit subglobose, orange-red, about 10 mm. diameter.

Tracyi Ashe. Fruit scarlet, subglobose, 10–15 mm. diameter.

urbica Sarg. Fruit pyriform, becoming subglobose to short-oblong, crimson, pale dotted, 10–15 mm. diameter.

CC. Leaves of flowering branches broader, mostly rounded, truncate or cordate at base (on vigorous shoots cordate)

Ellwangeriana Sarg. Fruit short-oblong, crimson, 15–20 mm. diameter.

exclusa Sarg. Fruit oblong, rarely ovoid, bright cherry-red, dark dotted, about 13 mm. diameter.

Huntiana Sarg. Fruit obovoid, scarlet, dark dotted, 16–18 mm. diameter.

lauta Sarg. Fruit oblong or short-oblong, bright orange-red, dark dotted, 15–18 mm. diameter.

pennsylvanica Ashe. Fruit globular or oblong, dull red mottled with green, russet and scarlet, 15–20 mm. diameter.

Robesoniana Sarg. Fruit oblong, bright scarlet, pale dotted, 10–15 mm. diameter.

BB. Stamens about 20

C. Leaves of flowering branches mostly cuneate or rounded at base (on vigorous shoots truncate or subcordate)

amabilis Ashe. Fruit globose, bright scarlet or red, about 20 mm. diameter. Color of anthers unknown.

dispessa Ashe. Fruit pyriform or obovoid, bright cherry-red, 10–15 mm. diameter.

gravida Beadle. Fruit depressed-globose, red, 12–15 mm. diameter.

Leaves sometimes broader, subcordate on shoots.

induta Sarg. Fruit short-oblong, crimson or reddish-yellow, pale dotted, 18–20 mm. diameter.

lanuginosa Sarg. Fruit subglobose to short-oblong, dark crimson, pale dotted, 10–12 mm. (or larger) diameter. Leaves sometimes broader, subcordate on shoots.

Kelloggii Sarg. Fruit subglobose to short-oval, bright yellow with pale lenticles, about 20 mm. diameter.

quercina Ashe. Fruit subglobose, dark red, pale dotted, 10–15 mm. diameter.

radians Sarg. Fruit short-oblong or obovoid, crimson, pale dotted, 16–18 mm. diameter.

texana Buckley. Fruit obovoid to short-oblong, bright scarlet, 12–18 mm. diameter.

CC. Leaves of flowering branches broader, rounded to subcordate at base (on vigorous shoots cordate)

brachyphylla Sarg. Fruit subglobose, dull dark red, 10–14 mm. diameter.

corusca Sarg. Fruit short-oblong to obovoid, dark cherry-red, pale dotted, 12–15 mm. diameter.

dasyphylla Sarg. Fruit short-oblong, crimson, pale dotted, 18–20 mm. diameter.

Fulleriana Sarg. Fruit short-oblong, scarlet, pale dotted, about 12 mm. diameter.

gravida Beadle. Leaves sometimes narrowed at base. Fruit characters as given above.

lanuginosa Sarg. Leaves sometimes narrowed at base. Fruit characters as given above.

nupera Ashe. (?) Fruit globular or slightly oblong, dark red, 13–15 mm. diameter. Color of anthers unknown.

placens Sarg. Fruit obovoid, scarlet, 10–12 mm. diameter.

valens Ashe. (?) Fruit subglobose, bright red, 16–20 mm. diameter. Color of anthers unknown.

venosa Ashe. (?) Fruit globose or subglobose, bright red, 15–18 mm. diameter. Color of anthers unknown.

verna Ashe. (?) Fruit subglobose, dark red, 16–18 mm. diameter. Color of anthers unknown.

vesca Ashe. (?) Fruit subglobose, scarlet, 18–24 mm. diameter. Color of anthers unknown.

AA. Anthers white, cream or yellow

B. Stamens about 10

C. Leaves of flowering branches mostly narrowed or rounded at base (on vigorous shoots truncate to subcordate)

Arnoldiana Sarg. Fruit subglobose, bright crimson with pale lenticles, 15–20 mm. diameter.

champlainensis Sarg. Fruit ovoid to oblong, bright scarlet with pale lenticles, 15–20 mm. diameter. Leaves sometimes broader.

submollis Sarg. Fruit ovoid, bright orange-red, 15–20 mm. diameter.

Tracyi Ashe. Fruit subglobose, deep scarlet, 8–12 mm. diameter.

uvaldensis Sarg. Fruit subglobose to short-oblong, orange-red, 12–14 mm. diameter.

CC. Leaves of flowering branches broader, mostly rounded to subcordate at base (on vigorous shoots cordate)

champlainensis Sarg. Leaves often narrowed at base. Fruit characters as given above.

contortifolia Sarg. Fruit subglobose to short-oblong, bright cherry-red, about 15 mm. diameter.

Tatnalliana Sarg. Fruit globose, ovoid or rarely oblong, bright orange-red, pale dotted, 10–12 mm. diameter.

BB. Stamens about 20

C. Leaves of flowering branches mostly narrowed or rounded at base (on vigorous shoots truncate or subcordate)

amabilis Ashe. (?) Color of anthers unknown. Fruit characters as given above.

arkansana Sarg. Fruit oblong or rarely ovoid, bright crimson with dark lenticels, 13–18 mm. diameter.

Berlandieri Sarg. Fruit short-oblong to subglobose, scarlet, 10–16 mm. diameter.

canadensis Sarg. Fruit short-oblong to subglobose, crimson with pale lenticles, 10–15 mm. diameter.

cibaria Beadle. Fruit short-oblong to ovoid, red, 12–15 mm. diameter.

dumetosa Sarg. Fruit ovoid, scarlet, dark dotted, 10–15 mm. diameter.

invisa Sarg. Fruit short-oblong, orange-red, pale dotted, 10–12 mm. diameter.

limaria Sarg. Fruit oval to ovoid or short-oblong, crimson, pale dotted, 12–14 mm. diameter. Leaves sometimes broad-based.

meridionalis Sarg. Fruit short-oblong to subglobose, scarlet, 10–12 mm. diameter.

mollipes Sarg. Fruit ovoid, scarlet, pale dotted, 8–9 mm. diameter. Leaves sometimes broader-based.

obtecta Ashe. Fruit pyriform, dark red, 12–15 mm. diameter.

Ridgwayi Sarg. Fruit subglobose or short-oblong, scarlet, 10–12 mm. diameter.

sera Sarg. Fruit ovoid or short-oblong, dull dark red, pale dotted, 12–20 mm. diameter.

Treleasei Sarg. Fruit subglobose, crimson, pale dotted, 12–17 mm. diameter.

umbrosa Sarg. Fruit ovoid to subglobose, scarlet, pale dotted, about 10 mm. diameter.

- viburnifolia* Sarg. Fruit subglobose, bright canary-yellow, 15–20 mm. diameter.
- CC. Leaves of flowering branches broader, mostly rounded to subcordate at base (on vigorous shoots cordate)
- altrix* Ashe. Fruit globose, bright red, 13–18 mm. diameter.
- cibilis* Ashe. Fruit depressed-globose, dark red, 12–15 mm. diameter.
- declivitatis* Sarg. Fruit subglobose to short-oblong, dull red, 12–13 mm. diameter.
- lanigera* Sarg. Fruit ovoid, crimson, 10–11 mm. diameter.
- lasiantha* Sarg. Fruit oval or subglobose, crimson, pale dotted, about 12 mm. diameter.
- limaria* Sarg. Leaves sometimes narrowed at base. Fruit characters as given above.
- macrophylla* Sarg. Fruit oval, crimson, pale dotted, 15–16 mm. (or more) diameter.
- mollipes* Sarg. Leaves sometimes narrowed at base. Fruit characters as given above.
- mollis* Scheele. Fruit short-oblong to subglobose, scarlet, dark dotted, 20–25 mm. diameter.
- nupera* Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- nutans* Sarg. Fruit short-oblong to subglobose or slightly ovoid, orange-red with dark lenticles, 10–12 mm. diameter.
- redolens* Ashe. Fruit globose, dark red, 15–20 mm. diameter.
- valens* Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- venosa* Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- verna* Ashe. (?) Color of anthers unknown. Fruit characters as given above.
- vesca* Ashe. (?) Color of anthers unknown. Fruit characters as given above.

XI. Dilatatae

Anthers pink, rose or purple. Stamens about 20

A. Leaves of flowering branches mostly narrowed at base, glabrous or nearly so

durobrivensis Sarg. Fruit globose, crimson, pale dotted, about 16 mm. diameter.

hudsonica Sarg. Fruit subglobose, crimson, 16–18 mm. diameter.

suborbiculata Sarg. Fruit subglobose, dull red blotched with green, 12–15 mm. diameter.

AA. Leaves of flowering branches mostly truncate to cordate

B. Leaves at flowering time glabrous or nearly so; corymbs glabrous

coccinoides Ashe. Fruit subglobose, dark crimson, pale dotted, 15–20 mm. diameter.

BB. Leaves at flowering time short villous above; corymbs more or less villous
dilatata Sarg. Fruit subglobose, bright scarlet, 18–20 mm. diameter.

XII. *Rotundifoliae*

A. Anthers pink, rose or purple (AA see p. 114)

B. Stamens about 10 (5–10)

C. Corymbs at flowering time glabrous or nearly so

Bicknellii Egglest. Fruit globose, red, 10–12 mm. diameter.

cernua Ashe. Fruit oblong, crimson, 13–14 mm. diameter.

chateaugayensis Sarg. Fruit short-oblong to oval, scarlet, pale dotted, 8–10 mm. diameter.

dacrioidea Sarg. Fruit obovoid, orange-red, pale dotted, 10–12 mm. diameter.

Harryi Sarg. Fruit short-oblong to slightly obovoid, cherry-red, pale dotted, 8–9 mm. diameter.

propria Sarg. Fruit oblong or slightly ovoid, scarlet, pale dotted, 7–8 mm. diameter.

spissa Sarg. Fruit subglobose to oval, scarlet, pale dotted, 7–9 mm. diameter.

Williamsii Egglest. (?) Fruit globose, red, 8–10 mm. diameter. Color of anthers unknown.

CC. Corymbs at flowering time villous or pubescent

Fernaldii Sarg. Fruit obovoid, bright scarlet, 9–10 mm. diameter.

illuminata Sarg. (?) Fruit oval to subglobose, bright cherry-red, 7–8 mm. diameter. Color of anthers unknown.

Jonesae Sarg. Fruit oblong to oblong-obvoid, dark carmine red, punctate, 12–15 mm. diameter.

Kennedyi Sarg. Fruit slender-obvoid, crimson, 7–8 mm. diameter.

lemingtonensis Sarg. Fruit short-oblong, scarlet, pale dotted, 8–9 mm. diameter.

maligna Sarg. Fruit short-oblong, crimson, pale dotted, 9–10 mm. diameter.

praecoqua Sarg. Fruit subglobose, dark crimson, 10–15 mm. diameter.

Proctoriana Sarg. Fruit subglobose, crimson, pale dotted, 10–12 mm. diameter.

puberis Sarg. Fruit short-oblong, orange-red, pale dotted, 9–10 mm. diameter.

verrucalis Peck. Fruit subglobose to short-oblong, scarlet, 8–10 mm. diameter.

Websteri Sarg. Fruit short-oblong, bright cherry-red, pale dotted, 8–10 mm. diameter.

BB. Stamens about 20 (15–20)

C. Corymbs at flowering time glabrous or nearly so

Brainerdii Sarg. Fruit oblong, bright scarlet, 8–10 mm. diameter.

kingstonensis Sarg. Fruit subglobose, dark red, about 15 mm. diameter.

- meiophylla* Sarg. Fruit subglobose, dark red, about 10 mm. diameter.
Corymbs sometimes slightly villous.
- neobaxteri* Sarg. Fruit short-oblong, rich deep red, pale dotted, 10–12 mm. diameter.
- neosmithii* Sarg. Fruit subglobose, light green becoming russet, 16–17 mm. diameter.
- scopulorum* Sarg. Fruit short-oblong to slightly ovoid, orange-red, pale dotted, 8–10 mm. diameter.
CC. Corymbs at flowering time villous or pubescent
- Blanchardii* Sarg. Fruit short-oblong, dark cherry-red, 10–12 mm. diameter.
- Evansiana* Sarg. Fruit subglobose to short-oblong, scarlet, 10–13 mm. diameter.
- insolens* Sarg. Fruit short-oblong, scarlet, 8–10 mm. diameter.
- mansfieldensis* Sarg. Fruit broad-ovoid or subglobose, dark red, 8–10 mm. diameter.
- meiophylla* Sarg. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.
- AA. Anthers white, cream or yellow
B. Stamens about 10 (5–10)
C. Corymbs at flowering time glabrous or nearly so
- Dodgei* Ashe. Fruit depressed-globose, dull crimson or orange, 10–15 mm. diameter.
- crassifolia* Sarg. Fruit short-oblong, orange-red, pale dotted, 10–11 mm. diameter.
- grossa* Sarg. Fruit subglobose to short-oblong, dull orange-red, 8–10 mm. diameter.
- illuminata* Sarg. (?) Color of anthers unknown. Fruit characters as given above.
- mercerensis* Sarg. Fruit short-oblong, orange-red, 10–12 mm. diameter.
- repentina* Sarg. Fruit subglobose to slightly ovoid, red, 12–13 mm. diameter. Stamens sometimes 20.
- rotundata* Sarg. Fruit short-oblong, orange-red blotched with red and with pale lenticles, 15–20 mm. diameter.
- rotundifolia* Moench. Fruit subglobose, dark crimson, dark dotted, 8–10 mm. diameter. Corymbs villous in variety.
- varians* Sarg. Fruit subglobose, dark red blotched with green, pale dotted, 10–12 mm. diameter. Stamens 10–15.
- Williamsii* Egglest. (?) Color of anthers unknown. Fruit characters as given above.
CC. Corymbs at flowering time villous or pubescent
- Brunetiana* Sarg. Fruit oblong, bright clear red, pale dotted, 9–10 mm. diameter.
- caesariata* Sarg. Fruit oval or slightly ovoid, dark crimson, dark dotted, 7–12 mm. diameter.

coccinata Sarg. Fruit short-oblong to slightly obovoid, very dark crimson, pale dotted, 11–12 mm. diameter.

divergens Sarg. Fruit short-oblong to subglobose, scarlet, pale dotted, 4–5 mm. (or more) diameter. Stamens sometimes 18–20.

Faxonii Sarg. Fruit oblong, dark crimson with pale lenticles, 8–10 mm. diameter.

Jackii Sarg. Fruit ovoid to oblong, angled, dull dark red, pale dotted, 10–12 mm. diameter.

Keepii Sarg. Fruit oblong, bright clear red, white dotted, 9–10 mm. diameter.

praetermissa Sarg. Fruit short-oblong to ovoid, crimson, pale dotted, 8–9 mm. diameter.

subrotundifolia Sarg. Fruit short-oblong, crimson, pale dotted, 10–15 mm. diameter.

BB. Stamens about 20 (15–20)

C. Corymbs at flowering time glabrous or nearly so

inaudita Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, 12–14 mm. diameter.

Macauleyae Sarg. Fruit subglobose to short-oblong, dark crimson, pale dotted, 10–12 mm. diameter.

Margareta Ashe. Fruit subglobose, reddish or orange, about 10 mm. diameter. Corymbs sometimes slightly villous.

Maribella Sarg. Fruit short-oblong, crimson, pale dotted, 9–10 mm. diameter.

repentina Sarg. Stamens sometimes only 10. Fruit characters as given above.

varians Sarg. Stamens usually about 10. Fruit characters as given above.

CC. Corymbs at flowering time villous or pubescent

divergens Sarg. Stamens sometimes only 10. Fruit characters as given above.

Margareta Ashe. Corymbs usually glabrous. Fruit characters as given above.

noveboracensis Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, about 10 mm. diameter.

Oakesiana Egglest. Fruit pyriform to oblong, yellowish-red, about 10 mm. diameter.

XIII. Microcarpae

A. Anthers pink or rose; stamens about 20

apiifolia Michx. Fruit short-oblong, bright scarlet, 4–5 mm. diameter. Leaves pubescent, deeply lobed.

Phaenopyrum Medic. Fruit subglobose, scarlet, 4–6 mm. diameter. Leaves glabrous, slightly lobed.

AA. Anthers cream or yellow; stamens about 20

spathulata Michx. Fruit subglobose, bright scarlet, 4–5 mm. diameter. Leaves narrow, spatulate.

Youngii Sarg. Fruit globose to subglobose, scarlet, 5–7 mm. diameter.
Leaves broadly ovate or deltoid in outline.

XIV. Flavae

- A. Anthers pink, rose or purple (AA see p. 117)
- B. Stamens about 10
 - C. Leaves at flowering time glabrous or nearly so
 - alleghaniensis* Beadle. Fruit globular-pyriform, red, 8–12 mm. diameter.
 - frugiferens* Beadle. Fruit globose or subglobose, red, 9–13 mm. diameter.
Leaves sometimes slightly villous.
 - CC. Leaves at flowering time villous or pubescent
 - frugiferens* Beadle. Leaves usually glabrous or nearly so. Fruit characters as given above.
 - villaris* Beadle. (?) Fruit pyriform, yellow or orange-yellow flushed with red, 8–11 mm. diameter. Floral characters unknown.
 - BB. Stamens about 20 (15–20)
 - C. Leaves at flowering time glabrous or nearly so
 - adunca* Beadle. Fruit globose, red, 12–15 mm. diameter.
 - clara* Beadle. Fruit globose, orange-red, 9–12 mm. diameter. Leaves at flowering time sometimes slightly villous.
 - extraria* Beadle. Fruit subglobose or oval, red, 9–12 mm. diameter.
 - ignava* Beadle. Fruit globose or subglobose, red or orange-red, 8–12 mm. diameter.
 - limata* Beadle. Fruit globose, red, 10–13 mm. diameter.
 - quaesila* Beadle. Fruit subglobose or slightly pyriform, red and orange, 8–11 mm. diameter. Young leaves sometimes slightly villous.
 - sodalis* Beadle. (?) Fruit pyriform, orange-red, 8–12 mm. diameter.
Leaves sometimes slightly villous. Color of anthers unknown.
 - valida* Beadle. Fruit subglobose, color not given, 12–15 mm. diameter.
 - CC. Leaves at flowering time villous or pubescent
 - arrogans* Beadle. Fruit short-oval to slightly pyriform, red, 9–10 mm. diameter.
 - clara* Beadle. Leaves usually nearly glabrous at flowering time. Fruit characters as given above.
 - colonica* Beadle. (?) Fruit pyriform, orange-red, 10–13 mm. diameter.
Color of anthers unknown.
 - condigna* Beadle. Fruit subglobose or pyriform, red or orange and greenish, 6–8 mm. diameter.
 - consanguinea* Beadle. Fruit globose to depressed-globose, red, 9–12 mm. diameter.
 - egregia* Beadle. Fruit subglobose, orange or flushed with red, 10–12 mm. diameter.
 - flava* Ait. Leaves at flowering time sometimes nearly glabrous. Fruit characters as given above.
 - illudens* Beadle. (?) Fruit pyriform, yellow, 8–11 mm. diameter. Color of anthers unknown.

invicta Beadle. (?) Fruit pyriform, color not given, 6–8 mm. diameter.
Color of anthers unknown.

lepidia Beadle. (?) Fruit subglobose, orange or orange-red, 7–11 mm. diameter. Color of anthers unknown.

Michauxii Pers. (?) Fruit subglobose, red or ruddy, 8–12 mm. diameter.
Color of anthers unknown.

munda Beadle. (?) Fruit pyriform, color not given, 7–9 mm. diameter.
Leaves sometimes nearly glabrous. Color of anthers unknown.

pexa Beadle. Fruit globose, color not given, 7–9 mm. diameter. Color
of anthers unknown.

quaesita Beadle. Leaves sometimes glabrous or nearly so. Fruit char-
acters as given above.

segnis Beadle. (?) Fruit globose, red, 8–12 mm. diameter. Color of
anthers unknown.

sodalis Beadle. (?) Leaves sometimes nearly glabrous. Color of anthers
unknown. Fruit characters as given above.

tristis Beadle. Fruit short-oval, red or orange-red, 10–12 mm. diameter.

vicenda Beadle. Fruit pyriform, yellow or flushed with red, 10–12 mm.
diameter.

yadkinensis Ashe. (?) Fruit pyriform or subglobose, dull red or orange,
7–11 mm. diameter. Color of anthers unknown.

AA. Anthers white, cream or yellow

B. Stamens about 10 (5–10)

C. Leaves at flowering time glabrous or nearly so

impar Beadle. Fruit oval, red, 10–13 mm. diameter. Stamens some-
times 15 or more.

CC. Leaves at flowering time villous or pubescent

aprica Beadle. Fruit globose, red or orange-red, 9–14 mm. diameter.

mira Beadle. Fruit subglobose, color not given, 10–12 mm. diameter.
Stamens sometimes 15–20.

BB. Stamens about 20

C. Leaves at flowering time glabrous or nearly so

abdita Beadle. Fruit globose, red, 10–15 mm. diameter. Leaves sometimes
slightly villous.

attrita Beadle. Fruit subglobose, yellow, splashed with red, 10–14 mm.
diameter. Leaves sometimes slightly villous.

calva Beadle. Fruit globose, yellow or orange-red, 7–10 mm. diameter.

impar Beadle. Stamens sometimes only 10. Fruit characters as given
above.

lacrimata Beadle. Fruit globose or subglobose, yellow or orange and
red, 8–10 mm. diameter.

sodalis Beadle. (?) Leaves at flowering time sometimes slightly villous.
Color of anthers unknown. Fruit characters as given above.

sororia Beadle. Fruit globose, red or red and yellow, 12–18 mm. diameter.
Leaves usually somewhat villous.

teres Beadle. Fruit oblong, red, 8–10 mm. diameter.

CC. Leaves at flowering time villous or pubescent
abdita Beadle. Leaves sometimes glabrous or nearly so. Fruit characters
as given above.

adusta Beadle. Fruit short-pyriform, orange-red, spotted and streaked
with red, 9–11 mm. diameter.

alabamensis Beadle. Fruit oval, red, 10–15 mm. diameter.

amica Beadle. Fruit subglobose, orange, blotched with red, 10–13 mm.
diameter.

anisophylla Beadle. Fruit globose or subglobose, orange or orange and
red, 6–8 mm. diameter.

annosa Beadle. Fruit subglobose or oval, orange-red or red and orange,
10–12 mm. diameter.

arguta Beadle. Fruit globose, red, 7–9 mm. diameter.

attrita Beadle. Leaves usually glabrous or nearly so. Fruit characters
as given above.

audens Beadle. Fruit pyriform, orange-yellow flushed with red, 9–12
mm. diameter.

colonica Beadle. (?) Color of anthers unknown. Fruit characters as given
above.

cirrata Beadle. Fruit globose or subglobose, red, 7–9 mm. diameter.

compitalis Beadle. Fruit subglobose or oval, red, 10–13 mm. diameter.

constans Beadle. Fruit subglobose or slightly pyriform, bright orange-
red, 10–12 mm. diameter.

crocea Beadle. Fruit pyriform, yellow to russet-red, 9–12 mm. diameter.

cullasagensis Ashe. Fruit globular or slightly oblong, dark orange,
mottled with orange-red and crimson, 10–14 mm. diameter.

curva Beadle. Fruit pyriform, orange or orange and red, 6–9 mm. diam-
eter.

dapsilis Beadle. Fruit globose or subglobose, yellow or orange and red,
10–15 mm. diameter.

dispar Beadle. Fruit subglobose or oval, red, 7–10 mm. diameter.

dolosa Beadle. Fruit globose or short-oval, yellow or orange, 9–12 mm.
diameter.

egens Beadle. Fruit subglobose or slightly oval, orange-red or orange and
red, 9–11 mm. diameter.

exilis Beadle. Fruit globose or subglobose, red or ruddy, 5–7 mm. diam-
eter.

florens Beadle. Fruit globose, orange-red, 10–15 mm. diameter.

floridana Sarg. Fruit ovoid or short-oblong, bright orange-red, pale
dotted, 7–9 mm. diameter.

frugalis Beadle. Fruit subglobose, red, 7–9 mm. diameter.

furtiva Beadle. Fruit pyriform, orange or orange and red, 7–9 mm.
diameter.

galbana Beadle. Fruit globose, red, 9–13 mm. diameter.

- geniculata Ashe. (?) Fruit pyriform, lemon-yellow or orange mottled with red, 7-12 mm. diameter. Color of anthers unknown.
- illudens Beadle. (?) Color of anthers unknown. Fruit characters as given above.
- incana Beadle. Fruit pyriform, orange-yellow or orange and red, 8-9 mm. diameter.
- inopina Beadle. Fruit pyriform, yellow or orange-yellow and red, 8-11 mm. diameter.
- inops Beadle. Fruit globose, orange or orange and red, 10-14 mm. diameter.
- insidiosa Beadle. Fruit oval or subglobose, red, 9-12 mm. diameter.
- integra Beadle. Fruit globose, red, 10-15 mm. diameter.
- invicta Beadle. (?) Color of anthers unknown. Fruit characters as given above.
- lanata Beadle. Fruit globose, color not given, 8-11 mm. diameter.
- lassa Beadle. Fruit pyriform, orange-red, 8-10 mm. diameter.
- laxa Beadle. Fruit subglobose or short-oblong, red, 7-9 mm. diameter.
- lepidia Beadle. (?) Color of anthers unknown. Fruit characters as given above.
- meridiana Beadle. Fruit pyriform, orange, yellow or yellow flushed with red, 7-9 mm. diameter.
- Michauxii Pers. (?) Color of anthers unknown. Fruit characters as given above.
- mundula Beadle. (?) Color of anthers unknown. Fruit characters as given above.
- panda Beadle. Fruit globose or depressed-globose, orange-red or orange tinged with red, 10-15 mm. diameter.
- pexa Beadle. (?) Color of anthers unknown. Fruit characters as given above.
- polla Beadle. Fruit subglobose to oval, orange-yellow flushed with red, 9-12 mm. diameter.
- Ravenelii Sarg. Fruit globose to short-oblong, bright orange-red, dark dotted, 10-12 mm. diameter.
- recurva Beadle. Fruit pyriform or short-pyriform, red, 7-9 mm. diameter.
- sesima Beadle. Fruit pyriform, orange-red, blotched with red, 9-11 mm. diameter.
- imosa Beadle. Fruit short-pyriform, yellow or orange-yellow and red, 9-12 mm. diameter.
- egnis Beadle. Color of anthers unknown. Fruit characters as given above.
- enta Beadle. Fruit globose, red, 10-14 mm. diameter.
- odalis Beadle. (?) Color of anthers unknown. Fruit characters as given above.
- ororia Beadle. Leaves sometimes nearly or quite glabrous. Fruit characters as given above.

versuta Beadle. Fruit short-pyriform, orange or greenish-yellow and red, 6–9 mm. diameter.

vicana Beadle. Fruit pyriform or oblong-pyriform, yellow or orange, blotched with red, 10–14 mm. diameter.

yadkinensis Ashe. (?) Color of anthers unknown. Fruit characters as given above.

XV. Bracteatae

Anthers yellow; stamens about 20 (15–20); corymbs at flowering time villous or pubescent

Ashei Beadle. Fruit subglobose or short-oval, bright red, 10–14 mm. diameter.

Harbisonii Beadle. Fruit globose, red, 10–13 mm. diameter.

XVI. Triflorae

Anthers white, cream or yellow in this group

A. Stamens about 10

austromontana Beadle. Fruit globose, bright red, 12–15 mm. diameter.

AA. Stamens about 20

conjugens Sarg. Fruit subglobose, orange-red, about 10 mm. diameter.
triflora Chapman. Fruit globose, red, 12–15 mm. diameter.

XVII. Uniflorae

A. Anthers pink, rose or purple; stamens about 20

arenicola Ashe. (?) Fruit pyriform, orange or orange-red, 8–10 mm. diameter. Color of anthers unknown.

bisulcata Ashe. (?) Fruit pyriform, yellow or greenish-yellow, 10–12 mm. diameter. Color of anthers unknown.

Croomiana Sarg. Fruit ovoid, yellow or greenish-yellow, 8–9 mm. diameter.

rhodella Ashe. Fruit pyriform, orange and red, 8–12 mm. diameter.

raleighensis Ashe. (?) Fruit globular, yellow or greenish, 7–9 mm. diameter. Color of anthers unknown.

AA. Anthers white, cream or yellow; stamens about 20

arenicola Ashe. (?) Color of anthers unknown. Fruit characters as given above.

armentalis Beadle. Fruit globose, red, about 10 mm. diameter.

bisulcata Ashe. (?) Color of anthers unknown. Fruit characters as given above.

Brittonii Egglest. Fruit globose or short-pyriform, reddish-brown, 8–12 mm. diameter.

choriophylla Sarg. Fruit subglobose to slightly ovoid, orange-red, 10–12 mm. diameter.

Earlei Ashe. Fruit pyriform, yellow or greenish-yellow, 7–9 mm. diameter.

gregalis Beadle. Fruit pyriform, red or ruddy, 10–14 mm. diameter.

raleighensis Ashe. (?) Color of anthers unknown. Fruit characters as given above.

Smithii Sarg. Fruit short-oblong, orange or orange-green, about 12 mm. diameter.

trianthophora Sarg. Fruit ovoid, light orange-red, pale dotted, 8-12 mm. diameter.

uniflora Moench. Fruit subglobose, yellow or greenish-yellow, 10-14 mm. diameter.

XVIII. Aestivales

Anthers red, rose or purple in this group

A. Young leaves and branchlets glabrous or nearly so

aestivalis T. & G. Fruit short-oblong or subglobose, scarlet, 12-15 mm. diameter.

fruticosa Sarg. Fruit subglobose, light orange-red, pale dotted, about 10 mm. diameter. Young leaves sometimes slightly villous.

monantha Sarg. Fruit subglobose or short-oblong, cherry-red, 8-10 mm. diameter. Young leaves sometimes slightly villous.

AA. Young leaves and branchlets villous

fruticosa Sarg. Young leaves usually nearly glabrous. Fruit characters as given above.

monantha Sarg. Young leaves sometimes slightly villous. Fruit characters as given above.

opaca Hook. Fruit depressed-globose, scarlet, pale dotted, 15-20 mm. diameter.

rufula Sarg. Fruit subglobose, scarlet, about 12 mm. diameter.

XIX. Brachyacanthae

Anthers yellow; stamens about 20

brachyacantha Sarg. & Engelm. Fruit subglobose or ovoid, bright blue (straw-color in var. *leucocarpa* Sarg.), 12-15 mm. diameter.

saligna Greene. Fruit globose, blue-black, 5-8 mm. diameter.

XX. Douglassianae

A. Anthers pink, rose or purple

B. Stamens about 10 (5-10)

C. Corymbs and leaves glabrous or nearly so at flowering time

cerronis Nelson. Fruit globose, red, 8-10 mm. diameter.

erythropoda Ashe. Fruit subglobose, red or orange and red, 7-8 mm. diameter.

Douglasii Lindl. Fruit short-oblong, black (chestnut-color in f. *badia* Sarg.), 10-12 mm. diameter. Corymbs sometimes slightly villous. Stamens sometimes 20.

CC. Corymbs and leaves villous at flowering time

Douglasii Lindl. Corymbs usually glabrous. Fruit characters as given above. Stamens sometimes 20.

rivularis Nutt. Fruit short-oblong, crimson, pale dotted, becoming black, about 10 mm. diameter.

BB. Stamens about 20

Douglasii Lindl. Stamens sometimes 5–10. Fruit characters as given above.

AA. Anthers white, cream or yellow; stamens about 10 (5–10); corymbs and leaves villous at flowering time

colorado Ashe. Fruit subglobose, bright red, 7–8 mm. diameter.

columbiana Howell. Fruit ovoid, scarlet, 8–12 mm. diameter.

Piperi Britton. Fruit spherical or nearly so, coral-red, about 12 mm. diameter.

XXI. Anomalae

Anthers pink, rose or purple in this group

A. Stamens about 10 (5–10)

B. Corymbs at flowering time glabrous or nearly so

asperifolia Sarg. Fruit oblong, bright scarlet, about 10 mm. diameter. Corymbs sometimes slightly villous.

Brockwayae Sarg. Fruit short-oblong to slightly ovoid, dark vinous-purple, 10–12 mm. diameter.

cyclophylla Sarg. Fruit subglobose to short-oblong, orange-red, pale dotted, about 10 mm. diameter.

Dunbarii Sarg. Fruit subglobose, crimson, dark dotted, 12–14 mm. diameter. Corymbs sometimes slightly villous.

Egglestonii Sarg. Fruit oblong, orange becoming crimson, pale dotted, about 10 mm. diameter. Corymbs sometimes slightly villous.

floridula Sarg. Fruit short-oblong to ovoid, crimson, 7–8 mm. diameter.

improvisa Sarg. Fruit ovoid-oblong, orange-red, 8–9 mm. diameter.

inopinata Sarg. Fruit subglobose, dark red, pale dotted, 7–9 mm. diameter.

Corymbs sometimes slightly villous.

Knieskerniana Sarg. Fruit short-oblong, scarlet, pale dotted, 10–11 mm. diameter.

scabrida Sarg. Fruit subglobose, scarlet, 8–12 mm. diameter. (Anthers described as pale yellow, but are pink or purple in specimens.)

BB. Corymbs at flowering time villous

affinis Sarg. Fruit short-oblong, oval or slightly ovoid, orange-red with pale lenticles, 9–10 mm. diameter.

asperifolia Sarg. Corymbs usually glabrous or nearly so. Fruit characters as given above.

Dunbarii Sarg. Corymbs often glabrous or nearly so. Fruit characters as given above.

Egglestonii Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.

honesta Sarg. Fruit subglobose, cherry-red, pale dotted, 7–8 mm. diameter.

Ideae Sarg. Fruit short-oblong to subglobose, crimson, pale dotted, about 10 mm. diameter.

inopinata Sarg. Corymbs sometimes nearly glabrous. Fruit characters as given above.

misella Sarg. Fruit short-oblong, crimson, pale dotted, about 10 mm. diameter.

pinguis Sarg. Fruit short-oblong, crimson, pale dotted, 8–10 mm. diameter.

repulsans Sarg. Fruit short-oblong to subglobose, orange-red, pale dotted, 9–10 mm. diameter.

AA. Stamens about 20 (15–20)

C. Corymbs at flowering time glabrous or nearly so

brachyloba Sarg. Fruit short-oblong to ovoid, orange-red with pale lenticles, 8–10 mm. diameter.

Coleae Sarg. Fruit subglobose, bright orange-red, pale dotted, 8–10 mm. diameter.

errata Sarg. Fruit short-oblong, dark red with pale lenticles, 10–12 mm. diameter.

fallsiana Sarg. Fruit subglobose, scarlet, pale dotted, 14–15 mm. (or less) diameter.

putata Sarg. Fruit oval, orange-red, dark dotted, about 8 mm. diameter.

Seelyana Sarg. Fruit short-oblong, dull orange-red, about 7 mm. diameter.

urbana Sarg. Fruit short-oblong, crimson, pale dotted, 9–10 mm. diameter.

CC. Corymbs at flowering time villous

Saundersiana Sarg. Fruit short-oblong, ovoid or depressed-globose, orange-red, with pale lenticles, about 10 mm. diameter.

shirleyensis Sarg. Fruit subglobose to short-oblong, orange-red, about 10 mm. diameter.

virilis Sarg. Fruit short-oblong to ovoid, scarlet, dark dotted, 8–9 mm. diameter.

XXII. Macracanthae

A. Anthers pink, rose or purple (AA see p. 127)

B. Stamens about 10 (BB see p. 125)

C. Corymbs at flowering time glabrous or nearly so

bristolensis Sarg. Fruit oval, bright scarlet, pale dotted, 10–12 mm. diameter.

cupulifera Sarg. Fruit globose to short-oblong, scarlet, dark dotted, about 10 mm. diameter. Corymbs sometimes slightly villous.

diaphora Sarg. Fruit ovoid, crimson, pale dotted, 13–15 mm. diameter. Corymbs sometimes slightly villous.

divida Sarg. Fruit short-oblong to subglobose, crimson with pale lenticles, 10–12 mm. diameter.

dumicola Sarg. Fruit subglobose, dark red, pale dotted, about 10 mm. diameter. Corymbs sometimes slightly villous.

laetifica Sarg. Fruit subglobose to ovoid, crimson, pale dotted, 10–14 mm. (or less) diameter. Stamens sometimes 15–20.

- nuda* Sarg. Fruit short-oblong to subglobose, bright red, about 10 mm. diameter.
- Peckietta* Sarg. Fruit ovoid, red or greenish-red, 9–11 mm. diameter.
- CC. Corymbs at flowering time villous or pubescent
- aquilonaris* Sarg. Fruit oblong, dull greenish-red, 8–10 mm. diameter.
- baccata* Sarg. Fruit subglobose, orange-red, pale dotted, 7–8 mm. diameter.
- Balkwillii* Sarg. Fruit short-oblong, orange-red blotched with yellow, 7–8 mm. diameter.
- Beckiana* Sarg. Fruit subglobose to ovoid, crimson, pale dotted, about 10 mm. diameter.
- Chapmanii* Ashe. Fruit globose or subglobose, bright red, 8–10 mm. diameter.
- coloradensis* Nelson. Fruit globose, bright red, about 8 mm. diameter.
- conspicua* Sarg. Fruit subglobose to short-oblong, crimson, 8–10 mm. diameter. Stamens sometimes 20.
- cupulifera* Sarg. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.
- Deweyana* Sarg. Fruit subglobose to short-oblong, scarlet with pale lenticles, 10–12 mm. diameter.
- diaphora* Sarg. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.
- dumicola* Sarg. Corymbs sometimes glabrous. Fruit characters as given above.
- fallax* Ashe. (?) Fruit depressed-globose, crimson, mottled with green and olive, 13–16 mm. diameter. Color of anthers unknown.
- fertilis* Sarg. Fruit subglobose, short-oblong or rarely ovoid, dark red, pale dotted, about 10 mm. diameter.
- flammea* Sarg. Fruit short-oblong to subglobose, orange-red, pale dotted, 10–11 mm. diameter.
- fulgens* Sarg. Fruit subglobose, crimson, pale dotted, 10–12 mm. diameter.
- illinoiensis* Ashe. Fruit globose, scarlet, dark dotted, 8–11 mm. diameter.
- insperata* Sarg. Fruit ovoid, light orange-red, 8–10 mm. diameter.
(This species was described as having 20 stamens, probably through error.)
- integriloba* Sarg. Fruit subglobose, bright scarlet, with pale lenticles, 10–12 mm. diameter.
- laurentiana* Sarg. Fruit oblong, dark crimson, 8–10 mm. diameter.
- obscura* Sarg. Fruit ovoid, light orange-red, pale dotted, 9–10 mm. diameter.
- ogdensbergensis* Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, 9–11 mm. diameter.
- prinoides* Sarg. Fruit globose or depressed-globose, scarlet, dark dotted, 8–9 mm. diameter.
- prunifolia* Pers. Fruit subglobose, dull crimson, 10–15 mm. diameter.

- rhombifolia* Sarg. Fruit subglobose, bright clear red, 8–9 mm. diameter.
- Robinsonii* Sarg. Fruit oblong to slightly ovoid, bright scarlet, dark dotted, 7–8 mm. diameter.
- rupicola* Sarg. Fruit subglobose to short-oblong, bright orange-red, 9–11 mm. diameter.
- sheridana* Nelson. (?) Fruit spherical, scarlet, 8–9 mm. diameter.
- spinulosa* Sarg. Fruit short-oblong, orange-red, 5–6 mm. diameter.
- BB. Stamens about 20 (15–20)
- C. Corymbs at flowering time glabrous or nearly so
- Heleneae* Sarg. Fruit short-oblong or ovoid, scarlet, 8–10 mm. diameter.
- laetifica* Sarg. Stamens sometimes 15 or more. Fruit characters as given above.
- menandiana* Sarg. Fruit subglobose, scarlet, pale dotted, about 12 mm. diameter. Corymbs sometimes slightly villous.
- micrantha* Sarg. Fruit subglobose, orange-red, pale dotted, 7–8 mm. diameter.
- neofluvialis* Ashe. Fruit subglobose, greenish-orange or flushed with red, 6–10 mm. diameter. (Stamens described as 10–15, but apparently about 20.)
- pisifera* Sarg. Fruit subglobose, crimson, pale dotted, 7–8 mm. diameter. Corymbs sometimes slightly villous.
- propixa* Sarg. Fruit subglobose to ovoid, crimson, pale dotted, 7–8 mm. diameter.
- radiosa* Sarg. Fruit subglobose, bright cherry-red, about 10 mm. diameter.
- rutila* Sarg. Fruit subglobose, orange-red, pale dotted, 8–10 mm. diameter. Corymbs sometimes slightly villous.
- simulata* Sarg. Fruit short-oblong, dull orange-red, 7–9 mm. diameter. Corymbs sometimes slightly villous.
- spinea* Sarg. Fruit subglobose to short-oblong, dark red, dark dotted, 6–7 mm. diameter.
- stenophylla* Sarg. Fruit short-oblong to ovoid, orange-red, pale dotted, 8–9 mm. diameter. Corymbs sometimes slightly villous.
- vaga* Sarg. Fruit ovoid or oval, orange-red, 6–7 mm. diameter. Corymbs sometimes slightly villous.
- CC. Corymbs at flowering time villous or pubescent
- admiranda* Sarg. Fruit subglobose to short-oblong or ovoid, bright cherry-red, pale dotted, 12–14 mm. diameter.
- ardua* Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, 12–14 mm. diameter.
- calpodendron* Medic. (?) Fruit pyriform or ellipsoid, red or orange-red, 6–8 mm. diameter.
- Calvinii* Sarg. Fruit oval to ovoid, scarlet, pale dotted, 8–10 mm. diameter.
- conspicua* Sarg. Stamens 10–20, according to description. Fruit characters as given above.

- diversa Sarg. Fruit oval or slightly ovoid. crimson, pale dotted, 6–7 mm. diameter.
- efferata Sarg. Fruit subglobose, scarlet, pale dotted, 8–9 mm. diameter.
- ensifera Sarg. Fruit oval, bright orange-red, pale dotted, 8–9 mm. diameter.
- finitima Sarg. Fruit short-oblong to subglobose, orange-red, pale dotted, about 10 mm. diameter.
- frutescens Sarg. Fruit short-oblong, scarlet, pale dotted, 7–8 mm. diam.
- Gaultii Sarg. Fruit subglobose to short-oblong, scarlet becoming dark crimson, 12–14 mm. diameter.
- gemmosa Sarg. Fruit subglobose to short-oblong, scarlet, 12–14 mm. diameter.
- globosa Sarg. Fruit globose to short-oblong, bright orange-red, dark dotted, 7–8 mm. diameter.
- hispidula Sarg. Fruit ovoid, orange-red, 7–8 mm. diameter
- honeoyensis Sarg. Fruit short-oblong or slightly ovoid, scarlet, pale dotted, 8–10 mm. diameter.
- longispina Sarg. Fruit subglobose, scarlet, pale dotted, 8–10 mm. diam.
- manandiana Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.
- missouriensis Ashe. Fruit pyriform or becoming subglobose, deep scarlet, 9–10 mm. diameter.
- obesa Ashe. (?) Fruit globose or slightly oblong, scarlet, 8–9 mm. diameter. Color of anthers unknown.
- pisifera Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.
- pubifolia Ashe. Fruit subglobose to short-oblong, scarlet or orange-red, 6–8 mm. diameter.
- rutila Sarg. Corymbs sometimes nearly or quite glabrous, Fruit characters as given above.
- saeva Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, 8–10 mm. diameter.
- scabera Sarg. Fruit subglobose to slightly ovoid, bright cherry-red, dark dotted, about 10 mm. diameter.
- simulata Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.
- sonnenbergensis Sarg. Fruit subglobose or short-oblong, crimson, about 10 mm. diameter.
- spinifera Sarg. Fruit subglobose to slightly ovoid, scarlet, pale dotted, 10–12 mm. diameter.
- stenophylla Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.
- succulenta Schrad. Fruit globose, bright scarlet, 6–8 mm. diameter.
- tomentosa L. Fruit ovoid, rarely subglobose, dull orange-red, 5–10 mm. diameter.

- vaga* Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.
- Vailiae* Britton. Fruit subglobose, yellowish-green becoming red, 8–10 mm. diameter.
- vegeta* Sarg. Fruit subglobose, scarlet, pale dotted, 8–10 mm. diameter.
- AA. Anthers white, cream or yellow
- B. Stamens about 10 (5–10)
 - C. Corymbs at flowering time glabrous or nearly so
- chadfordiana* Sarg. Fruit subglobose, scarlet, pale dotted, 12–14 mm. diameter.
- delectabilis* Sarg. Fruit ovoid, scarlet, pale dotted, 8–9 mm. diameter. Corymbs sometimes slightly villous.
- Doddsii* Ramaley. Fruit pyriform, bright red, about 9 mm. diameter.
- Emersoniana* Sarg. Fruit globose to subglobose, scarlet, pale dotted, 12–14 mm. diameter.
- glabrata* Sarg. Fruit subglobose, crimson blotched with yellow, pale dotted, 8–9 mm. diameter.
- Handya* Sarg. Fruit subglobose, bright scarlet, pale dotted, 10–12 mm. diameter.
- peramoena* Sarg. Fruit short-oblong to ovoid, light scarlet blotched with yellow, pale dotted, 8–10 mm. diameter.
- picta* Sarg. Fruit subglobose, scarlet, pale dotted, 8–9 mm. diameter.
- spatiosa* Sarg. Fruit oval to globose or depressed-globose, scarlet blotched with yellow, pale dotted, 12–14 mm. diameter.
- venustula* Sarg. Fruit subglobose, crimson, pale dotted, about 10 mm. diameter.
- CC. Corymbs at flowering time villous or pubescent
- agaia* Sarg. Fruit subglobose to short-oblong, dark crimson, pale dotted, 10–12 mm. diameter.
- armigera* Sarg. Fruit short-oblong to ovoid, crimson, pale dotted, 8–10 mm. diameter.
- coloradoides* Ramaley. Fruit globose, red or purplish, about 9 mm. diam.
- crux* Ashe. Fruit subglobose, bright scarlet, about 13 mm. diameter.
- debilis* Sarg. Fruit short-oblong, scarlet, 9–11 mm. diameter.
- delectabilis* Sarg. Corymbs sometimes glabrous or nearly so. Fruit characters as given above.
- Doddsii* Ramaley. Corymbs sometimes nearly glabrous. Fruit characters as given above.
- fallax* Ashe. (?) Color of anthers unknown. Fruit characters as above.
- ferentaria* Sarg. Fruit subglobose to short-oblong, crimson, pale dotted, about 12 mm. diameter.
- flagrans* Sarg. Fruit subglobose, dark crimson, 8–10 mm. diameter.
- fulgida* Sarg. Fruit subglobose, dark crimson, pale dotted, 10–12 mm. diameter.
- hystricina* Ashe. Fruit subglobose, bright scarlet, about 10 mm. diameter.

- macracantha* Lodd. Fruit globose, crimson, 5–6 mm. diameter.
- microsperma* Sarg. Fruit short-oblong to subglobose, crimson, pale dotted, 7–10 mm. diameter.
- mollicula* Sarg. Fruit subglobose to short-oblong, bright orange-red, pale dotted, 10–12 mm. diameter.
- Laneyi* Sarg. Fruit subglobose, dark orange-red, pale dotted, about 6 mm. diameter.
- pellucidula* Sarg. Fruit subglobose, scarlet, pale dotted, 12–14 mm. diameter.
- peramoena* Sarg. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.
- pertomentosa* Ashe. Fruit subglobose, red or yellowish (becoming dark red), 6–9 mm. diameter.
- praeclara* Sarg. Fruit subglobose to ovoid, crimson, 8–10 mm. diameter.
- pudens* Sarg. Fruit short-oblong, green or yellowish-red, 8–9 mm. diam.
- Searsii* Sarg. Fruit subglobose, crimson, pale dotted, 8–10 mm. diameter.
- stratfordensis* Sarg. Fruit oval, greenish-yellow mottled with crimson, 7–8 mm. diameter.
- BB. Stamens about 20 (15–20)
- C. Corymbs at flowering time glabrous or nearly so
- opica* Ashe. Fruit depressed-globose, crimson, about 12 mm. (or less) diameter. Corymbs sometimes finely pubescent.
- tanophylla* Sarg. Fruit short-oblong, crimson, 7–10 mm. diam.
- vernans* Ashe. Fruit globose, bright scarlet, 9–12 mm. diameter. Corymbs sometimes slightly villous.
- CC. Corymbs at flowering time villous or pubescent
- ambrosia* Sarg. Fruit short-oblong, crimson, dark dotted, 8–12 mm. diameter.
- carrollensis* Sarg. Fruit ellipsoidal, dark red, 6–7 mm. diameter.
- comans* Sarg. Fruit short-oblong to subglobose, orange-red, 7–8 mm. diameter.
- ferta* Sarg. Fruit subglobose to short-oblong or ovoid, scarlet, pale dotted, 12–14 mm. (or usually less) diameter.
- obesa* Ashe. (?) Color of anthers unknown. Fruit characters as above.
- opica* Ashe. Corymbs sometimes nearly or quite glabrous. Fruit characters as given above.
- structilis* Ashe. Fruit oblong or pyriform, orange or reddish-orange dotted with dark green and russet, 7–10 mm. diameter.
- truculenta* Sarg. Fruit subglobose, dark red, pale dotted, 7–8 mm. diameter.
- venulosa* Sarg. Fruit subglobose to ovoid or short-oblong, orange-red, 7–8 mm. diameter.
- vernans* Ashe. Corymbs usually glabrous, according to description. Fruit characters as given above.



BHL

Biodiversity Heritage Library

Palmer, Ernest J. 1925. "Synopsis of North American Crataegi." *Journal of the Arnold Arboretum* 6(1-2), 5-128. <https://doi.org/10.5962/p.185187>.

View This Item Online: <https://www.biodiversitylibrary.org/item/33585>

DOI: <https://doi.org/10.5962/p.185187>

Permalink: <https://www.biodiversitylibrary.org/partpdf/185187>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.