

Evidence for aboveground activity of *Zambian Molerats* (*Cryptomys*, *Bathyergidae*, *Rodentia*)

By A. SCHARFF and O. GRÜTJEN

Abt. Allgemeine Zoologie, Universität-GH Essen, Essen

Receipt of Ms. 29. 11. 1996

Acceptance of Ms. 09. 02. 1997

By definition, subterranean mammals spend most of their lives in underground burrows, where they find their food and reproductive partners, and most of them come to the surface only incidentally. The physical and biotic uniqueness of subterranean ecotope (which is dark, stable, monotonous, hypoxic, hypercapnic, etc.) leads to an adaptive convergence at all levels of organizations (cf., e.g. NEVO and REIG 1990). Consequently these mammals have become unique subjects for studies of (among others) sensory biology and physiology of metabolism.

Particularly *Zambian mole-rats* of the genus *Cryptomys* (*Bathyergidae*, *Rodentia*) have been very intensively studied within the last decade concerning their hearing and acoustic communication (CREDNER et al. 1996), orientation in space (BURDA et al. 1990; LINDENLAUB et al. 1995), and physiology of metabolism (MARHOLD and NAGEL 1995; BENNETT et al. 1994).

While all these studies have elucidated interesting adaptations to an underground life style, they have been carried out in laboratories, and the central question remains as for how strictly are these animals confined to underground life. Do they also, incidentally or even regularly, come into contact with daylight, aboveground sensory cues or climate? There are only few direct (GENELLY 1965; OATLEY and ANSELL 1966; SHEPPE 1973) and indirect observations (e.g. skulls in owl pellets) indicating aboveground activity in mole-rats and related species (cf. DE GRAAFF 1981).

During a six-months' field study in Zambia (April–October 1996) we found some indirect evidence that mole-rats of the genus *Cryptomys* spend more time outside the burrow system than generally assumed. We studied common mole-rats (*Cryptomys spec.*, 2 n = 68) in Lusaka and giant mole-rats (*Cryptomys mechowii*) in Ndola. We excavated burrow systems in different areas and caught the animals. Although we never observed mole-rats aboveground, local people and mole-rat hunters reported on mole-rats moving on the surface particularly during early morning hours.

Altogether we excavated 12 nests. Three nests consisted completely of plastic material (nylon stockings, pieces of plastic bags, and sacks). Three further nests contained a mixture of plastic materials, dry grass and root fibres. These six nests were situated near cultivated fields and human buildings. The six remaining nests contained a mixture of grass and root fibres. They were situated far away from human settlements. It seemed that mole-rats preferred plastic materials (if available) to grass or roots. It is obvious that mole-rats must have collected most of their nest material aboveground.

In a food chamber of a colony near Lusaka we found two seeds of the tree *Parinari curatellifolium* (*Chrysobalanaceae*). Also this finding indicates that mole-rats may even sometimes forage aboveground.

We found two small colonies of giant mole-rats (3 and 4 animals) in a "dambo" (a seasonally inundated area) near Ndola. Compared to normal habitats (e.g. cultivated fields), the burrow systems from dambos were small (50 m² versus >300 m²). The burrow systems were surrounded by areas where the level of ground water was near or above the surface (at the end of the dry season), so it was impossible for the mole-rats to reach this area underground. Most probably a breeding pair reached the dambo aboveground and established a new colony. This is supported by the finding that both families consisted of two adult animals (male and female) and one or two subadult individuals. Normally, giant mole-rats live in large colonies of even about 40 animals.

These findings indicate that mole-rats of the genus *Cryptomys* obviously spend more time outside their burrows than assumed so far. Most of the aboveground activity occurs probably at night.

Acknowledgement

This work was supported by a travel grant of the DAAD (Deutscher Akademischer Austauschdienst) to A. S.; and was carried out within the framework of the research project "Behavioural ecology and systematics of Zambian mole-rats" (Prof. Dr. H. BURDA, Abt. Allgemeine Zoologie, Universität-GH Essen). We thank Prof. Dr. H. BURDA for comments on the manuscript and Dr. P. M. S. PHIRI (University of Zambia) for the identification of seeds and plants.

References

- BENNETT, N. C.; AGUILAR, G. H.; JARVIS, J. U. M.; FAULKES, C. G. (1994): Thermoregulation in three species of Afrotropical subterranean mole-rats (Rodentia: Bathyergidae) from Zambia and Angola and scaling within the genus *Cryptomys*. *Oecologia* **97**, 222–227.
- BURDA, H.; MARHOLD, S.; WESTENBERGER, T.; WILTSCHKO, W.; WILTSCHKO, R. (1990): Magnetic compass orientation in the subterranean rodent *Cryptomys hottentotus* (Bathyergidae, Rodentia). *Experientia* **46**, 528–530.
- Credner, S.; Burda, H.; Ludescher, F. (1996): Acoustic communication underground: Vocalization characteristics in subterranean social mole-rats (*Cryptomys* sp., Bathyergidae). *J. Comp. Physiol. A*. (in press).
- DE GRAAFF, G. (1981): The rodents of Southern Africa. Durban, Pretoria: Butterworth.
- GENELLY, R. E. (1965): Ecology of the common mole-rat (*Cryptomys hottentotus*) in Rhodesia. *J. Mammalogy* **46**, 647–665.
- Lindenlaub, T.; Burda, H.; Nevo, E. (1995): Convergent evolution of the vestibular organ in the subterranean mole-rats, *Cryptomys* and *Spalax*, as compared with aboveground rat, *Rattus*. *J. Morphol.* **224**, 303–311.
- Marhold, S.; Nagel, A. (1995): The energetics of the common mole-rat *Cryptomys*, a subterranean eusocial rodent from Zambia. *J. Comp. Physiol. B* **164**, 636–645.
- NEVO, E.; REIG, O. A. (1990): Evolution of subterranean mammals at the organismal and molecular levels. New York: Alan R. Liss.
- OATLEY, T. B.; ANSELL, W. F. H. (1966): *Cryptomys mechowii* above ground. *Puku* **4**, 193–194.
- SHEPPE, W. A. (1973): Notes on Zambian rodents and shrews. *Puku* **7**, 167–190.

Authors' address: Dipl.-Biol. ANDREAS SCHARFF, OLIVER GRÜTJEN, Abt. Allgemeine Zoologie, FB 9; Universität-GH Essen, D-45117 Essen



Scharff, Andreas and Grütjen, Oliver. 1997. "Evidence for aboveground activity of Zambian Molerats (*Cryptomys*, *Bathyergidae*, *Rodentia*)." *Zeitschrift für Säugetierkunde : im Auftrage der Deutschen Gesellschaft für Säugetierkunde e.V* 62, 253–254.

View This Item Online: <https://www.biodiversitylibrary.org/item/163239>

Permalink: <https://www.biodiversitylibrary.org/partpdf/192298>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: Deutsche Gesellschaft für Säugetierkunde

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.