## Butterflies in a light trap (Lepidoptera: Papilionoidea)

Stanislav ABADJIEV

Samenvatting. Dagvlinders in een lichtval (Lepidoptera: Papilionoidea) De vangst van twee dagvlindersoorten - Lasiommata roxelana (CRAMER, 1777) en Quercusia quercus (LINNAEUS, 1758) - in een lichtval tijdens een nachtvangst in de oostelijke Rhodopen (Zuid-Bulgarije) op 7 juli 1992, wordt vermeld.

Résumé. Papillons diurnes dans un piège lumineux (Lepidoptera: Papilionoidea) La capture de deux espèces de papillons diurne - *Lasiommata roxelana* (CRAMER, 1777) et *Quercusia quercus* (LINNAEUS, 1758) - dans un piège lumineux lors d'une chasse nocturne dans les Rhodopes orientales (sud de la Bulgarie) le 7 juillet 1992, est rapportée.

Abadjiev, S.: Block 415, Entrance G, Flat 80, Druzhba 2, BG-1582 Sofia, Bulgaria.

In July 1992 my fellow-colleague and friend Stoyan BESCHKOW and I went on a lepidopterological trip to the Eastern Rhodopes in South Bulgaria. On 7th July we reached a small chalet near the town of Madjarovo. The place is situated at about 50 m close to the Arda River. Around was a mixed deciduous forest with predominance of *Quercus* spp.

On the same evening we switched on a 400 W Hg lamp. At 11.30 local time I observed a specimen of a butterfly fluttering near the lamp. I was very astonished to see that it was a worn male of *Lasionmata roxelana* (CRAMER, 1777). The night was very humid and with no wind whatsoever. Some time later at 0.40 I saw another species flying around the lamp. This time it was a female of *Quercusia quercus* (LINNAEUS, 1758).

This reminded me of a similar observation of collecting *Kanetisa circe* (FABRICIUS, 1775) in a light trap in Andorra (ROCHE 1990). I have only heard about the penumbral butterflies in the Tropics, but it was abnormal to see our European species flying at night. It is very interesting that the two species I collected must be absolutely nocturnal. What is it? What prompted them to do this? I am still not sure that this is characteristic for the behaviour of European species. Maybe there is an alternate explanation. I think that most likely their eyesight perception had mixed the sunlight with the strong light of the mercury lamp.

## Reference

Roche, P., 1990. A satyrid butterfly in a light trap. - Entomologist's Rec. J. Var. 102: 114.



Abadjiev, Stanislav. 1993. "Butterflies in a light trap (Lepidoptera: Papilionoidea)." *Phegea* 21, 27–27.

View This Item Online: <a href="https://www.biodiversitylibrary.org/item/207230">https://www.biodiversitylibrary.org/partpdf/192896</a>

Holding Institution Natural History Museum Library, London

**Sponsored by** Natural History Museum Library, London

**Copyright & Reuse** Copyright Status: In Copyright. Digitized with the permission of the rights holder Rights Holder: Vlaamse Vereniging voor Entomologie License: <u>http://creativecommons.org/licenses/by-nc-sa/4.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.