

The female of *Cochylimorpha salinarida* (Lepidoptera: Tortricidae, Tortricinae)

Anton Cox & Hugo van der Wolf

Summary. The female genitalia of the Tortricid moth *Cochylimorpha salinarida* Groenen & Larsen, 2003 are described and illustrated.

Samenvatting. Het vrouwtje van *Cochylimorpha salinarida* (Lepidoptera: Tortricidae, Tortricinae)

Het vrouwelijk genitaal van de Tortricidae mot *Cochylimorpha salinarida* Groenen & Larsen, 2003 wordt beschreven en afgebeeld.

Résumé. La femelle de *Cochylimorpha salinarida* (Lepidoptera: Tortricidae, Tortricinae)
Les génitalia femelles du Tortricidae *Cochylimorpha salinarida* Groenen & Larsen, 2003 sont décrits et figurés.

Zusammenfassung. Dass Weibchen von *Cochylimorpha salinarida* (Lepidoptera: Tortricidae, Tortricinae)

Die weibliche Genitalien von *Cochylimorpha salinarida* Groenen & Larsen, 2003 (Tortricidae) werden beschrieben und abgebildet.

Key words: Tortricidae – *Cochylimorpha salinarida* – female genitalia – Spain.

Cox, A. L.: Generaal Gavinstraat 11, NL 6585 WL Mook, Netherlands.

van der Wolf, H. W.: Wermersland 22, NL 5673 PT Nuenen, Netherlands.
h.vanderwolf@onsnet.nu.

Introduction

Several years ago *Cochylimorpha salinarida* Groenen & Larsen, 2003 was described in this journal from 3 male specimens collected by Mr. J. Wolschrijn (Twello, Netherlands) in the Spanish Province of Alicante (Groenen & Larsen 2003). It is now possible to describe the female from some specimens collected in the same region by the first author of this paper.

Material examined: "Espagne, Alicante, La Marina, 11–12.X.2005, leg. A. Cox", 2♂, 1♀ gen. prep. Wf. 9919, 1♀; "Espagne, Alicante, Royales, 12–14.X. 2005, leg. A. Cox", 2♂, 1♀ gen. prep. Wf. 9918, 1♀. (in collections Cox and van der Wolf).

Diagnosis. The female adults do not differ from the male ones. The wingspan of the male and female specimens varies between 15 and 21 mm.

Female genitalia (Fig. 1): Papillae anales stretched; apophyses long, stout; antrum weakly sclerotised; sterigma oblong, wider than long; ductus bursae short; corpus bursae as in *C. perturbatana* (Kennel, 1900), but first section oval, weakly sclerotised, with a curved sclerite, second section sack-like, transparent; ductus seminalis emerging from median part of corpus bursae.

Distribution: The locality Royales is about 16 km SW of the type locality La Marina. Unlike the type locality, which is on the coast, Royales is about 8 km land inwards.

Biology: The hostplant is still unknown.

Remark: The shape of the female genitalia confirms that the species is most closely related to *C. perturbatana* (Razowski 1970: plate 127, fig. 80).




Fig. 1. *Cochylimorpha salinarida* Groenen & Larsen, female genitalia, gen. prep. Wf. 9919.

References

- Groenen, F. & Larsen, K. 2003. *Cochylimorpha salinarida* sp.n. (Lepidoptera, Tortriciae: Tortricinae), a new Cochylid moth from Spain. — *Phegea* **31**(1): 11–14.
- Razowski, J. 1970. Cochylidae. – in Amsel, H. G., Gregor, F. & Reisser, H. (Eds), *Microlepidoptera Palaearctica* Vol. 3. — Verlag Georg Fromme & Co., Wien. 528 pp., 161 pls.



Cox, Anton and Wolf, Hugo W. van der. 2001. "The female of *Cochylimorpha salinarida* (Lepidoptera: Tortricidae, Tortricinae)." *Phegea* 36, 21–22.

View This Item Online: <https://www.biodiversitylibrary.org/item/193363>

Permalink: <https://www.biodiversitylibrary.org/partpdf/193139>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.