


ON *HELICINA PTEROPHORA*, N.SP., FROM GUATEMALA.

By E. R. SYKES, B.A., F.L.S., etc.

Read 8th November, 1901.

Testa depressa, conoidea, tenuiuscula, sub lente dense malleata, cornea, infra suturas et ad peripheriam zona brunnea notata, apice obtusulo; anfr. $5\frac{1}{2}$, plano-convexiusculi, ultimus non descendens, acute carinatus; apertura subtriangularis, intus ad carinam castanea, callum tenuissime circumscriptum emittens; peristoma album, expansum, reflexiusculum, ad carinam valde rostratum, margine columellari sinuato, basali reflexo. Operculum ignotum. Diam. max. 22, min. 12 mm.; alt. 8 mm.

*Helicina pterophora*, n.sp.

Hab.—Guatemala (coll. Boucard).

This striking shell recalls in form *H. rhyncostoma*, Shuttl., but may be readily separated by the difference in sculpture, the present species not being spirally striate, and also by its larger size. From *H. rostralis*, Morel., its shape will readily distinguish it, while *H. oxyrhynca*, Crosse & Debeaux, appears to be a much smaller and more conic species.



Sykes, E. R. 1902. "ON HELICINA PTEROPHORA, N.SP., FROM GUATEMALA."
Proceedings of the Malacological Society of London 5, 20–20.

View This Item Online: <https://www.biodiversitylibrary.org/item/52297>

Permalink: <https://www.biodiversitylibrary.org/partpdf/202781>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.