

Collection and selection of natural enemies of twospotted spider mites for biological control

D.R. GILLESPIE and D.J.M. QUIRING

**PACIFIC AGRI-FOOD RESEARCH CENTRE, P.O. BOX 1000,
AGASSIZ, BC, V0M 1A0**

M. GREENWOOD

**DEPARTMENT OF BIOLOGICAL SCIENCES, SIMON FRASER UNIVERSITY,
BURNABY, BC, V5A 1S6**

ABSTRACT

Natural enemies of the twospotted spider mite, *Tetranychus urticae* (Koch) were surveyed (at three locations) in the lower Fraser Valley, British Columbia. Our objective was to identify predators that might be used for the biological control of spider mites in greenhouse tomato crops. Three to four pots of bean plants infested with spider mites were exposed each week at each of three locations from early April through to October, in both 1992 and 1993. Predators were collected from these trap plants, identified and tested as potential predators of spider mites. Twenty-two species were collected, more than half of them predatory Hemiptera. Two species, *Feltiella acarisuga* (Vallot) (Diptera: Cecidomyiidae), and *Stethorus punctum picipes* (Casey) (Coleoptera: Coccinellidae) were specialist predators of spider mites. *Feltiella acarisuga* was the most promising candidate for use in biological control of spider mites (on tomatoes) based on the frequency with which it occurred on trap plants, its cosmopolitan distribution, and its monophagous feeding habits.

Key words: biological control, British Columbia, greenhouse, tomato, *Tetranychus urticae*, *Feltiella acarisuga*, *Therodiplosis persicae*, predators.

INTRODUCTION

Biological control of twospotted spider mites, *Tetranychus urticae* (Koch) (Acari: Tetranychidae), by the predatory mite *Phytoseiulus persimilis* (Dosse & Bravenboer) (Acari: Phytoseiidae) is common on cucumber and pepper crops in greenhouses in British Columbia (Costello *et al.* 1992) and elsewhere (Lenteren & Woets 1988). Twospotted spider mites are also pests of greenhouse-grown tomatoes, and biological control using *P. persimilis* has been described (Hussey & Scopes 1985). Outbreaks of twospotted spider mites are common on greenhouse grown tomato crops in B.C. but attempts at biological control with *P. persimilis* have generally failed. The increase of populations of *P. persimilis* on tomato crops is reduced by mortality of motile stages through entanglement on glandular hairs on stems and leaf petioles (Haren *et al.* 1987), and by reduced life span and fecundity of adults due to contact with tomato leaves (Gillespie & Quiring 1994).

As one of many possible solutions to this problem, a different natural enemy could be used either to replace or to supplement *P. persimilis*. Importation of exotic predators should be done cautiously, however, and potential endemic natural enemies should be screened first. Consequently, the predators of spider mites were surveyed at three locations in the lower Fraser Valley, BC, using trap plants baited with spider mites. We report here the results of that survey, and of the initial screening process used to select candidate species for further study.




Figure 1. Map of the Lower Fraser Valley, B.C. Showing sample locations: 1. Green Timbers Nursery, Surrey, B.C.; 2. Agriculture and Agri-Food Canada Sub-Station, Abbotsford; 3. Agriculture and Agri-Food Canada Research Station, Agassiz.

METHODS AND MATERIALS

Ten to twelve lima bean (*Phaseolus lunatus* L.) seeds were planted in compost mix in standard "8 inch" plastic pots, weekly from early April to late September of 1992 and 1993. After three to four weeks in a greenhouse at 22°C the primary leaves had completed growth, and the first secondary leaves were maturing. The plants were then lightly infested with approximately 200 adult and immature twospotted spider mites per pot, and taken to three different locations. At each location (Fig. 1) three to four pots were placed in 30 cm diam., plastic saucers, filled with water, under a 90 x 90 x 90 cm, plastic-roofed shelter with open sides. These plants were left unattended for one week.

After each week the trap plants were replaced with pots of fresh plants. The pots of exposed plants were carefully placed in plastic tote boxes with tight fitting, screened lids, and returned to the laboratory. Any obvious herbivores were removed, and known predators of spider mites were preserved and catalogued. Predators with unknown prey preferences and species whose role in the food chain was not obvious (eg. nymphs of some Hemiptera) were provided with spider mites on pieces of bean leaf for 24 h. Any insects that did not feed on spider mites were discarded. Species of unknown identity were reared to adults on a diet of spider mites and preserved for identification.

At site 1, in Surrey, trap plants were placed on mowed grass at the interface between a large open grass field, and a mature second-growth coniferous forest. At site 2, in Abbotsford, plants were in a mixed grass and clover fallow field. The site was surrounded by a windbreak of deciduous and coniferous trees, and much of the surrounding land was dedicated to raspberry culture. At site 3, in Agassiz, plants were among long grass at the interface between grass pasture and an extensive blackberry (*Rubus* spp.) thicket. The wild area behind the blackberry was mixed deciduous and coniferous forest.

RESULTS AND DISCUSSION

More than 1500 specimens, representing 22 species of predator in seven orders and 14 families were collected from trap plants in 1992 and 1993 (Table 1). These species were either known predators of spider mites, or were observed to feed on them in the laboratory.

Most of the species (12 of 22) were Hemiptera and could be classed as generalist predators, feeding on spider mites and other small arthropods (see tables and citations in Chazeau 1985).

With the exception of *Heterotoma meriopterum* (Scopoli) (Hemiptera: Miridae), all hemipterans were collected as adults. First instar nymphs of *H. meriopterum* were found on trap plants on several occasions. These were reared to adults on a diet of spider mites. It appears that adults of this species might lay eggs in or near to spider mite colonies.

Of the ten non-Hemipterous species, seven can be classed as generalist predators. Three species appeared to specialize as predators of spider mites. These were *Amblyseius* sp. (Acari: Phytoseiidae), *Feltiella acarisuga* (Vallot) (Diptera: Cecidomyiidae) and *Stethorus punctum-picipes* (Casey) (Coleoptera: Coccinellidae). Of these three, only *F. acarisuga* was collected at all three sites, and in virtually every month of the survey; it was the commonest predator. *Feltiella* spp. are predators of spider mites in many agricultural crops and are often considered the most important predators (Oatman *et al.* 1985, Pickett & Gilstrap 1986, Sharaf 1984). Moreover, *F. acarisuga* is virtually cosmopolitan (Gagne, 1995). In contrast, the other specialist feeder, *S. punctum picipes*, is restricted to western North America (Gordon 1985), and only 13 specimens were collected on 3 occasions.

Our objective was to select one or more natural enemies that might replace *P. persimilis* as a biological control for spider mites on greenhouse tomato crops. *Phytoseiulus persimilis* is ineffective on tomato plants, partly because motile stages are trapped on glandular hairs on stems and leaf petioles (Haren *et al.* 1987). Other phytoseiids and predators that walk between leaves to locate prey might also get trapped. Therefore trap plants were placed in water-filled saucers, in part to exclude such predators. This strategy was apparently successful, as only two species of predacious mites were found on trap plants. In a survey of spider mites on strawberry plants in the Fraser Valley, Raworth (1990) collected seven species of predacious Acari. All of the arthropod species found in this survey were collected either as winged adults, or as eggs, and first instars from eggs deposited by winged adults. The adults might therefore be able to avoid glandular hairs on stems and petioles when searching for prey.

The use of trap plants baited with target pest species and placed in areas where that pest is at a low density has been advocated as a way to select natural enemies for use in classical biological control (Pschorn-Walcher 1977). The natural enemies collected on such plants can presumably locate and feed on prey at low population densities. In this study, we placed trap plants baited with spider mites where spider mites were not obviously in outbreak, but might have been present naturally. At all sites, plants were placed in or near areas of mixed vegetation, separated from cultivated crops. Although spider mite density was not determined in surrounding vegetation, the lack of obvious damage on local plants presumably indicated low numbers. We therefore assumed that any natural enemies, particularly specialist natural enemies, that appeared on trap plants originated from nearby spider mite colonies that were not in outbreak or from distant outbreaks.

One of the desirable characteristics of natural enemies used for inundative biological control in greenhouses is their ability to locate hosts readily (Lentern & Woets 1988). Predators that discover spider mites on trap plants have either done so by chance or by some, presumably semiochemical, cues. It can be argued that those species that are known specialist predators of spider mites, and that also occur with regularity on trap plants probably have an efficient system for locating prey. In this context, *F. acarisuga* is an obvious candidate because of its consistently high numbers.

Table 1.

Species of predatory insects collected on trap plants baited with twospotted spider mite, *Tetranychus urticae* (Koch).

Species	Common Name	Type	Stage Collected ¹	# Collected ²
Acari: Anystidae				
<i>Anystis agilis</i> Banks	predatory mite	generalist	I, A	54
Phytoseiidae				
<i>Amblyseius</i> sp.	predatory mite	generalist	I,A	23
Hemiptera: Miridae				
	plant bugs			31
<i>Heterotomma meriopterum</i> (Scopoli)		generalist	I	
<i>Dicyphus</i> sp.		generalist	A	
<i>Deraeocoris brevis</i> Knight		generalist	A	
<i>Campyloneura virgula</i> (H.-S.)		generalist	A	
Lygaeidae				
	big-eyed bugs			
<i>Geocoris</i> sp		generalist	A	1
Anthocoridae				
	flower bugs			35
<i>Orius tristicolor</i> (White)		generalist	A	
<i>Orius minutus</i> (L.)		generalist	A	
<i>Anthocoris antevoleus</i> White		generalist	A	
Berytidae				
	stilt-legged bugs			
<i>Neides muticus</i> (Say)		generalist	A	1
Nabidae				
	damsel bugs			33
<i>Nabis alternatus</i> Parshley		generalist	A	
<i>Nabis roseipennis</i> Reuter		generalist	A	
<i>Nabis rufusculus</i> Reuter		generalist	A	
Diptera: Cecidomyiidae				
<i>Feltiella acarisuga</i> Vallot	predatory gall midge	specialist	E, I	1224
Syrphidae				
<i>Scavaea pyrastris</i> (L.)	flower fly	“generalist”	I	1
Thysanoptera: Aeolothripidae				
<i>Aeolothrips fasciatus</i> (L.)	thrips	generalist	A	32
Neuroptera: Chrysopidae				
<i>Chrysopa</i> sp.	green lacewing	generalist	I	40
Hemerobiidae				
<i>Wesmaelius</i> sp.	brown lacewing	generalist	E, I	3
Dermaptera: Forficulidae				
<i>Forficula auricularia</i> (L.)	european earwig	generalist	A	
Coleoptera: Coccinellidae				
<i>Stethorus punctum picipes</i> (Casey)	spider mite destroyer	specialist	E, I, A	13
<i>Coccinella</i> sp.	ladybird beetle	generalist	A	1

¹ stage collected on plants. E = egg, I = immature, A = adult

² total number collected at 3 sites during 2 years

A second important criterion for inundative biological control agents is that they should be able to complete their life history on the prey (Lentern & Woets 1988). Those predators found on infested baited trap plants as eggs and immatures probably meet this criterion. *Feltiella acarisuga*, *Wesmaelius* sp., and *Stethorus punctum-picipes* all occurred as both eggs and immatures. In addition, we have seen *F. acarisuga* among spider mite colonies in greenhouses, apparently invading naturally. Our preliminary test release of *F. acarisuga* on tomato plants in a greenhouse became established and reproduced. On the basis of this evidence *F. acarisuga* was selected for further investigation as a potential candidate for inundative biological control of spider mites.

ACKNOWLEDGEMENTS

We thank Darren Hiebert, Corrine Kikegawa and Tanya Lenton for technical assistance, and the B.C. Greenhouse Vegetable Research Council for funding that assisted in the conduct of this work. Insects were identified by M.D. Schwartz, I.M. Smith, E. Maw, and J.R. Vockeroth, all of Eastern Cereals and Oilseeds Research Centre, Agriculture and Agri-Food Canada, Ottawa, ON. This is contribution number 568 from the Pacific Agri-Food Research Centre, Agriculture and Agri-Food Canada, Agassiz, BC.

REFERENCES

- Chazeau, J. 1985. Predaceous Insects. pp. 211-246 IN Helle, W., Sabelis, M.W. Eds. Spider Mites. Their Biology, Natural Enemies and Control. Elsevier. Vol. 2. Amsterdam, Oxford, New York, Tokyo.
- Costello, R., Elliot, D.P., Gilkeson, L.A., and Gillespie, D.R. . 1992. Integrated control of greenhouse pests. Edition. B.C. Ministry of Agriculture, Fisheries and Food. Victoria, B.C. 19 pp
- Gillespie, D.R., Quiring, D.M.J. 1994. Reproduction and longevity of the predatory mite *Phytoseiulus persimilis* (Acari: Phytoseiidae) and its prey, *Tetranychus urticae* (Acari: Tetranychidae) on different host plants. Journal of the Entomological Society of British Columbia. 91: 3-8
- Gordon, R.D. 1985. The Coccinellidae (Coleoptera) of America North of Mexico. Journal of the New York Entomological Society. 93: 1-912.
- Haren, R.J.F.van, Steenhuis, M.M., Sabelis, M.W., Ponti, O.M.B.de, 1987. Tomato stem trichomes and dispersal success of *Phytoseiulus persimilis* relative to its prey *Tetranychus urticae*. Experimental and Applied Acarology. 3: 115-121
- Hussey, N.W., Scopes, N.E.A. 1985. Greenhouse Vegetables (Britain). pp. 285-297. IN Helle, W., Sabelis, M.W. Eds. Spider Mites. Their Biology, Natural Enemies and Control. Elsevier. Amsterdam, Oxford, New York, Tokyo.
- Lenteren, J.C. van, Woets, J. 1988. Biological and integrated pest control in greenhouses. Annual Review of Entomology. 33: 239-269
- Oatman, E.R., M.E. Bagley, and G.R. Platner. 1985. Predators of the two-spotted spider mite on strawberry. California Agriculture 39 (1/2): 9-12.
- Pickett, C.H., and F.E. Gilstrap. 1986. Natural enemies associated with spider mites (Acari: Tetranychidae) infesting corn in the High Plains region of Texas. Journal of the Kansas Entomological Society 59: 524-536.
- Pschorn-Walcher, H. 1977. Biological control of forest insects. Annual Review of Entomology. 22: 1-22
- Raworth, D.A. 1990. Predators associated with the twospotted spider mite, *Tetranychus urticae*, on strawberry at Abbotsford, BC, and development of non-chemical mite control. Journal of the Entomological Society of British Columbia. 87: 59-67
- Sharaf, N.S. 1984. Studies on natural enemies of tetranychid mites infesting eggplants in the Jordan Valley. Zeitschrift für Angewandte Entomologie 98: 527-5



Gillespie, David Roy, Quiring, Donald, and Greenwood, M. 1997. "Collection and selection of natural enemies of twospotted spider mites for biological control." *Journal of the Entomological Society of British Columbia* 94, 7–12.

View This Item Online: <https://www.biodiversitylibrary.org/item/181001>

Permalink: <https://www.biodiversitylibrary.org/partpdf/213830>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: Entomological Society of British Columbia

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.