Clarifying the nomenclature of *Crateva trifoliata* (Capparaceae)

I.M. Turner

Research Associate, Royal Botanic Gardens, Kew Richmond, Surrey TW9 3AE, U.K.

ABSTRACT. Capparis trifoliata Roxb., the supposed basionym of 'Crateva adansonii subsp. trifoliata (Roxb.) Jacobs' and 'Crateva trifoliata (Roxb.) B.S.Sun', is shown to represent an illegitimate renaming of Crateva religiosa G.Forst. However, Jacobs effectively excluded the nomenclatural type of Capparis trifoliata, thus avoiding creating a superfluous name, and instead described Crateva adansonii subsp. trifoliata Jacobs validated by the citation of the Latin description of Crateva erythrocarpa Gagnep. Sun also excluded the type of Crateva religiosa, but his effective transfer of C. adansonii subsp. trifoliata to specific rank does not have priority over Crateva erythrocarpa Gagnep., which is therefore the correct name for this taxon when it is considered a distinct species. The name Crateva nurvula Buch.-Ham. is formally lectotypified.

Keywords. Capparis, China, Crateva, India, nomenclature, Roxburgh, typification

Introduction

William Roxburgh named an Indian species Capparis trifoliata. The name was included in Roxburgh's Hortus Bengalensis (Roxburgh, 1814: 41) but as no description was included the name remained invalid. A brief description was eventually published in the second edition of Flora Indica (Roxburgh, 1832: vol. 2, p. 571), 17 years after Roxburgh had died. Jacobs (1964), in a revision of the genus Crateva, included Capparis trifoliata within the wide-ranging (seasonally dry tropical Africa and Asia) and variable Crateva adansonii DC. In order to categorise the variation within the species, Jacobs recognised a series of five subspecies separable on leaf characters. These consisted of Crateva adansonii subsp. adansonii in Africa, C. adansonii subsp. odora (Buch.-Ham.) Jacobs in India, C. adansonii subsp. trifoliata (Roxb.) Jacobs in Indochina, C. adansonii subsp. formosensis Jacobs in China and Taiwan and C. adansonii subsp. axillaris (C.Presl) Jacobs in the Philippines and Java. While some recent works have favoured splitting up Crateva adansonii, others have maintained Jacobs's system (Chayamarit, 1991; Liu & Liao, 1996; Philcox, 1996). However, there is a nomenclatural problem with *Capparis trifoliata* Roxb., upon which Jacobs evidently based C. adansonii subsp. trifoliata. In the original validation in Flora Indica, Roxburgh cited 'Crateva religiosa Willd. 2. 853' in synonymy. This is a reference to Willdenow's Species Plantarum. In turn, Willdenow referred to Vahl (1794: 62) and Forster (1786: 203). The latter element (also referred to by Vahl) relates to the original publication of Crateva religiosa G.Forst. Roxburgh, Willdenow and Vahl, in their somewhat confused view, also all referred to the Hortus Malabaricus element 'niirvala' (Van Rheede tot Draakestein, 1682: 49-50, t. 42) - now considered to represent a separate species. Roxburgh stated that his species was native to India and certainly included what Jacobs later referred to as Crateva adansonii subsp. odora as shown by a Roxburgh specimen from William Hooker's herbarium now at Kew and Roxburgh's Icones no. 157 also at Kew. Jacobs (1964: 199) referred to a Roxburgh specimen in the Herbarium of the East India Company as the holotype of Capparis trifoliata. This does seem to be a specimen of Crateva adansonii subsp. trifoliata. It is mounted with another Roxburgh specimen but Jacobs made it clear that he was referring to the specimen annotated '1' on the sheet. The origins of the specimen are unknown. Nomenclaturally, Capparis trifoliata Roxb. represents a superfluous renaming of Crateva religosa and, as no type was indicated by Roxburgh, it is typified by the type of C. religiosa (McNeill et al., 2012: Art. 7.5). Therefore Jacob's subspecies' name also appears to be based on Crateva religiosa. Similarly so does 'Crateva trifoliata (Roxb.) B.S.Sun' which has been used in recent Chinese floras (Sun, 1999; Zhang & Tucker, 2008).

However, apparently unwittingly, Jacobs (1964) effectively excluded the type of *Crateva religiosa* from *C. adansonii* subsp. *trifoliata* by recognising *Crateva religiosa* as a taxon distinct from *C. adansonii* subsp. *trifoliata* in the same paper. Such exclusion by implication is permitted under the code (McNeill et al., 2012: Art. 52.2; cf. Ex. 7). While Jacobs provided a description in English of *Crateva adansonii* subsp. *trifoliata* this is not sufficient to validate a new taxon at this date. However Jacobs also cited *Crateva erythrocarpa* Gagnep. in synonymy. This was published by Gagnepain with a Latin description. Therefore *Crateva adansonii* subsp. *trifoliata* is validated by the Latin description of *C. erythrocarpa*.

Sun (1999) also seemed to base *Crateva trifoliata* on *Capparis trifoliata* Roxb. However, Sun excluded *Crateva religiosa* G.Forst. from the flora of China, noting that the application of the name to the species used to make fish lures in Taiwan and the Ryukyus was erroneous. Therefore, like Jacobs, Sun excluded Roxburgh's nomenclatural type and effectively created a new combination at species rank based on Jacobs's subspecies. But this combination only has priority at species rank from the date of Sun's publication. Therefore the correct name at species rank is *Crateva erythrocarpa* Gagnep.

Crateva adansonii subsp. *trifoliata* Jacobs, Blumea 12: 199 (1964). – *Crateva trifoliata* (Jacobs) B.S.Sun, Fl. Reipubl. Popularis Sin. 32: 489 (1999). – TYPE: *W. Roxburgh* s.n. (EIC 6972C, specimen annot. 1) (holotype K-W [barcode no. K001126439]).

Crateva erythrocarpa Gagnep., Bull. Soc. Bot. France 55: 322 (1908). – TYPE: Vietnam, montagne de Chaudoc, July 1876, F.-J. Harmand 608 (lectotype P [barcode no. P05427398], designated by Jacobs (1964: 199)).

Crateva religiosa G.Forst., Diss. Pl. Esc. 45–46 (1786). – *Capparis trifoliata* Roxb., Fl. Ind. 2: 571 (1832), *nom. illeg.*, *superfl.* – TYPE: ?India, *G. Forster* s.n. (lectotype K [barcode no. K000651081], designated by Jacobs (1964: 191)).

Roxburgh, Willdenow and Vahl all cited the Van Rheede element 'niirvala' in synonymy. This has also been considered to be a separate species, *Crateva nurvala* Buch.-Ham. Jacobs (1964: 194) effectively lectotypified the name with a Buchanan-Hamilton specimen in the Wallich Herbarium (K-W). However, as Nicolson et al. (1988) have noted, Buchanan-Hamilton did not consider his own specimens to be the true *Crateva nurvala* but a variety of it. Therefore Jacobs's typification is incorrect. Nicolson et al. (1988) stated 'However it appears that Van Rheede's illustration should be taken as the type.' It is questionable whether this is a valid typification – the opening of the sentence casts doubt making it unclear that the authors are accepting their typification. I therefore formally typify the name here. Note that after the type of a Loureiro name was refound *Crateva nurvala* was included in synonymy of *Crateva magna* (Lour.) DC. (Jacobs, 1976).

Crateva magna (Lour.) DC., Prodr. 1: 243 (1824). – Capparis magna Lour., Fl. Cochinch. 1: 331 (1790). – TYPE: Cochinchina, J. Loureiro s.n. (lectotype BM [barcode no. BM000629693], designated by Jacobs (1976: 822)).

Crateva nurvala Buch.-Ham., Trans. Linn. Soc. London 15: 121 (1827), as 'nürvala'. – TYPE: Van Rheede, Hort. Malab. 3: t. 42 (1682) (lectotype, designated here).

ACKNOWLEDGEMENTS. I am very grateful to Prof. J. McNeill (E) for guidance on nomenclatural matters and Wong Wei San (SUTD) for Chinese translation. Comments from two anonymous reviewers helped improve the paper.

References

Chayamarit, K. (1991). Capparaceae. In: Smitinand, T. & Larsen, K. (eds) *Flora of Thailand* 5(3): 241–271. Bangkok: Royal Forest Department.

Forster, J.R. (1786). Florulae Insularum Australium Prodromus. Goettingen: Joann. Christian Dieterich.

Jacobs, M. (1964). The genus Crateva. Blumea 12: 177-208.

Jacobs, M. (1976). Capparaceae. In: Van Steenis, C.G.G.J. (ed) *Flora Malesiana* 7(4): 822. Leyden: Noordhoff International Publishing.

Liu, T.-S. & Liao, J.-C. (1996). Capparaceae. In: Huang, T.-C. (ed) *Flora of Taiwan*, 2nd edition, 2: 734–744. Taipei: Editorial Committee of the Flora of Taiwan.

- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Prud'homme van Reine, W.F., Smith, G.F., Wiersema, J.H. & Turland, N.J. (2012). *International Code of Nomenclature for algae, fungi, and plants (Melbourne Code)*. Regnum Vegetabile 154. Königstein: Koeltz Scientific Books.
- Nicolson, D.H., Suresh, C.R. & Manilal, K.S. (1988). An Interpretation of Van Rheede's Hortus Malabaricus. Regnum Vegetabile 119. Königstein: Koeltz Scientific Books.
- Philcox, D. (1996). Capparaceae. In: Dassanayake, M.D. (ed) *A Revised Handbook to the Flora of Ceylon* 10: 23–50. Rotterdam: A.A. Balkema.
- Roxburgh, W. (1814). Hortus Bengalensis. Serampore: Mission Press.
- Roxburgh, W. (1832). Flora Indica, vol.1–3. Calcutta: W. Thacker & Co.
- Sun, B. (1999). Capparaceae (excluding *Cleome*). In: Wu, C. (ed) *Flora Reipublicae Popularis Sinicae* 32: 484–531. Beijing: Science Press.
- Vahl, M. (1794). Symbolae Botanicae, vol. 3. Copenhagen: Nicolaus Möller et filius.
- Van Rheede tot Draakestein, H.A. (1682). *Horti Indici Malabarica pars Tertia*. Amsterdam: Van Someren & Van Dyck.
- Zhang, M. & Tucker, G.C. (2008). Capparaceae. In: Wu, Z., Raven, P.H. & Hong, D. (eds) *Flora of China* 7: 433–450. Beijing: Science Press.



Turner, I. M. 2015. "Clarifying the nomenclature of Crateva trifoliata (Capparaceae)." *The Gardens' bulletin, Singapore* 67(2), 403–406. https://doi.org/10.3850/s2382581215000320.

View This Item Online: https://www.biodiversitylibrary.org/item/223140

DOI: https://doi.org/10.3850/s2382581215000320

Permalink: https://www.biodiversitylibrary.org/partpdf/229530

Holding Institution

Singapore Botanic Gardens, National Parks Board Singapore

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: http://creativecommons.org/licenses/by-nc-sa/4.0/

Rights: https://biodiversitylibrary.org/permissions

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.