Parasitic Diatoms Inside Antarctic Sponges

GIORGIO BAVESTRELLO^{1,*}, ATTILIO ARILLO², BARBARA CALCINAI², RICCARDO CATTANEO-VIETTI², CARLO CERRANO², ELDA GAINO³, ANTONELLA PENNA⁴, AND MICHELE SARÀ²

¹Istituto di Scienze del Mare—Università di Ancona, Via Brecce Bianche, I-60131 Ancona, Italy; ²Dipartimento per lo Studio del Territorio e delle sue Risorse—Università di Genova, Viale Benedetto XV, 5 I-16132 Genova, Italy; and ³Dipartimento di Biologia animale ed Ecologia—Università di Perugia, Via Elce di Sotto I-06123 Perugia, Italy; and ⁴Istituto di Biochimica—Università di Urbino, I-60118, Italy

Antarctic sponges may host large populations of planktonic and benthic diatoms. After settling on the sponge, these diatoms enter its body through pinacocytes (1) and form, there, large mono- or pauci-specific assemblages. Yet the total amount of carbohydrates in the invaded sponge tissue is inversely correlated with that of chlorophyll-a. We suggest, therefore, that endobiont diatoms utilize the products of the metabolism of their host as an energy source. This is the first evidence indicating that an endobiotic autotrophic organism may parasitize its animal host. Moreover, this unusual symbiotic behavior could be a successful strategy that allows the diatom to survive in darkness.

Heterotrophic bacteria, autotrophic cyanobacteria, zoochlorellae, and zooxanthellae are common symbionts in Porifera, where they may actually constitute most of the sponge tissue. Sponges can also harbor fungal populations, the significance of which has been poorly investigated (2). The association of sponges with autotrophic symbionts occurs mainly in tropical (3) and temperate waters (4); it is a fruitful strategy, allowing the sponges to utilize the symbionts as a food source complementary to filter feeding (5, 6).

Diatoms in sponge tissues have rarely been recorded (7), but such associations are widespread in Antarctic waters (1). In fact, most sponges from Terra Nova Bay (Ross Sea, Antarctica) host large populations of both planktonic and benthic diatoms, and these microorganisms are actively taken up by the pinacoderm and thus enter the sponge choanosome (1). This could be an adaptive supplementation of the food supply in the Antarctic ecosystem, because diatoms produce extracellular polysaccharides (8, 9) that sponges might use as energy substrates. On the other hand, the integrity of the diatom shells, the presence of cytoplasm inside the shells, and the formation of large mono- or pauci-specific assemblages inside a single sponge suggest that diatoms are able to complete their entire biological cycle inside the sponge body.

The aim of this study was to investigate the energy source that symbiotic diatoms might provide to Antarctic sponges. To this end, we evaluated the variation in the total carbo-hydrate content of sponges as a function of the biomass of endobiotic diatoms; biomass was estimated from chloro-phyll-*a* concentration.

We examined 39 specimens representing 17 species of sponges (Table 1). The material was collected at Terra Nova Bay (Ross Sea), between 100 and 120 m depth, during the XIII Italian Antarctic Expedition (Austral Summer 1997–1998). The sponges were prepared for ultrastructural study as follows. Immediately after collection, small pieces from each specimen were fixed for 2 h in 2.5% glutaraldehyde in artificial seawater (ASW). The fixed tissues were rinsed, stored in ASW, and then dehydrated in a graded series of ethanol concentrations. Critical point drying was achieved with a CO₂ Pabish CPD 750 apparatus. Samples were examined with a Philips EM 515 scanning electromicroscope, and several species of diatoms were observed (Fig.

Received 29 June 1999; accepted 5 November 1999.

^{*} To whom correspondence should be addressed. E-mail: bavestrello@ popcsi.unian.it

Table I

Chlorophyll-a concentration and carbohydrate content for the sponge specimens analyzed

Specimens	Chlorophyll- <i>a</i> (µg/g dwaf)	Total sugars (mg/g dwaf)
Avociella nidificata	27.29 ± 2.5	8.12 ± 0.2
Dendrilla membranosa	29.13 ± 3.6	5.43 ± 0.4
	30.47 ± 2.1	11.40 ± 2.1
	31.65 ± 1.8	7.01 ± 0.6
	21.34 ± 1.9	16.29 ± 2.8
	7.31 ± 1	22.56 ± 1.8
	9.96 ± 0.9	27.06 ± 1.4
	14.43 ± 2.9	9.59 ± 1.1
Ectyodorix ramilobosa	8.47 ± 1.2	19.90 ± 3.1
	20.70 ± 3.1	6.91 ± 1.8
Gellius sp.	17.05 ± 2.4	7.25 ± 2
Gellius sp. 1	37.10 ± 6.9	6.91 ± 1.5
Haliclona dancoi	16.67 ± 2.4	9.92 ± 2.4
	15.05 ± 2.6	21.28 ± 3.2
	15.14 ± 0.9	8.10 ± 1.4
	21.85 ± 1.9	15.92 ± 1.9
Homaxinella balfourensis	14.37 ± 3.1	12.73 ± 2.1
Inflatella belli	5.75 ± 1.6	24.30 ± 3.1
	2.21 ± 0.8	24.41 ± 3.6
Isodictya conulosa	16.48 ± 1.6	12.29 ± 2.2
	13.59 ± 2	9.96 ± 1
Isodictva erinacea	11.86 ± 1.8	15.25 ± 2.8
Microxina benedeni	11.75 ± 3.2	11.21 ± 1.2
Phorbas glaberrima	18.22 ± 2.9	13.13 ± 2.8
	19.04 ± 4	12.73 ± 4.1
Pseudosuberites antarcticus	44.56 ± 3.5	3.32 ± 0.9
Pseudosuberites nudus	27.79 ± 5.6	10.23 ± 1.4
	45.71 ± 5.9	6.23 ± 0.8
	9.13 ± 1.7	17.71 ± 2.5
	11.88 ± 2.1	12.13 ± 1.3
	24.53 ± 4.8	9.23 ± 1.9
	25.65 ± 4.9	5.93 ± 1.4
	19.74 ± 3.6	7.41 ± 2.8
	11.54 ± 2.2	10.48 ± 2.3
Suberites caminatus	32.94 ± 3.8	8.91 ± 1.7
	6.35 ± 2.1	21.08 ± 3.4
	17.25 ± 5.7	7.11 ± 1.5
Tedania charcoti	16.67 ± 3.6	7.11 ± 0.9

Values are the mean of three replicates \pm SD; dwaf, dry weight ash free.

1). Among pennate diatoms, *Fragilariopsis curta* (Fig. 1a), *Fragilariopsis* sp. (Fig. 1a, b), *Achnanthes* sp. (Fig. 1c), and *Pseudogomphonema* sp. were the most recurrent species; centric diatoms belonging to the genera *Porosira* (Fig. 1c), *Coscinodiscus*, and *Rhizosolenia* were rarer. For transmission electron microscopy, fragments of selected fixed material were postfixed in 1.0% OsO₄ in ASW at 4°C. After repeated rinsing, these fragments were dehydrated in ethanol and embedded in araldite. Ultrathin sections were cut with a Reichert ultramicrotome equipped with a diamond

blade; these sections were gathered on copper grids, stained with uranyl acetate and lead citrate for contrast, and examined with a Philips 300 transmission electron microscope. Micrographs (TEM) of *Fragilariopsis curta* provide evidence that the cytoplasm and thylacoids are intact (Fig. 1d).

The concentrations of chl-*a* in the various sponge specimens were measured and are listed in Table 1. These values differed widely, ranging from 2.2 μ g/g in *Inflatella belli*, to 45.7 μ g/g in *Pseudosuberites nudus*. High values were found also in *Pseudosuberites antarcticus*, *Gellius* sp. 1., *Suberites caminatus*, and *Dendrilla membranosa*. Strong differences emerged when different specimens of the same species were compared (*e.g.*, from 7.3 to 31.6 μ g/g in *D. membranosa*; and from 9.1 to 45.7 μ g/g in *P. nudus*). This variation suggests that diatom uptake is fortuitous, depending primarily upon exposure of the sponge surface to diatoms settling upon it, rather than to some species-specific entrapping mechanism.

Carbohydrate in the sponge tissues (Table 1) ranged between 3.3 mg/g in *Pseudosuberites antarcticus* and 27.1 mg/g in *Dendrilla membranosa*. As with the chl-*a* concentration, carbohydrate content varied markedly when different specimens of the same species were compared.

For each specimen, we plotted carbohydrate content as a function of chl-*a* concentration and obtained the inverse correlation that is evident in Figure 2a. When we considered only those specimens belonging to a single species, this same trend was confirmed. In fact, in *Dendrilla membranosa* (Fig. 2b) and *Pseudosuberites nudus* (Fig. 2c), the total sugar content of the sponge tissue decreased as the concentration of chl-*a* increased.

The original aim of this work was to demonstrate that diatoms living inside the sponge tissue might represent a source of food for Antarctic sponges. This hypothesis seemed reasonable because diatoms produce large amounts of polysaccharides (8, 9). But our data show a clear inverse relationship between the amount of chl-*a* and of carbohydrate in sponge tissue. This relationship suggests that, as the population of diatoms within the sponge increases, it consumes more of the carbohydrate or related compounds produced by the sponge. This report, therefore, provides the first indication that diatoms in the field can utilize animal metabolic products as an energy source.

Like those in Antarctic sponges, most heterotrophic diatoms are pennate and occur in benthic habitats, where light is absent or limited and organic substrates are in great supply; such habitats strongly favor heterotrophic growth (10). Furthermore, several diatom species have developed active transport systems that bind to a limited number of organic substrates (glucose, lactate, and glutamate) and that respond to conditions of low light or darkness (11, 12); these diatoms have evolved highly sophisticated and ecologically significant mechanisms for



Figure 1. Diatoms living inside Antarctic sponges. (a) Several specimens of *Fragilariopsis curta* (arrows) and a *Fragilariopsis* sp. disposed under the exopinacoderm of *Tedania charcoti* (Scale bar, 50 μ m. (b) *Fragilariopsis* sp. (arrows) embedded inside the same sponge (scale bar, 50 μ m). (c) Large centric *Porosira* sp. (arrows) and numerous pennate *Achnanthes* sp. (arrow heads) in *Pseudosuberites antarcticus* (scale bar, 100 μ m). (d) Transmission electron micrograph of *Fragilariopsis curta* inside the sponge body showing intact cytoplasm (c) and thylocoid (t) (scale bar, 10 μ m). s, sponge spicules.

facultative heterotrophy. Mixotrophy, a combination of autotrophy and heterotrophy, is widespread among certain protists and, in particular, among flagellates such as haptophytes (13), dinoflagellates (14, 15, 16, 17), cryptophytes (18), and diatoms (19, 20, 21). Previous documentation of mixotrophy in diatoms was obtained by observations *in vitro*.

The relationship between myxotrophic diatoms and Antarctic sponges can be considered as a sort of symbiosis shifting to parasitism; *i.e.*, the algae can get nourishment, not only by photosynthesis, but also by utilizing organic carbon derived from their host. Although some Antarctic diatoms are considered among the most shade-adapted microalgae (23), the mixotrophy assumed in this report explains the surprising presence of dense diatom populations inside sponges living at 100-120 m depth, where the level of irradiance is very low (24).

Finally, the mechanism that causes a sponge to actively incorporate diatoms is unclear. However, some studies (22) have revealed a tendency of demosponges to incorporate



Figure 2. Concentration of chl-*a* as a function of the total amount of carbohydrate. Data from all the sponge specimens tested (a), from different specimens of *Dendrilla membranosa* (b), and from *Pseudosuberites nudus* (c). Each point represents a sponge specimen. The analyses were carried out on three samples of sponge tissue from each specimen. Samples were cylindrical cores of choanosomal tissue, 1 cm in diameter, 2 cm in height. Cores were longitudinally cut, and the two half-portions were tested to evaluate the chl-*a* concentration (as an estimate of diatom population biomass), as well as the total amount of carbohydrate. One subsample was extracted in 90% acetone. The chl-*a* was then assayed by spectrofluorimetry with excitation at 430 nm and emission measured at 668 nm. On the other subsample, the total carbohydrates were assayed: the method was based on the colorimetric reaction of sugars with phenol and sulphuric acid, and D(+) glucose was used as the standard (26).

siliceous particles, and the diatoms, with their siliceous shells, may thus "deceive" the sponge.

In conclusion, our data widen the ecological role of diatoms which, by exploiting the niche provided by sponges—one of the most important Antarctic benthic organisms—illustrate a strong functional link between plankton and benthos (25).

Literature Cited

- Gaino, E., G. Bavestrello, R. Cattaneo-Vietti, and M. Sarà. 1994. Scanning electron microscope evidence for diatom uptake by two Antarctic sponges. *Polar Biol.* 14: 1–4.
- Sarà, M., G. Bavestrello, R. Cattaneo-Vietti, and C. Cerrano. 1998. Endosymbiosis in sponges: relevance for epigenesis and evolution. *Symbiosis* 25: 57–70.
- Wilkinson, C. R. 1980. Cyanobacteria symbiotic in marine sponges. Pp. 553–563 in *Endocytobiology, Endosymbiosis and Cell Biology*. W. Schwemmler and H. E. A. Schenk, eds. Walter De Gruyter, New York.
- Sarà, M. 1971. Ultrastructural aspects of the symbiosis between two species of the genus *Aphanocapsa* (Cyanophyceae) and *Ircinia variabilis. Mar. Biol.* 11: 214–221.
- Wilkinson, C. R. 1983. Net primary productivity in coral reef sponges. Science 219: 410–412.
- Arillo, A., G. Bavestrello, B. Burlando, and M. Sarà. 1983. Metabolic integration between symbiotic cyanobacteria and sponges: a possible mechanism. *Mar. Biol.* 117: 159–162.
- Cox, G., and A. W. D. Larkum. 1983. A diatom apparently living in symbiosis with a sponge. Bull. Mar. Sci. 33: 943–945.
- Myklestad, S., O. Holm-Hansen, K. M. Värum, and B. Volcani. 1989. Rate of release of extracellular aminoacids and carbohydrates from the marine diatom *Chaetoceros affinis*. J. Plankton Res. 11: 763–773.
- Myklestad, S. 1995. Release of extracellular products by phytoplankton with special emphasis on polysaccharides. *Sci. Total Environ.* 165: 155–164.
- Hellebust, J. A., and J. Lewin. 1997. Heterotrophic nutrition. Pp. 169–197 in *The Biology of Diatoms*. D. Werner, ed. University of California Press, Berkeley.
- White, A. W. 1974. Growth of 2 facultatively heterotrophic marine centric diatoms. J. Phycol. 10: 292–300.
- Lewin, J. C., and J. A. Hellebust. 1976. Heterotrophic nutrition of marine pennate diatom *Nitzschia angularis* var. *affinis. Mar. Biol.* 36: 313–320.
- Tillmann, U. 1998. Phagotrophy by a plastidic haptophyte, Prymnesium patelliferum. Aquat. Microb. Ecol. 14: 155–160.
- Graneli, E., D. M. Anderson, P. Carlsson, and S. Y. Maestrini. 1997. Light and carbon uptake by *Dinophysis* species in comparison to other photosynthetic and heterotrophic dinoflagellates. *Aquat. Microb. Ecol.* 13: 177–186.
- Legrand, C., E. Graneli, and P. Carlsson. 1998. Induced phagotrophy in the photosynthetic dinoflagellate *Heterocapsa triquetra*. *Aquat. Microb. Ecol.* 15: 65–75.
- Skovgaard, A. 1996. Mixotrophy in *Fragilidium subglobosum* (Dinophyceae): growth and grazing responses as functions of light intensity. *Mar. Ecol. Prog. Ser.* 143: 247–253.
- Tiselius, P., and M. Kuylenstierna. 1996. Growth and decline of a diatom spring bloom: phytoplankton species composition, formation of marine snow and the role of heterotrophic dinoflagellates. J. Plankton Res. 18: 133–155.
- Gervais, F. 1997. Light dependent growth, dark survival and glucose uptake by cryptophytes isolated from a freshwater chemocline. *J. Phycol.* 33: 18–25.
- Azam, F., and B. E. Volcani. 1974. Role of silicon in diatom metabolism VI. Active transport of germanic acid in the heterotrophic diatom *Nitzschia alba. Arch. Microbiol.* 101: 1–8.
- Darley, W. M., B. B. Wimpee, and C. T. Ohlman. 1981. Heterotrophic and photoheterotrophic utilization of lactate by the diatom *Cylindrotheca fusiformis. Br. Phycol. J.* 16: 423–428.

- Tan, C. K., and M. R. Johns. 1996. Screening of diatoms for heterotrophic eicosapentaenoic acid production. J. Appl. Phycol. 8: 59-64.
- Bavestrello, G., A. Arillo, U. Benatti, C. Cerrano, R. Cattaneo-Vietti, L. Cortesogno, L. Gaggero, M. Giovine, M. Tonetti, and M. Sarà. 1995. Quartz dissolution by the sponge *Chondrosia reniformis* (Porifera, Demospongiae). *Nature* 378: 374–378.
- Palmisano, A. C., J. B. SooHoo, D. C. White, G. A. Smith, G. R. Stanton, and R. H. Burckle. 1985. Shade-adapted benthic diatoms beneath Antarctic sea ice. *J. Phycol.* 21: 664–667.
- Vanucci, S., and M. Innamorati. 1997. Available radiation profiles recorded in the Ross Sea. Pp. 281–290 in *Rossmize 1993–95 Data Report I.* F. M. Faranda, L. Guglielmo and P. Povero, eds. Lang Arti Grafiche, Genoa, Italy.
- Marcus, N. H., and F. Boero. 1998. The importance of benthicpelagic coupling and the forgotten role of the life cycles in coastal aquatic systems. *Limnol. Oceanogr.* 43: 763–768.
- Dubois, M., K. A. Gilles, J. K. Hamilton, P. A. Rebers, and F. Smith. 1956. Colorimetric method of determination of sugars and related substances. *Anal. Chem.* 28: 350–356.



Bavestrello, Giorgio et al. 2000. "Parasitic diatoms inside antarctic sponges." *The Biological bulletin* 198, 29–33. <u>https://doi.org/10.2307/1542801</u>.

View This Item Online: https://doi.org/10.2307/1542801 Permalink: https://www.biodiversitylibrary.org/partpdf/2440

Holding Institution MBLWHOI Library

Sponsored by MBLWHOI Library

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: University of Chicago License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.