

BRITISH FUNGI.

By M. C. COOKE.

(Continued from p. 88.)

Cortinarius (Phlegmacium) triumphans. *Fr.* "Magnificent Cortinarius."

Pileus fleshy, convex, then plane, viscid, even, at first spotted with evanescent darker adpressed scales; stem solid, clavate, concentrically squamose beneath the ring; gills emarginate, quite entire, crowded, pallid, then clay-coloured.—*Fr. Epicr. p. 256. B. & Br. Ann. N. H., No. 1263. Cortinarius sublanatus. Hussey Myc. Illus. ii. t. 22.*

In woods.

A large species. Pileus yellowish.

Cortinarius (Myxadium) pluvius. *Fr.* "Pluvial Cortinarius."

Pileus rather fleshy, globose, then convex, smooth, viscid, hygrophanous; stem stuffed, then hollow, slender, nearly equal, soft, white, becoming pallid; gills decurrent, then seceding, white, then ochraceous.—*Fr. Sys. Myc. i., 36. Fr. Epicr. p. 277. B. & Br. Ann. N. H., No. 1267. Secr. No. 265.*

In woods. Lea. Sept.

Pileus $\frac{1}{2}$ -1 in., stem sometimes short, sometimes 3 in. long, 2-3 lines thick. Gills at length free, pallid.

Cortinarius (Dermocybe) anthracinus. *Fr.* "Fiery-gilled Cortinarius."

Pileus innato-fibrillose, brownish cinnamon; stem fibrilloso-striate, pallid; gills very much crowded, narrow, fiery-yellow, blood-red when rubbed.—*Fr. Epicr. p. 288. B. & Br. Ann. N. H., No. 1269.*

In a wood. Coed Coch.

Certainly different from *C. sanguineus*. The Welsh plant exactly accords with a drawing from Fries. *B. & Br.*

Stem 2 in. long, 1-2 lines thick.

Cortinarius (Dermocybe) orellanus. *Fr.* "Red fleshed Cortinarius."

Pileus fleshy, obtusely umbonate, villosa-squamulose or fibrillose, tawny orange, flesh reddish; stem solid, firm, nearly equal, striato-fibrillose, tawny; gills adfixed, broad, rather distant, at length opaque.—*Fr. Epicr. p. 288. B. & Br. Ann. N. H. No. 1270.*

On the ground in a wood. Coed Coch. Oct.

With *C. cinnamomeus*, to which it is nearly related, but very distinct.

Cortinarius (Telamonia) bivelus. *Fr.* "Twin-veiled Cortinarius."

Pileus fleshy, convex, then plane, obtuse, smooth, or silky at the margin, brick red, becoming tawny; stem firm, rather bulbous, dirty whitish; spurious ring fugacious; gills adnexed, scarcely crowded, bright tawny cinnamon.—*Fr. Epicr. p. 292. B. & Br. Ann. N. H. No. 1271. Fr. Obs. ii. 58. Fr. Sys. Myc. i. 215.*

In woods. Coed Coch. Oct.

Pileus soft, bibulous, moist, but not truly hygrophanous.



Cooke, M. C. 1873. "British fungi [cont.]." *Grevillea* 1(7), 110–111.

View This Item Online: <https://www.biodiversitylibrary.org/item/30137>

Permalink: <https://www.biodiversitylibrary.org/partpdf/246390>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.