

wet season had served to stimulate the conglomerated gonidial granules into renewed activity, and produced (along with the consequent maceration) the modified thallus in question.

Had time and space permitted, I should have liked to have enlarged somewhat more on this subject; meanwhile, I shall content myself with describing a genuine parasite on *S. bispore*, viz:—

Lecidea epiphorbia, *Stn.*—Apothecia resemble, externally and internally, those of *L. Parmeliarum*, except that the paraphyses are neither thickened nor darker coloured at their apices, the spores are colourless, or present in a few instances, a faint tinge of yellow, and the reaction on the gelatina hymenia by means of iodine, shows a deep vinous red without any preceding cærulescent tint, instead of being negative, as in *L. Parmeliarum*. This lichen bears the same relationship to *L. Parmeliarum* that *L. solorinaria* does to *L. oxyspora*.

BRITISH FUNGI.

By the EDITOR.

(Continued from Page 176.)

The number of species of Fungi found in this country since the conclusion of our series of descriptions in the last volume of this Journal has rendered it incumbent upon us, in fulfilment of promise, to continue the descriptions of these additions as a supplement to the "Handbook of British Fungi."

Agaricus (Lepiota) granulatus. *Batsch.* var. **Carcharias.** *Fr.*

Pileus umbonate, clad with darker evanescent granules; stem sub-bulbous, and as well as the patent ring squamulose with granules; gills adnexed, white.—*Fries, Ep.*, p. 18. *Pers. Ic. Pict.*, t. 5, f. 1. *Smith, Journ. Bot.*, 1873, p. 335 = var. *incarnato-albida.* *Fr. Mon. Hym. Suec.*, p. 29.

On grass under old Scotch firs.

Agaricus (Armillaria) aurantius. *Schæff.*

Pileus fleshy, convex then plane, obtuse, innato-squamulose, viscid; stem solid, equal, concolorous, guttate above, clad with orange scales, sub-annulate; gills adnexed with a tooth, crowded, white.—*Fries Epicr.*, p. 21. *Sys. Myc. i.*, 39. *Schæffer, Ic.* 27. *B. & Br. Ann. N.H.*, 1335.

In pine woods. *Forres, N. B.*

Varying a good deal in the nature and frequency of the scales.

Agaricus (Clitocybe) tuba. *Fr.*

White, pileus fleshy, thin, convex, then plane, umbilicate, moist, shining with a whitish silky lustre, margin even; stem equal, soon hollow and compressed, naked above, gills attenuated and decurrent,



Cooke, M. C. 1874. "British fungi [cont.]." *Grevillea* 2(19), 108–109.

View This Item Online: <https://www.biodiversitylibrary.org/item/30137>

Permalink: <https://www.biodiversitylibrary.org/partpdf/246401>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.