

## Symbolae ad Mycologiam Fennicam.

### Pars XXV.

Auctore

P. A. Karsten.

~~~~~  
(Societati exhibitum die 3 Martii 1888).

**Lactarius lateritioroseus** Karst. Symb. ad Myc. Fenn. XXIV, p. 14, emend.

Pileus demum repandus, saepe inaequalis, in squamulas in disco minores, granuliformes, marginem versus majores secedens, usque ad 11 cm. latus, carne grumosa, usque ad 1 cm. crassa, ad marginem admodum tenuis. Stipes fartus, nonnumquam demum basi cavus, inaequalis, deorsum solito incrassatus, curvatus vel flexuosus, rarius strictus, subtiliter subflocculosus, tandem pileo concolor, at pallidior, usque ad 7 cm. longus et 2 cm. crassus. Lamellae demum flavescentes. — Pileus subinde obsolete zonatus, fragilis. Lamellae subdistantes, usque ad 5 mm. latae. Odor fere nullus.

**Bjerkandera chionea** (Fr.) Karst.

In Lapponia rossica prope vicum Knjäscha-guba, die 9 Aug. 1861 legimus.

**Bjerkandera simulans** Karst. in Rev. myc. 1888, Avril.

Valde imbricata, conrescens. Pileus carnosus, lentus, dimidiatus, triqveter, irregularis, laevis, glaber, azonus, pallescens vel albus, siccitate subferruginascens, intus albus 1—3 cm. latus. Pori minuti, rotundi, inaeqvales, laceri, tenues, e

pallido subferruginascentes, nudi. Sporae cylindraceae, curvulae, hyalinae, 4—5 mmm. longae, 1—1,5 mmm. crassae.

In cortice *Piceae excelsae* prope Vasa, m. Majo 1864.

Quoad formam cum *Inonoto radiato* (Sow.) omnino convenit.

***Poria separabilis* Karst. n. sp.**

Effusa, subtus glabra, immarginata, siccitate rimosa, ambitu concolore. Pori membrana tenuissima, molli, pallida vel albida, sicca secedente impositi, perexigui, rotundi, aequales, pertenuēs, circiter 0,4 mm. alti, a membrana facile separabiles, subfusci.

Ad corticem laevem, semivivum *Betulae* in paroecia Tammela semel obvia.

***Cyphella terrigena* Karst. n. sp.**

Receptaculum membranaceum, campanulatum, subinde in stipitem brevissimum porrectum, laeve, fere album, albo-puberulum, 0,2—0,3 mm. altum. Hymenium laeve, lutescente album. Basidia clavulato-cylindracea, longit. 20—25 mmm., crassit. 4—5 mmm. Sporae ellipsoideo-sphaeroideae, longit. 6—7 mmm., crassit. 5—6 mmm. Cystidia nulla.

Supra terram nudam in horto Mustialensi, m. Julio 1869.

***Corticium calotrichum* Karst. in Rev. myc. 1888, Avril.**

Oblongatum vel effusum, immarginatum, adnatum, subtus adpresse fibrillosum ambituque byssinum, demum membranaceum, subsecedens et glabrescens. Mycelium e hyphis ramosis, articulatis, apice obtuso-rotundatis, albis, 6—12 mmm. crassis contextum. Hymenium tenue, ceraceum, laeve, contiguum, rarius siccitate rimosum, albido-alutaceum vel sordide flavidum. Cystidia nulla. Sporae sphaeroideae, molles, flavescēte hyalinae, parcae, 6—10 mmm. diam.

Supra corticem vetustum *Alni incanae* in regione Mustialensi pluries m. Aug.—Nov. lectum.

***Corticium confluens* Fr.**

Var. **triviale** Karst. in Rev. myc. 1888, Avril.

Rotundatum vel effusum, adglutinatum, immarginatum, ambitu primitus subfimbriatum, album, dein glabrescens.

Hymenium ceraceum, rigescens, laeve, contiguum, rarissime (siccum) rimosum, livido-caesium, siccum albido-alutaceum seu flavescens, nudum. Cystidia nulla. Sporae sphaeroideae vel subsphaeroideae, dilutissime flavescens vel subhyalinae 6—9 mmm. diam. vel 7—9 longi, 6—8 mmm. crassi.

Supra corticem, rarius lignum *Betulae*, *Alni*, *Sambuci racemosae*, *Aceris* et *Populi* frequens usque ad Mare-glaciale.

Num hoc varietas tantum sit, ut omnis analogia sva-det, in medio relinqvimus.

Var. **subcalceum** Karst. in Rev. myc. 1888, Avril.

Rotundatum vel effusum, confluens, adglutinatum, immarginatum, ambitu simile vel primitus tenuissime subbyssinum, concolor. Hymenium ceraceum, rigescens, laeve, contiguum vel saepius rimoso-partitum, nudum, albidum. Cystidia nulla. Sporae sphaeroideae, ditutissime flavescens vel subhyalinae, 6—9 mmm. diam., parcae.

Supra corticem et lignum *Betulae* circa Mustiala, fere per annum.

A var. *padino* Karst., cui proxime accedit, hymenio vulgo rimoso, albidiores ambituque similari vel mox glaucescente concolori discrepat.

**Tromera microtheca** Karst n. sp.

Apothecia superficialia, sessilia, gregaria, solitaria, rarissime in caespitulos sociata (2—3 insimul), nuda, nigra, opaca, subrugulosa, aperta, plana, orbicularia vel ovalia, tenuiter marginata, vix 0,3 mm. lata. Asci clavati, obtusissimi, jodo dilute caerulescentes, longit. 36—40 mmm., crassit. 10—12 mmm., myriospori. Sporae oblongatae vel elongatae, rectae, hyalinae, longit. 2—3,5 mmm., crassit. 1—1,5 mmm. Paraphyses haud bene discretas. Hymenium jodo fulvescens.

In ligno vetusto abietino ad Mustiala, m. Aug. 1870.

A sequente, cui proxime accedit, praecipue ascis minoribus, sporis longioribus apotheciisque fere minoribus tenuiusque marginatis recedit.

**Tromera ligniaria** Karst. n. sp.

Apothecia superficialia, solitaria, raro in caespitulos sociata (plerumque 3 insimul), sessilia, rarius basi constricta, aperta, concaviuscula, demum plana vel convexiuscula, margine obtuso, demum subimmarginata, nuda, atra, rugulosa, latit. 0,3 mm. Asci clavati, obtusi, jodo dilute caerulescentes, longit. 80—100 mmm., crassit. 12—14 mmm., myriospori. Sporae ellipsoideo-sphaeroideae vel sphaeroideae, hyalinae, longit. 1,5—2 mmm., crassit. 1,5 mmm. vel diam. 1,5 mmm. Paraphyses filiformes, 1 mmm. crassae.

Ad lignum abietinum vetustum prope Mustiala, m. Martio 1872. — An cum priore *Tympanidi* adscribenda?

**Mollisia Artemisiae** (Lasch) Karst. *Peziza Artemisiae* Lasch. in Rab. Herb. Myc. N:o 335; Handb. I, p. 344.

Perithecia erumpentia, e sphaeroideo — hemisphaerico-applanata, scabrosella, subolivaceo-cinerea vel cinerea, disco pallidiore, margine primitus albicante, siccitate nigrescentia.

Ad caules aridos *Artemisiae vulgaris* in regione Mustialensi, m. Majo et Junio. Sterilis tantum lecta.

**Amerosporium Sedi** Karst. in Rev. myc. 1888, Avril.

Cupulae superficiales, gregariae, atrae, setis simplicibus, rigidis, erectis, continuis, atris, circiter 220 mmm. longis obsessae, circa 0,5 mm. latae. Conidia fusoides-elongata, continua, curvula, eguttulata, hyalina, longit. 24—26 mmm., crassit. 2,5—3 mmm.

In foliis langvescentibus *Sedi Telephii* prope Aboam, m. Sept. 1887, leg. Onni Karsten.

**Cytosporina stellulata** Sacc. Syll. III, p. 602 in ligno vetusto *Ulm*i ad Aboam m. Aprili 1861 semel a nobis lecta.

Sporulae filiformes, arcuatae, circiter 30 mmm. longae et 1 mmm. crassae.

**Rhabdospora pleosporoides** Sacc. \***Rh. longior** Karst. n. subsp.

Maculae nullae. Pyrenia sparsa, cladogena, mox nuda,

rotundata vel oblongata, applanata vel saepius depressa, ostiolo papillato, circiter 0,4 mm. lata. Sporulae filiformes, rectae, continuae, multiguttulatae, hyalinae, 50—70 mmm. longi.

In caulibus aridis *Cerefolii sylvestris* prope Mustiala, m. Aprili 1866.

**Leptosporium mycophilum** Karst. n. sp.

Sporodochia effusa, mollia, tenuissima, aurantiaca, ambitu subbyssino, albo. Conidia oblongata, continua, recta, hyalina, longit. 3—4 mm., crassit. 1—1,5 mmm.

In specie indescripta Myxomycetum in regione Aboënsi, Merimasku, m. Aug. 1860.

**Helminthosporium arbusculoides** Peck.

Caespites effusi, atri, velutini. Hyphae fasciculatae, rigidae, simplices, subflexuosae, saepe basi decumbentes, nigrae, articulatae, usque ad 250 mmm. longae, 7—8 mmm. crassae. Conidia ellipsoidea vel oblongata, 3-septata, fuliginea, semipellucida, unicoloria, longit. 20—25 mmm., crassit. 8—10 mmm.

In cortice et ligno vetusto *Betulae* prope Mustiala et Vasa, m. Aprili—Julio.

**Botrytis campotricha** Sacc. Syll. IV, p. 117, var. **fennica** Karst.

Caespituli pulvinati, minuti, confluentes, albi. Hyphae parce septatae, ramosae, 4—6 mmm. crassae, ramis vulgo rectis. Conidia ellipsoidea vel ovalia vel suboblonga, eguttulata, hyalina, longit. 5—8 mm., crassit. 2—3,5 mmm.

Ad lignum *Pini sylvestris* alicubi in Fennia.

Nudo oculo lignum videtur farina adpersum.

**Monilia arctica** Karst. n. sp.

Caespituli effusi, velutini, e sordido helvoli. Hyphae septatae, longiusculae, 8—10 mmm. crassae. Conidia breve catenulata, ovalia, saepissime distincte apiculata indeqve isthmico-conjuncta, hyalina vel demum, dilutissime flavida, endochromate granuloso, longit. 20—30 mmm., crassit. 10—13 mmm.

Supra lignum *Betulae* vetustum juxta sinum Kolaënsem Lapponiae rossicae, m. Julio 1861 legimus.

**Tolypomyria fungicola** Karst. n. sp.

Caespituli effusi vel subcirculares, tomentosuli, albi. Hyphae vage ramosae, flexuosae, articulatae, intricatae, 8—10 mmm. crassae. Glomeruli rotundati, mucosi. Conidia ellipsoidea vel ovalia, subinde sphaeroideo-ellipsoidea, hyalina, 2—4 mmm. longa, 1—2 mmm. crassa.

In *Hansenia velutina* putrescente in paroecia Tyrvis, m. Sept. 1859.

**Oospora Clavariarum** Karst. n. sp.

Caespituli effusi, ambientes, mucedinei, albidi. Catenulae breves, simplices, ex hyphis brevissimis (40—50 mmm. longis) oriundae. Conidia (saltem ultima) sphaeroidea, episporio crasso, verruculoso, hyalina, diam. 18—36 mmm.

Supra Clavariam putrescentem prope Mustiala, m. Aug. 1866.

**Trichosporium fuscum** (Link.) Sacc. Syll. IV, p. 289 in ligno vetusto prope vicum Lapponiae rossicae, Knjäscha-guba, m. Aug. 1861 legimus.

Fuscum. Hyphae 4—6 mmm. crassae, dilute fuligineae et pellucidae (sub.micr.). Conidia ellipsoidea vel ovalia, dilute fuliginea pellucidaque (sub.micr.), longit. 8—14 mmm., crassit. 6—7 mmm. — *Trichosporium fuscescens* Cook. et Harkn. verisimiliter huc pertinet.

**Hormiscium septonema** (Preuss.) Karst. var. **betulinum** Karst.

Syn. *Hormiscium curvatum* (Peck.) Sacc. var. *betulinum* Karst. Symb. ad Myc. Fenn. XX, p. 97.


Caespites late effusi, atri. Catenae conidiorum rarissime ramosae, 60—90 mmm. longae. Conidia subcuboidea, 4 mmm. longae, 6 mmm. crassae.

In cortice *Betulae* prope Vasa, m. Aug. 1867.

**Torula obducens** Karst. n. sp.

Ramos late ambiens et crustam tenuem, atram, effusam formans. Hyphae steriles decumbentes, articulatae, hyalinae. Catenulae ut plurimum decemsporae, simplices vel ramosae, curvatae vel flexuosae, saepe connatae. Conidia primitus hyalina, ellipsoidea vel sphaeroidea, dein sphaeroidea, fuliginea, pellucida (sub. micr.), diam. 4—8 mmm., sat facile secedentes.

Supra corticem semivivum *Populi tremulae* in regione Mustialensi, m. Januario 1882.





Karsten, P. A. 1888. "Symbolae ad mycologiam Fennicam. XXV." *Meddelanden af Societas pro Fauna et Flora Fennica* 16, 20–26.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/38482>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/246688>

**Holding Institution**

MBLWHOI Library

**Sponsored by**

MBLWHOI Library

**Copyright & Reuse**

Copyright Status: NOT\_IN\_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.