
Saurauia homotricha (Actinidiaceae), a New Species from Honduras and Nicaragua

Amy Pool

Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri 63166-0299, U.S.A.

ABSTRACT. *Saurauia homotricha*, a new species from montane forests of western Honduras and Nicaragua, is described and illustrated. It is placed in the Central American series *Gymnogynae* Buscalioni and is most similar to *Saurauia rubiformis* Vatke, from which it differs most markedly in the type, length, and distribution of trichomes on the leaves and the abaxial surface of the sepals.

The genus *Saurauia* consists of approximately 300 species from the Neotropics (excluding the Antilles) and tropical Asia. Twenty-five species have been recognized for Mexico and Central America; 22 are treated in Hunter's (1966) revision of the Mexican and Central American species, with 2 more Mexican species described by Keller and Breedlove (1981) and 1 more from Honduras described by Soejarto (1985). An additional species of *Saurauia* from Honduras and Nicaragua was recognized during preparation of a treatment of the Actinidiaceae for the *Flora de Nicaragua* and is here described. The pubescence terminology employed is that described and illustrated in Hunter's revision.

***Saurauia homotricha* A. Pool, sp. nov. TYPE:** Nicaragua. Nueva Segovia: Río Achuapa, al sur del Cerro Mogotón, 1500 m, 12 June 1975 (buds and flowers), *Atwood & Neill 16* (holotype, MO; isotype, HNMN?). Figure 1.

Species nova *Saurauiae rubiformis* Vatke similis, sed ab ea foliis supra inter venulas setulosis trichomatibus 1–1.5 mm longis infra inter venulas hirsutis atque stellato-pilosis, sepalorum partibus in alabastro expositis plumuloso-setulosis trichomatibus 1.75–2 mm longis differt.

Small tree, ca. 5 m tall, the young branches densely covered with shaggy-hirsute to shaggy-setose trichomes; trichomes to 2 mm long, shaggy mainly at base, golden to rusty-brown, mixed with minute, appressed-stellate trichomes. Leaves alternate, aggregated at tips of branches. Petioles 1.5–3.5 cm long, terete, indumentum as on young branches. Leaf blades oblanceolate, 19–28 cm long, 7–11 cm wide, short-acuminate at apex, cuneate to rounded at base, serrulate, with 16–29

pairs of secondary veins; tertiary veins more prominent on abaxial surface than higher order venation; adaxial surface abundantly to sparingly covered with setose trichomes 1–1.5 mm long; abaxial surface abundantly covered with a mixture of hirsute and stellate trichomes with long spreading arms, with primary and secondary veins densely covered by shaggy-setose trichomes. Inflorescence a narrow, axillary thyrse with 9–20 flowers, 3.5–14 cm long, peduncle 1.5–6 cm long, indumentum as on young branches; flowers actinomorphic, 14–18 mm diam.; sepal aestivation quincuncial, sepals elliptic to oblanceolate, 7–9 mm long, 4–4.5 mm wide, obtuse at apex, non-ciliolate, adaxial surfaces densely covered with minute, appressed-stellate trichomes, abaxial surface of outer two sepals densely covered with 1.75–2 mm long, shaggy-setose trichomes throughout, abaxial surface of imbricate sepal with exterior half (half exposed in bud) densely covered with 1.75–2 mm long, shaggy-setose trichomes and interior half densely covered with minute, appressed-stellate trichomes, abaxial surface of two inner sepals medially with dense 1.75–2 mm long, shaggy-setose trichomes and laterally and marginally densely covered with minute, appressed-stellate trichomes; petals 5, connate at base, oblong, 6–7 mm long, 3–5 mm wide, obtuse at apex, white; stamens 20–26, filaments adnate to base of corolla, 1.5–2 mm long, pubescent at base, trichomes filiform, anthers 2–2.5 mm long; ovary globose, glabrous, 5-locular, styles 5, 4–5 mm long. Fruit a berry, globose, 5-sulcate, ca. 5 mm diam., glabrous.

Soejarto (1969) suggested that many species of *Saurauia* have dimorphic flowers: one form functionally staminate with short styles and high pollen fertility and the other form functionally pistillate with long styles and low pollen fertility. A short-styled form of *Saurauia homotricha* is not currently known.

Hunter (1966) recognized four series in *Saurauia* for Mexico and Central America: *Gymnogynae* Buscalioni, *Gynotrichae* Buscalioni, *Oreophilae* Buscalioni, and *Laevigatae* Buscalioni. *Saurauia homotricha* belongs to the series *Gymnogynae*, which (as emended by Hunter) is characterized by leaves




Figure 1. *Saurauia homotricha* A. Pool. —a. Flowering branch. —b. Calyx, abaxial surface. Sepal position starting at 12 o'clock and moving clockwise: imbricate, inner, outer, inner, outer. —c. Leaf, abaxial surface. —d. Leaf, adaxial surface.

with tertiary veins prominent abaxially, sepals densely pubescent and non-ciliolate, and ovaries and fruits glabrous. Within this series, three of the species recognized by Hunter resemble *S. homotricha* in having the adaxial surface of all the sepals completely and densely covered with minute, appressed-stellate trichomes. Hunter's key to these species may be amended to include *S. homotricha* as follows:

- 1a. Adaxial surface of leaves abundantly sericeous; abaxial surface between veins with dendroid and long-armed stellate trichomes; longer trichomes of sepals shaggy-strigose . . . *S. villosa* DC. (Mexico)
- 1b. Adaxial surface of leaves abundantly to sparsely covered with setose, shaggy-setose or cluster-type trichomes; abaxial surface between veins without dendroid trichomes, the stellate trichomes with medium to very short arms or, if arms longer, then unbranched trichomes also present; longer trichomes of sepals shaggy-setose.
 - 2a. Adaxial surface of leaf abundantly to sparsely covered with setose trichomes 1–1.5 mm long, surface smooth to touch; abaxial surface between veins abundantly covered with mixture of hirsute and stellate trichomes; abaxial surface of outer sepals homotrichous, trichomes 1.75–2 mm long *S. homotricha* (Nicaragua and Honduras)
 - 2b. Adaxial surface of leaf with scattered, shaggy-setose trichomes less than 1 mm long (often only base of hair present), and/or cluster-type trichomes, surface scabrous; abaxial surface between veins with sparse to scattered, stellate to radiate trichomes; abaxial surface of outer sepals heterotrichous with longer trichomes to 1 mm long (often only base of trichomes present) mixed with dense, minute-appressed stellate trichomes.
 - 3a. Abaxial surface of leaf with sparse, medium-armed stellate trichomes (the longest arms ca. 0.2–0.3 mm long); flowers 9–13 mm diam.; inflorescence usually more than 70-flowered . . . *S. pittieri* J. Donnell Smith (Costa Rica and Panama)
 - 3b. Abaxial surface of leaf with scattered, radiate to short-armed stellate trichomes (the longest arms ca. 0.1 mm long); flowers 15–22 mm diam.; inflorescence usually less than 60-flowered
 - . . . *S. rubiformis* Vatke (Guatemala, Honduras, Costa Rica, and Panama)

The South American species of *Saurauia* most similar to *Saurauia homotricha* belong to the series *Macrophyllae* Buscalioni, as emended in Soejarto's revision of the South American species (1980). The two species most similar to *S. homotricha* are *Saurauia herthae* Sleumer and *S. prainiana* Buscalioni. They both differ from *S. homotricha* in having the larger trichomes of the sepals, peduncles, and petioles strigose. In addition, the abaxial leaf surface

of *S. herthae* has predominately stellate trichomes, with the unbranched trichomes restricted to the veins. In *S. prainiana*, the area between the veins has a combination of setose and radiate trichomes; in *S. prainiana* var. *prainiana* the trichomes are predominately radiate and in *S. prainiana* var. *pastasana* (Diels) Soejarto, they are predominately setose.

Specimens of *Saurauia homotricha* have been distributed under a variety of names: *Saurauia villosa*, *S. veraguasensis* Seemann, and *S. cf. scabrida* Hemsley. *Saurauia homotricha* is distinguished from *S. villosa* in the preceding key. *Saurauia veraguasensis* (= *Saurauia montana* Seemann) is distinguished from *S. homotricha* by its densely pubescent ovary and its sepals with adaxial surfaces submarginally glabrous, the abaxial surface of the two outer sepals heterotrichous with shaggy-strigose trichomes mixed with minute, appressed-stellate trichomes, and the two inner sepals ciliolate. *Saurauia scabrida* is distinguished from *S. homotricha* by its ciliolate sepals, with the adaxial surfaces submarginally glabrous (rarely pubescent for apical half to three-quarters of length) and the abaxial surface of the two outer sepals heterotrichous, with shaggy-strigose to shaggy-setose trichomes less than 1 mm long mixed with minute, appressed-stellate trichomes.

Paratypes. HONDURAS. **Intibucá:** Barranca Yamaranguila, cerca de Yashse, 1500 m, 12 abr. 1956 (buds), *Molina R. 6513* (US). **Comayagua:** in summit forest, cloud zone, range above El Achiote, above plains of Siguatepeque, 1850 m, 1 Aug. 1936 (buds), *Yuncker et al. 6261* (MO). **Morazán:** slopes of Cerro Uyuca (near El Zamorano), ca. 1800 m, 10 Oct. 1975 (buds, flowers, fruit), *Pilz & Pilz 1354-b* (MO).

Acknowledgments. I thank W. D. Stevens, Michael Grayum, and Djaja Soejarto for their advice and encouragement, Roy Gereau for assistance in preparing the Latin description, Barbara Alongi for providing the illustration, and the curators at US and F for specimen loans.

Literature Cited

- Hunter, G. E. 1966. Revision of Mexican and Central American *Saurauia* (Dilleniaceae). Ann. Missouri Bot. Gard. 53: 47–89.
- Keller, B. T. & D. E. Breedlove. 1981. Two new species of *Saurauia* (Actinidiaceae) from Mexico. Syst. Bot. 6: 65–73.
- Soejarto, D. D. 1969. Aspects of reproduction in *Saurauia*. J. Arnold Arbor. 50: 180–193.
- . 1980. Revision of South American *Saurauia* (Actinidiaceae). Fieldiana, Bot., n.s. 2: i–141.
- . 1985. *Saurauia molinae*, a new species of Actinidiaceae from Central America. Ann. Missouri Bot. Gard. 72: 878–880.



Pool, Amy. 1998. "Saurauia homotricha (Actinidiaceae), a new species from Honduras and Nicaragua." *Novon a journal of botanical nomenclature from the Missouri Botanical Garden* 8, 280–282. <https://doi.org/10.2307/3392020>.

View This Item Online: <https://www.biodiversitylibrary.org/item/14668>

DOI: <https://doi.org/10.2307/3392020>

Permalink: <https://www.biodiversitylibrary.org/partpdf/24897>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.