

my views as to the temperate character of that flora. But I have also got some plants from the northern part of the island which are quite new. You will probably be interested to learn, that amongst them there occurs a most beautiful specimen of Comptonia acutiloba Brong., which is known from the Miocene strata of Europe. As this species is most nearly allied to C. argenteifolia it affords an interesting proof that this latter probably is a descendant of the former, which may have passed to America over the old land about Bering's Strait.

Yours very truly
A. G. Mathernt.

Geological Survey Office
Stockholm

27 Feb. 1884

My dear Professor Gray,

Very many thanks for your kind letter of 18th Dec. 1883 and please excuse that I have not answered on it long ago. But I wished to be able to state positively, if I should be able to go to America this year or not, and I am sorry to say, that this will not be possible. I have many researches to make in the field this summer, which cannot be postponed, as I have already been absent during two summers (1882 and 1883) in



Nathorst, A. G. 1884. "Nathorst, Alfred G. Feb. 27, 1884." *Asa Gray correspondence*

View This Item Online: <https://www.biodiversitylibrary.org/item/224661>

Permalink: <https://www.biodiversitylibrary.org/partpdf/251924>

Holding Institution

Harvard University Botany Libraries

Sponsored by

Arcadia 19th Century Collections Digitization/Harvard Library

Copyright & Reuse

Copyright Status: Public domain. The Library considers that this work is no longer under copyright protection

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.