

Wheat Cal July 31, 90

Mr Tereno Watson

Dear Sir

Yours received of date July 25, and I am obliged for facts contained. As regards *C. pub-
chellus* I agree with you in naming spotted var
var *maculosus*. It may be true that in dried specimen
it is difficult to distinguish between variations of
C. Nuttallii. In blooming plants a very marked difference
is apparent between *C. Nuttallii* & *C. Leichtlinii*, although
morphologically they be very near alike. It seems to me however
that if plants of the two species grown side by side
are so different in appearance that any one would notice
it, and if blooming plants of the one are all very simi-
lar, while the other varies greatly, but does not approach
the former. I should say there was room for a variety,
if not a species. However I hope to bloom a consider-
able number of bulbs of both next year, and we can
study further on the subject. An odd feature in
Calochorti met my eye this season. It is in *C. luteo-*
(venosus) *ovatus*. I found on an elevated pass
numerous flowers of very brilliant coloring. Some of
these had the ordinary lunata gland, but in some



Purdy, Carl. 1890. "Purdy, Carl July 31, 1890 [to S. Watson]." *Asa Gray correspondence*

View This Item Online: <https://www.biodiversitylibrary.org/item/225468>

Permalink: <https://www.biodiversitylibrary.org/partpdf/252374>

Holding Institution

Harvard University Botany Libraries

Sponsored by

Arcadia 19th Century Collections Digitization/Harvard Library

Copyright & Reuse

Copyright Status: Public domain. The Library considers that this work is no longer under copyright protection

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.