

N. A. Flora, says "radical leaves on slender petioles, broadly rhomboidal." In my plant the radical leaves are always linear-spatulate - the caudine pair is very variable - usually at first united into a pinnatifid leaf, but presenting numerous gradations between that and oblong lanceolate or almost linear and sometimes starting below the middle of the stem - I have mature seeds and shall try to cultivate it more at large next year.

*Helianthus rigidus* - I meant to have written *H. hirsutus*, as I know *H. rigidus* quite well -

With many thanks for your kindness

I am truly yours

G. Vasey

Ringwood  
Prof. A. Gray  
Dear Sir,

Your  
inst. was duly received  
sheets of Silliman's  
Botany from the Public  
This Garden Botany is  
which has been needed  
work to the use of the  
whose principal reliance  
is upon the Garden & C

The Additions are of  
though I apprehend there  
in giving the names of  
plants - thus, *Artemisia*  
give the name of E Hall,  
it should be J. J. Hale,  
from Mr Hale, and I have  
not collected any in Wisconsin  
thing occurs under Lygodes



BHL

# Biodiversity Heritage Library

Vasey, George. 1849. "Vasey, George Nov. 28, 1862." *George Vasey letters to Asa Gray*

**View This Item Online:** <https://www.biodiversitylibrary.org/item/268647>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/253436>

**Holding Institution**

Harvard University Botany Libraries

**Sponsored by**

Arcadia 19th Century Collections Digitization/Harvard Library

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.