

Liquor Musci esculenti Odorem habet Spir. Tart.	Liqu. farin. secalis eundem,	Spirit. panis secalis, magis acidum	Spir. panis triticei eundem
Cum Spir. Vitr. Nitri, Salis, solut. alum. Acet. mutatio nulla	nulla	nulla,	nulla
Cum Mercur. sublim. turbidus & segnis principitab.	idem	idem,	turbidus ut Opal.
Luna, Sacch. Sat. Vitr. Vener. & Solut. Vener. nihil	nihil,	nihil,	nihil
Vitr. Mart. solut. obscuro-ferrugineus	idem	nigridus	obscuro-ferrua gineus
Solut. Mart. turbidus & niger	nigridus	ater	idem
Sal. Tart. tantillum fermentavit	nihil novi	nihil	parum alter- rationis
Solut. Resin. ferrugineus	granosus con- gulatus	turbidus	turbidus
Scor. Reg. Antimon. principiat. cum massa rubicunda	ferruginens c. foetore	ferrugineus	turbid. & fer- rug. c. foetore
Calx viva. nihil	nihil	nihil	nihil
Sulph. sol. c. Calce viva albus cum foetore	idem	idem	idem
Spirit. Sal. arm. parum ferment. obscurior	subturbidus	idem	nihil
Syrup. Viol. ruber	rubicundus	rubicundus	rubicundus
Lacmus cinnaberinus	idem	idem	idem
Solut. Gallar. nihil	nihil	nihil	nihil

Ex quo maxima eorum convenientiam intelliges.

Nomina, a Botanicis musco huic attributa, & Lin-
naei

nei studio collecta, Regiaeque Academiae exhibita sunt sequentia:

Muscus quidam crispæ lactucæ similis, minor & per ambitus tenuiter aculeatus. *Cord. Sylv.* 221.

Pulmonariæ alterum terrestre genus. *Camer. epit.* 283.
fig. 2. hist. 363. *fig. 2.*

Muscus catharticus. *Borrich. in Adis Haffn.* I. p. 126.
descr. absque figur. Sibbald. Scot. 2. p. 39.

Muscus Islandicus Bartholini. *Rai. hist.* 114.

Muscofungus terrestris, superne cinereus islandicus.
Mons. hist. 3. p. 632.

Muscus pulmonarius terrestris sanguineus. *Breyn. in Act.*
Imperial. 3. p. 439. c. f.

Coralloides cornua damæ referens. *Tournef. elem.* 442.
inst. 565. *paris.* 423.

Muscus terrestris intybaceus crispus duriusculus. *Cuppan.*
Suppl. 1. p. 248.

Muscus Islandicus purgans Bartholini. *Rai. syn.* 2.
p. 333. *opp.* 30.

Lichen terrestris pinetorum rigidus foliorum eryngii æmulus, basi purpurascente substantia ad fuscum vergente.
Rupp. Jen. I. p. 341. & 2. p. 297.

Lichen terrestris, foliis eryngii. *Buxb. cent.* 2. p. II.
t. 6. f. 1.

Lichen pulmonarius minor angustifoliis spinis tenuissimis
cel. *Mich. gen.* 85. t. 44. f. 4.

Lichen foliis oblongis laciniatis, marginibus conniventibus ciliatis. *Lin. flor. Lap.* 445. *obs. Stall. betv.* 75. *Roy Lugd.* 508.

Lichenoides rigidum eryngio folio referens. *Dill. musc.*
209. t. 28. f. III. *descr.* & *fig. opt.* *Rai. syn.* 3. p. 77.

Muscus est genuinus, & per totam Sveciam, in plagiis sterilibus, in sylvis abietinis, & inter junipreta occurrit. Hodie omnibus notissimus, & usus ejus pro pulce frequentissimus est, quam pthisi laborantibus optimam esse credunt. Hinc ab aliis muscus pulmonarius dicitur.

De alia ejus virtute recenter detecta, vide *Viri Spæring Schediasma*, de curatione mulieris, cuius uterus hydatidum plenus erat, supra pag. 273. expositum.



Linne

, Carl von. 1762. "Nomina, a Botanicis musco huic adtributa, & Linnaei studio collecta, Regiæque Academiæ exhibita sunt sequentia." *Analecta transalpina* 1, 328–329.

View This Item Online: <https://www.biodiversitylibrary.org/item/238177>

Permalink: <https://www.biodiversitylibrary.org/partpdf/259907>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.