

NOTE ON THE PREVALENCE OF INTESTINAL WORMS IN DOGS IN CAMBRIDGE.

BY GEORGE H. F. NUTTALL AND CYRIL STRICKLAND.

IN the course of investigations conducted in Cambridge upon dogs infected with *Piroplasma canis*, a number of animals have been incidentally examined for intestinal parasites. In an unbroken series of 24 dogs, which died (March—September 1907) from piroplasmosis, the animals were autopsied immediately after death and the intestine examined by slitting it open along its whole length. We briefly record the results of these examinations, for the reason that similar reports do not appear to have hitherto been published in this country. All of the dogs harboured worms. Only three species of worms were encountered: *Ascaris mystax*, *Dipylidium caninum* and *Taenia serrata*.

The parasites were distributed as follows:—

				No. of dogs har- bouring each species
<i>A. mystax</i>	17
<i>D. caninum</i>	14
<i>T. serrata</i>	4

The species occurred either alone or associated with each other:

10 dogs harboured *A. mystax*.

5 " " *D. caninum*.

5 " " *A. mystax* and *D. caninum*.

2 " " *D. caninum* and *T. serrata*.

2 " " *A. mystax*, *D. caninum* and *T. serrata*.

The numbers of *A. mystax* varied from 1 to 113 per dog.

" " " *D. caninum* " " 1 to 102 " "

" " " *T. serrata* " " 1 to 8 " "

The total number of each species recovered from the 24 dogs was

A. mystax 296.

D. caninum 312.

T. serrata 17.

Intestinal worms in dogs

The following table contains details regarding the numbers and species of worms found in each of the dogs examined:—

No. of dog	Species of worm	No. of worms found	No. of dog	Species of worm	No. of worms found
1	<i>A. mystax</i>	4	13	<i>A. mystax</i>	3
2	{ <i>D. caninum</i>	1	14	<i>D. caninum</i>	1
	{ <i>T. serrata</i>	3	15	<i>D. caninum</i>	82
	{ <i>A. mystax</i>	2	16	<i>A. mystax</i>	2
3	{ <i>D. caninum</i>	2	17	{ <i>D. caninum</i>	30
	{ <i>T. serrata</i>	1		{ <i>A. mystax</i>	6
4	<i>A. mystax</i>	5		{ <i>D. caninum</i>	1
5	<i>A. mystax</i>	1	18	{ <i>A. mystax</i>	45
6	<i>A. mystax</i>	48		{ <i>T. serrata</i>	5
7	<i>A. mystax</i>	4	19	<i>D. caninum</i>	1
8	<i>D. caninum</i>	7	20	{ <i>D. caninum</i>	53
9	{ <i>D. caninum</i>	20		{ <i>A. mystax</i>	7
	{ <i>T. serrata</i>	8	21	<i>A. mystax</i>	5
10	{ <i>D. caninum</i>	7	22	<i>A. mystax</i>	31
	{ <i>A. mystax</i>	2	23	{ <i>A. mystax</i>	113
11	{ <i>A. mystax</i>	4		{ <i>D. caninum</i>	4
	{ <i>D. caninum</i>	1	24	<i>D. caninum</i>	102
12	<i>A. mystax</i>	14			



Nuttall, George Henry Falkiner and Strickland, Cyril. 1908. "Note on the Prevalence of Intestinal Worms in Dogs in Cambridge." *Parasitology* 1(3), 261–262. <https://doi.org/10.1017/s0031182000003504>.

View This Item Online: <https://www.biodiversitylibrary.org/item/249184>

DOI: <https://doi.org/10.1017/s0031182000003504>

Permalink: <https://www.biodiversitylibrary.org/partpdf/264018>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.