

Crustacea Amphipoda: Hyperiidea from MUSORSTOM cruises

Wolfgang ZEIDLER

South Australian Museum
North Terrace, Adelaide
Australia 5000

ABSTRACT

Recent French expeditions to the Philippines, New Caledonia and Makassar Strait (Indonesia) have resulted in a small collection of hyperiid amphipods representing 11 species. All are tropical or warm temperate species. *Megalanceola stephensi*, a rare species, is represented by 8 specimens including some very large females and two males; only one male specimen has been recorded previously. *Paratyphis promontorii* is a new record for the south western Pacific.

RÉSUMÉ

Crustacea Amphipoda : Hyperiidea récoltés lors des campagnes MUSORSTOM

Des expéditions françaises récentes aux Philippines, en Nouvelle-Calédonie et dans le détroit de Makassar (Indonésie) ont permis de rassembler une petite collection d'Amphipodes hyperiens renfermant 11 espèces. Toutes sont des espèces d'eaux tropicales ou tempérées chaudes. *Megalanceola stephensi*, espèce rare, est représentée par 8 spécimens comprenant quelques très grandes femelles et deux mâles; jusqu'à présent un seul mâle de cette espèce avait été récolté. *Paratyphis promontorii* est signalé pour la première fois du Pacifique sud-ouest.

INTRODUCTION

This paper is a report on the hyperiid amphipods collected by recent French expeditions to Makassar Strait, Indonesia (CORINDON 2, 1980), the Philippines (MUSORSTOM 1-3, 1976, 1980, 1985) and New Caledonia (BIOCAL, 1985; CHALCAL 2, 1986; MUSORSTOM 4-5, 1985, 1986). More details regarding these expeditions are provided by FOREST (1981, 1985, 1989) for the Philippines, MOOSA (1984) for Indonesia and RICHER DE FORGES (1990)

for New Caledonia. Although the collection is very small and consists mainly of common tropical or warm temperate species it does contain two species of note. *Megalanceola stephensi*, a rare species, is represented by eight specimens including some very large females and two males, only one specimen of which has been recorded previously (SHOEMAKER, 1945), providing the opportunity to present here more complete illustrations (Fig. 1). *Paratyphis promontorii* is a new record for the south western Pacific.

There are no previous studies of hyperiids dealing specifically with the area covered by the present collections. However, some "historical" expeditions have passed through the general area and provide information on hyperiids collected e.g. "Challenger" (STEBBING, 1888), "Dana" (FAGE, 1960; SHIH, 1969) and "Siboga" (PIRLOT, 1930, 1938). Additional information on the Physosomata, from some parts of the study area, is provided by VINOGRADOV (1960) and ZEIDLER (1990).

In the following station list, the depths given refer to bottom depths, the specimens having been caught during bottom trawling or, being pelagic animals, most likely when the net was coming up. The precise depth of capture is therefore unknown. The systematic arrangement of families follows that of BOWMAN and GRÜNER (1973) except for Brachyscelidae which like VINOGRADOV *et al.* (1982), I have recognised as a separate family. The genera and species are arranged alphabetically. For each species only the original reference and synonymy is given. Specimen length was measured along a lateral parabolic line drawn from the anterior extremity of the head through the middle of the body to the posterior limit of the telson. All specimens are in the collection of the Muséum national d'Histoire naturelle, Paris.

LIST OF STATIONS

MUSORSTOM 1. Philippines.

Station 47. — 25.03.1976, 13°40.7'N, 120°30.0'E, 757-685 m : *Phronima sedentaria*.

CORINDON 2. Indonesia.

Station 201. — 30.10.1980, 01°10.2'S, 117°06.1'E, 21 m : *Platyscelus armatus*.

Station 231. — 04.11.1980, 0°04.9'N, 119°47.8'E, 980-1080 m : *Megalanceola stephensi*.

Station 281. — 08.11.1980, 01°57.5'S, 119°02.0'E, 1120-1150 m : *Megalanceola stephensi*.

MUSORSTOM 2. Philippines.

Station 50. — 27.11.1980, 13°37.0'N, 120°33.0'E, 810-820 m : *Megalanceola stephensi*.

Station 54. — 27.11.1980, 13°59.5'N, 120°09.3'E, 174-170 m : *Vibilia propinqua*.

MUSORSTOM 3. Philippines.

Station 95. — 01.06.1985, 13°56.0'N, 119°59.0'E, 865 m : *Phronima sedentaria*.

Station 106. — 02.06.1985, 13°47.0'N, 120°30.0'E, 640-668 m : *Megalanceola stephensi*.

Station 114. — 02.06.1985, 13°34.0'N, 120°29.0'E, 1000-1040 m : *Megalanceola stephensi*.

Station 129. — 05.06.1985, 11°43.0'N, 121°46.0'E, 1350 m : *Phronima sedentaria*.

Station 135. — 05.06.1985, 11°58.0'N, 122°02.0'E, 486-551 m : *Phronima sedentaria*.

BIOCAL. New Caledonia.

Station 4. — 11.08.1985, 21°16.0'S, 166°40.0'E, 2340 m : *Brachyscelus cruscum*, *Paratyphis promontorii*, *Prinno latreillei*, *Streetsia challenger*.

Station 5. — 11.08.1985, 21°16.0'S, 166°44.0'E, 2340 m : *Phronima sedentaria*.

Station 42. — 30.08.1985, 23°46.0'S, 167°13.0'E, 380 m : *Phronima sedentaria*.

Station 59. — 02.09.1985, 23°56.0'S, 166°41.0'E, 2650 m : *Platyscelus serratus*, *Vibilia propinqua*.

Station 62. — 02.09.1985, 24°19.0'S, 167°49.0'E, 1395-1410 m : *Phrosina semilunata*.

Station 74. — 04.09.1985, 22°14.0'S, 167°29.0'E, 1300-1475 m : *Phronima sedentaria*, *Scina crassicornis*.

Station 104. — 08.09.1985, 21°31.0'S, 166°21.0'E, 375-450 m : *Scina crassicornis*.

MUSORSTOM 4. New Caledonia.

Station 236. — 02.10.1985, 22°11.3'S, 167°15.0'E, 495-550 m : *Phronima sedentaria*.

Station 238. — 02.10.1985, 22°13.0'S, 167°14.0'E, 500-510 m : *Phronima sedentaria*.

MUSORSTOM 5. New Caledonia.

Station 335. — 15.10.1986, 20°03.24'S, 158°45.35'E, 315 m : *Phrosina semilunata*.

Station 340. — 16.10.1986, 19°48.5'S, 158°40.9'E, 675-680 m : *Phrosina semilunata*.

Station 363. — 19.10.1986, 19°47.9'S, 158°44.3'E, 700-685 m : *Phronima sedentaria*.

Station 382. — 21.10.1986, 19°37.39'S, 158°43.29'E, 580 m : *Phronima sedentaria*.

Station 389. — 22.10.1986, 20°44.95'S, 160°53.67'E, 500 m : *Phronima sedentaria*.

CHALCAL 2. New Caledonia.

Station 3. — 30.10.1986, 23°39.03'S, 167°43.11'E, 424 m : *Phronima sedentaria*.

SPECIES LIST

Scina crassicornis (Fabricius, 1775)

Megalanceola stephensi (Chevreux, 1920)

Vibilia propinqua Stebbing, 1888

Phronima sedentaria (Forskål, 1775)

Phrosina semilunata Risso, 1822

Prinno latreillei Stebbing, 1888

Brachyscelus cruscolum Bate, 1861

Streetsia challenger Stebbing, 1888

Paratyphis promontori Stebbing, 1888

Platyscelus armatus (Claus, 1879)

Platyscelus serratulus Stebbing, 1888

SYSTEMATIC ACCOUNT

Infraorder PHYSOSOMATA Pirlot, 1929

Superfamily SCINOIDEA Bowman & Gruner, 1973

Family SCINIDAE Stebbing, 1888

***Scina crassicornis* (Fabricius, 1775)**

Astacus crassicornis Fabricius, 1775 : 415.

Hyperia cornigera Milne-Edwards, 1830 : 387.

Clydonia gracilis Dana, 1853 : 834, pl. 55, fig. 6 a,b.

Tyro atlantica Bovallius, 1885 : 14.

Tyro sarsii Bovallius, 1885 : 15.

Scina edwardsi Garbowski, 1896 : 103, pl. 1, fig. 2; pl. 3, figs 19-33; pls 4-7; pl. 8, figs 97-109.

Scina crassicornis var. *bermudensis* Shoemaker, 1945 : 228, fig. 31.

MATERIAL EXAMINED. — **New Caledonia.** BIOCAL : stn 74, 1300-1475 m : 1 ♀ 12.4 mm. — Stn 104, 375-450 m : 1 ♀ 14 mm (approx. - damaged).

REMARKS. — This is a relatively common, cosmopolitan species occurring approximately between the polar circles (BRUSCA, 1978). It is often found near the surface (VINOGRADOV, 1957) particularly at night (THURSTON, 1976).

Superfamily LANCEOLOIDEA Bowman & Gruner, 1973

Family LANCEOLIDAE Bovallius, 1887

Megalanceola stephensi (Chevreux, 1920)

Fig. 1

Lanceola sp. Stebbing, 1888 : 1308, fig. 27.*Lanceola stephensi* Chevreux, 1920 : 4-7, figs 1-3.*Megalanceola terrae-novae* Pirlot, 1935 : 2, figs 1-4.

MATERIAL EXAMINED. — **Indonesia.** CORINDON 2 : stn 231, 980-1080 m : 1 ♀ 95 mm. — Stn 281, 1120-1150 m : 1 ♀ 90 mm.

Philippines. MUSORSTOM 2 : stn 50, 810-820 m : 1 ♀ 71 mm, 1 ♂ 53 mm.

MUSORSTOM 3 : stn 106, 640-668 m : 1 ♀ 68 mm, 1 ♂ 56 mm (approx. - damaged). — Stn 114, 1000-1040 m : 2 ♀ 68 and 81 mm.

REMARKS. — This appears to be a rare species as it has only been recorded in the literature on 5 previous occasions. From the Banda Sea STEBBING (1888), near the Azores (CHEVREUX, 1920), near Newfoundland (PIRLOT, 1935), near Bermuda (SHOEMAKER, 1945) and from the "Southern Ocean" (HERRING, 1981). Regarding STEBBING's record (as *Lanceola* sp.) the figures accompanying STEBBING's remarks, made by WILLEMOES SUHM, are rather crude but I am in agreement with SHOEMAKER in that they represent this species. The material from CORINDON 2 confirms the presence of this species in Indonesian waters.

The present collection is significant for the presence of very large females (up to 95 mm) and two males. Previously the largest female, recorded by SHOEMAKER (1945), measured only 73 mm, not taking into account the bent nature of the anterior part of the animal. SHOEMAKER also recorded the first and only known male specimen (48 mm) of this species and provided some illustrations. As SHOEMAKER's illustrations are limited I have provided more detailed drawings of one of the males for future reference. The males differ from the females mainly in the shape and size of the first antenna which is relatively larger in males. Mature males also have distinct penial projections on the ventral surface of pereonite 7 adjacent to the seventh pereopod.

Infraorder PHYSOCEPHALATA Bowman & Gruner, 1973

Superfamily VIBILIOIDEA Bowman & Gruner, 1973

Family VIBILIIDAE Dana, 1852

Vibilia propinqua Stebbing, 1888

Figs 2-3

Vibilia propinqua Stebbing, 1888 : 1279, pl. 147.*Vibilia milnei* Stebbing, 1888 : 1284, pl. 148A.*Vibilia* sp. Stebbing, 1888 : 1285, pl. 148B, figs C, D.

MATERIAL EXAMINED. — **Philippines.** MUSORSTOM 2 : stn 54, 174-170 m : 1 ♀ 6.3 mm.

New Caledonia. BIOCAL : stn 59, 2650 m : 1 ♀ 8.0 mm.

REMARKS. — The genus *Vibilia* is in a state of considerable taxonomic confusion and a revision is long overdue. VINOGRADOV *et al.* (1982) recognise 17 valid species and provide a useful key but specific determination,




FIG. 1. — *Megalanceola stephenseni* (Chevreux, 1920) : A, lateral view of ♂ 53 mm, MUSORSTOM 2, Stn 50 (scale = 5.0 mm); B-L, ♂ 58 mm, MUSORSTOM 3, Stn 106 (scales = 1.0 mm); B, urosome; C, upper lip; D, lower lip; E, first maxilla; F, second maxilla; G, mandible, inner face; H, mandible, outer face; I, maxilliped, outer face; J, maxilliped, inner face; K, first antenna, outer face; L, first antenna, inner face; M-N, outer and inner face of first antenna, ♀ 68 mm, MUSORSTOM 3, Stn 106.




FIG. 2. — *Vibilia propinqua* Stebbing, 1888, ♀ 6.5 mm, BIOCAL, Stn 59 : A, lateral view; B, urosome; C, first antenna; D, second antenna. (Scale A = 1.0 mm; B-D = 0.2 mm).

except for a few distinctive species, is still very difficult. The present specimens most closely resemble *V. propinqua* and are most likely this species but because there is some doubt I have figured one of the specimens for future reference.

A relatively common species, widely distributed in tropical and temperate regions.

Superfamily PHRONIMOIDEA Bowman & Gruner, 1973

Family PHRONIMIDAE Dana, 1852

Phronima sedentaria (Forskål, 1775)

Cancer sedentarius Forskål, 1775 : 95.

Phronima custos Risso, 1816 : 121, pl. 2, fig. 3.




FIG. 3. — *Vibilia propinqua* Stebbing, 1888, ♀ 6.5 mm, BIOCAL, Stn 59 : A-G, first to seventh pereopods (Scale = 0.2 mm); H, maxilliped, inner face; I, first maxilla; J, second maxilla; K, mandible (palp twisted). (Scale H-K = 0.1 mm).

- Phronima borneensis* Bate, 1862 : 318, pl. 51, fig. 3.
Phronima novaezealandiae Powell, 1875 : 294, pl. 21, figs 1,2.
Phronima spinosa Bovallius, 1887 : 25.
Phronima tenella Stebbing, 1888 : 1354, pl. 161A.
Phronima affinis Vosseler, 1901 : 20, pl. 1, figs 12-16.

MATERIAL EXAMINED. — **Philippines.** MUSORSTOM 1 : stn 47, 757-685 m : 1 ♀ 30.0 mm.

MUSORSTOM 3 : stn 95, 865 m : 1 ♀ 45.6 mm. — Stn 129, 1350 m : 1 ♀ 36.0 mm. — Stn 135, 486-551 m : 1 ♀ 30.4 mm.

New Caledonia. BIOCAL : stn 5, 2340 m : 1 ♀ 18.0 mm. — Stn 42, 380 m : 1 ♀ 25.4 mm. — Stn 74, 1300-1475 m : 1 ♀ 27.2 mm.

MUSORSTOM 4 : stn 236, 495-550 m : 1 ♀ 34.5 mm. — Stn 238, 500-510 m : 2 ♀ 29.7 and 30.2 mm.

MUSORSTOM 5 : stn 363, 700-685 m : 2 ♀ 23.7 and 28.0 mm. — Stn 382, 580 m : 1 ♀ 32.6 mm. — Stn 389, 500 m : 1 ♀ 38.5 mm.

CHALCAL 2 : stn 3, 424 m : 2 ♀ 20.0 and 31.8 mm.

REMARKS. — This is the most common and largest species of the family; widely distributed in tropical and temperate regions. It is often found in association with salps and pyrosomes and its ability to build and live in "barrels" made from these organisms is well documented in the literature (LAVAL, 1978).

Family PHROSINIDAE Dana, 1853

Phrosina semilunata Risso, 1822

- Phrosina semilunata* Risso, 1822 : 245.
Dactylocera nicaeensis Milne-Edwards, 1830 : 393.
Phrosina longispina Bate, 1862 : 320, pl. 5, fig. 7.
Phrosina pacifica Stebbing, 1888 : 1430.
Phrosina australis Stebbing, 1888 : 1431.

MATERIAL EXAMINED. — **New Caledonia.** BIOCAL : stn 62, 1395-1410 m : 1 ♀ 15.5 mm.

MUSORSTOM 5 : stn 335, 315 m : 1 ♀ 13.1 mm. — Stn 340, 675-680 m : 1 ♀ 14.8 mm.

REMARKS. — A very common, cosmopolitan species favouring tropical and temperate regions.

Primno latreillei Stebbing, 1888

Primno latreillei Stebbing, 1888 : 1445-1447, pl. 179A.

MATERIAL EXAMINED. — **New Caledonia.** BIOCAL : stn 4, 2340 m : 1 ♀ 5.5 mm.

REMARKS. — The genus *Primno* was revised by BOWMAN (1978) who recognised 4 species previously lumped as *P. macropa*. Two additional species were described later, one by BOWMAN (1985) and the other by SHEADER (1986). The present specimen is readily identified as typical *P. latreillei*, distinct from the very closely related species *P. johnsoni* Bowman, 1978.

The distribution of this species is difficult to determine due to past confusion with *P. macropa* but BOWMAN (1978) records it from the central North Pacific, southeast Gulf of Guinea, eastern Mediterranean and the Red Sea and I have recorded it from eastern Australia (ZEIDLER, 1978).

Superfamily PLATYSCELOIDEA Bowman & Gruner 1973

Family BRACHYSCHELIDAE Stephensen, 1923

Brachyscelus crusculum Bate, 1861

? *Orio Zancleus* Natale, 1850 : 12.

Brachyscelus crusculum Bate, 1861 : 7, pl. 2, figs 1,2.

Thamyris antipodes Bate, 1862 : 335, pl. 50, fig. 4.

? *Thamyris lycaeoides* Claus, 1887 : 60, pl. 21, figs 1,2.

? *Thamyris mediterranea* Claus, 1887 : 60, pl. 16, figs 11-18.

Brachyscelus acuticaudatus Stebbing, 1888 : 1555-1556, pl. 197C.

Brachyscelus stebbingi Boone, 1935 : 226-230, pls 67, 68.

MATERIAL EXAMINED. — New Caledonia. BIOCAL : stn 4, 2340 m : 1 ♀ 7.7 m.

REMARKS. — The genus *Brachyscelus* is in a state of considerable taxonomic confusion and a revision is long overdue. According to MADIN and HARBISON (1977) at least 17 species have been described, most of which are synonyms of *B. crusculum*. VINOGRADOV *et al.* (1982) recognised only 4 species. *B. crusculum* is the most common species of the genus being widely distributed in tropical and temperate regions.

Family OXYCEPHALIDAE Bate, 1861

Streetsia challengerii Stebbing, 1888

Streetsia challengerii Stebbing, 1888 : 1603-1606, pl. 207.

? *Oxycephalus pronoides* Bovallius, 1887 : 37.

Streetsia stebbingi Chevreux, 1900 : 161, pl. 18, fig. 4.

Streetsia washingtoni Senna, 1902 : 15, pl. 2.

Streetsia sabauda Colosi, 1918 : 218, pl. 17, figs 5-9.

Streetsia gaussi Spandl, 1927 : 184, fig. 17.

MATERIAL EXAMINED. — New Caledonia. BIOCAL : stn 4, 2340 m : 1 ♀ 11.0 mm (approx. - damaged; ovigerous).

REMARKS. — This is a very common, widely distributed species recorded mainly from tropical regions. FAGE (1960) gives more details regarding the biology of this species.

Family PLATYSCELIDAE Bate, 1862

Paratyphis promontorii Stebbing, 1888

Paratyphis promontorii Stebbing, 1888 : 1476-1479, pl. 209D.

? *Paratyphis theelii* Bovallius, 1887 : 47.

MATERIAL EXAMINED. — New Caledonia. BIOCAL : stn 4, 2340 m : 1 ♀ 3.5 mm.

REMARKS. — Four species of *Paratyphis* are currently recognised (VINOGRADOV *et al.*, 1982) but specific determination is difficult and the genus is in need of revision. The best distinguishing character of *P. promontorii* appears to be the form of the gnathopods.

This is a relatively rare species known mainly from the Atlantic in tropical and temperate areas. STEBBING (1888) and DICK (1970) record it from around South Africa and SPANDL (1927) records it from the Red Sea (misidentified as *P. maculatus*). According to VINOGRADOV *et al.* (1982) it has also been recorded from the south eastern Pacific Ocean but I have been unable to verify this from the literature. The present specimen is the first record from the south-west Pacific.

This species is usually found near the surface; 0-150 m (THURSTON, 1976).

Platyscelus armatus (Claus, 1879)

Eutyphis armatus Claus, 1879 : 10.

Eutyphis inermis Claus, 1887 : 37.

MATERIAL EXAMINED. — Indonesia. CORINDON 2 : stn 201, 21 m : 1 ♀ 15.0 mm.

REMARKS. — This species is easily distinguished from its congeners by the lateral process on the fifth and sixth coxal plates. It is a relatively rare species known from widely scattered records from tropical and warm temperate regions.

Platyscelus serratulus Stebbing, 1888

Eutyphis serratus Claus, 1879 : 11.

Platyscelus serratulus Stebbing, 1888 : 1470-1471.

Platyscelus dubius Shoemaker, 1925 : 51, figs 20, 21.

MATERIAL EXAMINED. — New Caledonia. BIOCAL : stn 59, 2650 m : 1 ♀ 4.4 mm.

REMARKS. — This is a widely distributed species occurring in tropical and subtropical regions.

ACKNOWLEDGEMENT

I am very grateful to Dr Alain CROSNIER (ORSTOM and Muséum national d'Histoire naturelle, Paris) and Mr Michel SEGONZAC (Centre national de tri d'Océanographie biologique, Plouzané) for forwarding specimens to me for study.

REFERENCES

- BATE, C.S., 1861. — On the morphology of some Amphipoda of the Division Hyperina. *Ann. Mag. nat. Hist.*, Ser. 3, 8 (43) : 1-16, pls 1, 2.
- BATE, C.S., 1862. — Catalogue of the specimens of Amphipodous Crustacea in the collection of the British Museum. 399 pp., pls 1-58. London.
- BOONE, L., 1935. — Scientific results of the world cruise of the yacht "Alva", 1931, William K. Vanderbilt, commanding. Crustacea: Anomura, Macrura, Euphausiacea, Isopoda, Amphipoda and Echinodermata. *Bull. Vanderbilt Mar. Mus.*, 6 : 1-264, pls 1-96.

- BOVALLIUS, C., 1885. — On some forgotten genera among the Amphipodous Crustacea. *Bih. Svenska Vetensk. Akad. Handl.*, **10** (14) : 1-17, 1 pl.
- BOVALLIUS, C., 1887. — Systematical list of the Amphipoda Hyperiidæ. *Bih. Svenska Vetensk. Akad. Handl.*, **11** (16) : 1-50.
- BOWMAN, T.E., 1978. — Revision of the pelagic amphipod genus *Primno* (Hyperiidæ: Phrosinidae). *Smithson. Contrib. Zool.*, (275) : 1-23.
- BOWMAN, T.E., 1985. — The correct identity of the pelagic amphipod *Primno abyssalis* (Hyperiidæ: Phrosinidae). *Proc. biol. Soc. Wash.*, **98** (1) : 121-126.
- BOWMAN, T.E., & GRÜNER, H.-E., 1973. — The families and genera of Hyperiidæ (Crustacea: Amphipoda). *Smithson. Contrib. Zool.*, (146) : 1-64.
- BRUSCA, G.J., 1978. — Contributions to the knowledge of hyperiid amphipods of the family Scinidae from near Hawaii, with a description of a new species, *Scina hawaiiensis*. *Pacif. Sci.*, **32** (2) : 281-292.
- CHEVREUX, E., 1900. — Amphipodes provenant des campagnes de l'Hirondelle. *Résult. Camp. sci. Monaco*, **16** (1-4) : 1-195, pls 1-18.
- CHEVREUX, E., 1920. — Révision des Lanceolidæ provenant des campagnes de S.A.S. le Prince de Monaco. *Bull. Inst. océanogr., Monaco*, (363) : 1-12.
- CLAUS, C., 1879. — Der Organismus der Phronimiden. *Arb. zool. Inst. Univ. Wien*, **2** : 59-146, pls 1-8.
- CLAUS, C., 1887. — Die Platysceliden. 77 pp., 25 pls. Vienna, Alfred Hölder.
- COLOSI, G., 1918. — Oxicephalidi. - Raccolte planctoniche fatte dalla R. Nave "Liguria" 1903-1905, II, fascicule 3. *Crostacei. Pubbl. R. Ist. Stud. super. Firenze*, **2** (3) : 207-224, pl. 17.
- DANA, J.D., 1852. — On the classification of the Crustacea Choristopoda or Tetracapoda. *Am. J. Sci. Arts, Ser. 2*, **14** (41) : 297-316.
- DANA, J.D., 1853. — Crustacea, Part II. U.S. Expl. Exped. **14**: 689-1618. (Pls 1-96 published 1855).
- DICK, R.I., 1970. — Hyperiidæ (Crustacea: Amphipoda) keys to South African genera and species, and a distribution list. *Ann. S. Afr. Mus.*, **57** (3) : 25-86.
- FABRICIUS, J.C., 1775. — Systema Entomologiae, sistens Insectorum Classes, Ordines, Genera, Species, adjectis Synonymis, Locis, Descriptionibus, Observationibus. Flensburgi et Lipsiae, 832 pp.
- FAGE, L., 1960. — Oxycephalidae, amphipodes pelagique. *Dana Rep.*, (52) : 1-145.
- FOREST, J., 1981. — Compte rendu et remarques générales. In : Résultats des Campagnes MUSORSTOM. I. Philippines (18-28 Mars 1976), **1** (1). *Mém. ORSTOM*, (91) : 9-50.
- FOREST, J., 1985. — La campagne MUSORSTOM II (1980). Compte rendu et liste de stations. In : Résultats des Campagnes MUSORSTOM I et II. Philippines (1976, 1980), **2** (1). *Mém. Mus. natn. Hist. nat.*, (A), **133** : 7-30.
- FOREST, J., 1989. — Compte rendu de la Campagne MUSORSTOM 3 aux Philippines (31 mai - 7 juin 1985). In : J. FOREST (ed.), Résultats des Campagnes MUSORSTOM, **4** (1). *Mém. Mus. natn. Hist. nat.*, (A), **143** : 9-23.
- FORSKÅL, P., 1775. — Descriptiones Animalium, Avium, Amphibiorum, Piscium, Insectorum, Vermium; quae in Itinere orientali observavit Petrus Forskål. Prof. Haun. Post Mortem Auctoris edidit. Carsten Niebuhr. Adjuncta est Materia Medica Kahirina atque Tabula Maris Rubri Geographica. Hafniae : 1-9 + i-xxxiv + 1-164, 1 map.
- GARBOWSKI, T., 1896. — Zoologische Ergebnisse IX, Hyperienartige Amphipoden des Mittelmeeres. Monographisch bearbeitet auf Grund des während der fünf Expeditionen S.M. Schiffes "Pola" gesammelten Materiales (1890-1894). I. Die Sciniden. (Bericht Commission Tiefsee-Forschung XX). *Denkschr. K. Akad. Wiss. Wien, Math.-naturwiss. Kl.*, **63** (2) : 89-116, 9 pls.
- HERRING, P.J., 1981. — Studies on bioluminescent marine amphipods. *J. mar. biol. Ass. U.K.*, **61** (1) : 161-176.
- LAVAL, P., 1978. — The barrel of the pelagic amphipod *Phronima sedentaria* (Forsk.) (Crustacea: Hyperiidæ). *J. exp. mar. Biol. Ecol.*, **33** : 187-211.
- MADIN, L.P. & HARBISON, G.R., 1977. — The associations of Amphipoda Hyperiidæ with gelatinous zooplankton - I. Associations with Salpidae. *Deep Sea Res.*, **24** : 449-463.

- MILNE EDWARDS, H., 1830. — Extrait de recherches pour servir à l'histoire naturelle des Crustacés Amphipodes. *Annls Sci. nat.*, **20** : 353-399, pls 10-11.
- MOOSA, M.K., 1984. — Report on the Corindon Cruises. *Mar. Res. Indonesia*, (24) : 1-6, figs 1-2.
- NATALE, G. DE., 1850. — Descrizione zoologica d'una nuova specie di Plojaria ed alcuni Crostacei del porto di Messina con poche considerazioni generali sulla natura delle appendici aculeiformi delle piante e degli animali. Messina : 1-31, appendix 2 pp., pls 1-2.
- PIRLOT, J. M., 1929. — Les Amphipodes Hypérides. - Résultats zoologiques de la croisière atlantique de l' "Armauer Hansen" (1922). *Mém. Soc. r. Sci. Liège, Ser. 3*, **15** (2) : 1-196.
- PIRLOT, J. M., 1930. — Les Amphipodes de l'expédition du "Siboga". Première Partie, Les Amphipodes Hypérides (à l'exception des Thaumtopsidae et des Oxycephalidae). *Siboga-Exped.*, Monogr. **33** : 1-55.
- PIRLOT, J. M., 1935. — Un grand Amphipode Hypéride, nouveau comme genre et comme espèce. *Bull. Inst. océanogr., Monaco*, (681) : 1-8.
- PIRLOT, J. M., 1938. — Amphipodes de l'expédition du Siboga. 2. Les Amphipodes littoraux. Addendum. Les Amphipodes Hypérides. *Siboga-Exped.*, Monogr. **33** : 329-388.
- POWELL, L., 1875. — Description of a new crustacean *Phronima novae-zealandiae*. *Trans. Proc. N. Z. Inst.*, **7** : 294-295.
- RICHER DE FORGES, B., 1990. — Les campagnes d'exploration de la faune bathyale dans la zone économique de la Nouvelle-Calédonie [Explorations for bathyal fauna in the New Caledonian economic zone]. In : A. CROSNIER (ed.), *Résultats des Campagnes MUSORSTOM*, **6** (1). *Mém. Mus. natn. Hist. nat.*, **145** : 9-54.
- RISSE, A., 1816. — Histoire naturelle des Crustacés des environs de Nice. Paris. Librairie Grecque Latine Allemande. 175 pp., 3 pls.
- RISSE, A., 1822. — Mémoire sur quelques nouveaux Crustacés observés dans la mer de Nice. *J. Phys. Chim. Hist. nat.*, **95** : 241-248.
- SENNA, A., 1902. — Le esplorazione abissali nel Mediterraneo del R. Piroscapo "Washington" nel 1881. I. Nota sugli Oxycefalidi. *Bull. Soc. ent. ital.*, **34** : 10-32, pl. 2.
- SHEAHER, M., 1986. — *Primno evansi* sp. nov. (Amphipoda: Hyperiidea) from the eastern North Atlantic. *Jnl. nat. Hist.*, **20** : 975-980.
- SHIH, C.T., 1969. — The systematics and biology of the family Phronimidae (Crustacea: Amphipoda). *Dana Rep.*, (74) : 1-100.
- SHOEMAKER, C.R., 1925. — The Amphipoda collected by the United States Fisheries Steamer "Albatross" in 1911, chiefly in the Gulf of California, Scientific Results of the Expedition to the Gulf of California. *Bull. Am. Mus. nat. Hist.*, **52** : 21-61.
- SHOEMAKER, C.R., 1945. — The Amphipoda of the Bermuda Oceanographic Expeditions, 1929-1931. *Zoologica*, N.Y., **30** : 185-266.
- SPANDL, H., 1927. — Die Hyperiidien (exkl. Hyperiidea Gammaroidea und Phronimidae) der Deutschen Südpolar-Expedition 1901-1903. *Dt. Südpolar-Exped. 1901-1903*, **19**, Zoologie **11** : 145-287, pl. 10.
- STEBBING, T.R.R., 1888. — Report on the Amphipoda collected by H.M.S. CHALLENGER during the years 1873-1876. *Rep. scient. Res. Voy. 'Challenger'*, Zool., **29** : xxiv + 1-1737, pls. 1-210.
- STEPHENSEN, K., 1923. — Crustacea Malacostraca, V. Amphipoda I. *Danish Ingolf-Exped.*, **3**(8) : 1-100.
- THURSTON, M.H., 1976. — The vertical distribution and diurnal migration of the Crustacea Amphipoda collected during the SONDA cruise, 1965. II. The Hyperiidea and general discussion. *J. mar. biol. Ass. U.K.*, **56** : 383-470.
- VINOGRADOV, M.E., 1957. — Hyperiid (Amphipoda) of the north western Pacific Ocean. 1. Tribe Hyperiidea Physosomata. (In Russian.) *Trudy Inst. Okeanol. Akad. Nauk SSSR*, **20** : 186-227. (English translation : *Mar. Biol.* : 149-184. Washington D.C. : American Institute of Biological Sciences, 1959.)
- VINOGRADOV, M.E., 1960. — Hyperiidea Physosomata of the tropical Pacific Ocean. (In Russian.) *Trudy Inst. Okeanol. Akad. Nauk SSSR*, **41** : 198-247.
- VINOGRADOV, M.E., VOLKOV A.F. & SEMENOVA, T.N., 1982. — Amphipody - Giperiidy (Amphipoda: Hyperiidea) Mirovogo Okeana. *Akademiia Nauk SSSR, Opredeliteli po Faune SSSR*, **132** : 1-492.

- VOSSELER, J., 1901. — Die Amphipoden der Plankton-Expedition. 1. Thiel. Hyperiidea 1. *Ergebn. Plankton-Exped.*, 2 : 1-129, pls 1-13.
- ZEIDLER, W., 1978. — Hyperiidea (Crustacea: Amphipoda) from Queensland waters. *Aust. J. Zool.*, Suppl. Ser. (59) : 1-93.
- ZEIDLER, W., 1990. — Pelagic amphipods, Infraorder Physosomata (Crustacea: Amphipoda: Hyperiidea) from the CSK International Zooplankton Collection (western North Pacific), with the description of four new species of *Scina*. *Publs Seto mar. biol. Lab.*, 34 (4/6) : 167-200.

Crustacea Isopoda : Deep-sea Chaetiliidae (Valvifera) from New Caledonia and the Philippines

Gary C.B. POORE

Department of Crustacea, Museum of Victoria
Swanston Street, Melbourne, Victoria 3000
Australia

ABSTRACT

Four species of *Stegidopsis* Poore (*S. pinnata* Poore, *S. longipes* sp. nov., *S. carinata* sp. nov., and *S. jerdoni* sp. nov.) are reported from deep water in the region of New Caledonia. The last of these is unusual in several respects having a complete set on the palm of the first large pereopod, subequal unipodal rami, a single like median process and a habitus different from all other species of the genus. *Symmetrus philippinensis* sp. nov. is described from the Philippines. One third species in this north western Pacific.

The 14 species of *Chaetiliidae* known from the Western Pacific (Japan, Philippines, New Caledonia, Australia and New Zealand) are tabulated.

RÉSUMÉ

Crustacés Isopoda : Chaetiliidae (Valvifera) des eaux profondes de Nouvelle-Calédonie et des Philippines.

Quatre espèces de *Stegidopsis* Poore (*S. pinnata* Poore, *S. longipes* sp. nov., *S. carinata* sp. nov., et *S. jerdoni* sp. nov.) ont été récoltées dans les eaux profondes de la Nouvelle-Calédonie. La dernière espèce est présente des caractères inhabituels, ayant une série sous dentée sur le propode, des rami unipodales subégales, un processus médian en forme de pointe et un habitus différent de celui de toutes les autres espèces de genre. *Symmetrus philippinensis* sp. nov. est décrit des Philippines et est la troisième espèce de ce genre du Pacifique nord-ouest.

Une liste des 14 espèces des Chaetiliidae du Pacifique occidental (Japon, Philippines, Nouvelle-Calédonie, Australie et Nouvelle-Zélande) est donnée.

POORE, G.C.B., 1991. — Crustacea Isopoda : Deep-sea Chaetiliidae (Valvifera) from New Zealand and the Philippines. In : A. Chiriac (ed.), Résultats des campagnes MUSORSTOM, Volume 9. Mémoires de la Société de Biogéographie, Paris 1990, 2 : 85-113-127-11.



Zeidler, Wolfgang. 1991. "Crustacea Amphipoda: Hyperiidea from Musorstom cruises." *Mémoires du Muséum national d'histoire naturelle* 152(1), 125–137.

View This Item Online: <https://www.biodiversitylibrary.org/item/266010>

Permalink: <https://www.biodiversitylibrary.org/partpdf/283046>

Holding Institution

Muséum national d'Histoire naturelle

Sponsored by

Muséum national d'Histoire naturelle

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Muséum national d'Histoire naturelle

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.