

Establishment of a new shrimp family Chlorotocellidae for four genera previously assigned to Pandalidae (Decapoda, Caridea, Pandaloidae)

Tomoyuki Komai¹, Tin-Yam Chan², Sammy De Grave³

¹ Natural History Museum and Institute, Chiba, 955-2 Aoba-cho, Chuo-ku, Chiba 260-8682, Japan

² Institute of Marine Biology and Center of Excellence for the Oceans, National Taiwan Ocean University, 2 Pei-Ning Road, Keelung 20224, Taiwan

³ Oxford University Museum of Natural History, Parks Road, Oxford, UK

<http://zoobank.org/86895CA3-596A-4015-8350-82EEF10F9885>

Corresponding author: Tin-Yam Chan (tychan@mail.ntou.edu.tw)

Academic editor: Kristina von Rintelen ♦ Received 8 May 2019 ♦ Accepted 17 June 2019 ♦ Published 3 July 2019

Abstract

A new caridean shrimp family, Chlorotocellidae, is established to accommodate four genera previously assigned to Pandalidae, viz., *Chlorotocella* Balss, 1914 (type genus), *Chlorocurtis* Kemp, 1925, *Anachlorocurtis* Hayashi, 1975, and *Miopandalus* Bruce, 1983, which represents the sister clade to a clade consisting of all other pandalid genera (including the two genera previously assigned to Thalassocarididae) in a recent comprehensive phylogenetic analysis of Pandaloidae. Diagnoses are provided for the new family and its constituent genera, and a comparison with Pandalidae is provided, for which a new diagnosis is given.

Key Words

Anachlorocurtis, ASR analysis, *Chlorotocella*, *Chlorocurtis*, *Miopandalus*

Introduction

The caridean family Pandalidae Haworth, 1825 is predominantly composed of cold-water and deep-sea taxa, including several species of commercial importance (Holthuis 1980). An extensive multimarker molecular phylogenetic analyses of the family and the closely related Thalassocarididae by Liao et al. (2019) clarified that Thalassocarididae, represented by two genera, *Thalassocaris* Stimpson, 1860 and *Chlorotocoides* Kemp, 1925, are deeply nested within Pandalidae, and that four genera, *Chlorotocella* Balss, 1914, *Chlorocurtis* Kemp, 1925, *Anachlorocurtis* Hayashi, 1975, and *Miopandalus* Bruce, 1983, comprise the sister clade to the remaining clade consisting of all other pandalid genera. These four genera are represented by small-sized species inhabiting shallow subtidal waters in tropical to subtropical and often associated with cnidarians (Hayashi and Miyake 1968; Hayashi 1975; Bruce 1983; Okuno and Yokota 1995; Minemizu 2000, 2013; Kato and Okuno 2001; Kawamoto and Okuno 2003; Humann and DeLoach

2010; Horká et al. 2014; Anker and De Grave 2016) in sharp contrast to the ecologies of most Pandalidae. To incorporate the documented phylogenetic pattern into the formal classification of Caridea, together with considerations on the morphological distinctness and ecological traits of these taxa, we hereby propose a new family, Chlorotocellidae fam. nov. for these four genera and redefine the family Pandalidae.

Materials and methods

The morphological data assembled following an examination of the literature and direct examination of specimens of relevant taxa formed the basis of the phylogenetic analysis by Liao et al. (2019) and can be found in the online supplementary material for that study. Aside from the family diagnosis, diagnoses are provided for each genus in the new family; to shorten diagnoses, synapomorphies for the family are not repeated in generic diagnosis; autapomorphies are in bold italics.

Illustrations showing diagnostic characters are given for *Chlorotocella* (*C. gracilis* Balss, 1914) and *Chlorocurtis* (*C. jactans* (Nobili, 1904)), as no published modern illustrations are available in easily accessible literature for those taxa. The three species of *Anachlorocurtis* and the monotypic *Miropandalus* have been well illustrated in their respective type description, and thus, no additional figures are presented.

Details of specimens used for preparation of drawings are listed below. These specimens are deposited in the Laboratory of Marine Zoology, Faculty of Fisheries, Hokkaido University, Hakodate, Japan (HUMZ), National Research Institute for Far Seas Fisheries, Shizuoka, Japan (NRIFS), and the Natural History Museum and Institute, Chiba, Japan (CBM).

Chlorotocellidae fam. nov.

Chlorocurtis jactans (Nobili, 1904): CBM-ZC 11596, 1 ovigerous female (cl 1.3 mm), Uehara, Iriomote Island, Yaeyama Islands, Ryukyu Islands, sea grass beds, 0.5–1 m at low tide, 18 July 2007, dip net, coll. T. Komai.

Chlorotocella gracilis Balss, 1914: CBM-ZC 12534, 1 male (cl 5.0 mm), TRV “Toyoshio-maru”, 2001-6 cruise, stn 4, W of Tanegashima Island, Ohsumi Islands, 30°33.50'N, 130°53.30'E, 47 m, 26 May 2001, dredge, coll. T. Komai; HUMZ-C 1556, 1 male (cl 4.1 mm), Tosa Bay, 40 m, 22 May 1960, coll. M. Toriyama.

Pandalidae Haworth, 1825

Chlorotocus novaezealandiae (Borradaile, 1916): NRIFS 578, 1 male (cl 17.0 mm), New Zealand, no other data.

Heterocarpus ensifer A. Milne-Edwards, 1881: HUMZ-C 255, 1 male (cl 33.0 mm), no data.

Pandalus montagui Leach, 1814: CBM-ZC 3422, 1 transitional male (cl 10.5 mm), Texel, The Netherlands, 8 April 1991.

Systematics

Infraorder Caridea Dana, 1852

Superfamily Pandalioidea Haworth, 1825

Family Pandalidae Haworth, 1825

Type genus. *Pandalus* Leach, 1814, by original designation.

Composition. *Atlantopandalus* Komai, 1999, *Austropandalus* Holthuis, 1950, *Bitias* Fransen, 1990, *Chelonika* Fransen, 1997, *Chlorotocoides* Kemp, 1925, *Chlorotocus* A. Milne-Edwards, 1882, *Dichelopandalus* Caullery, 1896, *Dorodotes* Bate, 1888, *Heterocarpus* A. Milne-Edwards, 1881, *Heteronika* Hendrickx, 2019, *Notopandalus* Yaldwyn, 1960, *Pandalina* Calman, 1899, *Pandalus* Leach, 1814, *Pantomus* A. Milne-Edwards, 1883, *Peripandalus* de Man, 1917, *Plesionika* Bate, 1888, *Proclestes* Bate, 1888, *Pseudopandalus* Crosnier, 1997, and *Thalassocaris* Stimpson, 1860.

Diagnosis. Rostrum well developed, usually ventrally with teeth or rows of setae. Thoracic sternites 6–8 each with paired conspicuous prominences, teeth or protuberances (Fig. 3A). Pleomere 6 posterolateral process usually terminating in small tooth. Telson with longitudinal row of spiniform setae located on dorso-lateral ridges. Eyestalks subpyriform or kidney-shaped, cornea distinctly longer and wider than eyestalk. Antennular stylocerite with proximolateral projection, distally acuminate or rounded; article 2 usually with minute spiniform setae (Fig. 6A); outer flagellum with distal portion (distal to aesthetasc-bearing portion) usually well developed, consisting of numerous articles. Article 1 of mandibular palp with prominent expansion on inner distal margin (Fig. 6B). Maxilliped 2 with podobranch. Maxilliped 3 with or without exopod. Pereopod 1 fingers minute or completely reduced. Pereopod 2 subequal or unequal; basis with small process on lateral surface (Fig. 6C); carpal articulation greatly variable, but never tri-articulated. Arthrobranches usually present on maxilliped 3 and pereopods.

Chlorotocellidae fam. nov.

<http://zoobank.org/CBE45390-043A-46E8-9743-7F5B167255AD>

Type genus. *Chlorotocella* Balss, 1914, by present designation.

Composition. *Chlorotocella* Balss, 1914 (two species), *Chlorocurtis* Kemp, 1925 (monotypic), *Anachlorocurtis* Hayashi, 1975 (three species) and *Miropandalus* Bruce, 1983 (monotypic).

Diagnosis. Rostrum, if present, without teeth or fringe of setae on ventral margin (Figs 1A, 4A). Thoracic sternites without conspicuous ornamentation, such as prominences, teeth or protuberances (Fig. 3B). Pleomere 6 posterolateral process rounded or truncate (Figs 1C, 4B). Telson with dorsolateral spiniform setae located adjacent to lateral margins (Figs 1D, 4C). Eyestalks subcylindrical, cornea distinctly shorter than eyestalk (Figs 1E, 4D). Antennular stylocerite devoid of proximolateral projection, distally obliquely truncate, bi- or tridentate (Figs 1A, F, 4E); outer flagellum with distal portion (distal to aesthetasc-bearing portion) reduced, consisting only of few articles (Figs 1A, 4E). Maxilliped 2 without podobranch (Figs 1L). Maxilliped 3 with no exopod (Figs 2A, 5A). Pereopod 1 fingers absent (Figs 2B, 5B, C). Pereopod 2 always subequal; basis without small process on lateral surface of basis; carpus consistently divided into three articles (Figs 2C, 5D, 6D). Arthrobranches always absent from maxilliped 3 and pereopods.

Remarks. Characters differentiating Chlorotocellidae fam. nov. and Pandalidae are summarized in Table 1, with the character states of Chlorotocellidae fam. nov. being synapomorphic against Pandalidae (see Liao et al. 2019).




Figure 1. *Chlorotocella gracilis* Balss, 1914, male (cl 5.0 mm), CBM-ZC 12534. **A.** Carapace, antennule and antenna, lateral view (left eye removed); **B.** Anterior part of carapace, lateral view; **C.** Pleon, lateral view; **D.** Telson, dorsal view; **E.** Left eye, dorsal view; **F.** Left antennular peduncle, dorsal view; **G.** Left antenna, ventral view; **H.** Left mandible, outer view; inset, palp, outer view; **I.** Left maxillule, outer view (coxal endite missing); **J.** Left maxilla, outer view; **K.** Left maxilliped 1, outer view; **L.** Left maxilliped 2, outer view; **M.** Endopod of left pleopod 1, ventral view.




Figure 2. *Chlorotocella gracilis* Balss, 1914, male (cl 5.0 mm), CBM-ZC 12534. Left thoracic appendages in lateral view. **A.** Maxilliped 3; **B–F.** Pereopods 1–5, respectively.




Figure 3. Thoracic sternites 6–8 and coxae (cx) of pereopods 3–5, ventral view. **A.** *Chlorotocus novaezealandiae* (Borradaile, 1916), male (17.0 mm), NRIFS 578 (arrows indicating prominences on sternites 6–8); **B.** *Chlorotocella gracilis* Balss, 1914, male (cl 4.1 mm), HUMZ-C 1556.

Table 1. Comparison of diagnostic characters between Chlorotocellidae fam. nov. and Pandalidae. Character states of Chlorotocellidae are apomorphic against those of Pandalidae.

	Chlorotocellidae fam. nov.	Pandalidae
1	Rostrum, if present, without teeth or fringe of setae on ventral margin (Figs 1A, 4A).	Rostrum usually well developed, ventral margin armed with a few or series of teeth accompanied by row(s) of short setae.
2	Thoracic sternites without conspicuous ornamentation, such as keels, teeth or protuberances (Fig. 3B).	Thoracic sternites 6 and 7 each with paired prominences on either side of median keel; thoracic sternite 8 with transverse carina bearing submedian spines or teeth (Fig. 3A).
3	Pleomere 6 without posteroventral tooth; posterolateral process rounded or truncate (Figs 1C, 4B).	Pleomere 6 with small posteroventral tooth; posterolateral process terminating in small tooth.
4	Telson with dorsolateral spiniform setae located adjacent to lateral margins (Figs 1D, 4C).	Telson with dorsolateral spiniform setae located on dorsolateral ridges.
5	Eyestalks subcylindrical, cornea distinctly shorter than eyestalk (Figs 1E, 4D).	Eyestalks pyriform or subpyriform, cornea distinctly wider than eyestalk.
6	Antennular stylocerite distally obliquely truncate, bi- or tridentate, devoid of proximolateral projection (1F, 4E).	Antennular stylocerite acuminate or rounded, usually having small proximolateral projection (Fig. 6A).
7	Outer antennular flagellum with distal portion (distal to thickened aesthetasc-bearing portion) reduced, consisting of few articles (Figs 1A, 4E).	Outer antennular flagellum with distal portion (distal to aesthetasc-bearing portion) long and slender, consisting of 10 or more articles.
8	Maxilliped 2 without podobranch (Fig. 1L).	Maxilliped 2 with podobranch.
9	Maxilliped 3 without exopod (Figs 2A, 5A).	Maxilliped 3 with or without exopod.
10	Pereopod 1 fingers absent (Figs 2B, 5B, C).	Pereopod 1 fingers minute or completely reduced.
11	Pereopods 2 always subequal; basis without small process on lateral surface of basis; carpus consistently divided into 3 articles (Figs 2C, 5D, 6D).	Pereopods 2 subequal or unequal; basis without small process on lateral surface of basis; division of carpus highly variable, but never 3-articulated.
12	Arthrobranchs absent from maxilliped 3 and pereopods.	Arthrobranchs usually present on maxilliped 3 and pereopods 1–4.

Supplementary figures of diagnostic characters can be found in Hayashi (1975: figs 1–3), Bruce (1983: figs 1–5), Hayashi (2007a: figs 538, 539, 542a–f), Hayashi (2007c: figs 557–559a–e), Horká et al. (2014: figs 1–8), and Ahyong (2015: figs 9, 10).

Amongst these characters, the division of the carpus of pereopod 2 and quite possibly the absence of ventral rostral teeth can readily be used to differentiate the two families, although determination of their polarity is not straightforward. In Chlorotocellidae the pereopod 2 car-




Figure 4. *Chlorocurtis jactans* (Nobili, 1904), ovigerous female (cl 1.3 mm), CBM-ZC 11596. **A.** Carapace, lateral view; **B.** Pleon, lateral view; **C.** Telson, dorsal view; **D.** Left eye, dorsal view; **E.** Left antennule, dorsal view (inner flagellum damaged); **F.** Right antennular peduncle, distal 2 articles, mesial view; **G.** Left antenna, ventral view (flagellum missing); **H.** Left mandible, outer view; **I.** Left maxillule, outer view; **J.** Left maxilla, outer view; **K.** Left maxilliped 1, outer view; **L.** Left maxilliped 2, outer view (epipod broken off).




Figure 5. *Chlorocurtis jactans* (Nobili, 1904), ovigerous female (cl 1.3 mm), CBM-ZC 11596, left thoracic appendages in lateral view (except for C). **A.** Maxilliped 3; **B.** Pereopod 1; **C.** Same, propodus, mesial view; **D–G.** Pereopods 2–5, respectively.

pus is consistently divided into three articles, whereas in Pandalidae, the number of the carpal articles is quite variable according to taxa, but none are tri-articulate (cf. Komai 1994). The absence of ventral rostral teeth is also reported in three taxa of the pandalid genus *Plesionika* (Chace 1985) but with doubt (see Chace 1985; Komai et al. 2005; Hayashi 2009; Komai 2011; Li and Chan 2013). Such a similarity, if really present, can be resulted from homoplasy (Liao et al. 2019).

According to the Ancestral State Reconstruction (ASR) analysis by Liao et al. (2019), Pandalidae is characterized by the following synapomorphic features: (1) second article of the antennular peduncle with a few minute spiniform setae on the dorsodistal margin (Fig. 6A); (2) mandibular palp consisting of three articles (Fig. 6B); and (3) basis of pereopod 2 bearing a small process on the lateral surface (Fig. 6C). In these regards, Chlorotocellidae shows the following plesiomorphic states: (1) mandibular palp tends to be reduced, being absent or consisting of two articles at most (Figs 1H, 4H); (2)

second article of the antennular peduncle is unarmed on the dorsodistal margin (Figs 1A, F, 4E); and (3) basis of pereopod 2 being unarmed (Fig. 6D).

Nevertheless, an assessment of the polarity of the development of the mandibular palp is fraught with difficulty and heavily dependent on outgroup selection. In Caridea in general, however, a reduction of the mandibular palp is considered to be derived (e.g., Christoffersen 1987, 1989), as compared to the well-developed palp in Dendrobranchiata and most other Decapoda.

Furthermore, the other two characters are subject to reversal within Pandalidae (Komai 1994; Liao et al. 2019). In species of *Thalassocaris* and *Chlorotocoides* (previously in their own family Thalassocarididae, but now considered part of Pandalidae), the second article of the antennular peduncle is devoid of spiniform setae and the basis of the second pereopods unarmed (Komai 1994).

In addition to the three aforementioned characters, the possession of a rounded laminar expansion at the inner distal angle of the first article of the mandibular palp




Figure 6. **A, B.** *Pandalus montagui* Leach, 1814, transitional male (cl 10.5 mm), CBM-ZC 3422; **C.** *Heterocarpus ensifer* A. Milne-Edwards, 1881, male (cl 33.0 mm), HUMZ-C 255; **D.** *Chlorotocella gracilis* Balss, 1914, male (cl 4.1 mm), HUMZ-C 1556. **A.** Left antennular peduncle, dorsal view; **B.** Left mandible, outer view; inset, palp, lateral view; **C.** Basis of left pereopod 2; **D.** Basis of right pereopod 2.

(Fig. 6B, inset) might be synapomorphic to Pandalidae (Komai 1994; Liao et al. 2019), although a secondary loss of this structure is observed in *Thalassocaris* and *Chlorotocoides* (Komai 1994; Liao et al. 2019). It is impossible to evaluate the homology of this character for the taxa assigned to Chlorotocellidae, because in those taxa, the mandibular palp only comprises two articles (*Chlorotocella*) or is absent (*Chlorocurtis*, *Anachlorocurtis* and *Miopandalus*), and the homology of the articles has not been established in taxa with different numbers.

Genus *Chlorotocella* Balss, 1914

Chlorotocella Balss 1914: 33; Holthuis 1955: 118, 127; 1993: 263, 266; Hayashi 2007a: 150.

Type species. *Chlorotocella gracilis* Balss, 1914.

Diagnosis. Rostrum elongate, very slender, gently upturned, exceeding far beyond distal margin of antennal scaphocerite, dorsally armed with two teeth around rostral base (one postrostral); ventral margin unarmed (Fig. 1A). Carapace without projections on dorsal midline; **supraorbital tooth present; suborbital lobe prominent, longer than antennal tooth, distally rounded, slightly constricted at base;** pterygostomial tooth moderately small (Fig. 1A, B). Pleomeres 1–6 dorsally rounded; **pleomeres 4 and 5 each with pair of posterolateral teeth; pleomere 5 with deep transverse groove near posterodorsal margin;** pleuron with small posteroventral

tooth (Fig. 1C). **Pleomere 6 with minute postero-medial tooth;** posteroventral angle with minute tooth (Fig. 1C). **Telson with additional anterior pair of spiniform setae located more mesial to other lateral series of spiniform setae;** posterior margin narrow, slightly produced medially, with two pairs of unequal spiniform setae (Fig. 1D). Eye with ocellar spot (nebenauge) (Fig. 1E). **Antennular peduncle article 1 armed with tooth on dorsodistal margin** (Fig. 1A, F). Mandible with two-articulated palp (Fig. 1H). **Maxillule palp without distal outer lobule** (Fig. 1I). Maxilla with short, moderately slender endopod (Fig. 1J). Maxilliped 1 with coxal and basal endites well developed, both with row of setae on mesial margin; exopodal flagellum well developed (Fig. 1K). Maxilliped 2 endopod with dactylus located at distal portion of propodus; exopod well developed (Fig. 1L). Pereopod 1 fingers completely reduced (Fig. 2B). Pereopods 3–5 propodi each with closely spaced, short to long spiniform setae in distal 0.2; carpi each with few spiniform setae on lateral surface; meri usually with spiniform setae arranged in two rows; ischia each with spiniform seta on ventral surface in pereopods 3 and 4 (Fig. 2D–F). Male pleopod 1 endopod without appendix interna (Fig. 1M).

Composition. *Chlorotocella gracilis*; *C. spinicaudus* (H. Milne Edwards, 1837).

Distribution. Indo-West Pacific, South Australia; shallow subtidal to 60 m; free living in algal-rich habitats or facultatively associated with gorgonarians and hydroids.

Remarks. At present, two species are assigned to *Chlorotocella* (De Grave and Fransen 2011), viz., *C. gracilis* (type species) and *C. spinicaudus*. Holthuis (1995) clarified that *Hippolyte spinicaudus* H. Milne Edwards, 1837 was a senior subjective synonym of *Pandalus leptorhynchus* Stimpson, 1860. In addition, a third taxon, which was placed in the synonymy of *C. spinicaudus* by De Grave and Fransen (2011), *Pandalus (Parapandalus) leptorhynchus* var. *gibber* Hale, 1924, was described from Gulf St Vincent, South Australia, characterized mainly by the prominently crested tergite of pleomere 3 (see Hale 1927). This taxon has been seldom mentioned in more recent literature. Ledoyer (1984) illustrated a specimen with a weakly crested tergite from Nouméa (New Caledonia), which he assigned to *C. gracilis*, but left it open as to whether this should be a distinct species or merely a “forme *gibber*” of *C. gracilis*. In contrast, Poore (2004) treated the taxon as a distinct species, *C. gibber* (Hale), noting it was restricted to the Gulf St Vincent (South Australia).

Because no modern descriptions are available for *C. spinicaudus*, the above generic diagnosis is largely based on *C. gracilis* and the summary information available on the other species. It seems possible that Hale’s (1924) taxon might be distinct from *C. gracilis* and *C. spinicaudus* as it is characteristic by having a highly crested tergite of the pleomere 3 (Hale 1924: pl. 4, fig. 6; 1927: fig. 35). Re-assessment of the taxonomic status of *C. spinicaudus* and *Pandalus (Parapandalus) leptorhynchus* var. *gibber* will be necessary to fully clarify the taxonomy of the genus.

Genus *Chlorocurtis* Kemp, 1925

Chlorocurtis Kemp 1925: 272, 279; Holthuis 1955: 118, 127; 1993: 263, 265; Hayashi 2007b: 248.

Type species. *Chlorocurtis miser* Kemp, 1925.

Diagnosis. Rostrum short but well developed, directed forward, reaching midlength of article 1 of antennular peduncle; **dorsal margin crested**, with five to seven teeth including two or three postrostral; ventral margin unarmed (Fig. 4A). Carapace without conspicuous projections on dorsal midline; no supraorbital tooth; suborbital lobe absent; pterygostomial tooth moderately small (Fig. 4A). Pleomeres 1–5 dorsally rounded; **pleomeres 1–3 with long, erect setae on dorsal surface**; pleomeres 4 and 5 each without pair of posterolateral teeth; pleomere 5 without deep transverse groove near posterodorsal margin, pleuron rounded posteriorly; pleomere 6 without posteromedian tooth, posteroventral angle unarmed (Fig. 4B). Telson posterior margin rather broad, convex, with three pairs of unequal spiniform setae (Fig. 4C). Eye without ocellar spot (nebenauge) (Fig. 4D). Antennular peduncle article 1 unarmed on dorsodistal margin; stylocerite obliquely truncate distally, distolateral angle terminating in tooth, distomesial angle subacute or blunt; outer

flagellum shorter than peduncle, distal portion reduced to single article (Fig. 4E). **Short, club-like, modified setae present at ventrodistal margin of article 2 of antennular peduncle (one seta) and distal margin of antennal scaphocerite (two setae)** (Fig. 4F, G.). Mandible without palp (Fig. 4H). Maxillule palp with well-developed distal outer lobule (Fig. 4I). Maxilla with short, distally tapering endopod (Fig. 4J). Maxilliped 1 with coxal and basial endites well developed, both with row of setae on mesial margin; exopodal flagellum well developed (Fig. 4K). Maxilliped 2 endopod with dactylus located at distal portion of propodus; exopod well developed (Fig. 4L). **Pereopods 3–5 propodi broadened distally, oblique flexor distal margins each with short rows of narrowly spaced long spiniform setae flanking field of short setae, forming prehensile structure together with dactylus folded back**; carpi without spiniform setae on lateral surface; meri without spiniform setae; ischia without spiniform seta on ventral surface (Fig. 5E–G). Male pleopod 1 endopod without appendix interna.

Composition. Monotypic.

Distribution. Indo-West Pacific, intertidal to 10 m; seagrass beds.

Remarks. *Chlorocurtis* was originally established for *Chlorocurtis miser* by Kemp (1925).

Later, Holthuis (1947) synonymized *Chlorocurtis miser* with *Virbius (?) jactans* (Nobili, 1904) without any argumentation, although clearly correct. This synonymy has since been widely adopted (e.g., Holthuis 1955; Ledoyer 1968, 1984; Bruce 1976; Hayashi 2007b; Holthuis 1993; De Grave and Fransen 2011; Gan and Li 2018).

Genus *Anachlorocurtis* Hayashi, 1975

Anachlorocurtis Hayashi 1975: 173; 2007a; 147; Holthuis 1993: 263; Horká et al. 2014: 12.

Type species. *Anachlorocurtis commensalis* Hayashi, 1975.

Diagnosis. Rostrum short, ascending in adults, reaching midlength of article 1 of antennular peduncle, terminating in acute tip or obliquely truncate distally with irregular dentition; dorsal and ventral margins usually unarmed. Carapace without supraorbital tooth; dorsal midline with two prominent processes, anterior one postrostral, irregularly denticulate anteriorly, posterior one cardiac in position, directed forward, acuminate; suborbital lobe absent; pterygostomial angle rounded, unarmed. Pleomeres 1–5 dorsally rounded; pleomeres 4 and 5 each without pair of posterolateral teeth. Pleomere 5 without deep transverse groove near posterodorsal margin; pleuron rounded posteriorly. Pleomere 6 without posteromedian tooth; posteroventral angle without tooth. Telson posterior margin truncate or rounded, with five pairs of unequal spiniform

setae. Eye without ocellar spot (nebenauge); **cornea with papilla-like tubercle**. Antennular peduncle article 1 unarmed on dorsodistal margin; stylocerite obliquely truncate distally, both distal angles dentate; outer flagellum shorter than peduncle, distal portion reduced to one or two articles. Mandible without palp. Maxillule palp with well-developed distal outer lobule bearing apical seta. Maxilliped 1 with coxal and basial endites poorly developed, narrow; exopodal flagellum absent. Maxilliped 2 endopod with dactylus located at mesial portion of propodus or fused to propodus; exopod absent. Articulation between carpal article 1 and 2 of pereopod 2 strongly oblique. Pereopods 3–5 propodi slightly narrowing distally, with few widely spaced minute spiniform setae on flexor margin; carpi without spiniform setae on lateral surface; meri of pereopods 3 and 4 each with one spiniform seta distolaterally and one minute spiniform seta at midlength of ventral surface. Male pleopod 1 endopod with small rounded lateral lobe far exceeded by well-developed appendix interna.

For illustrations see Hayashi (1975: figs 1–3), Hayashi (2007a: figs 538, 539, 542a–f); Horká et al. (2014: figs 1–8) and Ah Yong (2015: figs 9, 10).

Composition. *Anachlorocurtis commensalis*, *A. occidentalis* Horká, De Grave & Ďuriš, 2014, and *A. australis* Ah Yong, 2015.

Distribution. Indo-West Pacific, shallow subtidal to 40 m; associated with antipatharian corals.

Genus *Miopandalus* Bruce, 1983

Miopandalus Bruce, 1983: 482; Holthuis 1993: 263, 269; Hayashi 2007c: 585.

Type species. *Miopandalus hardingi* Bruce, 1983.

Diagnosis. **Rostrum absent.** Carapace without supra-orbital tooth; **dorsal midline with two very prominent, erect processes, anterior one postrostral, tapering, posterior one cardiac in position, slightly curved, anteriorly, blunt**, suborbital lobe absent; pterygostomial angle rounded or angular. **Pleomere 1 with prominent protuberance; pleomere 3 with triangular crest on posterior half of dorsal midline**; pleomeres 4 and 5 each without pair of posterolateral teeth; pleomere 5 without deep transverse groove near posterodorsal margin; pleuron rounded posteriorly. Pleomere 6 without posteromedian tooth; posteroventral angle unarmed. Telson posterior margin rounded, with several short spiniform setae. Eye without ocellar spot (nebenauge); cornea without papilla-like tubercle. Antennular peduncle article 1 unarmed on dorsodistal margin; **stylocerite subtruncate distally, bi- or tridentate**; outer flagellum shorter than peduncle, distal portion completely reduced. Mandible without palp. Maxillule palp with well-developed distal outer lobule, without apical seta. Maxilliped 1 with coxal and

basial endites poorly developed, narrow; endopod stout; exopodal flagellum absent. Maxilliped 2 endopod with dactylus fused to propodus; exopod absent. Articulation between carpal article 1 and 2 of pereopod 2 strongly oblique. Pereopods 3–5 propodi narrowing distally, with few minute spiniform setae on flexor margin; carpi without spiniform setae on lateral surface; meri of pereopods 3 and 4 unarmed. Male pleopod 1 endopod with small rounded lateral lobe far exceeded by well-developed appendix interna.

For illustrations, see Bruce (1983: figs 1–5) and Hayashi (2007c: figs 557–559a–e).

Composition. Monotypic.

Distribution. West Pacific, subtidal to 58 m; associated with antipatharian corals.

Acknowledgements

This work was supported by grants from the Ministry of Science and Technology, Taiwan, R.O.C., and the Center of Excellence for the Oceans (National Taiwan Ocean University), which is financially supported by The Featured Areas Research Center Program within the framework of the Higher Education Sprout Project by the Ministry of Education in Taiwan, R.O.C.

References

- Ah Yong ST (2015) Decapod Crustacea of the Kermadec Biodiscovery Expedition 2011. *Bulletin of the Auckland Museum* 20: 405–442.
- Anker A, De Grave S (2016) An updated and annotated checklist of marine and brackish caridean shrimps of Singapore. *Raffles Bulletin of Zoology* 34 (Supplement): 343–454.
- Balss H (1914) Ostasiatische Decapoden II. Die Natantia und Reptantia. *Abhandlungen der Mathematisch-Physikalischen Klasse der Königlich Bayerischen Akademie der Wissenschaften* 10 (Supplement 2): 1–101. [pl. 1]
- Bate SC (1888) Report on the Crustacea Macrura collected by the Challenger during the years 1873–76. Report on the Scientific Results of the Voyage of H.M.S. “Challenger” during the Years 1873–1876 24: 1–942. [pls 1–157]
- Borradaile LA (1916) Crustacea. Part I. Decapoda. *British Antarctic (“Terra Nova”) Expedition, 1910. Natural History Report, Zoology* 3: 75–110.
- Bruce AJ (1976) A report on a small collection of shrimps from the Kenya National Marine Parks at Malindi, with notes on selected species. *Zoologische Verhandlungen* 145(1): 1–72.
- Bruce AJ (1983) *Miopandalus hardingi*, new genus, new species, a bizarre commensal pandalid shrimp from the Marshall Islands. *Journal of Crustacean Biology* 3: 482–490. <https://doi.org/10.2307/1548149>
- Caullery M (1896) Résultats scientifiques de la Campagne du “Caudan” dans le Golfe de Gascogne. Aout-Septembre 1885. *Crustacés schizopodes et décapodes. Annales de l’Université de Lyon* 2: 365–419.

- Calman WT (1899) On the British Pandalidae. The Annals and Magazine of Natural History (Series 7) 3: 27–39. <https://doi.org/10.1080/00222939908678072>
- Chace Jr FA (1985) The caridean shrimps (Crustacea: Decapoda) of the *Albatross* Philippine Expedition, 1907–1910, Part 3: families Thalassocarididae and Pandalidae. Smithsonian Contributions to Zoology 411: 1–143. <https://doi.org/10.5479/si.00810282.411>
- Christoffersen ML (1987) Phylogenetic relationships of hippolytid genera, with an assignment of new families for the Crangonoidea and Alpheoidea (Crustacea, Decapoda, Caridea). Cladistics 3(4): 348–362. <https://doi.org/10.1111/j.1096-0031.1987.tb00898.x>
- Christoffersen ML (1989) Phylogeny and classification of the Pandaloidae (Crustacea, Caridea). Cladistics 5: 259–274. <https://doi.org/10.1111/j.1096-0031.1989.tb00489.x>
- Crosnier A (1997) Crustacea Decapoda: *Pseudopandalus curvirostris*, genre et espèce nouveaux (Pandalidae) de Nouvelle-Calédonie. In: Crosnier A (Ed.) Résultats des Campagnes MUSORSTOM, vol. 18. Mémoires du Muséum National d'Histoire Naturelle 176: 169–176.
- De Grave S, Fransen CHJM (2011) Carideorum catalogus: the recent species of the dendrobranchiate, stenopodidean, procarididean and caridean shrimps (Crustacea: Decapoda). Zoologische Mededelingen 85: 195–589.
- de Man JG (1917) Diagnoses of new species of macrurous decapod Crustacea from the Siboga-Expedition. Zoologische Mededelingen 3: 279–284.
- Fransen CHJM (1990). *Bitias stocki*, a new genus and new species of pandalid shrimp (Crustacea, Decapoda, Caridea) in the eastern Atlantic Ocean. Beaufortia 41: 67–73.
- Fransen CHJM (1997) Crustacea Decapoda: *Chelonika macrochela*, a new genus and new species of pandalid shrimp (Caridea) from new Caledonian waters. In: Crosnier A (Ed.) Résultats des Campagnes MUSORSTOM, vol. 18. Mémoires du Muséum National d'Histoire Naturelle 176: 177–185.
- Gan Z, Li X (2018) Four new records of caridean shrimp (Crustacea: Decapoda: Caridea) from the East China Sea and South China Sea. Acta Oceanologica Sinica 37 (10): 212–217. <https://doi.org/10.1007/s13131-018-1325-y>
- Hale HM (1924) The flora and fauna of Nuys Archipelago and the Investigator group. No. 16. The Crustacea. Transactions and Proceedings of the Royal Society of South Australia 48: 67–73. [pls 4, 5]
- Hale HM (1927) The Crustaceans of South Australia. Part 1. South Australian Government Printer, Adelaide, 201 pp.
- Haworth AH (1825) A new binary arrangement of the macrurous Crustacea. The Philosophical Magazine and Journal 65: 183–184. <https://doi.org/10.1080/14786442508628417>
- Hayashi K (1975) *Anachlorocurtis commensalis* gen. nov., sp. nov. (Crustacea, Decapoda, Pandalidae), a new pandalid shrimp associated with antipatharian corals from central Japan. Annotationes Zoologicae Japonenses 48: 172–182.
- Hayashi K (2007a) Prawns, shrimps and lobsters from Japan (152). Family Pandalidae – genera *Anaculorocurtis* and *Chlorotocella*. Aquabiology 169: 147–153. [in Japanese]
- Hayashi K (2007b) Prawns, shrimps and lobsters from Japan (153). Family Pandalidae – genera *Chlorotocella*, *Chlorocurtis* and *Heterocarpus*. Aquabiology 170: 248–255. [in Japanese]
- Hayashi K (2007c) Prawns, shrimps and lobsters from Japan (156). Family Pandalidae – genera *Miopandalus* & *Pandalopsis* 1. Aquabiology 173: 585–590. [in Japanese]
- Hayashi K (2009) Prawns, shrimps and lobsters from Japan (166). Family Pandalidae – genus *Plesionika* (4). Aquabiology 183: 425–430 [in Japanese].
- Hayashi K, Miyake S (1968) Three caridean shrimps associated with a medusa from Tanabe Bay, Japan. Publications of the Seto Marine Biological Laboratory 16(1): 11–19. <https://doi.org/10.5134/175493>
- Hendrickx ME (2019) Redescription of the rare shrimp *Heterocarpus nesisi* (Burukovsky, 1986) (Crustacea: Caridea: Pandalidae) rediscovered off western Mexico, with the proposal of a new genus. Zootaxa 4565: 49–60. <https://doi.org/10.11646/zootaxa.4565.1.3>
- Holthuis LB (1947) The Decapoda of the Siboga Expedition. Part IX. The Hippolytidae and Rhynchocinetidae collected by the Siboga and Snellius expeditions with remarks on other species. Siboga Expeditie 39a(8): 1–100.
- Holthuis LB (1950) The Decapoda of the Siboga Expedition. Part X. The Palaemonidae collected by the Siboga and Snellius expeditions with remarks on other species. I. Subfamily Palaemoninae. Siboga Expeditie 39a(9): 1–268.
- Holthuis LB (1955) The recent genera of the caridean and stenopodidean shrimps (Class Crustacea, Order Decapoda, Supersection Natantia) with keys for their determination. Zoologische Verhandlungen 26: 1–157.
- Holthuis LB (1980) FAO Species Catalogue. Vol. 1. Shrimps and prawns of the world: an annotated catalogue of species of interest to fisheries. FAO Fisheries Synopsis 125 (1): 1–271.
- Holthuis LB (1993) The Recent Genera of the Caridean and Stenopodidean Shrimps (Crustacea, Decapoda) with an Appendix on the Order Amphionidacea. Nationaal Natuurhistorisch Museum, Leiden, 328 pp. <https://doi.org/10.5962/bhl.title.152891>
- Holthuis LB (1995) Notes on Indo-Pacific Crustacea III to IX. Zoologische Mededelingen 69(13): 139–151.
- Horká I, De Grave S, Ďuriš Z (2014) A new species of shrimp of the genus *Anachlorocurtis* Hayashi, 1975 from the Red Sea, with range extension of *A. commensalis* Hayashi, 1975 (Crustacea, Decapoda, Pandalidae). ZooKeys 407: 9–28. <https://doi.org/10.3897/zookeys.407.7457>
- Humann P, DeLoach N (2010) Reef Creatures Identification Tropical Pacific. New World Publications, Jacksonville, Florida, 497 pp.
- Kato S, Okuno J (2001) Shrimps and Crabs of Hachijo Island. TBS Britannica, Tokyo, 158 pp. [in Japanese]
- Kawamoto T, Okuno J (2003) Shrimps and Crabs of Kume Island, Okinawa. Hankyu Communications, Tokyo, 174 pp. [in Japanese]
- Kemp S (1925) Notes on Crustacea Decapoda in the Indian Museum. XVII On various Caridea. Records of the Indian Museum 27: 249–342.
- Komai T (1994) Phylogeny and cladistic classification of the superfamily Pandaloidae (Crustacea: Decapoda: Caridea). Unpublished doctoral thesis, Hokkaido University.
- Komai T (1999) A revision of the genus *Pandalus* (Crustacea: Decapoda: Caridea: Pandalidae). Journal of Natural History 33: 1265–1372. <https://doi.org/10.1080/002229399299914>
- Komai T (2011) Deep-sea shrimps and lobsters (Crustacea: Decapoda: Dendrobranchiata and Pleocyemata) from the Sagami Sea and Izu Islands, Central Japan. Memoirs of the National Museum of Natural Science 47: 279–337.
- Komai T, Chan T-Y, Hanamura Y, Abe Y (2005) First record of the deep-water shrimp, *Plesionika williamsi* Forest, 1964 (Decapoda, Caridea, Pandalidae) from Japan and Taiwan. Crustaceana 78: 1001–1012. <https://doi.org/10.1163/156854005775197307>

- Leach WE (1813–1814) Crustaceology. In: Brewster D (Ed.) The Edinburgh Encyclopædia. A. Balfour, Edinburgh, 383–437.
- Ledoyer M (1968) Les Caridea de la frondaison des herbiers de phanérogames de la région de Tuléar (République Malgache). Recueil des Travaux de la Station Marine d'Endoume 8: 63–123.
- Ledoyer M (1984) Les Caridea (Crustacea: Decapoda) des herbiers de phanérogames marines de Nouvelle-Calédonie (région de Nouméa). Zoologische Verhandlungen 211: 1–58.
- Li X, Chan T-Y (2013) Pandalid shrimps (Crustacea, Decapoda, Caridea) collected from the Philippines PANGLAO 2005 deep-sea expedition. In: Ahyong ST, Chan T-Y, Corbari L, Ng PKL (Eds) Tropical Deep-Sea Benthos Vol. 27. Mémoires du Muséum national d'Histoire naturelle 204: 129–154.
- Liao Y, Ma K-Y, De Grave S, Komai T, Chan T-Y, Chu K-H (2019) Systematic analysis of the caridean superfamily Pandaloidea (Crustacea: Decapoda) based on molecular and morphological evidence. Molecular Phylogenetics and Evolution 134: 200–210. <https://doi.org/10.1016/j.ympev.2019.02.006>
- Milne Edwards H (1837) Histoire naturelle des crustacés: comprennent l'anatomie, la physiologie et la classification de ces animaux. Librairie Encyclopédique de Roret, Paris, 531 pp. <https://doi.org/10.5962/bhl.title.16170>
- Milne-Edwards A (1881) Description de quelques crustacés macroures provenant des grandes profondeurs de la Mer des Antilles. Annales des Sciences Naturelles (Série 6) 11: 1–15.
- Milne-Edwards A (1882) Rapport sur les travaux de la commission charge par M. le Ministre de l'Instruction Publique d'étudier la faune sous-marine dans les grandes profondeurs de la Méditerranée et de l'Océan Atlantique. Archives Des Missions Scientifiques et Littéraires (Série 3) 9: 1–56.
- Milne-Edwards A (1883) Recueil de figures de crustacés nouveaux ou peu connus. Paris, 3 pp. [44 pls.]
- Minemizu R (2000) Marine Decapod and Stomatopod Crustaceans Mainly from Japan. Bun-ichi, Tokyo, 344 pp. [in Japanese]
- Minemizu R (2013) Coral Reef Shrimps of Indo-West Pacific. Bun-ichi, Tokyo, 144 pp. [in Japanese]
- Nobili G (1904) Diagnoses préliminaires de vingt-huit espèces nouvelles de stomatopodes et Decapodes Macroures de la mer Rouge. Bulletin du Muséum Histoire Naturelle 10: 228–238.
- Okuno J, Yokota M (1995) Record of a bizarre shrimp, *Miopandalus hardingi* Bruce, 1983 from Izu Peninsula, Central Japan. I.O.P. Diving News: 7 (1): 2–3. [in Japanese with English abstract]
- Poore GCB (2004) Marine Decapod Crustacea of Southern Australia. A Guide to Identification. CSIRO Publishing, Collingwood, Victoria, 574 pp. <https://doi.org/10.1071/9780643092129>
- Stimpson W (1860) Prodrômus descriptionis animalium evertetorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladore Ringgold et Johanne Rodgers Ducibus, observavit et descripsit. Pars VIII, Crustacea Macrura. Proceedings of the Academy of Natural Sciences of Philadelphia 1860: 22–47.
- Yaldwyn JC (1960) Biological results of the Chatham Islands 1954 Expedition. Decapoda Natantia. New Zealand Department of Scientific and Industrial Research Bulletin 139: 13–53.



Komai, Tomoyuki, Chan, Tin-Yam, and De Grave, Sammy. 2019.
"Establishment of a new shrimp family Chlorotocellidae for four genera previously assigned to Pandalidae (Decapoda, Caridea, Pandalioidea)."
Zoosystematics and evolution 95(2), 391–402.
<https://doi.org/10.3897/zse.95.35999>.

View This Item Online: <https://www.biodiversitylibrary.org/item/276109>

DOI: <https://doi.org/10.3897/zse.95.35999>

Permalink: <https://www.biodiversitylibrary.org/partpdf/291842>

Holding Institution

Museum für Naturkunde, Berlin

Sponsored by

Museum für Naturkunde, Berlin

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Copyright held by individual article author(s).

License: <https://creativecommons.org/licenses/by/4.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.