

THE FOSSIL INSECTS OF AUSTRALIA

P.A. JELL

Jell, P.A. 2004 16 08: The fossil insects of Australia. *Memoirs of the Queensland Museum* 50(1): 1-124. Brisbane. ISSN0079-8835.

A taxonomic catalogue of the known fossil insects of Australia provides illustrations of at least one species of every genus so far identified. Every known species is recorded with the age, rock unit (formation) and location (sedimentary basin). Some synonymies and updated taxonomic placements are included but this has not been an exhaustive review of taxonomy. The principal sources for fossil insects are described as the Upper Permian Boolaroo Formation in the Newcastle Coal Measures adjacent to Lake Macquarie, the Hawkesbury Sandstone at Brookvale in Sydney, the Mt Crosby and Blackstone Formations in the Upper Triassic Ipswich Coal Measures of southeastern Queensland and the Lower Cretaceous Koonwarra Fossil Bed in South Gippsland, Victoria. Other minor sites are discussed and potential for further research is outlined. *Austroscarites* is introduced to replace the preoccupied *Scarites* Hamilton not Fabricius. □ *Fossil insects, Australia, Permian, Triassic, Cretaceous, catalogue.*

Peter A. Jell, Queensland Museum, PO Box 3300, South Brisbane, Queensland 4101, Australia; 1 August 2004.

The living insect fauna of Australia is obviously abundant and diverse to inhabitants and visitors alike. The importance of the insect fauna to our way of life and our economy has been documented by many people since the first arrival of Europeans. Assuming living diversity is an indication of the insect fauna at any instant of the geological past and in view of the known evolution of insects during geological time we can project an enormous number of species that must have inhabited this continent since the origin of insects during the early Palaeozoic. However, the majority of these were almost certainly never fossilised and have left no record. Marine faunas have a better fossilization potential than terrestrial ones. It is estimated that overall less than 1% of living species at any one instant in time will be fossilized and collectable to enter the palaeontological literature. Thus the known fossil insect record of more than 400 species in approximately 250 genera is only a small fraction of the insect species that have lived in Australia.

In each edition of 'The Insects of Australia' a chapter was devoted to their fossil history. Riek (1970c) and Kukalova-Peck (1991) gave good overviews of the fossil history of Australia and fitted it into the broader story of insect evolution across the globe. The aim of this publication is to complement those accounts with a comprehensive listing of the fossil insect species described from Australia along with illustration of at least one species from each genus.

GEOLOGICAL SETTING OF FOSSIL INSECT SITES

Compared to the number of Australian fossil sites or to the number of insect fossil sites worldwide only a few sites around Australia have yielded fossils of insects (Fig. 1). These sites provide imperfect data for only a very few instants of geological history (Fig. 2).

HELLYER GORGE SITE IN UPPER CARBONIFEROUS WYNYARD TILLITE.

The earliest known insect from Australia comes from the Hellyer Gorge district of northwestern Tasmania where the single, almost complete specimen, with 50mm wing span, was found in a 1m fossiliferous band of varve-like sediments within the Wynyard Tillite. Tillites occur in the formation both above and below the finegrained fossiliferous section. The co-occurring fossil plant, *Botrychiopsis plantiana*, and Stage 1 microflora indicate a Late Carboniferous age and the primitive insect structure suggests early Late Carboniferous. The sedimentary environment and paucity of associated insects indicate deposition in cold conditions close to glacial ice. Riek (1976) erected the Psychroptilidae and Suborder Neosecoptera as monotypic taxa for this insect, *Psychroptilus burrettiae* Riek, 1976 within the Order Megasecoptera. He outlined the difficulty of classifying this species by indicating features it shares with the Megasecoptera (wing venation except Sc ends abruptly on the costal margin just beyond the midlength of the wing) others it shares with the Palaeodictyoptera (wings




FIG. 1. Distribution of main fossil insect sites. 1=UCarboniferous Wynyard Tillite at Hellyer Gorge; 2=UPermian Newcastle Coal Measures at Belmont just south of Newcastle; 3=MTriassic (Anisian) Brookvale Shale in Hawkesbury Sandstone near Manly in suburbs of Sydney; 4=MTriassic (Anisian) Ashfield Formation in Wianamatta Group at St Peters near Newtown, suburban Sydney; 5=UTriassic (Carnian) Mt Crosby Formation in lower Ipswich Coal Measures just north of Ipswich; 6=UTriassic (Carnian) Blackstone Formation at top of Ipswich Coal Measures on Denmark Hill in suburban Ipswich; 7=LJurassic Talbragar Fish Bed in the Purlawaugh Formation near Gulgong; 8=LCretaceous Cockleshell Gully Formation in Hill River area, northern Perth Basin; 9=LCretaceous (Aptian) Koonwarra Fossil Bed on the South Gippsland Highway near Tarwin in the Gippsland Basin; 10=LCretaceous Wallumbilla Fm near Richmond, Great Artesian Basin; 11=UCretaceous Winton Formation near Winton, Great Artesian Basin; 12=Palaeocene Redbank Plains Formation east of Ipswich; 13=Late Oligocene/Early Miocene Upper Site in Riversleigh Limestones north of Mount Isa.

broad at the base) and others not previously known in either of those orders. By comparison to the Northern Hemisphere faunas the Carboniferous insect fauna of Australia is virtually unknown.

BELMONT INSECT FAUNA IN UPPER PERMIAN NEWCASTLE COAL MEASURES. The Upper Permian insects from the Newcastle Coal Measures described mainly by Tillyard came from an area of 10-20km² between the town of Belmont and Warners Bay on the eastern side

of Lake Macquarie some 10km southwest of Newcastle. A sand and conglomerate fan was probably braiding through coal swamps along the southeastern flank of the Hunter Trough in the Late Permian, sourcing its clastic material primarily from higher land to the north and northwest (Engel, 1966). The widespread and persistent insect beds, which are fine grained volcanoclastic units (tuffs) associated with plenty of plant detritus mainly *Glossopteris* leaves and *Phyllothea*, were apparently deposited during a brief, episodic event and fairly rapidly buried.

The insects were very likely a wing-float concentration, with the wings remaining buoyant until the waterbody was choked with ash, possibly enhanced by storm action during or just after the eruption. Adjacent coal seams, conglomerates and tuff beds indicate a dynamic environment with swamp conditions prevailing when and where short term quiet conditions became established. Details of stratigraphic relationships in the Belmont area are provided by Hawley & Brunton (1995) and Little et al. (1996). Rare insects have been found in the Lambton Formation adjacent to the Dudley Coal Seam (*Permoscarta mitchelli* Tillyard, 1918b, *Mitchelloneura permiana* Tillyard, 1921b and *Lophioneura ustulata* Tillyard, 1921b) at the base of the Newcastle Coal Measures but the major horizon is within the Boolaroo Formation in the upper half of the coal measures. As with many coal measure stratigraphies there is an extremely fine subdivision to reflect the various coal seams but these units are often of restricted extent so the listing below refers to the Newcastle Coal Measures as the provenance of the insects from this area. This fauna is rich in Homoptera, Psocoptera, Neuroptera, Mecoptera but lacks blattoids or palaeoptera.

MIDDLE TRIASSIC HAWKESBURY SANDSTONE AT BROOKVALE, SYDNEY. The Middle Triassic (Anisian) Hawkesbury Sandstone in the Sydney Basin envelops a lens of fine mudstone referred to as the Brookvale Shale lens (about 8m thick) which was best exposed in the Brookvale Quarry on Beacon Hill, Brookvale, near Manly, north of Sydney Harbour where it yielded many fish, many plants (*Dicroidium zuberi* flora), amphibians, unionid bivalves, crustaceans, insects, trace fossils and some enigmatic fossils. This mudstone lens may be interpreted as a lacustrine interval on a quartz sandstone coastal plain and from the fauna a freshwater environment is most probable. The insect fauna includes the large titanopteran, *Clatrotitan*, a mecopteroid, a protorthopteroid, orthopteroids and a stonefly.

MIDDLE TRIASSIC ASHFIELD FORMATION AT ST PETERS, SYDNEY. The Ashfield Shale (40-60m thick in the Wianamatta Group) overlies the Mittagong Formation (up to 15m thick) which in turn overlies the Hawkesbury Sandstone in the area immediately west and southwest of Sydney known as the Cumberland Basin. Outcrop of the Ashfield Shale defines the shape of the basin. Siltstone and




FIG. 2. Stratigraphic succession of important known insect sites.

clay from low in the formation was extensively quarried in the St Peters district for manufacture of house bricks. An extensive fauna of bivalves, fish, a shark, labyrinthodonts, an isopod, the insects and plants has been collected from these quarries and indicates a freshwater environment. Its age is Middle Triassic (late Anisian) based mainly on the floral components of the fauna. The insects are predominantly beetle elytra, a large blattodea and the large titanopteran, *Mesotitan*,

but nothing has been added to the work of Tillyard (1916). Tillyard (1918b) described another Ashfield insect fauna from a railway cutting at Glenlee, 6.5km south of Campbelltown on the main southern line with a very similar assemblage to that from St Peters i.e. dominated by beetle elytra, the same blattodean genus and a mecopteroid rather than the large titanoptera.

UPPER TRIASSIC MT CROSBY FORMATION, IPSWICH COAL MEASURES AT MT CROSBY, NORTH OF IPSWICH, SOUTH-EASTERN QUEENSLAND. The Mount Crosby Formation, the oldest in the Ipswich Coal Measures occupies an area north of Ipswich adjacent to the Brisbane River of some 30km². It is 20-30m thick and consists mainly of polymictic, poorly sorted conglomerate but has 3 shale units, up to 6m thick near the bottom, middle and top of the unit. The upper two shale units are fossiliferous with prolific plant and insect remains. The conglomerates of the formation indicate rapid erosion from an adjacent highland (the Daguiar Block) and little transport. The shale units indicate restricted quiet lacustrine conditions at various times on this piedmont fan. The insect fauna is dominated by Blattodea and Homoptera with Mecoptera, Diptera, Orthoptera, Neuroptera, Coleoptera, Odonata, Plecoptera, and Hymenoptera represented.

UPPER TRIASSIC BLACKSTONE FORMATION, IPSWICH COAL MEASURES AT DENMARK HILL, IPSWICH, SOUTH-EASTERN QUEENSLAND. At Denmark Hill very close to the centre of Ipswich the upper part of the Ipswich Coal Measures underlies an urban park reserve. In 1890 insects were first found in this locality and much geological interest followed. The richly fossiliferous horizon is a finely arenaceous siltstone about 15cm thick assigned to the Blackstone Formation, the youngest formation of the Ipswich Coal Measures. Coal seams are known above and below the fossiliferous sequence indicating a lacustrine environment. Plant fossils are prolific and among the fauna freshwater bivalves and estherid crustaceans accompany the insects while dinosaur footprints are known in the formation. Dunstan (1916) gave a detailed account of the small quarry in which the insects were collected. The fauna is dominated by beetle elytra, Blattodea and Homoptera and includes fewer Orthoptera, Phasmatodea, Neuroptera, and Mecoptera.

LOWER CRETACEOUS (APTIAN) KOONWARRA FOSSIL BED NEAR LEONGATHA, SOUTH GIPPSLAND, VICTORIA. During road widening excavations on the South Gippsland Highway between Koonwarra and Tarwin council workers recognised fossil fish in the spoil. A short succession of varved lake deposits is within a thick fluvialite succession, the Korumburra Group (=Strzelecki Group). The lacustrine sequence contains some beds with abundant fish (Waldman, 1971) and other beds with numerous insects, plants and a few other arthropods, some annelid worms (Jell & Roberts, 1986) and a few feathers (Talent et al., 1966), the earliest indications of birds in Australia. The insects include many aquatic larvae and numerous terrestrial adults. By analogy with present day lakes in western Victoria it was possible to identify representatives of the insect fauna that lived in the lake, a fauna of larvae that lived in a feeder stream and insects that lived on the surrounding land and then fell onto the surface of the water (Jell & Duncan, 1986). Mayfly nymphs from the stream fauna and dipteran and beetle larvae from the lake dominate the aquatic fauna while Hemiptera, Mecoptera, hymenopterans, beetles, flies and blattodea are common among the winged adults. The beds are dated palynologically (Dettmann, 1986) as Aptian and a fission track date (Drinnan & Chambers, 1986) of 116±5Ma has been obtained.

PALEOCENE REDBANK PLAINS FORMATION NEAR IPSWICH, SOUTHEASTERN QUEENSLAND. The Tertiary Booval Basin which overlies the Mesozoic Ipswich Basin extends over 50km² in the Ipswich-Bundamba-Dinmore area. The Palaeocene Redbank Plains Formation consists of about 70m of mudstone, claystone, shale and fine sandstone which were deposited in a freshwater lake in structural depressions on the eroding upper surface of the Ipswich Basin sediments. The fossil content includes fish, turtles, crocodiles, ostracodes, cladocerans, insects and plants. Riek (1952a) noted two different faunas: one at Redbank Plains in a hardened band of ferruginous mudstone is dominated by Homoptera and Coleoptera with a few Diptera, Mecoptera, Neuroptera, Orthoptera and Hemiptera and the other, at Dinmore in compact clays and clay shales, contains Orthoptera, Isoptera, Megaloptera, Homoptera and possibly Odonata.

POTENTIAL FOR FUTURE FOSSIL INSECT RESEARCH IN AUSTRALIA

CARBONIFEROUS. The nonmarine mostly Upper Carboniferous Joe Joe Formation in the Drummond Basin of central Queensland contains finegrained varved siltstones and mudstones in its upper part. In the varved units at a number of localities are arthropod tracks closely comparable to those which occur with the only known Australian Carboniferous insect in the Wynyard Tillite in Tasmania. Although no evidence of fossil insects in the Joe Joe Formation exists it is clear that the right sorts of environments existed and some further prospecting in the fine lacustrine sediments could be productive.

UPPER TRIASSIC BEDS IN NORTHERN ESK GRABEN. North of the Ipswich area in southeastern Queensland extends the Esk Trough, a NNW/SSE trending structurally controlled intermontane basin of mainly Middle Triassic (Anisian) sedimentation. The northern extent of this structure is not obvious and the more widespread sediments of this age in the Murgon-Gayndah area are probably northern deposits in the same large lake of that time. The whole belt from Wivenhoe to Gayndah had a dynamic environment with active degradation of adjacent highlands as indicated by abundant and widespread conglomerates and conglomerate/sandstone/shale alternation vertically and laterally. Tuffs throughout and larva flows in the northern areas indicate contemporary volcanism. Through this large area of deposition, plant fossils are common with floras dominated by *Dicroidium*. A few insects have been collected from the Esk Formation in the Wivenhoe/Bellevue area (*Steinhardtia* and a proto-elytroptera in the QM collection), an extensive collection of insects, dominated by blattodeans and beetles from the Mondure Formation in the region north of Murgon to Windera is in the Queensland Museum and insects are known (Lambkin, 1988) in the Gayndah Formation near Gayndah. The insects in these three occurrences are Middle Triassic and thus older than the Denmark Hill and Mt Crosby faunas and of comparable age to the Brookvale and Ashfield faunas in Sydney. There is a very significant scope for further work in this region with the promise of significant results in piecing together Australian insect faunas of that age.

MESOZOIC INSECTS IN WESTERN AUSTRALIA. Two records of insects in the Mesozoic of Western Australia are in the Lower Jurassic Cockleshell Gully Formation (Riek, 1968c) and the Cretaceous (Santonian) Gingin Chalk (Etheridge, 1913).

Riek (1968c) described the tenebrionid beetle, *Mesothoris westraliensis* and noted two other indeterminate beetle elytra and a poorly preserved cockroach forewing possibly allied to *Austroblatulla*. Riek (1968c) speculated that the sediments were Triassic based on the ranges of various insect groups in Australia. However, Balme (1964a) determined an Early Jurassic age palynologically and the enclosing sediments are now considered part of the Cattamarra Coal Measures Member of the Cockleshell Gully Formation. The material was collected at 3 different localities in the Hill River region inland from Jurien Bay so the occurrence is not a restricted one. These insects occur in nonmarine finegrained shales and siltstones associated with widespread subsurface coal seams, an environment very conducive to insect fossilization. Riek (1968c) indicated that although they were too poorly preserved for identification, numerous other insects occurred in the same collection. There is every indication that a more comprehensive insect fauna remains to be found in this rock unit. Since our knowledge of Australian Jurassic faunas is almost non-existent, any further results from this source could be extremely significant.

Etheridge (1913) illustrated a beetle elytron and indicated a myriapod sclerite in a collection of fossils sent to him from One Tree Hill, just north of Gingin. The small quarry on top of this hill is in the Upper Cretaceous Gingin Chalk, a white, friable slightly glauconitic chalk. It is a marine unit with foraminifera, corals, crinoids, bivalves, ammonites and other typical marine fossils. This is an extremely unlikely environment in which to find terrestrial arthropods fossilized and Etheridge's report must be examined cautiously. Apparently, no further insects or myriapods have been collected from this unit and my efforts in the same quarry produced no semblance of them so I am very skeptical of the 1913 report.

JURASSIC TALBRAGAR FISH BED NEW SOUTH WALES. The celebrated Talbragar Fish Beds in the Lower Jurassic Purlawaugh Formation between Gulgong and Mudgee in central New South Wales are best known for their



FIG. 3. *Cicada? lowei* Etheridge & Oliff, 1890; AMF35725; 20mm long. The only insect so far described from the Talbragar Fish Bed.

prolific fish and plant fossils but it has long been known that they contain a few insects. These fine cherty shales were deposited in a relatively small body of water near the southern margin of a very large freshwater lake that extended north into Queensland and west almost to South Australia. In 1889 soon after the fossil site was found Charles Cullen was sent by the Geological Survey of New South Wales to make a collection of the fish and plants. Among the very many specimens he procured was a single insect specimen. Etheridge & Oliff (1890) erected *Cicada? lowei* for this specimen (Fig. 3). Riek (1970c) noted this specimen as coming from the Jurassic but considered its identification as a cicada to be wrong. However, because he did not show any identified Jurassic occurrences of insects from Australia (Riek, 1970c, table 8.1) it may be assumed he considered the specimen unidentifiable. Other rare insects from the Talbragar Fish Beds are known to be in private collections and a study of this fauna is required as the only available Australian Jurassic insect fauna.

CRETACEOUS OF WESTERN QUEENSLAND. The marine Lower Cretaceous Walumbilla Formation of the Richmond-Hughenden area in the Great Artesian Basin has yielded several dragonfly wings of one species. The first of these was the first fossil insect recorded from Australia (Woodward, 1884). These are quite rare among a very diverse marine fauna and prospecting for these specific forms would be an extremely time costly exercise. They are more likely to be found coincidentally by people collecting the marine fauna and usually at different times.

During excavations to collect sauropod dinosaurs in the Upper Cretaceous, fluvatile, nonmarine Winton Formation near Winton, western Queensland, three insect specimens have been collected. One is a dragonfly wing, the same or similar to that from the marine Walumbilla Formation and the other two are incomplete mecopteroid specimens. The insects are relatively rare in fine shales with prolific plant fossils. These sediments weather rapidly on exposure and the fossils were only found because fresh sediment was being excavated on a considerable scale to build a dam. Further excavation in these sediments in the same area is likely to produce more specimens and although the rate of discovery could be expected to be erratic and not guaranteed its continuation could be highly beneficial as the second Cretaceous site in Australia. The Winton Formation outcrops over wide areas of western Queensland on top of the marine sequence in the Artesian Basin and since it is now known to contain at least one fossil locality, further prospecting, especially of sites with fine lacustrine sediments and/or plant fossils should be undertaken.

TERTIARY LIMESTONES AT RIVERSLEIGH, NORTHWESTERN QUEENSLAND. Riversleigh, northwestern Queensland, has been made a household word in vertebrate palaeontology by the extensive etching of Tertiary limestones which has liberated diverse assemblages of bones from numerous Oligocene to Pleistocene horizons. Limestone from the Late Oligocene/early Miocene Upper Site yielded some 3D insect remains mentioned in preliminary lists in Archer et al. (1989, 1991). Duncan, Briggs & Archer (1998) studied the mechanism of preservation of these Riversleigh insects in which the cuticle has been replaced by phosphate and offered a preliminary taxonomy of the specimens used in their study. They referred to 4 different species of beetles (including a

curculionid, a hydrophyloid and a cupidoid), a trichopteran larva and a juliid millipede and noted that other insects are present in the fauna. Co-occurring taxa are one, possibly two, species of ants, an hemipteran, possibly a cicada, and millipedes. The specimen illustrated by Archer et al. (1991) as a millipede represents the same taxon as that described as the tail assembly of a trichopteran larva by Duncan et al. (1998). Discussions with various entomologists have unearthed a variety of different opinions as to the identification of several of these taxa. The Upper Site which yielded the faunule is interpreted as a shallow lime-rich pool in a rain forest where anaerobic conditions may have intermittently occurred. Etching of the Riversleigh limestones is continuing and a detailed study of these insects preserved in such a rarely seen way is necessary as soon as possible.

FOSSIL INSECT THAT IS A CRUSTACEAN

Austrolestidion duaringae Tillyard, 1916 was erected for two specimens (GSQ120) in a core of Tertiary sediments from the Daringa Bore in central Queensland that were identified as zygopterid dragonfly larvae. Rozefelds (1985) provided a good photograph of the specimens and suggested that it probably represents a single

juvenile parastacid crustacean. The chelate appendages are confirmation of this identification. However, Riek (1970b), Kukalova-Pek (1991) and Carpenter (1992) still maintained it amongst the Zygoptera, apparently not having examined the specimen and unaware of Rozefelds' paper.

INSECT THAT IS A FAKED FOSSIL

Austrodictya Tillyard, 1922c, with type species *A. corbouldi* Tillyard, 1922c, was based on a wing fragment (distal third) preserved in a crystal of selenite (or gypsum). The specimen came from the Mount Elliott Mine in Precambrian terrane in northwestern Queensland. It was collected 87m (260 feet) below the surface in the copper lode being worked in the mine. Tillyard went to great pains to impress on his readers that the wing fragment was completely embedded in the crystal and could not possibly have been mechanically inserted. Tillyard considered it an archaic orthopteran quite unlike anything he had seen in the living or fossil fauna. He even sent a drawing of the venation to Mr A.N. Caudell, well known orthopterist of the day at the US National Museum, who concurred that it resembled nothing in the living fauna.

The specimen which had been forwarded to the Geological Survey of New South Wales, Sydney in about 1912 by Mr Corbould, Manager of the Mt Elliott Mine, remains in the Mines Department collections to this day (MMF25954). It next attracted attention in 1952. Harry S. Ladd, U.S. National Museum, Washington DC who was compiling a major work on the variety of types of fossilization of different fossils enquired about it as an unique specimen. Mr H.F. Whitworth, Curator of the Geological & Mining Museum set out to photograph the specimen for Ladd. He found the face of the crystal somewhat scratched over time and decided it would be best photographed under water.

To his surprise, Mr Whitworth found that when he immersed it in water, air bubbles formed around the wing. This suggested a cavity. On further investigation he found a slit along the side and concluded that the wing fragment must have been inserted between the flexible laths of the crystal. Whitworth related how he had carried out the same exercise of spreading the laths with the blade of a penknife and inserting a wing. Indeed I have done the same thing and it is perfectly achievable.

Whitworth took Tillyard's specimen to Mr A. Musgrave, entomologist at the Australian



FIG. 4. *Austrolestidion duaringae* Tillyard, 1916, GSQ120; body 20mm long. A juvenile parastacid crustacean.

Museum, who identified the wing as belonging to *Terpandrus horridus*, commonly known as the Great Gum-Tree Grasshopper and well known in northern Australia.

Clearly, someone had inserted the wing fragment into the crystal as a joke, but the joke went on too far and thus we have *Austrodictya corbouldi* as a junior synonym of *Terpandrus horridus* and one generic name is removed from the catalogue of Australian fossil insects.

INSECT TRACES

LEAF MINES As they are traces of insect activity, mention should be made of a series of papers by Andrew Rozefelds (1988a, 1988b; Rozefelds & Sobbe, 1987) on leaf mines in Triassic to Eocene leaves. Mesozoic examples are rarely recorded and probably not common. Further, the inference that the Late Jurassic to Early Cretaceous *Pachypteris* leaf mine is the work of a lepidopteran makes it a very early record for this order and the only record of a lepidopteran mining a gymnosperm. It has already raised expressions of doubt from the Northern Hemisphere (Zherikhin in Rasnitsyn & Cooke, 2002:360).

TAXONOMIC PLACEMENT

Taxonomic nomenclature applied to fossil insects is not yet generally agreed between all palaeoentomologists. The contents of many groups vary widely among authors (cf. the family classification of the Mecoptera between Willman (1989b), Carpenter (1992) and Novokshonov in Rasnitsyn & Quicke (2002)). In the taxonomic compilation herein I have worked from the original placement and applied what appear to me to be sensible amendments. In many cases I have inserted notes in [] to indicate some history of the taxon but this is not exhaustive and I acknowledge that many of the taxa are relatively poorly known and in need of careful revision. Part of the aim of this work is to collate the entire known fauna to promote some of the necessary revisionary work.

In the case of our earlier paper on the Koonwarra Fossil Bed fauna (Jell & Duncan, 1986) we made something of the same statement, acknowledging our shortcomings across all the insect orders and hoping that specialists would be able to improve on these placements. I am delighted to be able to report here on the work of Ansoorge (1996), Borkent (1997), Hamilton

(1992) and Nauman (1993) who have improved our knowledge of various taxa from that fauna.

A couple of areas of taxonomy are worthy of further discussion.

TINDALE'S THEORY OF PRIMITIVE LEPIDOPTERA IN THE TRIASSIC OF QUEENSLAND. Tindale (1945) erected *Eoses triassica*, on a specimen from the Upper Triassic Mt Crosby Formation, as a member of the Lepidoptera within a new suborder, the Eoneura. This claim had considerable significance as it would be the earliest known lepidopteran by a long way. Riek (1955) discussed this claim at length and concluded that *Eoses triassica* is a junior synonym of *Mesochorista proavita* Tillyard, 1916 from Denmark Hill (Riek (1955) identified numerous specimens of this species in material from Mt Crosby) and belongs to the Mecoptera. Tindale (1980) replied by stating further evidence for why he considered *Eoses* to be a lepidopteran and at the same time erected *Eocorona iani* as another lepidopteran in the Eoneura from the same Mt Crosby horizon. Sukhacheva (1982:9) considered *Eocorona* to be a junior synonym of *Prorhyacophila* Riek, 1955 within the Trichoptera. Thus there is considerable refutation of the ideas espoused by Tindale and apparently no other worker has supported, in print, Tindale's assertions of Triassic lepidopterans. Carpenter (1992) noted Tindale's work, discussed Riek's point and considered them unlikely to be lepidopterans. On that basis I have accepted the synonymies of these two genera.

MESOTITAN. *Mesotitan giganteus* Tillyard, 1916, the type species by monotypy, is based on a single large specimen, GSQI22a from brick pits in the Ashfield Formation (previously the Wianamatta Shales) at St Peters, Newtown, Sydney. *Clatrotitan* McKeown, 1937 (see below) was also assigned to the Mesotitanidae and several recent accounts (notably Sharov (1968), Carpenter (1992) and Gorochoff & Rasnitsyn in Rasnitsyn & Quicke (2002)) have considered the two genera synonymous. On the otherhand Riek (1954c) considered that *Mesotitan* (i.e. *M. giganteus*) 'is a quite distinct genus [from *Clatrotitan*] most probably referable to the Homoptera'. The type specimen of *M. giganteus* is a difficult specimen to interpret and has never been adequately described or understood in the published literature. A preliminary examination indicates that an argument can be mounted to support Riek's claim of separate generic status



FIG. 5. *Mesotitan giganteus* Tillyard, 1916, holotype, GSQI22a, 15cm maximum width. The central axis of the body is more or less up and down the vertical crack. This is apparently the counterpart and we are looking up at the underside of the wings. If any part of the body is preserved it would be expected on the opposing part which has not been located. On the left side, wing at 45° to the axis is the hindwing with the posterior cubital and anal areas still folded together with main veins extremely close together in a bundle just overlapping onto and concealing the proximal parts of the subcostal and costal areas of the forewing. The forewing is the long wing down the axis, probably in the position of rest. On the right side the two wings lie on top of each other at a very small angle as deduced by veins of the underlying forewing being pressed through the hindwing. The outer marginal or costal area of the forewing has flexed along the subcostal vein and been turned in under the subcostal area. This is why it appears the oblique crossveins of the outer part of the two wings are in two different directions.

for *Mesotitan* and *Clatrotitan* but there seems good reason to classify them in the one family. However, the considerable explanation and description of the specimen that will be necessary to fully understand this taxon is outside the scope of the present work and will be reserved for another publication. It is sufficient to say *Mesotitan* and *Clatrotitan* belong to the same family, currently classified under Titanoptera.

PERMOCEPHALUS & PERMOCAPITUS. Evans (1943a, 1943b) described these two genera as the heads of homopteran bugs. Although Evans (1957) later entertained some doubt about the specimens he continued to describe them as heads of Homoptera. Riek (1971) showed these specimens to be thoracic segments of small insects. Although he provided a detailed discussion of the possible relationships of these




FIG. 6. *Permocephalus knighti*; holotype, AMF39865; 5mm long.

thoracic fragments he did not formally assign them to any group. The closest comparison he made was with recent Symphyta: Hymenoptera and with Neuroptera. In this state of uncertainty the two genera are not assigned within the following compilation. *Permocephalus knighti* Evans, 1943a (Fig. 6) based on the holotype AMF39865 and *Permocapitus globulus* Evans, 1943b based on holotype AMF40078 both from

the Upper Permian Newcastle Coal Measures at Belmont, near Newcastle in the Sydney Basin.

CATALOGUE FORMAT AND REPOSITORIES

In the catalogue of Australian fossil insects that follows I have attempted to include all genera so far described along with most specimens in open nomenclature but the latter may not be entirely comprehensive. They are organised taxonomically by Orders and then in families arranged usually alphabetically under the order. There is no particular order to genera within the family headings. I have not attempted to diagnose any suprageneric taxa. Each generic entry has the "Name Author, date [type species]" on the first line. The next line has a diagnosis, usually quoted from the original work, or abbreviated where possible. After the diagnosis comes a list of "species Author, date; Age; Geological Formation (i.e. the rock unit); Sedimentary Basin or nearest town". Ages are abbreviated thus: - L, M, U for Lower, Middle and Upper, respectively; Carb=Carboniferous, Perm= Permian, Trias=Triassic, Jur=Jurassic, Cret= Cretaceous. Fm = Formation, FB=Fossil Bed, Sst=Sandstone, CM=Coal Measures. Sydney, Ipswich, Gippsland, etc are sedimentary basins.

Abbreviations for repositories are Australian Museum, Sydney (AMF), Geological Survey of Queensland, Brisbane (GSQI), Museum of Victoria (NMVP), Melbourne University Geology Department (MUGD), the Natural History Museum, London (BMNHIn), Queensland Museum, Brisbane (QMF), University of Queensland (UQF and UQC), University of Sydney, Geology Department (USGD). The Geological Survey of Queensland and University of Queensland collections have now been transferred to the Queensland Museum; the Melbourne University Geology Department collection is now in the Museum of Victoria; and the Sydney University collection is in the Australian Museum.

Order EPHEMEROPTERA (mayflies)

Family **BAETIDAE** Leach, 1815

Cleon Leach, 1815

Larva: small, shrimp-like; head very small; cerci short.

emmavillensis Riek, 1954b; Pliocene;
Vegetable Ck, Emmaville, NSW.

Family **LEPTOPHLEBIIDAE** Banks, 1900

Atalophlebia Eaton, 1881 [*Ephemera australis* Walker, 1853]

Nymph dorsoventrally depressed, with widely flattened femora; head rectangular; eyes small, posterolateral, protruding slightly at margin, well separated medially; thorax slightly wider than head; insertions of foreleg close together, of mid and hindlegs well apart; femora 3 times as long as wide; tibiae not strongly flattened, not quite as long as femora; abdominal segments gradually increasing in length but narrowing, with 9th segment as long as wide; caudal cerci as long as abdomen.

culleni (Etheridge & Olliff, 1890); Plio;
Vegetable Ck; Emmaville, N.S.W.

Family **SIPHONURIDAE** Banks, 1900

Promirara Jell & Duncan, 1986 [*cephalota*]

Nymph only. Similar to living *Mirawara* but with mouthparts less highly adapted for predation. Abdomen with 7 pairs of subequal gills, first pair not reduced. Cerci with hairs on the lateral as well as the median margin but hairs on lateral margins shorter and not extending to the base of cercus.

cephalota Jell & Duncan, 1986; LCret;
Koonwarra FB; Gippsland.

Australurus Jell & Duncan, 1986 [*plexus*]

Nymph only. Head elongate; mandibles large; antenna long; face above antennal insertions ridged. Legs long, thin; tibia shorter than tarsus; tarsal claw apparently long. Mesonotum raised, produced caudally; pronotum large. Abdomen with 7 pairs of unmodified lamellar gills; gill structure imperfectly known but each gill with a strengthened longitudinal ridge at or close to fore margin, possibly with a fibrillar tuft at base; segments 1-9 strongly produced posterolaterally into spines. Caudal filaments with dense fringes of fine hairs except on outer margins of each lateral filament.

plexus Jell & Duncan, 1986; LCret; Koonwarra
FB; Gippsland.

Cleon emmavillensis,
AMD115, x5



Atalophlebia culleni,
AMD113, x5




Promirara cephalota;
large nymph NMVP
102472B (above), x3;
immature nymph
MUGD3757 (right), x9.





Australurus plexus; nymphs; MUGD3754 (far left), x8; NMVP102450 (left), x6; NMVP48610 (top, above), x8; NMVP103120 (above), x4.



Dulcimanna sculptor; NMVP102507 (above centre) and AMF66758 (above left); x3. Line drawing (above right) of Jell & Duncan, 1986, fig. 2H



***Dulcimanna* Jell & Duncan, 1986 [*sculptor*]**

Nymph only, large, elongate body 10 times as long as wide. Head rounded, relatively small, with large bulging dorsal eyes, with entire margin on labrum. Legs with femora of approximately equal length from fore to hind legs, with tibiae and unsegmented tarsi becoming longer to hind leg, with strong curved relatively long tarsal claw. Abdomen with flat lobate gills on segments 1-7, with gently convex internal margins not spinose posterolaterally. Caudal filaments equal, with dense comb of fine long hairs along lateral margins except on outer edges of cerci.

sculptor Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

***Siphonuridae* gen. nov. Jell & Duncan, 1986**

Nymph only. Head rounded; mouthparts conspicuous; eye small; pleural spines at least on segment 9. Caudal filaments moderately long, slightly more than half length of abdomen.

gen. et sp. nov. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Siphonuridae gen. et sp. nov., NMVP103210, x6. Line drawing (left) of Jell & Duncan, 1986, fig. 2E.

Order MEGASECOPTERA


Family **PSYCHROPTILIDAE** Riek, 1976

Psychroptilus Riek, 1976 [*burrettae*]

Hindwing slightly broader than forewing; Sc ending on costa about 2/3 wing length from base; Rs 3-branched; MA, CuA and CuP simple; MP 2-branched;

burrettae Riek, 1976; UCarb; Wynyard Fm; ETas.

[Riek originally assigned this species to the Megasecoptera noting that it combined features of that order and the Palaeodictyoptera. Subsequently it has been assigned to the Palaeodictyoptera by Carpenter (1992) and Rasnitsyn & Quicke (2002) among others.]





Psychroptilus burrettae,
UTGD94563, 50mm
wing span. Wing
venation (left) from Riek,
1976, fig. 1.



Mesomantidion queenslandicum; GSQI1; x1.2.
Line drawing from Tillyard, 1916, pl. 5, fig. 2.



Mesophlebia antinodalis; GSQI3; x2. Line drawing
from Tillyard, 1916, pl. 4, fig. 2.



Peraphlebia tetrastichia; NMVP103212; x10.;
immature nymph.

Order ODONATA (dragonflies)

Family **MESOMANTIDIIDAE** Tillyard, 1916

Mesomantidion Tillyard, 1916

[*queenslandicum*]

Forewing: large, elongate; C weak, reaching margin near base; precostal area narrow, crossed by series of simple crossveins; Sc long, simple; R strong, simple;



subcostal area large, with numerous irregular cross veins; M branching strongly; M1, M2 and M3 separating at level of end of C; M1 and M2 branching again distally; Cu2 2-branched; A1 present; cross veins in single rows.

queenslandicum Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

[Riek (1956:110) considered this specimen may not represent an insect but offered no alternative].

Family **MESOPHLEBIIDAE** Tillyard, 1916

Mesophlebia Tillyard, 1916 [*antinodalis*]


Hindwing: nodus distinct, transverse behind Sc, not reaching R; pterostigma well formed parallelogram with lower side thickened; M1 arching up close to pterostigma; M3-branched; M4 very distant from Cu, separated by single row of long parallel cross veins; cross veins numerous between all veins; M4 distally indistinct, distally a zigzag division between cells.

antinodalis Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

Triassophlebia Tillyard, 1922b [*stigmata*]


Wing with pterostigma elongate; costal veinlets beyond nodus numerous; M1 only slightly converging towards R; M1a arising from M2; no oblique vein between M2 and Ms; M4 not arching downwards.

stigmata Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.




Triassophlebia stigmata; AMF29267; x4 (left).
Line drawing from Tillyard, 1922b, text-fig. 76.

Peraphlebia Jell & Duncan, 1986 [*tetrastichia*]
R3 meeting R2 at level of nodal, continuing in smooth curve to costal margin; with wide area between R3 and R4, with brace vein between R1 and R2 oblique and strong.
tetrastichia Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.




Peraphlebia tetrastichia; mature nymph; MUGD 3731; x3.




Peraphlebia tetrastichia; NMVP 103204; x2 (far left). Line drawing from Jell & Duncan, 1986, fig. 9B.

Family **AESCHNIDIIDAE** Handlirsch, 1906
Aeschnidiopsis Tillyard, 1918a [*flindersiensis*]
Triangles with strongly curved distal side; arculus not connected to CuP.
flindersiensis Woodward, 1884; LCret; Walumbilla Fm; Artesian.



Aeschnidiopsis flindersiensis, UQF3162; 55mm long. Line drawing (left) of Riek, 1954d, text-fig. 1




Family **PERMAESCHNIDAE** Martynov, 1931
Polytaxineura Tillyard, 1935b [*stanleyi*]
Forewing: C and Sc areas narrow, towards nodus; C and R areas beyond nodus almost without cross veins; Cu2 and A1 with an anterior convexity about 0.3 of length from base; cubital space with 4 cross veins before the anterior convexity; polygonal network distally and posteriorly.
stanleyi Tillyard, 1935b; UPerm; Newcastle CM; Sydney.

Family **PROTOMYRMELEONTIDAE** Handlirsch, 1906

Triassagrion Tillyard, 1922b [*australiense*]
Forewing: pterostigma short, twice as long as wide; R1, M1, M1a and M2a finishing close together near apex; 24 cross veins between C and R1 after Sc finished and before pterostigma; M with 7 major branches at margin; Cu simple and zigzag; A1 zigzag distally and with only one row of cells behind it; triangular areas between M2a and M2b and between Ms and M3 with irregular polygonal veins.
australiense Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.





Polytaxineura stanleyi; S343; 40mm long. Line drawing (above) of Tillyard, 1935b, text-fig. 1.



Triassagrion australiense; GSQI290a; x3. Line drawing from Tillyard, 1922b, text-fig. 77.



Triassolestes epiphleboides; GSQI205a; x3. Line drawing (right) from Tillyard, 1918d, text-fig. 11.



Coenagrionid indet.; NMVP102508; x7.

Family **TRIASSOLESTIDAE** Tillyard, 1918d

Triassolestes Tillyard, 1918d [*epiphleboides*]

Wing: M 4-branched; M4 and Cu1 slightly divergent; single row of cells between Cu1 and Cu2 basally, irregular polygons distally; Cu2 curving back distally to margin; one row of cells between Cu2 and margin increasing to 2 distally



epiphleboides Tillyard, 1918d; UTrias; Blackstone Fm; Ipswich.

Family **COENAGRIONIDAE** Kirby, 1890

Coenagrionid indet.

Body long, thin, with moderately expanded gills and antenna, with pedicel not markedly lengthened; mask with medial projection; gills with rounded apex

coenagrionid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Perrisophlebia multiseriata; GSQI203; x3. Line drawing from Tillyard, 1918d, text-fig. 13.


Family **UNCERTAIN**

Perissophlebia Tillyard, 1918d

[*multiseriata*]

Wing with an irregular double row of cells between C and R beyond pterostigma; between R and M1 at same level an irregular triple row of cells proximally then a more regular double row; 3 cell rows below M1 a strong convex sector.

multiseriata Tillyard, 1918d; UTrias; Blackstone Fm; Ipswich.



Samaruura Brauer, Redtenbacher & Ganglbauer, 1889 [*gigantea*]

Nymph. long slender abdomen, long thin legs, lamellar gills.

Samarura sp. Rozefelds, 1985; UTrias; Aberdare Conglomerate; Ipswich.

Samarura sp.; QMF12996; 10mm long.

Order PERLARIA (stoneflies)

Family **EUSTHENIIDAE** Tillyard, 1921

Mesonotoperla Riek, 1954c [*sinuata*]

Rs simple (if forked only as terminal twigging); Sc extending almost to apex; Cu1 2-branched; M 2-branched; m-cu1 and cu1-cu2 well separated; no r-m over basal half.

sinuata Riek, 1954c; MTrias; Hawkesbury Sst; Sydney.



Mesonotoperla sinuata, AMF35877, x3 (upper right). Line drawing from Riek, 1954c, fig. 5.

Stenoperlidium Tillyard, 1935c [*permianum*] [= *Antitaxineura* Tillyard, 1935c; type *anomala*]

Forewing: very narrow; Sc short, ending just beyond midlength in short fork connecting with both C and R1; costal veinlets only moderately developed; Rs with 4 pectinate branches distally; M 2-branched, forking just before midlength; Cu1 strong, 3-branched distally; cu1-m strong; Cu2 simple.

permianum Tillyard, 1935c; UPerm; Newcastle CM; Sydney.

triassicum Riek, 1956; UTrias; Blackstone Fm; Ipswich.

anomala Tillyard, 1935c; UPerm; Newcastle CM; Sydney.

Family **GRIPOPTERYGIDAE** Enderlein, 1909

Eodinotoperla Jell & Duncan, 1986





[*duncanae*]

Nymph with typical anal gill tufts, long, thin, with small tenth abdominal segment, with expanded femora, with long fine cerci. Adult wing with small terminal branch on Rs, with fewer crossveins.

duncanae Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.




Eodinotoperla duncani; adult NMVP32285, x6 (right); nymph NMVP27040, x6 (far right); line drawing from Jell & Duncan, 1986, fig. 15A.




Stenoperlidium permianum; B77; 22.5mm long. Line drawing (2nd above) of Tillyard, 1935b, text-fig. 1. Nymph, AMF40701 (3rd above); x4.





Austroidelia perplexa, AMF39179, 17mm long.
Line drawing (above) from Riek, 1954c, fig. 1.



Mesacridites elongata, AMF30970; 55mm long.
Line drawing (above) from Riek, 1954c, fig. 2.



Mesorthopteron locustoides; GSQ172; 7mm long.
Line drawings (right) from Tillyard, 1916, pl. 2,
figs 5, 6.

Order PROTORTHOPTERA

Family IDELIIDAE Zalessky, 1929

Austroidelia Riek, 1954c [*perplexa*]

Forewing: Sc with numerous anterior branches; Rs arising near base; M forked before origin of Rs; M 3-branched; M3+4 simple; Cu1 sigmoidally curved near base, with many pectinate branches posteriorly, of which proximal 3 or 4 do not reach wing margin; Cu2 straight, simple; anals short, straight.

perplexa Riek, 1954c; MTrias; Hawkesbury Sst; Sydney.

Mesacridites Riek, 1954c [*elongata*]

Forewing: Costal area only slightly expanded; Sc short, ending well before apex; R1 simple; Rs with numerous pectinate branches.


elongata Riek, 1954c; MTrias; Hawkesbury Sst; Sydney.

Family MESORTHOPTERIDAE Tillyard, 1916

Mesorthopteron Tillyard, 1916 [*locustoides*]

Forewing: narrow, densely veined, weakly thickened, forming weak tegmen; Sc strong, straight, sending series of oblique cross veins into costal space; costal space wide; R strong, parallel to Sc, giving off Rs; M weaker, giving off 3 branches before origin of Rs, with branches running forward from main stem; Cu 2-branched. Hindwing: not thickened, with less dense venation; Cu unbranched; A1 and A2 present.

locustoides Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.



Order BLATTARIA (cockroaches)

Family MESOBLATTINIDAE Handlirsch, 1906

Austroblattula Tillyard, 1919c [*ipsviciensis*]
Tegmen; small; Sc fused with R basally, straight behind humeral area; R curved basally, straight for most of its length, curving forward distally; anal area very broad; anal veins apparently unforked.

ipsviciensis Tillyard, 1919c; UTrias;
Blackstone Fm; Ipswich.



Samaroblatta Tillyard, 1919c [*reticulata*]
Tegmen: with wide humeral area, as long as clavus; Sc simple, strong, with or without weaker branch onto humeral area; R sigmoidally curved; anal area well defined; forkings of A, if any, as terminal twiggings.

reticulata Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

triassica Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

jonesi Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

blatteloides Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

intercalata Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

Triassoblatta Tillyard, 1919c [*typica*]
Tegmen: Sc reduced; R strongly developed, sending numerous veinlets across the costal area, almost reaching apex; M dividing dichotomously and often; Cu similarly branched, curved posteriorly; Y-vein on clavus.

typica Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

insignita Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

intermedia Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.

tasmanica Riek, 1962; Trias; New Town Fm; Hobart, Tas.

jonesi Tillyard, 1919c; UTrias; Blackstone Fm; Ipswich.



denmeadi Tillyard, 1937; MTrias; Mt Crosby Fm; Ipswich.



Austroblattula ipsviciensis; GSQI105; 7.5mm long. Line drawing (left) of Tillyard, 1919c, text-fig. 39.



Samaroblatta reticulata; GSQI155; 10.5mm long. Line drawing (above) of Tillyard, 1919c, text-fig. 34.




Triassoblatta typica; GSQI180; 11.5mm long. Line drawing (above) of Tillyard, 1919c, text-fig. 31.



Triassoblatta jonesi; UQC1861; 7mm long.



Samaroblatta reticulata; GSQI14; 12mm long.





Methana? sp., nymph, NMVP102532 (below left), x6; wings NMVP27045 (above), x5; incomplete adult NMVP103231 (below right), x6.



Family MYLACRIDAE

Austromylacrites Tillyard, 1916 [*latus*]
Forewing: broadly oval; numerous main veins diverging from base; all veins dichotomously branched; no cross veins.

latus Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.



Austromylacrites latus, GSQI8; 11mm long. Line drawing (above) from Tillyard, 1916, pl. 2, fig. 1.

Family BLATTIDAE Stephens, 1829

Methana Stål, 1877

Wing with long costal cell more than half wing length; Sc simple; clavus with markedly curved margin.

Methana? sp. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family INCERTAE SEDIS



Notoblattites Tillyard, 1916 [subcostalis]

Tegmen: large, broad, oval; Sc strong and almost to wing tip; costal space broad, with simple or once-branched cross veins; costal border slightly convex; R very strong, parallel to Sc; R with 3 posterior branches, posterior one unbranched, second 3-branched, anterior one 2-branched; M 4-branched.



subcostalis Tillyard, 1916; MTrias; Ashfield Fm; Sydney.

wianamattensis Tillyard, 1918b; MTrias; Ashfield Fm; Sydney.


mittelli Tillyard, 1918b; MTrias; Ashfield Fm; Sydney.



Notoblattites wianamattensis; AMF39324; 28mm long (right). Line drawing (right above) from Tillyard, 1918b, text-fig. 8.



Notoblattites subcostalis; GSQI24a; 46mm long. Line drawing (left) of Tillyard, 1916, pl. 6, fig. 1 [specimen obviously damaged after it had been drawn].




Order ISOPTERA(termites)

Family **MASTOTERMITIDAE** Desneux, 1904

Blattotermes Riek, 1952b [*neoxenus*]

Forewing: archdictyon strongly developed; Sc reduced, very short; M parallel to R, forking irregularly only in apical third; Cu1 with many pectinate branches, the basal one being further branched.

neoxenus Riek, 1952b; Palaeocene; Redbank Plains Fm; Ipswich.



Blattotermes neoxenus; UQF10431; 24mm long. Lime drawing from Riek, 1952b, fig. 1.

Order PROTELYTROPTERA

Family **PROTOCOLEIDAE** Tillyard, 1924

Austrelytron Kukalová, 1966 [*tillyardi*]

Forewing: small densely granulose, with isolated pointed tubercles in rows on the veins; costal expansion circular but not prominent; tapering from midlength; Sc with 2 or more parallel branches finishing well before apex; R sending several irregular branches towards Sc; Rs distinct, originating after midlength of wing; CuP long, concave forward; 3 unbranched anal veins; cross veins few, simple.

tillyardi Kukalová, 1966; UPerm; Newcastle CM; Sydney.

Phyllelytron Kukalová, 1966 [*folium*]

Forewing: tegminous, dark, densely granulose; proximal costal expansion circular, strongly projecting; tubercles distinct in anal area; apex curved back, only slightly prolonged, pointed; main veins not parallel; branches of R, M and CuA usually indistinct; Sc with 2 or more long branches with numerous veinlets; R with a series of weak irregular anterior branches; CuP long, branched; 4-5 anal veins broad, some directed anteriorly.

folium Kukalová, 1966; UPerm; Newcastle CM; Sydney.

petalon Kukalová, 1966; UPerm; Newcastle CM; Sydney.


granulatum Kukalová, 1966; UPerm; Newcastle CM; Sydney.

melinum Kukalová, 1966; UPerm; Newcastle CM; Sydney.



Protocoleus Tillyard, 1924 [*mitchelli*]

Forewing: tegminous, darkly pigmented, covered by flat regularly arranged tubercles (not on veins) and fine granules; apical part prolonged and tapering; main veins with almost parallel branches; Sc with several parallel branches almost to apex; R with series of long branches towards Sc; Rs similar to other branches of R; M and Cu with variable numbers of irregular branches; CuP long, branched; 5 anal veins; cross veins numerous and regular.


mitchelli Tillyard, 1924; LPerm; Newcastle CM; Sydney.





Austrelytron tillyardi; BMIn45525; 11mm long. Line drawing (above) of Kukalova, 1966, fig.5.




Phyllelytron folium; BMIn45364; 23mm long. Line drawing (above) of Kukalova, 1966, fig. 1





Protocoleus mitchelli; AMF41710; 21.5mm long (left). Line drawing of Tillyard, 1924, text-fig. 3.



Chanoselytron gingiva; BMNHIn45493; 7.5mm long (left). Line drawing of Kukalova, 1966, text-fig. 14.



Dermelytron conservativum; BMNHIn45725; 7.2mm long. Line drawing (above) of Kukalova, 1966, text-fig. 11.



Psychelytron progressivum; BMNHIn45900; 8mm long (above left). Line drawing of Kukalova, 1966, text-fig. 13.



Elytrathrix hirsuta; BMNHIn45503; 15mm long. Line drawing (above) of Kukalova, 1966, text-fig. 8.

Family DERMELYTRIDAE Kukalova, 1966

Chanoselytron Kukalová, 1966 [*gingiva*]

Forewing: convex, thin, narrow; costal expansion large, rounded, extending beyond base of wing proximally; posterior margin slightly concave in anal area; no veins evident.

gingiva Kukalová, 1966; UPerm; Newcastle CM; Sydney.

Dermelytron Kukalová, 1966 [*conservativum*]

Forewing: oval, little sclerotised; costal expansion projecting only a little; setae making a patch in subcostal area; apex directed posteriorly; CuP and one anal vein sometimes weakly indicated.

conservativum Kukalová, 1966; UPerm; Newcastle CM; Sydney.

pigmentatum Kukalová, 1966; UPerm; Newcastle CM; Sydney.

Psychelytron Kukalová, 1966 [*progressivum*]

Forewing: convex, thin, weakly sclerotised; apex directed anteriorly; posterior margin concave distally; CuP and A short, simple, not reaching margin.

progressivum Kukalová, 1966; LPerm; Newcastle CM; Sydney.

Family PERMOPHILIDAE Tillyard, 1924

Elytrathrix Kukalová, 1966 [*hirsuta*]

Forewing: densely granulose, with sparse indistinct tubercles distinct basally; setae bordering the yard in presence of conspicuous large tubercles in costal area; costal expansion projecting near base, with large tubercles; main veins strong in basal 1/3, absent distally.

hirsuta Kukalová, 1966; UPerm; Newcastle CM; Sydney.

Permophilus Tillyard, 1924 [*pincombei*]

Forewing: surface densely granulose, with isolated indistinct tubercles; narrowed and prolonged apically; costal expansion projection; main veins strong in basal third but completely lacking in rest.

pincombei Tillyard, 1924; UPerm; Newcastle CM; Sydney.

minor Tillyard, 1924; UPerm; Newcastle CM; Sydney.

hirtus Kukalová, 1966; UPerm; Newcastle CM; Sydney.

capulus Kukalová, 1966; UPerm; Newcastle CM; Sydney.

Family **PERMOFULGORIDAE** Tillyard, 1918b

Permofulgor Tillyard, 1918b [*belmontensis*]

Tegmen: delicate, broadest near base, narrow, elongate; anal veins strongly developed; A1 and A2 both 2-branched from near their bases; Sc less than half wing length; R, M and Cu united near base; oblique cross veins between R, M, Cu and A veins; anal Y-vein absent.

belmontensis Tillyard, 1918b; UPerm; Newcastle CM; Sydney.

indistinctus Tillyard, 1922a; UPerm; Newcastle CM; Sydney.

Family **STENELYTRIDAE** Kukalova, 1966

Labidelytron Kukalová-Pek, 1988 [*Stenelytron enervatum* Kukalová, 1966]

[replacement name for *Stenelytron* Kukalova]

Forewing: costal expansion subcircular; setae forming patch in subcostal area; apex directed anteriorly; Sc short not reaching margin; subcostal area broad; Rs originating beyond midlength, simple, weak; CuA short; CuP slightly concave forward; anal veins broad, 3-5, simple or forked.

enervatum (Kukalová, 1966); UPerm; Newcastle CM; Sydney.


Xenelytron Kukalová, 1966 [*ligula*]

Forewing: almost flat; costal expansion large, semicircular; R1 directed towards apex; Rs originating late, weak; M close to Cu, simple or forked; Cu long; 3-5 strong anal veins, simple or forked.


ligula Kukalová, 1966; UPerm; Newcastle CM; Sydney.




Permophilus pincombei; AMF19779; 21.5mm long. Line drawing (above) of Tillyard, 1924, text-fig. 1a.



Permofulgor belmontensis; AMF41758; 9.5mm long. Line drawing (above) of Tillyard, 1918b, text-fig. 3.




Labidelytron enervatum; BMNHIn45958; 14mm long. Line drawing (above) of Kukalova, 1966, text-fig. 9.



Xenelytron ligula; BMNHIn45526; 11.5mm long (right). Line drawing of Kukalova, 1966, text-fig. 10.





Elcanopsis sydneyensis; MMF15455; 7mm long.
Line drawing of Tillyard, 1918c, text-fig. 1.



Mesogryllacris giganteus; UQC1920; 60mm long.
Line drawing (right) of Riek, 1955, fig. 31.



Proparagryllacris crassifemur; AMF39251, x3.
Line drawing (right) of Riek, 1956, text-fig. 2.



Neohagla sinuata; UQC2216; x2.


Order ORTHOPTERA (grasshoppers, crickets)

Family ELCANIDAE Handlirsch, 1906

Elcanopsis Tillyard, 1918c [*sydneyensis*]

Sc closer to C distally than to R; wing with brown pigment along costal space; Rs with few branches wide apart; cross veins not numerous, wide apart.

sydneyensis Tillyard, 1918c; MTrias;
Hawkesbury Sst; Sydney.



Family PROPARGRYLLACRIDIDAE Riek, 1956

Mesogryllacris Riek, 1955 [*giganteus*]


Forewing: costal margin expanded at base; C strong, 0.25 in length; costal space with a number of costal veinlets; R1 with series of short branches to wing margin beyond Sc; Rs with series of pectinate branches over apical half; M 2-branched, forking in basal half of wing, not touching Rs; Cu 2-branched, not fused with posterior branch of M; Cu forking at same level as M.

giganteus Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

Proparagryllacris Riek, 1956 [*crassifemur*]

Forewing: Sc curving forward distally, 2 anterior branches; M arising from R close to base; stem of M fused with upper branch of Cu; M forking distally after separating from Cu; M and Cu both 2-branched.

crassifemur Riek, 1956; UTrias; Blackstone Fm; Ipswich.



Family HAGLIDAE Handlirsch, 1906b

Neohagla Riek, 1955 [*sinuata*]

Forewing: costal space expanded over basal half; Sc with oblique veinlets to margin between which are many cross veins; R1 almost parallel to Sc, with 2 or 3 oblique anterior branches towing margin; Rs diverging rapidly from R1; M3+4 and Cu fusing only after origin of CuA2; r1-rs sinuous.

sinuata Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

Prohagla Riek, 1954c [*superba*]

Forewing: C with numerous weak branches in expanded base of costal space; Sc extending towards apex with numerous anterior branches; Rs arising near middle, with a few pectinate branches; M 3-branched, forking early; M3+4 fused to CuA for short distance; M1 not sigmoidally curved; CuA1 multibranched; CuP arched away from CuA2. Hindwing: costal space narrower; no distinct costal vein; M 3-branched; m-cu present; Cu and anals branching from base.

superba Riek, 1954c; MTrias; Hawkesbury Sst; Sydney.


imperfecta Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

Family **LOCUSTOPSEIDAE** Handlirsch, 1906

Triassolocusta Tillyard, 1922b [*leptoptera*]


Forewing: very long and narrow; Sc less than half costal length, with shorter anterior branch; Rs 4-branched in pectinate series; M 3-branched from about level of origin of Rs; M3+4 forked distally; M5 short, connecting to Cu1; rs-m near base of Rs.

leptoptera Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.








Gryllacridoid indet., NMVP32281, x3 (left);
Tridactylid indet., NMVP103232, x6 (lower right).



Xenopterum crosbyi; UQC2066; 13mm long. Line drawing (above) of Riek, 1955, fig. 34.



Triassomantis pygmaeus; GSQI86a; 10mm long. Line drawing (above) of Tillyard, 1922b, text-fig. 73.



Gryllacridoid? indet.

Crumpled adult. Costal area markedly expanded; C, Sc and R parallel; C-Sc space pigmented; scattered crossveins over most of the wing

Gryllacridoid? indet. Jell & Duncan, 1985; LCret; Koonwarra FB; Gippsland.

Family TRIDACTYLIDAE Brunner, 1882

Tridactylid indet.

Head large, with vaulted upper area; eye rounded, moderately sized; antenna short, inserted close to anterior margin of head, with most segments longer than wide; pronotum as long as head. Foreleg with tibia not obviously expanded; tarsus with basal article shorter than middle. Middle leg with tibia not expanded, with tarsus 2-segmented, with first segment long. Hind leg with markedly expanded femur; tibia long, slender, with long ventral terminal spurs.

tridactylid indet.; LCret; Koonwarra FB; Gippsland.

Family TRIASSOMANTEIDAE Tillyard, 1922

Xenopterum Riek, 1955 [*crosbyi*]

Forewing: curved sigmoidally almost to apex; M1+2 separating from Rs before Rs gives rise to the anterior pectinate branches; M3+4 simple or branched near margin; CuA 1 and 2 series of close spaced, parallel branches, rarely branched near margin; CuA 3 giving off a series of posterior pectinate branches.

crosbyi Riek, 1955; UTrias; MtCrosby Fm; Ipswich.

Triassomantis Tillyard, 1922b [*pygmaeus*]

Forewing: costal margin straight; C short; all veinlets and cross veins parallel and oblique; Sc and R1 both turn sharply forward near their apices; Rs 4-branched; R3 straight; R2 running forward, parallel to R5; M 4-branched, all simple; Cu1 straight, with small terminal fork.


pygmaeus Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

Family TETTIGONIIDAE Krauss, 1902

Tettigoides Riek, 1952b [*pectinata*]

Elytron very narrow; Sc long; RS arising at midwing; M nearly parallel to R.

pectinata Riek, 1952b; Palaeocene; Redbank Plains Fm; Ipswich.



Tettigoides pectinata; UQF10640; 32mm long (left). Line drawing of Riek, 1952b, fig. 4

Order TITANOPTERA

Family MESOTITANIDAE Tillyard, 1925

Clatrotitan McKeown, 1937 [*andersoni*]

[=*Mesotitan scullyi* Tillyard, 1925c]

Forewing: broadest at midlength; precostal area apparently absent; stridulatory area about half wing width; Rs 6-branched; M 2-branched; cubito-median Y-vein well-developed.

scullyi Tillyard, 1925c; MTrias; Hawkesbury Sst; Sydney.

tillyardi McKeown, 1937; MTrias; Hawkesbury Sst; Sydney.

andersoni McKeown, 1937; MTrias; Hawkesbury Sst; Sydney.


[*Mesotitan* is discussed above]
[Riek (1954c) considered *tillyardi* and *andersoni* to be junior synonyms of *scullyi*]

Family UNCERTAIN (probably new)



Steinhardtia Jell & Lambkin, 1993 [*maryae*]

Hindwing: Large (>40mm long); primary R fork well beyond that of M; RA, RP, MA, MP, CuA, CuP1+2 and CuP3+4 all deeply dichotomously forked; RA with 4 major branches, RP with 6, MA with 8, MP with 2, CuA with 4, CuP1+2 with 3 and CuP3+4 with 5; strut from CuA to CuP1 rather than CuP1+2; cellular network between veins absent or not preserved.

maryae Jell & Lambkin, 1993; MTrias; Esk Fm; Ipswich.



Clatrotitan scullyi; AMF20270; 86.5mm long



Steinhardtia maryae; QMF25507; 41mm long.
Line drawing of Jell & Lambkin, 1993, fig.1C



Clatrotitan scullyi; USGD310; 131mm long.



Clatrotitan andersoni; AMF36274; 138mm long. Line drawing (above left) of McKeown, 1937, fig. 3.


Order PHASMATODEA (stick-insects)

Family AEROPLANIDAE Tillyard, 1918b

Aeroplana Tillyard, 1918b [*mirabilis*]

Wings very long, narrow; parallel veins strong, close together; Sc weak; R 2-branched, strong; Rs arising near base by double root; M 7-branched; R + M fused at base; Cu1 and Cu2 arising near base and diverging; A1 in front of anal field with irregular veining. Forewing: M forking at same level as first origin of Rs; anterior branch giving 3 branches dichotomously; posterior branches into 3, middle one dividing again. Hindwing: anterior branch of M 2-branched distally; posterior branch dividing early then anterior of these dividing into 3.

mirabilis Tillyard, 1918b; UTrias; Blackstone Fm; Ipswich.



Aeroplana mirabilis; GSQ1126; 41.5mm long. Line drawing of Tillyard, 1918b, pl.44, fig. 12.

Order THYSANOPTERA (thrips)

Family **LOPHIONEURIDAE** Tillyard, 1921

Austrocypha Tillyard, 1935a [*abrupta*]

Forewing: clavus well-developed, wedge-shaped; Cu2 running straight into margin distally; Sc absent; costal area wide; R1 curved forward distally; Rs, M and Cu1 arising close together but independently; Cu1 and Cu2 connected by a basal piece.

abrupta Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

baretti Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

Lophiocypha Tillyard, 1935a [*permiana*]

Forewing: Cu1 unbranched, sigmoidal; M 2-branched; clavus well-developed, with A1 present; distinct break in margin at its end on Cu2.

permiana Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

stanleyi Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

thysanella Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

maxima Davis, 1942; UPerm; Newcastle CM; Sydney.



Lophioneura Tillyard, 1921b [*ustulata*]

Forewing: small, elongate, oval, 3 times as long as wide; R and Cu1 high sharp strong ridges; M and Cu1 fused with R basally; R1 almost straight; Sc and R1 parallel; Rs weak, 2-branched; M 2-branched; Cu1 2-branched, anterior branch curved forward then back distally.



ustulata Tillyard, 1921b; UPerm; Newcastle CM; Sydney.

angusta Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

conjuncta Tillyard, 1926a; UPerm; Newcastle CM; Sydney.



Austrocypha abrupta; AMF39809; 2.6mm long. Line drawing (above) of fore and hindwings from Tillyard, 1935a, text-fig. 10.





Lophioneura ustulata; AMF39344; 5.7mm long. Line drawing (above) of Tillyard, 1921, text-fig. 3.



Edgariekia una; NMVP103242; body 2mm long. Wing venation (above right) of Jell & Duncan, 1986, fig. 18C.

Undacypha Vishniakova, 1981 [*fumida*] [= *Edgariekia* Jell & Duncan, 1986; type *una*] R, M and CuA with common stem; Rs and M 2-branched; CuA unforked, reaching posterior margin at marked break in smooth marginal curve but CuP marking off pigmented clavus; A very weak in clavus.

una Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.




Zoropsocus delicatulus; BMNHIn46399; 2.5mm long (left). Line drawing (above) of Tillyard, 1935a, text-fig. 6.

Zoropsocus Tillyard, 1935a [*delicatulus*]

Wings narrow at base; sides straight; apex rounded; Sc short; R1 simple; Rs 2-branched, finishing before apex; pterostigma chitinised, covering both R1 and R2+3.

delicatulus Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

stanleyi Davis, 1942; UPerm; Newcastle CM; Sydney.



Order PSOCOPTERA (psocids, booklice)


Family PSOCIDIIDAE Tillyard, 1926

Austropsocidium Tillyard, 1935a [*pincombei*]

Forewing: elongate, oval; C gently curved; apex well rounded; Sc ending on R₁; R well removed from C; Rs 2- or 3- branched, branches short; M 4-branched; r-m present; Cu₁ arising just below M, with long flat fork not connected to M; Cu₂ straight, weak, bounding small clavus with 2 veins; A₁ straight, close to Cu₂; A₂ slightly sigmoidal.

pincombei Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

stigmaticum Tillyard, 1935a; UPerm; Newcastle CM; Sydney.





Austropsocidium pincombei; AMF40318; 8mm long (below). Line drawings (above) of fore and hind wings of Tillyard, 1935a, text-figs 1A, 1B.




Megapsocidium Tillyard, 1935a [*australe*]

Forewing: R₁ and R_s connected by cross vein below the fork of R₁; R_s forking at this cross vein and again distally on upper branch; M₁+2 3-branched; M₃+4 unbranched.


australe Tillyard, 1935a; UPerm; Newcastle CM; Sydney.



Megapsocidium australe; BMNHIn46396; 6mm long. Line drawing (above) of Tillyard, 1935a, text-fig. 3



Stenopsocidium elongatum; BMNHIn46397; 6.5mm long. Line drawing (below) of Tillyard, 1935a, text-fig. 4.



Zygopsocus permianus; BMNHIn46398; 4.7mm long (below). Line drawing (above) of Tillyard, 1935a, text-fig. 5.

***Stenopsocidium* Tillyard, 1935a [*elongatum*]**

Wings narrow, elongate-oval; C nearly straight to end of R1; apex well rounded; Sc ending on R1 with branch veinlet to C also; R1 simple; Rs 2-branched; M 5-branched in forewing, 4 in hindwing; Cu1 with long flat fork, not connected with M at all; Cu2 faint; A1 strong, nearly straight.


elongatum Tillyard, 1935a; UPerm; Newcastle CM; Sydney.

Family ZYGOPSOCIDAE Tillyard, 1935a

***Zygopsocus* Tillyard, 1935a [*permianus*]**

Forewing: well rounded; Sc ending on R1 just before midlength; Rs arising near base; i-r beneath end of Sc; Rs forking just below this cross vein, with posterior branch again dividing into 3; M arising close to base; r-m present; M curving back sigmoidally distally, branching; Cu separate from M; Cu1 ending in strong posterior curve; Cu2 and A1 forming Y-vein finishing at margin near Cu1; A2 present; base of wing heavily chitinised; margin broadly chitinised all around.

permianus Tillyard, 1935a; UPerm; Newcastle CM; Sydney.




Order HEMIPTERA (bugs)

Family **APHIDIDAE** Latreille, 1802

?**Triassoaphis** Evans, 1956 [*cubitus*]

Forewing: very strong vein parallel to costal margin incorporating all veins basally except Sc; Sc terminating near midlength; Rs short; M 3-branched; Cu1 2-branched; anals incomplete.

cubitus Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.



Triassoaphis cubitus; UQC2235; 6mm long. Line drawing (above) of Evans, 1956, fig. 26.

Crosaphis Evans, 1971 [*anomala*]

Wings broad; M basally incorporated in same vein as R so that M1+2 and M3+4 diverge separately from their common stem with Rs. CuA 2-branched, meeting R proximally at an acute angle. Clavus absent; anal veins lacking.

anomala Evans, 1971; UTrias; Mt Crosby Fm; Ipswich.

Family **ARCHESCYTINIDAE** Tillyard, 1926

Austroscytina Evans, 1943b [*imperfecta*]


Hindwing: costal space narrow; R1a meets costal margin just beyond midlength; R1b not reaching apex; M 3-branched; M and Cu1 have separate origins.

imperfecta Evans, 1943b; UPerm; Newcastle CM; Sydney.


Bekkerscytina Evans, 1958 [*primitiva*]

Forewing: elongate, oval; costal space wide; R not parallel to costal border; Rs arises from R closer to junction with M rather than with R1; M 3-branched (M1, M2, M3+4); Cu1a not angulate and longer than straight portion of Cu1; Cu1b meets margin distal to Cu2; cu-m present; clavus small; 2 anal veins completely separate.


primitiva Evans, 1958; UPerm; Newcastle CM; Sydney.




Crosaphis anomala; QMF6508; 3.5mm long. Line drawing (above) of Evans, 1971, fig. 3A.




Austroscytina imperfecta; AMF40084; 8mm long. Line drawing (above) of Evans, 1943b, fig. 3.




Bekkerscytina primitiva; AMF47190; 7.8mm long. Line drawing (above) of Evans, 1958, fig. 4.



Eopsyllidium delicatulum; AMF39795; 2.8mm long. Line drawing (right) of Davis, 1942, fig. 5



Eoscytina migdisovae; AMF47185; 9mm long. Line drawing (above) of Evans, 1958, fig. 1.




Alotrifidus interruptus; UQC347; 5mm long. Line drawing (right) of Evans, 1956, fig. 16D.

Eopsyllidium Davis, 1942 [*delicatulum*]

Forewing: very small; Sc absent; R anteriorly convex before junction with M; R1a normal to R1b, running across base of well-developed pterostigma; Rs simple.

delicatulum Davis, 1942; UPerm; Newcastle CM; Sydney.



Eoscytina Evans, 1958 [*migdisovae*]

Forewing: expands distally; R parallels costal margin; Rs arises approx. midway between junction with M and R1 fork; Cu1 steeply bent at its point of apposition to R+M; Cu1a angulate, joined to M by cross vein; Cu1b meets margin distant from Cu2; clavus small; anal veins form a Y-vein.

migdisovae Evans, 1958; UPerm; Newcastle CM; Sydney.


incompleta Evans, 1958; UPerm; Newcastle CM; Sydney.

Family **CERCOPIDAE** Westwood, 1838

Alotrifidus Evans, 1956 [*interruptus*]

Costal margin arched; costal space wide; Rs arising from R nearer to the apex of the tegmen than its base; M and Cu1 a single vein proximally.


interruptus Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.



Triassoscarta Tillyard, 1919d [*subcostalis*]

Forewing: R unbranched; long series of transverse costal veinlets; M and Cu arising together by short stem; A2 finishing as distant from A1 as Cu2.

subcostalis Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.




Triassoscarta subcostalis; GSQ1116a; 8mm long. Line drawing (left) of Tillyard, 1919d, text-fig. 8.


Trifidella Evans, 1956 [*perfecta*]

Forewing: surface rugose; series of transverse accessory veins between R and C; costal margin not arched; Rs originating proximal from midlength, oblique; M and Cu1 fused basally; M 2-branched, each with apical fork; cross veins r, r-m and m-cu present.

perfecta Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.



Trifidella perfecta; UQC945; 7mm long. Line drawing (left) of Evans, 1956, fig. 16C.


**Tychicoloides** Evans, 1963 [*belmontensis*]

Forewing: long, narrow; surface rugose; anterior margin curved; Rs leaving R 0.33 of tegmen length from base; R+Rs linked by a cross vein; M1+2 and Rs in contact at 2 points; Cu1 of 2 parallel arms; clavus unknown.



belmontensis Evans, 1963; UPerm; Eleebena Fm; Sydney.





Tychicoloides belmontensis; AMF50110; 7.2mm long. Line drawing (above) of Evans, 1963, fig. 4.




Chiliocycla scolopoides; GSQI158a; 6.4mm long.
Line drawing (right) of Tillyard, 1919d, text-fig. 2.



Mesoscytina affinis; GSQI235; 6mm long. Line drawing (above) of Tillyard, 1919d, text-fig. 6



Eurymelidium australe; GSQI248a; 5mm long.
Line drawing (above) of Tillyard, 1919d, text-fig. 16.



Family CHILIOCYCLIDAE Evans, 1956

Chiliocycla Tillyard, 1919d [*scolopoides*]

Forewing: basal 0.6 of wing with strong sculpture of flat circular tubercles; R4+5 arising a little before half length; r-m above median cell; M 3-branched dichotomously; M1+2 joined to M3 by cross vein; Cu1 strong with distal fork; a cross vein joins Cu1a to M3+4 behind median cell.


scolopoides Tillyard, 1919d; UTrias; Mt Crosby and Blackstone Fm; Ipswich.

Mesoscytina Tillyard, 1919d [*australis*]

Forewing: Sc waved; R gives off first branch to costal margin 1/3 length from apex, then divided into 4 short branches; M branched at midlength; 2 crossveins from R4+5 and M1+2; cross veins between branches of M; Cu1 branched distally; Cu2 strong, straight.

australis Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.

affinis Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.



Mesoscytina australis; GSQI112a; wing venation from Tillyard, 1919d, text-fig. 5.

Family CICADELLIDAE Latreille, 1802

Eurymelidium Tillyard, 1919d [*australe*]

Forewing: Sc close to costal margin; R, M, Cu basally fused; R giving 1 costal veinlet before Rs then 2 more before apex; Rs incorporated with M1+2 in part; M branches about midlength of median cell and becomes stronger beyond the cell; Cu1 forked distally; A1 wavy; A2 encloses anal angle.


australe Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.

Mesothymbris Evans, 1956 [*perkinsi*]

Forewing: small posterior expansion distally; apex not evenly rounded; sides not parallel; R1 4- or more branched; Rs simple; M1+2 straight, parallel to margin; M3+4 curving sharply towards m-cu.

perkinsi Evans, 1956; UTrias; MtCrosby Fm; Ipswich.

woodwardi Evans, 1956; UTrias; MtCrosby Fm; Ipswich.



Mesothymbris perkinsi; UQC889; 7mm long (left). Line drawing of Evans, 1956, fig. 5d.



Austroscarites gen. nov. [= *Scarites* Hamilton, 1992 not Fabricius, [*femoratus*]

Head narrower than pronotum; frons large, inflated, with indications laterally of muscle impressions; clypellus short, broad; lora long, upper edge poorly defined but probably not reaching antennal pits. Antennae small, slender, set near frons well removed from eyes. Hind coxae large, transverse, not jointed by a mesal peg. Middle and hind femora of similar length, fore femur distinctly shorter. Hind tibia slender, 50% longer than femur. Abdomen robust; laterotergites wider than long; terminal 3 segments much narrower than basal segments

femoratus Hamilton, 1992; LCret; Koonwarra FB; Gippsland.


Triassocotis Evans, 1956 [*australis*]

Tegmen: long, narrow; R1 4-branched; Rs simple; M 4-branched; enclosed cell between M1+2 and M3+4.

australis Evans, 1956; UTrias; MtCrosby Fm; Ipswich.

stricta Evans, 1961; UTrias; MtCrosby Fm; Ipswich.

amplicata Evans, 1961; UTrias; MtCrosby Fm; Ipswich.




Triassocotis australis; UQC1554; 11mm long (right). Line drawing of Evans, 1956, fig. 5L.

Austroscarites femoratus; MUGD3578; 5mm long. Line drawing (right) of Jell & Duncan, 1986, fig. 28C.




Austroscarites femoratus; NMVP32283; 5mm long (right). Line drawing of Jell & Duncan, 1986, fig. 20D.





Triassojassus proavitus; GSQI191a; 5.8mm long (left). Line drawing of Tillyard, 1919d, text-fig. 18.



Homopterulum jelli; NMVP103103; x9. Line drawing (above) of Hamilton, 1992, fig.6.



Ligavena gracilipes; NMVP103350; 11mm long.

Triassojassus Tillyard, 1919d [*proavitus*]

Tegmen: costal margin arched basally; Sc absent; Cu1 leaving R+M near base; R+M fused for more than 0.25 length; R 3-branched; cross vein between R2+3 and R4+5 closing radial cell; r-m weak; M 3-branched; small median cell enclosed; crossveins between branches of M and Cu1 forming 4 cells; 3 cross veins m-cu; A1 wavy; A2 enclosing anal angle. **proavitus** Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.

Family **JASCOPIDAE** Hamilton, 1971

Homopterulum Hamilton, 1992 [*Cercopidium signoretti* Westwood, 1854]

Head narrower than pronotum; frons large, inflated, with indications laterally of muscle impressions; clypellus short, broad; lora long, upper edge poorly defined but probably not reaching antennal pits. Antennae small, slender, set near frons well removed from eyes. Hind coxae large, transverse, not jointed by a mesal peg. Middle and hind femora of similar length, fore femur distinctly shorter. Hind tibia slender, 50% longer than femur. Abdomen robust; laterotergites wider than long; terminal 3 segments much narrower than basal segments

jelli Hamilton, 1992; LCret; Koonwarra FB; Gippsland.

Family **LIGAVENIDAE** Hamilton, 1992

Ligavena Hamilton, 1992 [*gracilipes*]

Head much narrower than pronotum; frons with distinct muscle impressions; clypellus broad basally; lora not reaching antennal pits. Pronotum with divergent lateral margins. Tegmina with complex, weak, apical venation defining numerous apical cells. Hind wings nearly as long as tegmina. Abdomen large; laterotergites about as broad as long; pregenital sternite medially excavated; ovipositor slender, apex slightly exceeding pygofer. **gracilipes** Hamilton, 1992; LCret; Koonwarra FB; Gippsland.

prosboloides Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

dunstani Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.


Family **CIXIIDAE** Spinola, 1838

Mesocixioides Tillyard, 1922b [*termioneura*]
Forewing: R1 unbranched; R2+3 sends a series of veinlets to the costa distally; costal area broad; median cell complete, small, placed distally; Cu1 with small distal fork.

termioneura Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.


orthoclada Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

brachyclada Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.





Mesocixius Tillyard, 1919d [*triassicus*]
Forewing: widest near base; R divides dichotomously twice; R1 connected to costal margin by 2 veinlets; R2+3 branches into 2 simple veins; M 6-branched; M3+4 4-branched; closed median cell defined by strong m-cu; Cu1 forked before midlength; crossveins present between R1-R2, R4+5-M1+2, M-Cu1a.

triassicus Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.



Triassocixius Tillyard, 1919d [*australicus*]
Forewing: series of oblique costal veinlets; first 2 branchings of R close together; r4+5-m and r3-r4+5; M 2-branched; m-cu present.

australicus Tillyard, 1920; UTrias; Blackstone Fm; Ipswich.



Ligavena gracilipes; NMVP103350; line drawings of Hamilton, 1992, figs 9, 10.



Mesocixioides termioneura; GSQI88a; 12.5mm long. Line drawing (above left) of Tillyard, 1922a, text-fig. 82.





Mesocixius triassicus; GSQI215; 10mm long. Line drawing (left) of Tillyard, 1919d, text-fig. 11.



Triassocixius australicus; GSQI267a; 10.5mm long. Line drawing (left) of Tillyard, 1919d, text-fig. 12.



Dunstania pulchra; GSQI2a; 20mm long. Line drawing (right) of Tillyard, 1916, pl. 3, fig. 6.



Dunstaniopsis triassica; GSQI107a; 31.5mm long. Line drawing (above) of Tillyard, 1918e, text-fig. 18.




Paradunstania affinis; GSQI147; 17mm long. Line drawing (right) of Tillyard, 1918e, text-fig. 20.

Family DUNSTANIIDAE Tillyard, 1916

Dunstania Tillyard, 1916 [*pulchra*]

Forewing: corium pitted all over; distinct break (node) in costal margin; R1 weak, 2-branched; R1b ending well before apex; Rs at apex; 2 closed cells just beyond node in wing, one between M1+2 and M3+4 the other between 2 branches of R1+2 which divides then reunites.


pulchra Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.



Dunstaniopsis Tillyard, 1918e [*triassica*]

Hemelytron: smooth corium; membrane lightly pitted; node less prominent than in *Dunstania*; R1 2-branched, long, parallel; Rs simple; r-m present; M 3-branched; Cu1 2-branched.

triassica Tillyard, 1918e; UTrias; Blackstone Fm; Ipswich.



Paradunstania Tillyard, 1918e [*affinis*]

Hemelytron, larger than *Dunstania*. Crossveins basally between M1 and lowest branch of R absent; crossveins developed not far from dividing line between R1 and R2 and between R3 and R4+5; R4 and R5 unite distally; at same level M receives a curving branch from above.

affinis Tillyard, 1918e; UTrias; Blackstone Fm; Ipswich.

Family **DYSMORPHOPTILIDAE**

Handlirsch, 1906

Dysmorphoptiloides Evans, 1956 [*elongata*]



Forewing: evenly punctate, narrow margin; costal border emarginate near apex of R1; apex narrowly rounded; R parallels costal margin and connected to it by series of parallel veins; R1 3-branched; Rs may be separate from or incorporated with R1; M 4-branched apically, joined basally to Cu1; Cula longer than Culb; cross veins r, r-m, m-cu may be present.

elongata Evans, 1956;

UTrias; Mt Crosby Fm, Ipswich.

parva Evans, 1956;

UTrias; Mt Crosby Fm, Ipswich.



Dysmorphoptiloides elongata; UQC686; 16mm long. Line drawing (above) of Evans, 1956, fig. 17A.

Family **EOSCARTERELLIDAE** Evans, 1956

Belmontocarta Evans, 1958 [*perfecta*]



Forewing: rugose, expanding laterally; Sc short, curved, at base of R+M; R1a single; R1b branched, linked to Rs by oblique cross vein; M 4-branched; Cu1 basally parallel with R+M, joined to base of M by short cross vein; Cu1 evenly arched; Culb aligned with margin; r-m and m-cu present.

perfecta Evans, 1958; UPerm; Newcastle CM; Sydney.


Eoscarterella Evans, 1956 [*media*]

Forewing: rugose, expanded distally; R1 with >2 branches; Rs arises 0.3 of distance from base; R and M fused basally; M 4-branched; Cu1 not fused basally with M and proximally straight; Cula arched, longer than Culb.


media Evans, 1956; UTrias; MtCrosby Fm; Ipswich.



Belmontocarta perfecta; AMF47199; 10.5mm long. Line drawing (above) of Evans, 1958, fig. 5.





Eoscarterella media; UQC1110; 10.5mm long. Line drawing (left) of Evans, 1956, fig. 18C.



Eoscarteroides bryani; UQC172; 12mm long. Line drawing (above) of Evans, 1956, fig. 18A.



Mesojassus ipsviciensis; GSQI33; 6mm long. Line drawing (right) of Tillyard, 1916, pl. 2, fig. 7.



Eoscartoides Evans, 1956 [*bryani*]

Forewing: complete marginal border; R and M basally arched; R1a and R1b both forked; M 3-branched; M3+4 unbranched; Cu slightly arched basally, meeting Cu1b apically; Cu1a, Cu1b of equal length.


bryani Evans, 1956; UTrias; Mt Crosby Fm, Ipswich.

Family **EURYMELIDAE** Amyot & Serville, 1843

Mesojassus Tillyard, 1916 [*ipsviciensis*]

Forewing: R, M and Cu1 fused basally; Cu1 arising strongly; M arising weakly; R without longitudinal branches; 3 costal veinlets; M1+2 simple; closed cell formed by cross vein from M1+2 to M3+4; Cu1 forked distally; m-cu present; A1 wavy; A2 encloses anal angle.

ipsviciensis Tillyard, 1916; UTrias; Blackstone; Ipswich.



Family **HYLICELLIDAE** Evans, 1956

Eochiliocycla Davis, 1942 [*angusta*]

Forewing: Sc weak, parallel to base of R; R1 straight, ending at a reentrant angle on costal margin at half wing length; R1b weak; Rs simple, arising from R1 just before it bifurcates; M simple, arising from base of R; 2 rs-m; between these a cross vein (ir) runs to R1b; Cu arises separately from M and R, forks immediately; m-cu present; Cu1b sigmoidal; 1cu crossvein joins Cu2 to Cu1b; 3 strong anal veins arise separately from Cu and distally confluent.

angusta Davis, 1942; UPerm; Newcastle CM; Sydney.

Eochiliocycla angusta;


AMF39793; 5.7mm long. Line drawing (above left) of Davis, 1942, fig. 3.

Hylicella Evans, 1956 [*colorata*] [= *Hylicellites* Becker-Migdisova, 1962 (type *Hylicella reducta* Evans, 1956)]

Forewing: M complete or reduced and with M3+4 in part incorporated with Cu₁a; Cu₁ with a steep basal bend; cross veins rs-m and one between arms of M.

colorata Evans, 1956; UTrias; MtCrosby Fm; Ipswich.

reducta Evans, 1956; UTrias; MtCrosby Fm; Ipswich.



Hylicella colorata; UQC1841; 11mm long. Line drawing (above) of Evans, 1956, fig. 7A.

Family **IPSVICIIDAE** Tillyard, 1919d

Ipsvicia Tillyard, 1919d [*jonesi*]

Forewing: rugose, pitted, narrowing apically; lacking Rs; M and Cu₁ form single vein; Y-vein in clavus; costal fracture usually present from fork of R+Sc and M+Cu₁ to costal margin.

jonesi Tillyard, 1919b; UTrias; Blackstone Fm; Ipswich.


maculata Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.

acutipennis Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.


Ipsviciopsis Tillyard, 1922b [*elegans*] like *Ipsvicia* but with Sc as a sigmoidal vein separate from R and joined to it by 3 cross veins.

elegans Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

magna Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.



Ipsvicia jonesi; GSQ1122a; 14.2mm long (above). Line drawing of Tillyard, 1919d, text-fig. 13.




Ipsviciopsis elegans;
GSQ1178a;
12.5mm long
(right). Line
drawing(above)
of Tillyard,
1922b, text-fig.
85.





Mesogereon neuropunctatum;
GSQI19a; 23mm
long.



Mesogereon superbum; GSQI169; 44.5mm long.
Line drawing (right) of Tillyard, 1922b, text-fig.



Family MESOGEREONIDAE Tillyard, 1921

Mesogereon Tillyard, 1916 [*neuropunctatum*]

Forewing: R₁ and R_s long, parallel, simple; M 4-branched; long cross vein m-cu joins M₄; a closed cell above this cross vein; in the cell are remains of an archidictyon; Cu₁ weak basally, Cu_{1a} and Cu_{1b} simple; Cu₂ in deep groove, dividing off clavus; A₁ and A₂ simple; cross veins c-m and sometimes cu-a present.


neuropunctatum Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

superbum Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.


compressum Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

affine Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

shepperdi Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.



Mesogereon shepperdi;
GSQI97; 21mm
long (left). Line
drawing of
Tillyard, 1922b,
text-fig. 71.




Family PALAEONTINIDAE Handlirsch, 1906


Fletcheriana Evans, 1956 [*triassica*]

Forewing: costal space wider proximally, with several oblique veinlets; nodal line thickened between R_s and M; M 4-branched; nodal line crosses M at its first fork; m-cu sinuate; r-m present; Cu₁ sharply flexed basally.

triassica Evans, 1956; UTrias; MtCrosby Fm; Ipswich.



Fletcheriana triassica; forewing AMF39166 (above), 58mm long; hindwing AMF30971 (right), 40mm long. Line drawing (above) of Evans, 1956, fig. 21F.



Family **PARAKNIGHTIIDAE**
Evans, 1950

Paraknightia Evans, 1943b
[*magnifica*]

Pronotum with large well-developed pronotal expansions, narrowly produced anteriorly. Forewing: Cu1 proximally parallel to M and R; C thickened; Sc short, transverse; nodal line from margin to R-M fork; R parallel to C most of way; A1 and A2 parallel; rugose except smooth area close to apex.


magnifica Evans, 1943b;
UPerm; Newcastle CM; Sydney.

Family **PINCOMBEIDAE**
Tillyard, 1922


Pincombea Tillyard, 1922a
[*mirabilis*]

Sc absent; R, M and Cu1 meeting basally; r-m present; R 2-branched; M 3-branched; Cu1 2-branched.

mirabilis Tillyard, 1922a;
UPerm; Newcastle CM; Sydney.



Paraknightia magnifica; AMF40450; 17mm long. Line drawing of Evans, 1943b, fig. 24b





Pincombea mirabilis; AMF28464; 3mm long (right). Line drawing of Tillyard, 1922a, fig. 2.



Eupincombea Davis, 1942 [*postica*]

Hindwing: very small; Sc short, near costa; R strong for basal quarter then divided into 3 namely R, M and Cu1; R1 runs forward to anterior margin; Rs concave forward; M+Cu1 simple, slightly convex forward; Cu2 and A1 arise together then diverge immediately.


postica Davis, 1942; UPerm; Newcastle CM; Sydney.




Eupincombea postica; AMF39794; 2.6mm long. Line drawing (left) of Davis, 1942, fig. 4.




Protopincombea obscura; AMF40075; 4.2mm long.
Line drawing (above right) of Evans, 1943b, fig. 44.



Anomaloscytina incompleta (hindwing);
AMF40009; 10mm long.



Anomaloscytina metapteryx (hindwing);
AMF39792; 7.3mm long. Line drawing of Davis,
1942, fig. 2.



Austroprosbole maculata; AMF40048; 13mm long.
Line drawing (right) of Evans, 1943b, fig. 2

Protopincombea Evans, 1943b [*obscura*]

Tegmen: broad; clavus short, narrow; R parallel to costal margin; 2 cross veins from Rs to M; m-cu present; M 3-branched; M and Cu1 arise from same point on R but lack common stem; anals forming Y-vein.

obscura Evans, 1943b; UPerm; Newcastle CM; Sydney.

Family **PROSBOLIDAE** Handlirsch, 1906

Anomaloscytina Davis, 1942 [*metapteryx*]

Hindwing: Sc short but distinct; R1 with an articulation at half wing length where Rs originates from it; R1a normal to R1 and weak; Rs simple, with possible crossvein to R1b; M 2-branched, connected to Cu1a by possible crossvein; Cu1 arising from base of R; Cu2 connected by 1 oblique vein to base of Cu1 and by another to A1; A1+A2 straight.


metapteryx Davis, 1942; UPerm; Newcastle CM; Sydney.


incompleta Evans, 1943b; UPerm; Newcastle CM; Sydney.

Austroprosbole Evans, 1943b [*maculata*]

Forewing: long, broad, delicate; a break in curvature of the costal margin occurs at R1a; costal space wide; Rs bent and in close contact with M1; M 4-branched; clavus broad; A1 and A2 meet distally; anal margin not in alignment with rest of hind margin.

maculata Evans, 1943b; UPerm; Newcastle CM; Sydney.





Mesojassula Evans, 1956 [*marginata*]

Forewing: costal margin with marked medial depression; R1a, R1b and Rs present; M unbranched; Cu1 has 2 equal branches; marginal vein present.

marginata Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.

Mitchelloneura Tillyard, 1921b [*permiana*]


Forewing: Sc short, fused with R1, ending halfway along costal margin; R1 strong; Rs branching 0.25 of distance from apex; R3+4 joining R1; 2 cross veins from R4 to R2+3 and R1; M arising from R near base, 3-branched; M3+4 unbranched; Cu1a 2-branched; r-m and m-cu present.

permiana Tillyard, 1921b; UPerm; Newcastle Fm; Sydney.

Permodipthera Tillyard, 1926a [*robusta*]


Forewing: C strongly curved basally, then running fairly straight to R1a; apex well-rounded basal cell very small; Rs curved; nodal line anterior from fork in M; Cu1 strongly curved down to come close to Cu2.

robusta Tillyard, 1926a; UPerm; Newcastle CM; Sydney.







Belpsylla reticulata; AMF39992; 3.8mm long. Line drawing (right) of Evans, 1943b, fig. 42.



Clavopsyllidium minutum; AMF39800; 2.4mm long. Line drawing (above) of Davis, 1942, fig. 9.



Permopsyllidium mitchelli; AMF28020; 3.7mm long. Line drawing (right) of Tillyard, 1926a, text-fig. 25.



Family **PROTOPSYLLIDIIDAE** Carpenter, 1931

Belpsylla Evans, 1943b [*reticulata*]

Forewing: long, apically broad; R1a and R1b curve towards costa; 2 cross veins r-m; M r-branched, all straight; Cu1 narrowly forked; clavus with well-developed Y-vein.

reticulata Evans, 1943b; UPerm; Newcastle CM; Sydney.

Clavopsyllidium Davis, 1942 [*minutum*]

Very small. Forewing: Sc absent; Rs arising from R at 1/3 wing length; r-m absent; M+Cu1 arising from midlength of R; M 3-branched; Cu1a curved back, finishing closer to Cu1b than in most genera; Cu straight, narrow, independent of Cu1; A1+A2 with common origin, diverging then meeting again before margin.


minutum Davis, 1942; UPerm; Newcastle CM; Sydney.


Permopsyllidium Tillyard, 1926a [*mitchelli*]

Forewing: Sc absent; R+M scarcely arched; R 2-branched; Rs relatively short, curved; M 3-branched; r-m present; m-cu absent.

mitchelli Tillyard, 1926a; UPerm; Newcastle CM; Sydney.

affine Tillyard, 1926a; UPerm; Newcastle CM; Sydney.







Permopsyllidops stanleyi; AMF39799; 3mm long (right). Line drawing (above) of Davis, 1942, fig. 9

Permopsyllidops Davis, 1942 [*stanleyi*]

Forewing: Sc absent; R divides at 1/3 wing length; R1a normal to R1; Rs simple but drawn out medially at r-m; M+Cu arising at base of R; weak vein running from fork of M and Cu1 to R1 (original M+Cu1); M forked before m-cu; Cu1a curved back; Cu2 absent; A1 short, posteriorly directed.


stanleyi Davis, 1942; UPerm; Newcastle CM; Sydney.



Permopsylloides Evans, 1943b [*insolita*]

Tegmen: uniform width; costal space wide; R1a meeting costal margin beyond midlength; Rs curved forward; clavus wide, long; anal veins form a Y-vein.



insolita Evans, 1943b; UPerm; Newcastle CM; Sydney.



Permothea Tillyard, 1926a [*latipennis*]

Tegmen: pterostigmatic area flatly triangular, strongly chitinised; Rs curved near base, then straight; r-m present; m-cu absent; M 3-branched; Cu1 curving forward to touch M near base, 2-branched.


latipennis Tillyard, 1926a; UPerm; Newcastle CM; Sydney.




Permothea latipennis; AMF28022; 1.5mm long (right). Line drawing (above) of Tillyard, 1926a, text-fig. 27.

Permopsylloides insolita; AMF40727; 4mm long (part and counterpart). Line drawing (above) of Evans, 1943b, fig. 46.





Permotheella scytinopteroides; AMF39797; 3.8mm long. Line drawing (above) of Davis, 1942, fig. 7.



Protopsyllidium australe; AMF19790; 2.8mm long. Line drawing (right) of Tillyard, 1926a, text-fig. 24.



Protopsylllops minuta; AMF40005; 3.1mm long. Line drawing (right) of Evans, 1943b, fig. 37.

Permotheella Davis, 1942 [*scytinopteroides*]

Forewing: Sc short, faint, parallel to base of R; pterostigma and costal margin well-developed; R_s arising adaxial to midlength, curved forward distally; M+Cu₁ arising at 0.16 wing length; M forking into M₁+2 which is connected to R_s by r-m, then dividing into M₁, M₂ and M₃+4, the latter a single sinuous vein. Cu_{1a} curved; Cu₂ separating a distinct clavus with A₁ and A₂; A₁ and A₂ diverging near base, coalescing for half length.


scytinopteroides Davis, 1942; UPerm; Newcastle CM; Sydney.

Protopsyllidium Tillyard, 1926a [*australe*]

Tegmen: Sc absent; R+M arched; R straight to end of R_{1b}; pterostigma 2.5 times as long as the oblique R_{1a}; M and Cu₁ 2-branched; Cu₂ straight; no cross veins except a short cu-a basally.

australe Tillyard, 1926a; UPerm; Newcastle CM; Sydney.


sinuatum Davis, 1942; UPerm; Newcastle CM; Sydney.




Protopsylllops Evans, 1943b [*minuta*]

Tegmen: short, wide; R₁ close to costal margin, unbranched; R_s long; first branching of M close to junction of M and Cu₁; M 3-branched; Cu_{1b} parallel to Cu₂.

minuta Evans, 1943b; UPerm; Newcastle CM; Sydney.






Psocopsyllidium Davis, 1942 [*media*]

Forewing: Sc absent; Rs from R at wing midlength; M+Cu1 arising from R at 0.25 winglength; M and Cu separate early; M dividing same distance further; M1, M2 and M3+4 present; r-m between Rs and M1; Cu1a curved backwards; Cu2 and A1 arising together, diverging initially, then parallel and converging gently distally; cu2-r present; A2 short.
media Davis, 1942; UPerm; Newcastle CM; Sydney.



Psocopsyllidium media; AMF39796; 4.6mm long. Line drawing (left) of Davis, 1942, fig. 6.



Psocoscytina Davis, 1942 [*bifida*]

very small. Forewing: Sc short, weak, parallel to base of R; R1a curving strongly forward; several obscure veinlets in pterostigmatic area between R1a and R1b; Rs arising just before midlength of wing, branching distally; M+Cu1 arising together near base of R1; r-m strong; M forked just beyond r-m to M1+2 and M3+4; Cu1 2-branched; Cu2 joined to base of R by crossvein.

bifida Davis, 1942; UPerm; Newcastle CM; Sydney.





Psocoscytina bifida; AMF39790; 3.9mm long. Line drawing (above) of Davis, 1942, fig. 1.

Psyllidella Evans, 1943b


[*magna*]

Tegmen: broad; costal margin thickened, sinuate; R1 straight; Rs leaving R level with first fork of M.


magna Evans, 1943b; UPerm; Newcastle CM; Sydney.



Psyllidella magna; AMF40043; 5mm long. Line drawing (left) of Evans, 1943b, fig. 36.




Psyllidiana davisia; AMF40711, 3mm long. Line drawing (right) of Evans, 1943b, fig. 33.



Psyllidiana Evans, 1943b [*davisia*]
Tegmen: narrow; R1 and Rs straight; M 3-branched; Cula longer than basal portion of Cu1.
davisia Evans, 1943b; UPerm; Newcastle CM; Sydney.




Triassopsylla plecioides; AMF39331; 4.6mm long. Line drawing (right) of Tillyard, 1918b, text-fig. 16




Triassopsylla Tillyard, 1918b [*plecioides*]
Forewing: broad; apex rounded; R1 2-branched; Rs simple; M 3-branched; membrane smooth.


plecioides Tillyard, 1918b; MTrias; Ashfield Fm; Sydney.



Triassothea analis; UQC1590; 2.8mm long. Line drawing (above) of Evans, 1956, fig. 25F.



Triassothea Evans, 1956 [*analis*]
Forewing: Rs arising from R very soon after M+Cu; M 2-branched distally; Cu1 2-branched; clavus small; single anal vein.
analis Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.

**Tripsyllidium** Evans, 1956 [*wadei*]

R1a meeting costal margin near midlength; Rs ending at apex; M unbranched; no cross veins; clavus small; one anal vein only.

wadei Evans, 1956; UPerm; Newcastle CM; Sydney.

Family **SCYTINOPTERIDAE** Handlirsch, 1906

Crosbella Evans, 1956 [*elongata*]



Forewing: numerous terminal accessory veins and many supplementary cross veins.

elongata Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.

alata Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.



Tripsyllidium wadei; AMF30980; 3mm long. Line drawing (left) of Evans, 1956, fig. 25H.




Crosbella elongata; UQC2125; 12mm long. Line drawing (above) of Evans, 1956, fig. 5H.

Elliptoscarta Tillyard, 1926a [*ovalis*]

Forewing: broadly oval with wide costal space; R1 evenly branched; Rs arises closer to branch of R1 than to origin of M; M 5-branched; Cu1 3-branched; m-cu and r-m present.


ovalis Tillyard, 1926a; UPerm; Newcastle CM; Sydney.




Elliptoscarta ovalis; AMF41392; 5.6mm long. Line drawing (left) of Tillyard, 1926a, text-fig. 15.




Homaloscytina plana; AMF19783; 6.4mm long.
Line drawing (right) of Tillyard, 1926a, text-fig. 14.



Mesodipthera grandis; GSQ1213; 12mm long. Line drawing (above) of Tillyard, 1919d, text-fig. 7.



Mesonirvana abrupta; UQC1652; 9mm long (left).
Line drawing (right) of Evans, 1956, fig. 5E.



***Homaloscytina* Tillyard, 1926a [*plana*]**

Forewing: R1a almost transverse to mainstem of R1 that continues as R1b in line to margin; Rs arises well beyond midlength; apical margin bluntly rounded; pterostigma moderately long; M 4-branched; Cu1 2-branched, straight from basal cell to fork.

plana Tillyard, 1926a; UPerm; Newcastle CM: Sydney.

***Mesodipthera* Tillyard, 1919d [*grandis*]**


Forewing: R branches dichotomously just before the branching of M; branches of R and M branched irregularly distally; slightly impressed division of the wing in a line from end of R1 to end of clavus.

grandis Tillyard, 1919d; UTrias; Blackstone Fm; Ipswich.

***Mesonirvana* Evans, 1956 [*abrupta*]**

Forewing: R1 4-branched; M1+M2 of equal length; M3+4 single vein; Cu1 a straight line as far as m-cu; Cu1a and Cu1b apically parallel with each other.

abrupta Evans, 1956; UTrias; MtCrosby Fm; Ipswich.



Orthoscytina Tillyard, 1926 [*mittelli*]

Forewing: expanding distally; apex bluntly rounded; R+M arched above basal cell; R strong beyond cell, running forward to costal margin; R_{1b} with two accessory veins across long pterostigmatic area; M 3- to 5-branched; Cu_{1a} with series of accessory veins running to margin towards Cu_{1b}.

mittelli Tillyard, 1926; UPerm; Newcastle CM; Sydney.

quinquimedia Tillyard, 1926; UPerm; Newcastle CM; Sydney.

indistincta Tillyard, 1926; UPerm; Newcastle CM; Sydney.

subcostalis Tillyard, 1926; UPerm; Newcastle CM; Sydney.


irregularis Tillyard, 1926; UPerm; Newcastle CM; Sydney.

belmontensis Tillyard, 1926; UPerm; Newcastle CM; Sydney.


obliqua Tillyard, 1926; UPerm; Newcastle CM; Sydney.

pincombei Tillyard, 1926; UPerm; Newcastle CM; Sydney.

tetraneura Tillyard, 1926; UPerm; Newcastle CM; Sydney.



Orthoscytina mittelli; AMF28026; 10mm long. Line drawing (above) of Tillyard, 1926a, text-fig. 4.




Permobrachus magnus; AMF40029; 16mm long. Line drawing (above left) of Evans, 1943b, fig. 1.



Permoglyphis Tillyard, 1926 [*belmontensis*]

Forewing: C strong, with prominent node; evenly rounded apically; basal tegula swollen; Sc short, weak; basal cell long, narrow; faint nodal line to Cu₂; no strong R_{1a}; series of pterostigmatic veinlets present; R_s arising just before nodal line, distally forked; Cu₁ convex posteriorly, 2-branched; clavus triangular; A₁ long, gently sigmoidally curved; A₂ short, enclosing anal angle.



belmontensis Tillyard, 1926a; UPerm; Newcastle CM; Sydney.



Permoglyphis belmontensis; AMF28023; 9.4mm long. Line drawing (left) of Tillyard, 1926a, text-fig. 19.



Stenoglyphis kimblensis; AMF41207; 36mm long; Line drawing (above) of Evans, 1947, fig. 1.



Permojassus australis; forewing AMF19782; 6mm long (left upper); hindwing AMF19781(left lower). Line drawings (above) of Tillyard, 1926a, text-figs 2 and 3.

Permojassus Tillyard, 1926a [*australis*]

Forewing: Sc absent; R1b and pterostigmatic area short; M 4-branched; m-cu present; Cu1 curved forward close to M basally, 2-branched; Cu2 straight; A1 wavy; A2 enclosing anal angle. Hindwing: costal space wide; costal margin convexly rounded; M 2-branched.

australis Tillyard, 1926a; UPerm; Newcastle CM; Sydney.

dubius Tillyard, 1926a; UPerm; Newcastle CM; Sydney.

Stenoglyphis Evans, 1947 [*kimblensis*]

Tegmen: long, narrow, with circular basal lobe; M1 4-branched; M2, M3, M4 undivided; Cu1 short; Cu1a bent; reticulate cross veins between several main veins; clavus 0.3 wing length.

kimblensis Evans, 1947; UPerm; Newcastle CM; Sydney.

Stenoscytina australiensis; AMF28027; 7.2mm long. Line drawing (left below) of Tillyard, 1926a, text-fig. 13.

Stenoscytina Tillyard, 1926a [*australiensis*]
Tegmen: narrow; R curving forward to end of R1a; costa evenly curved from base to apex; Rs arising distally; M 4-branched; Cu1 curving down from basal cell close to Cu2; r-m and m-cu present.
australiensis Tillyard, 1926a; UPerm; Newcastle CM; Sydney.


Triassoscelis Evans, 1956 [*anomala*]

Tegmen: Rs 2-branched; R1 multibranched; M 4-branched; enclosed cell between Rs and R1b; r-m and m-cu present.

anomala Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.



Triassoscelis anomala; UQC976; 12mm long (counterpart - right). Line drawing (above right) of Evans, 1956, fig. 5F.



Triassoscytina incomplete; UQC1559; 10.8mm long (right). Line drawing of Evans, 1956, fig. 2B.

Triassoscytina Evans, 1956

[*incompleta*]

Forewing: complete venation like Homaloscytina but M dividing at same distance from wing base as origin of Rs; M dividing at same distance from wing base as origin of Rs.

incompleta Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.

Triassoscytinopsis Evans, 1956

[*stenulata*]

Forewing: parallel-sided; R1 with 4 or more parallel branches; Rs simple or 2-branched; M 4-branched; Cu1 with normal venation, with sharp proximal bend; anal veins separate for full length.


stenulata Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.

aberrans Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.

paranotalis Evans, 1956; MTrias; Hawkesbury Sst; Sydney.



Triassoscytinopsis paranotalis; AMF38265; 21mm long. Line drawing of Evans, 1956, fig. 5A.



Family **STENOVICHIIDAE** Evans, 1956

Apheloscyta Tillyard, 1922b [*mesocampta*]

Forewing: Rs originating close to apex; M with posterior convexity just beyond midlength; M and Cu 2-branched; m-cu, r-m and clavus present; A2 close to basal posterior margin, running close to the posterior margin of clavus.

mesocampta Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.



Apheloscyta mesocampta; GSQ198a; 10mm long. Line drawing (above left) of Tillyard, 1922b, text-fig. 78.




Palaeoviccia Evans, 1943b [*incerta*]

Forewing: broad, narrowing apically; R and R1b parallel costal margin; Rs short, straight; M 3-branched; M1+2 not branched; M and Cu1 fused basally; clavus broad, anals forming Y vein.

incerta Evans, 1943b; UPerm; Newcastle CM; Sydney.



Permagra distincta; AMF39866; 7mm long. Line drawing of Evans, 1943a, text-fig. 1.



Palaeoviccia incerta; AMF39999; 5.7mm long (above left). Line drawing of Evans, 1943b, fig. 26.

Permagra Evans, 1943a [*distincta*]


Forewing: costal margin thickened; Sc short and curved to meet R; R2+3 straight to apex; M 2-branched; Cu with 2 short branches.

distincta Evans, 1943a; UPerm; Newcastle CM; Sydney.

Permocentrus Evans, 1956 [*Permoscarta trivenulata* Tillyard, 1926a]

Forewing: 2 oblique veins between R1a and R1b; R1 linked to Rs by 3 cross veins; r-m present; M 2-branched; Cu1 not arched basally, separate from M.

trivenulatus Tillyard, 1926a; UPerm; Newcastle CM; Sydney.



Permocentrus trivenulatus; AMF19787; 6mm long. Line drawing (above right) of Tillyard, 1926a, text-fig. 17.




Permoscarta


Tillyard, 1918b [*mittelli*]

Tegmen: finely tuberculate; nodal line weak, evident near midlength; series of oblique costal veinlets; M unbranched; 2 m-cu; M leaving R near base, fused to Cu for short distance then diverging again; Cu straight, simple, parallel to A1.

mittelli Tillyard, 1918b; UPerm; Newcastle CM; Sydney.



Permoscarta mittelli; AMF39316; 6.4mm long (above right). Line drawing of Tillyard, 1918b, text-fig. 2a.




Permoviccia Evans, 1943b [*obscura*]

Tegmen: short, broad; Sc short, wide, transverse; R and R1b parallel costal margin; Rs widely curved; M and Cu1 fused for short distance basally; r-m and m-cu present.

obscura Evans, 1943b; UPerm; Newcastle CM; Sydney.




Permoviccia obscura; AMF40059; 4.8mm long. Line drawing (above) of Evans, 1943b, fig. 28.


**Stanleyana** Evans, 1943b [*pulchra*]

Tegmen: narrowing apically; R parallel to costal margin; M and Cu1 fused basally; M 3-branched; Cu1 2-branched; m-cu present.

pulchra Evans, 1943b; UPerm; Newcastle CM; Sydney.



Stanleyana pulchra;
AMF40016; 7mm
long. Line
drawing (above
left) of Evans,
1943b, fig. 25.



**Stenoviccia** Evans, 1943b [*angustata*]

Tegmen: narrowing apically; R parallel to costal margin; M and Cu1 fused close to R; M1 and M2 short; Cu1a twice length of Cu1b.

angustata Evans, 1943b; UPerm; Newcastle CM; Sydney.



Stenoviccia angustata;
AMF40008; 6mm long. Line
drawing (above left) of
Evans, 1943b, fig. 27.



Psyllid indet.; NMVP103240; 2.1mm long. Line
drawing (above) of Jell & Duncan, 1986, fig. 23D.

Family **PSYLLIDAE** Loew, 1879**Psyllid** indet.

Small forewing with broadly truncated apex; pterostigma long, slender; R 2-branched, with fork closer to base than apex; M, Cu 2-branched, with main stem joining just before common petiole joins basal vein, with 2 branches of M almost equal but 2 branches of Cu markedly different in length; clavus well developed, with angular change in posterior margin near wing base.

Psyllid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family
RICANIIDAE
Amyot & Serville,
1843



Scolypopites
Tillyard, 1923a
[*bryani*]

Forewing as in
Scolypopa but Sc
shorter, not as strong as
R, reaching only to a
little beyond midwing;
M and R not
completely fused at
base, only 1 series of
gradate veins.

bryani Tillyard,
1923a; Palaeocene;
Redbank Plains Fm;
Ipswich.



Scolypopites bryani; UQF7218; 15mm long. Line drawing (left) of Tillyard, 1923a, text-fig. 1.




FAMILY UNCERTAIN

Mesocicadella Evans, 1956
[*venosa*]

Forewing: series of oblique veinlets
proximal of separation of Rs from R1;
Rs and R1 with accessory veins, M
profusely branched; Cula with another
apical branch, arched forward.

punctata Evans, 1961; UTrias; Mt
Crosby Fm; Ipswich.



venosa Evans, 1956; UTrias; Mt
Crosby Fm; Ipswich.




Mesocicadella venosa; UQC844; 12mm long. Line drawing (above) of Evans, 1956, fig. 51



Knezouria unicus;
QMF18850;
13mm long.



Beaconsiella fennahi, hindwing; AMF30974; 17mm long. Line drawing (above) of Evans, 1963, fig. 5b.



Tricrosbia minuta; QMF6520; 3.2mm long (right). Line drawing (above) of Evans, 1971, fig. 1.

Knezouria Jell, 1993 [*unicus*]

Nymph: wings rudimentary; rostrum slender, extending beyond posterior margin of abdomen; frons large, diamond-shaped. Abdomen inverted beehive; 8 segments, with median longitudinal ridge, with paired areas of small tubercles on each segment. Terminal segment with anal and genital structures in midline, well forward of margin.

unicus Jell, 1993; UTrias; Blackstone Fm; Ipswich.

Order HETEROPTERA


Family FULGORIDAE

Beaconsiella Evans, 1963 [*fennahi*]

All principal veins multibranching; Sc separate from R proximally but joining with R soon after; Rs joined to M by crossvein prior to branching of either vein; Rs diverging from R near base of wing; Rs, M, Cu1 branch initially the same distance from the base of the hindwing; clavus unknown.

fennahi Evans, 1963 MTrias; Hawkesbury Sst; Sydney.

multivenata Evans, 1963; MTrias; Hawkesbury sst; Sydney.




Beaconsiella multivenata, forwing, line drawing of Evans, 1963, fig. 5a.

Tricrosbia Evans, 1971 [*minuta*]

Rs 2-branched, diverging from R slightly proximally to its centre; M1+2 unbranched; M3 and M4 apically separate, with a third vein arising from their common stem; an enclosed cell proximal to the initial branching of M, bounded laterally by pair of m-cua; CuA terminating proximally at base of claval suture, not associated with base of M; clavus with anal veins apically fused into a single vein.

minuta Evans, 1971; UTrias; MtCrosby Fm; Ipswich.



Family **PROGONOCIMICIDAE** Handlirsch, 1906

Actinoscytina Tillyard, 1926a [*belmontensis*]


Forewing: origins of Rs and R1a coincident and distal of midlength of wing; distal branches of M and Cu straight, slightly diverging; costal very wide and strong from base to R1a; R1a transverse; M2 running through middle of wing to margin in straight line; Cu1 concave towards M; M and Cu1 each with 3 distal branches meeting margin equidistance apart.

belmontensis Tillyard, 1926a; UPerm; Newcastle CM; Sydney.

Gelastocorid indet.

Nymph broadly elliptical; head transverse, short, with usual elongate mouthparts extending to rear of thorax; eye large, subrounded, lateral on head; pronotum with convex lateral margins, with several anterior projections adjacent to the eyes, rounded posterolaterally; mesonotum and metanotum subequal in length and only slightly shorter than pronotum. Abdomen with segments 1 and 2 very short medially, with segments thus chevron-shaped; anterior femur arising from the expanded coxa distally; 2nd and 3rd femora arising near midline.




Gelastocorid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Heterochterus Evans, 1971 [*timmsii*]

Forewing. Costal fracture well defined, 1/3 of length from base; R multi-branched, terminating near commencement of the apical appendix; M diverging from common stem with R at angle of costal fracture, with 2 equal branches; M and CuA linked by 3 crossveins; marginal vein meeting claval suture at an acute angle; clavus unknown.

timmsii Evans, 1971; UTrias; Mt Crosby Fm; Ipswich.




Actinoscytina belmontensis; AMF19788, 7mm long. Line drawing (above) of Tillyard, 1926a, text-fig. 16.




Gelastocorid indet., NMVP102519, 4mm long; ventral (above) and line drawing (left).



Heterochterus timmsii; QMF6473; 4.3mm long. Line drawing (left) of Evans, 1971, fig. 1.




Heterojassus membranaceus; QMF3701; 5.1mm long. Line drawing (above) of Evans, 1961, fig. 5B.



Heteronella marksei; QMF3699; 3.8mm long. Line drawing (above) of Evans, 1961, fig. 5A.



Heterojassus Evans, 1961 [*membranaceus*] membranous tegmina; Rs multibranched; M and Cu1 form single vein basally; elongate enclosed cell between M and Cu1 and wide appendix; all veins curved towards anterior corner of apex of tegmen. **membranaceus** Evans, 1961; UTrias; MtCrosby Fm; Ipswich.




Hexascytina transecta; BMNHIn44923; 2mm long. Line drawing (above) of Wootton, 1963, fig. 1.

Hexascytina Wootton, 1963 [*transecta*] Incomplete tegmen: 6 veins extend distally from the composite, transverse vein; transverse vein almost straight, running from anterior to posterior margin of remigium. **transecta** Wootton, 1963; UTrias; Mt Crosby Fm; Ipswich.

Heteronella Evans, 1961 [*marksei*] costal fracture in tegmen; venation in basal half only; in distal half indistinct lines curve towards costal margin; R proximally incorporated in same vein as M; apically R seems to be 3-branched and M is single vein; cu-m wide; 2 branches of Cu elongated. **marksei** Evans, 1961; UTrias; MtCrosby Fm; Ipswich.

Heteroscytina Evans, 1956 [*tillyardi*] Forewing: narrow apically; R1 2-branched or simple; Rs simple; r-m present. **tillyardi** Evans, 1956; UTrias; MtCrosby Fm; Ipswich.



Heteroscytina tillyardi; UQC691; 4mm long (left). Line drawing (above) of Evans, 1956, fig. 29F.

Platyscytinella Evans, 1956 [*paradoxa*]

Tegmen: R1a absent; r-m nearer apex than m-cu1; anterior branch of M straight to apex.

paradoxa Evans, 1956; UTrias; MtCrosby Fm; Ipswich.

Triassodoecus Evans, 1963 [*chinai*]

Pronotal paranota and tegmina large, overlapping apically; R widely separated from costal margin; Rs arising from R1b; M basally joined to R, linked to Rs by oblique cross vein; Cu1 joined to M by cross vein; claval veins forming a Y-vein.

chinai Evans, 1963; MTrias; Hawkesbury Sst; Sydney.



Triassodoecus chinai; AMF46398; 5mm long.



Triassodoecus chinai; line drawing of holotype, of Evans, 1963, fig. 6.

Microscytinella Wootton, 1963 [*radians*]


Composite transverse vein straight, just more than half wing length from the forewing base; M continued in straight line distal to transverse vein by common stem of its first two branches.

radians Wootton, 1963; M Trias; Mt Crosby Fm; Ipswich.


Triscytina Evans, 1956 [*rotundata*]

Tegmen: apex evenly rounded; R1a present; Rs+M1 fused apically; A1 adjacent to claval suture; A2 along hind margin; anals forming Y-vein.


rotundata Evans, 1956; UTrias; Mt Crosby Fm; Ipswich.



Platyscytinella paradoxa; UQC169; 3.2mm long. Line drawing (above) of Evans, 1956, fig. 29C.




Microscytinella radians; BMNHIn44920; 2.2mm long. Line drawing (above) of Wootton, 1963, fig. 2.



Triscytina rotundata; UQC2311; 3mm long. Line drawing (above) of Evans, 1956, fig. 29G.



Anthocorid indet.; NMVP103163; 2.4mm long.
Line drawing(right) of Jell & Duncan, 1986, fig.
23C.



Family **ANTHOCORIDAE** Amyot & Serville,
1843

Anthocorid indet.

Hemelytron with cuneus and embolium, membrane without basal cells, with large clavus and distinct vein close to posterior margin.

Anthocorid indet. Jell & Duncan, 1986;
LCret; Koonwarra FB; Gippsland.

Family **VELIIDAE** Dohrn, 1859

Veliid indet.

Dorsoventrally compressed; head triangular, with bulging lateral eyes, anteriorly obtuse; antennae long, thin, of few segments; legs all long, thin, with femora wider than tibiae, with hindleg longest; abdomen tapering, with wide doublure, with short segments anteriorly becoming longer posteriorly; 6th segment more than half as long as wide.

Veliid indet. Jell & Duncan, 1986; LCret;
Koonwarra FB; Gippsland.

Family **MESOVELIIDAE** Douglas & Scott,
1867

Duncanovelina Jell & Duncan, 1986 [*extensa*]

Head elongate; eyes near midlength of head; antenna of 4 long segments; thoracic segments relatively short, subequal, becoming slightly longer to posterior.

extensa Jell & Duncan, 1986; LCret;
Koonwarra FB; Gippsland.



Veliid indet.; NMVP103235; 3mm long



Duncanovelina extensa Jell & Duncan, 1986;
NMVP103237; 5mm long, with line drawing (above)
from Jell & Duncan, 1986, fig. 24B and
NMVP27044 (right); 5mm long



Family **TRIASSOCORIDAE** Tillyard, 1922b

Triassocoris Tillyard, 1922b [*myersi*]


Hemelytron: short, broad, smooth, shiny, dark; venation faint; 3 main veins on corium – R, M, Cu – fused basally for a short distance; R parallel to costal margin, some distance from it; one veinlet near origin of M then series to C near apex; a costal fracture apparently present.

myersi Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

scutulum Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

ovalis Tillyard, 1923b; UTrias; Blackstone Fm; Ipswich.



grandis Tillyard, 1923b; UTrias; Blackstone Fm; Ipswich.




Triassocoris myersi; GSQI140a; 5.8mm long. Line drawing of Tillyard, 1922, text-fig. 87.




Triassocoris ovalis; GSQI129a; 4.8mm long.



Ademosyne australiensis; GSQI12b; 3.4mm long. Line drawing of Dunstan, 1923, pl.1, fig. 3.



Ademosyne lata; GSQI132a; 2.2 mm, Dunstan, 1923, pl. 1, fig. 5



Ademosynoides obtusa; GSQI9b; 2.4mm long. Line drawing of Dunstan, 1923, pl. 2, fig. 17.

Order COLEOPTERA

Family **ADEMOSYNIDAE** Ponomarenko, 1968

Ademosyne Handlirsch, 1906 [*major*]

Elytron: punctate-striate, costate; without wide border; 8-11 costae.

major Handlirsch, 1906; UTrias; Blackstone Fm; Ipswich.

australiensis Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

olliffi (Handlirsch, 1906); UTrias; Blackstone Fm; Ipswich.

congener Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

cameroni Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

punctata Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

parva Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

intermedia Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

lata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

brevis Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

curvata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

ramocostata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

rugulosa Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

vittamargina Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

adunca Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

wianamattensis Tillyard, 1918b; UTrias; Blackstone Fm; Ipswich.

Ademosynoides Dunstan, 1923 [*minor*]

Elytron: impunctate-striate; 8-10 costae; borders narrow or moderately wide.

minor (Handlirsch, 1906); UTrias; Blackstone Fm; Ipswich.

obtusa (Tillyard, 1916); UTrias; Blackstone Fm; Ipswich.


angustata Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

alternata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

striatella Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

abnormis Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

magnifica Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.



Grammositus Dunstan, 1923 [*bilineatus*]

Elytron: apex blunt; basal margin long; humerus extended; only 2 rows of striae; elongate granules oriented longitudinally.

bilineatus Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

Permosyne Tillyard, 1924 [*belmontensis*]


Elytra with lateral margins not thickened; sutural margin narrow; both margins curved; striae plain or punctate, with 2nd, 3rd and 4th ending on 1st, with 8th ending on 9th; 9th stria (next to sutural margin) arising basally further from margin than from 8th stria; usually vestige of a short 10th stria in basal part of interval between 9th stria and margin.

belmontensis Tillyard, 1924; UPerm; Newcastle CM; Sydney.


affinis Tillyard, 1924; UPerm; Newcastle CM; Sydney.

mitchelli Tillyard, 1924; UPerm; Newcastle CM; Sydney.

pincombeae Tillyard, 1924; UPerm; Newcastle CM; Sydney.




Grammositus bilineatus; GSQI136a; 2.5mm



Permosyne belmontensis; AMF19777; 3.1mm long.
From Tillyard, 1924, text-fig. 2.



Permosyne pincombei; AMF19778; 3.3mm long.



Family **BUPRESTIDAE** Leach, 1815

Lobites Dunstan, 1923 [*tuberculata*]

Elytron: lateral margin bisinuate; sutural margin arcuate; surface with several lines of tuberculation, 3 lines of tubercles or smooth.


tuberculata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

trivittata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.


granulata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.



Lobites tuberculatus; GSQI342a; 5.8mm long. Line drawing (above) of Dunstan, 1923, pl. 5, fig. 41.



Mesostigmodera typica; GSQI61a; 15mm long. Line drawings (right) of Dunstan, 1923, pl.5, fig. 39.



Mesostigmodera Etheridge & Olliff, 1890
[*typica*]

Like *Lobites* but differing in ornamentation, border details and in lacking an incision on the disc.

typica Etheridge & Olliff, 1890; UTrias; Blackstone Fm; Ipswich.

Family **CANTHARIDAE** Thomson, 1864

Cantharid indet.

Head narrow; clypeal region produced forward, with convex anterior; mandibles elongate; pronotum much wider than head, with lateral margin concave anteriorly; elytra soft, elongate, not covering apical segments of abdomen; middle and forelegs similar, with small coxae not touching, expanded femur and long thin tibia; abdomen of 8 uniform segments except for 2 longer posterior ones.

Cantharid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **CERAMBYCIDAE** Latreille, 1802

Willcoxia Dunstan, 1923 [*magnopunctata*]

Elytron: 3 longitudinal and 2 oblique double rows of pits; lateral border wide, pronounced.

magnopunctata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.



Willcoxia magnopunctata; GSQI 261; 10mm long (left). Line drawing of Dunstan, 1923, pl.6, fig. 47.

Family **CURCULIONIDAE** Latreille, 1802

Etheridgea Handlirsch, 1906b [*australis*]



Elytron: small, distally pointed; 13 rows of minute tubercles of variable size and distance from one another.

australis Handlirsch, 1906b; UTrias; Blackstone Fm; Ipswich.

petrica Tillyard, 1916; MTrias; Ashfield Fm; Sydney.



Cantharid indet.; NMVP103331; 7mm long. Line drawing of Jell & Duncan, 1986, fig. 39A.



Etheridgea australis; GSQI31; 9mm long. Line drawing of Dunstan, 1923, pl. 7, fig. 60. [The type specimen is not readily identified in either GSQ or AM

Mesorhynchophora Tillyard, 1916 [*dunstani*]
Elytron: very broad base, tapering to narrow rounded apex; surface convex, with 4 converging striae distally separated by wide flat interstices.


dunstani Tillyard, 1916; MTrias; Ashfield Fm; Sydney.

Tillyardiopsis Dunstan, 1923 [*tuberculata*]
Elytron: sutural border narrow; lateral and humeral borders wide; surface granulate; short deep incision or line of close deep punctae at middle of lateral half.

tuberculata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

granulata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.


variotubercula Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.



Mesorhynchophora dunstani; GSQI27a; 15mm long. Line drawing of Dunstan, 1923, pl. 1, fig. 7




Tillyardiopsis variotubercula; GSQI66a; 6.5mm long.



Tillyardiopsis tuberculata; GSQI133; 7mm long (below). Line drawing of Dunstan, 1923, pl. 7, fig. 57.



Tillyardiopsis granulata; GSQI289a; 6.9mm long.



Family **DASCILLIDAE** Guérin-Ménéville, 1823

Apheloodes Dunstan, 1923 [*obliqua*]
Elytron: rugose lines concentric with lateral borders; sometimes with 2 oblique lines; low convexity.

obliqua Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.


rugosa Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.



Apheloodes obliqua; GSQI143a; 2mm long. Line drawing of Dunstan, 1923, pl. 6, fig. 54.



Leioodes plana; GSQI348; 1.8mm long. Line drawing (right) of Dunstan, 1923, pl. 6, fig. 53.



Reeveana major; GSQI251a; 5.4mm long. Line drawing (above) of Dunstan, 1923, pl. 5, fig. 38.




Tryoniopsis punctata; GSQI250b; 6mm long. Line drawing (right) of Dunstan, 1923, pl. 7, fig. 65.

Leioodes Dunstan, 1923 [*plana*]

Elytron: very small, highly convex; ornament very faint; lateral margin arcuate.

plana Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

pygmaea Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.



Family **DERMESTIDAE** Latreille, 1807

Reeveana Dunstan, 1923 [*minor*]

Elytron: distally tapered; widest point near base; subulate or cuneate overall; ornament very weak, concentric to absent.

major Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

intermedia Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.


minor Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.


Tryoniopsis Dunstan, 1923 [*punctata*]

Elytron: explanate, distally attenuate; borders narrow, well defined; costae low; striae shallow; almost flat.

punctata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

granulata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.





Dytiscid indet. 2; larva laterally compressed (top), NMVP103330, 6.5mm long; larva dorsally compressed (lower above), NMVP103302, 4.2mm long.

Family **DYTISCIDAE** Latreille, 1825

Dytiscid indet 1.

Larva with head longer than pronotum; mandibles strongly curved, opposed, single-tipped; 8th abdominal tergite drawn out into spine-like projection. Adult and elytron only tentatively assigned, rather indistinct.

Dytiscid. indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Dytiscid indet 2.

Larvae: Head subrectangular, with prominent Y-shaped ridge; antennae short, 3-4 segments; pronotum shorter than head, longer than succeeding segments, with truncated corners; abdomen 8-segmented, posteriorly pointed. Adult oval; antennae 10-segmented, becoming beaded; pronotum half body length; coxae large, bearing hairs along outer margin; tibia and tarsus covered with hairs; abdomen of 6 sternites, 1st longest; elytra with longitudinal ridges.

Dytiscid. indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Dytiscid indet 3.

Thorax 2nd = 3rd in length; body widest at rear of thorax; foreleg femur long narrow; mid leg femur heavier, expanding distally; hindleg long, stout, femur & tibia same length; tibia with strong lateral hairs on apex; tarsus 5-segmented, tapering distally to sharp tip, with long fine hairs from concave side and stout prominent lateral hairs from convex side.

Dytiscid. indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Dytiscid indet. 1; larva, MUGD3749, 9mm long (upper); immature adult (above left) and elytron (above right), NMVP103325, body 5.5mm long.




Dytiscid indet. 2; adult, NMVP103169, 4mm long, whitened with ammonium chloride on right.



Dytiscid indet. 3; NMVP 102527; 10mm long. Detail of distal part of hindleg at left.



Elaterium punctomarginum; GSQI200b; 7mm long.
Line drawing (right) of Dunstan, 1923, pl.5, fig. 43.



Elateridium angustius; AMF39328; 9mm long.



Elateridium transversum; GSQI159; 6.5mm long.
Line drawing (right) of Dunstan, 1923, pl. 4, fig.



Family ELATERIDAE Leach, 1815

Elaterium Westwood, 1854 [*pronaesus*]

Elytron: elongate disc with parallel costae; double lines of punctate striae or minute groups of punctae centrally on border.

punctomarginum Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

bipunctatum Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

Elateridium Tillyard 1918b [*wianamattense*]

Elytron: elongate, oval, tapering posteriorly; no sculpture evident except a slight roughness on the surface.

wianamattense (Tillyard, 1916); MTrias; Ashfield Fm; Sydney.

angustius Tillyard, 1918b; MTrias; Ashfield Fm; Sydney.

subulatum (Dunstan, 1923); UTrias; Blackstone Fm; Ipswich.

transversum (Dunstan, 1923); UTrias; Blackstone Fm; Ipswich.




Family HELODIDAE Leconte, 1862

Helodid indet.

Larva comparatively broad: Head small, subquadrate; antennae long, with long 1st segment, of many segments; eyes small, lateral, near midlength of head behind insertion of antenna; median groove on rear of head dividing forward, running to lateral margins just forward of eyes; thorax of subequal segments, with rounded lateral margins to segments, with fine raised ridge around entire margin of tergite. legs short, with large tibia, tarsus and tarsal claw; abdomen of 8 segments, 8th shorter than 7th.

Helodid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Helodid larva indet.; NMVP103308; 7mm long.
Line drawing of Jell & Duncan, 1986, fig. 37A.

Family **HYDROPHILIDAE** Latreille, 1802

Platycrossos Dunstan, 1923 [*tumidus*]

Elytron: punctate-striate; 8-9 costae; lateral border very wide; extended humeral area and border.

tumidus (Tillyard, 1916); UTrias; Blackstone Fm; Ipswich.

ligulatus Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

subtumidus Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.




Platycrossos tumidus; GSQI45b; 6.2mm long. Line drawing (left) of Dunstan, 1923, pl. 3, fig. 21.

Polysitum Dunstan, 1923 [*punctatus*]

Elytron: broadly ovate, uniformly granulate, otherwise like *Ademosyne*.

punctatus Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

minutus Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.




Polysitum punctatum; GSQI153b; 7mm long. Line drawing (left) of Dunstan, 1923, pl. 4, fig. 37.


Sheperdia Dunstan, 1923 [*quadrivittata*]

Elytron: banded sculpture with fine cross lining; apex acutely rounded; otherwise like *Ademosyne*.



quadrivittata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.



Sheperdia quadrivittata; GSQI130a; 11mm long. Line drawing (left) of Dunstan, 1923, pl. 4, fig. 32.



Simmondsia cylindrica; GSQI87a; 3mm long. Line drawing (right) of Dunstan, 1923, pl. 4, fig. 29.



Hydrophilid indet.; NMVP103314; 102522; 102768; x12.



Hydraenid indet.; NMVP 103319; 2.5mm long.



Hydrophilid larva indet.; NMVP 32284; 9mm long.



Gyrinid indet.; NMVP103311; 2.5mm long; whitened with ammonium chloride (left) and in normal light (right).



Simmondsia Dunstan, 1923 [*subpyriformis*]

Elytron: number of longitudinal ridges and connecting crossbars; enclosed squares depressed; lateral and sutural borders narrow; scutellary margin long.

subpyriformis Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

cylindrica Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

Hydrophilid indet.

Small body; head trapezoidal; antennae of 2 long basal segments, shorter broader 3rd segment and 3 short broad distal to the cupule; eyes tiny, posterolateral; pronotum coarsely pustulose, with large paratergal lobes; elytra with broad longitudinal ridges separated by narrow grooves with closely spaced pits along entire length; abdomen of 6 sternites.

Hydrophilid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Hydrophilid larva indet.

Large curved mandibles; thoracic segments subequal; abdomen tapering posteriorly

Hydrophilid larva indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **HYDRAENIDAE** Mulsant, 1844

Hydraenid indet.

Elongate adult; head longer than wide; antennae inserted anterolaterally, of 2 long basal segments and 5 bead-like apical sections; pronotum not expanded; 3rd thoracic segment very long; elytra long, thin; abdomen with 6 sternites.

Hydraenid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **GYRINIDAE** Latreille, 1810

Gyrinid indet.

Hind leg with greatly expanded coxa; tibia wide, expanding distally; basitarsus wider than long, relatively massive; 1st abdominal sternite with pair of circular central and margined depressions to receive posterior tip of hind coxa, much longer than succeeding sternites; 6th sternite as long as others; elytra with widely separated longitudinal ridges.

Gyrinid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **MALACODERMIDAE**

Metrorhynchites Tillyard, 1916 [*sydneiensis*]
Elytron: elongate, oval, with 3 strong longitudinal ribs, one parallel to inner margin and extending to tip, other two roughly parallel with outer margin and meeting first independently near apex.

sydneiensis Tillyard, 1916; MTrias; Ashfield Fm; Sydney.

grandis Tillyard, 1918b; MTrias; Ashfield Fm; Sydney.

Family **MORDELLIDAE** Latreille, 1802

Mordellid indet.

Body typically hump-shaped, with small high head tucked down below convex pronotum; elytra well sclerotised, with well-defined border and longitudinal striations.

Mordellid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **PSELAPHIDAE** Latreille, 1802

Pselaphid indet.

Head small; antenna 11-segmented, with basal segment twice as long as wide; becoming beaded distally; pronotum longer than head, convex; legs with forefemur greatly enlarged, club-shaped mid and hind femora; elytra short, over only abdomen anterior; abdomen short; segments short.

Pselaphid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **STAPHYLINIDAE** Latreille, 1802

Staphylinid indet.

Brachypterous;

head

subquadrate;

antennae

11-segmented,

expanding

apically, apical

segment larger than penultimate; pronotum with small anterolateral spine; elytra subrectangular, just reaching abdomen; abdomen with segments 2-7 subequal; segment 8 small, with indistinct cerci and terminalia.

Staphylinid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **TENEBRIONIDAE** Latreille, 1802

Adelidium Tillyard, 1918b [*cordatum*]

Elytron: broad, small, highly convex; sculpture of 9 slightly raised longitudinal striae, separated by wider interstices; striae punctate.

cordatum Tillyard, 1918b; MTrias; Ashfield Fm; Sydney.



Metrorhynchites grandis; AMF39330; 14.4mm long.



Mordellid indet.; NMVP103158; 3.7mm long.



Pselaphid indet.; NMVP103321; 2mm long.




Staphylinid indet.; MUGD3737B; 4mm long. Line drawing (left) of Jell & Duncan, 1986, fig. 35A




Adelidium cordatum; AMF39329; 4mm long. Line drawing (left) of Tillyard, 1918, text-fig. 14.






Mesothoris westraliensis; AMF52315; 7mm long.



Mesothoris tenuiclatrata; GSQI313a; 6.8mm long. Line drawing (right) of Dunstan, 1923, pl. 6, fig.



Ulomites willcoxi; GSQI50b; 11mm long. Line drawing of Dunstan, 1923, pl. 4, fig. 33.

Mesothoris Tillyard, 1916 [*clathrata*]

Elytron: elongate, almost parallel sided; sculpture areolate, formed by 9 broad flat longitudinal ridges intersected by transverse ridges; areolae subquadrate.

clathrata Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.

quadripartita Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

tenuiclatrata Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

grandis Dunstan, 1923; UTrias; Blackstone Fm; Ipswich.

westraliensis Riek, 1968; LJur; Cockleshell Gully Fm; Perth.



Ulomites Tillyard, 1916 [*willcoxi*]

Elytron: sides subparallel, base rounded, apex pointed; marginal and sutural borders narrow; sculpture of 7 distinct punctate striae; distance between punctures half interstitial width; marginal interstices becoming wider apically; interstices finely granulose.

willcoxi Tillyard, 1916; UTrias; Blackstone Fm; Ipswich.



Undetermined beetles from Koonwarra Fossil Bed (left) and Wianamatta Shale (right)



Order NEUROPTERA (lacewings)

Family **ARCHEOSMYLIDAE** Riek, 1953

Archeosmylus Riek, 1953 [*pectinatus*]

Forewing: Sc and R1 fused; pterostigma long, curved round towards apex; Rs 8- or 9-branched; Rs and M1+2 fused for short distance, with proximal portion of M1+2 almost transverse; Cu2 deeply forked; anals simple, with short pectinate branches near margin.

pectinatus Riek, 1953; UPerm; Newcastle CM; Sydney.

stigmatus Riek, 1955; UTrias; MtCrosby Fm; Ipswich.

costalis Riek, 1955; UTrias; MtCrosby Fm; Ipswich.

Family **PERMITHONIDAE** Tillyard, 1922

Permithone Tillyard, 1922a [*belmontensis*]

[=*Permosmylus* Tillyard, 1926b; type *P. pincombeae*]. Forewing: Sc fused to R1 distally; costal space expanded at base; 6-7 cross veins from R to Rs; Rs with only 4-5 pectinate branches; MP forking before Rs; cubito-median Y-vein distinct; Cu1 branched; Cu2 simple or with one distal fork; A1 pectinately branched; A2 and A3 branched, enclosing an irregularly celled area; cross veins between Cu1 and Cu2 simple or irregular.

Hindwing: costal space narrow; Cu1 branched pectinately; Cu2 simple; Cu1 touching or fused to M for short distance; no cubito-median Y-vein.

belmontensis Tillyard, 1922a; UPerm; Newcastle CM; Sydney.

oliarcoides Tillyard, 1926b; UPerm; Newcastle CM; Sydney.

neoxenus Riek, 1953; UPerm; Newcastle CM; Sydney.

pincombeae (Tillyard, 1926b); UPerm; Newcastle CM; Sydney.

Permopsychops Tillyard, 1926b



[*belmontensis*]

Forewing: Sc fused to R1 distally; costal veinlets simple, rarely branched, without cross veins; 3 r-rs; Rs with numerous pectinate branches; M forking near base before Rs; cubito-median Y-vein distinct; Cu1 2-branched; Cu2 simple; A1 forked near apex; 2A and 3A branched.

belmontensis Tillyard, 1926b; UPerm; Newcastle CM; Sydney. [= *Permithone venosa* Davis, 1943; UPerm; Newcastle CM; Sydney].




Archeosmylus pectinatus; AMF40264; 11mm long.
Line drawing (above) of Riek, 1953, text-fig. 66.





Permithone belmontensis; AMF28465; 9.4mm long.
Line drawing (above) of Tillyard, 1922a, text-fig. 6.




Permopsychops belmontensis; AMF39867; 15mm long, holotype of *P. venosa*.




Permorapisma biserialis; AMF28053; 20mm long. Line drawing (right) of Riek, 1953, text-fig. 48.



Euporismites balli; GSQI34; 22mm long. Line drawing (above) of Tillyard, 1916, pl. 3, fig. 1.



Lithosmylidia lineata; UQC2189 (part and cp); 20mm long. Line drawing (right) of Riek, 1955, fig. 22.



Permorapisma Tillyard, 1926 [*biserialis*]

Forewing: costal space enlarged near base; Sc fused with R1 distally; costal veinlets often branched and connected by numerous irregular cross veins; 10 or more r-rs; Rs with 9 pectinate branches (including MA), with dense end twigging; cubito-medial Y-vein distinct; Cu1 branched towards apex; 3 anal veins branched pectinately; Cu1 to Cu2 cross veins reticulate, forming 2-3 cells; similar reticulate cross veins between anals.

biserialis Tillyard, 1926b; UPerm; Newcastle CM; Sydney.

triserialis Tillyard, 1926b; UPerm; Newcastle CM; Sydney.

Family OSMYLIDAE Leach, 1815

Euporismites Tillyard, 1916 [*balli*]

Bordered by close veins (costal cross veins anteriorly, many branches of primary veins apically and of Cu and A veins posteriorly); Sc and R1 very close, parallel; Rs parallel to R1 giving off large series of pectinate branches; M widely separated from Rs; Cu1 and Cu2 not distinct, zigzag, curving posteriorly very early and then branching.

balli Tillyard, 1916; Palaeocene; Redbank Plains Fm; Ipswich.

Lithosmylidia Riek, 1955 [*lineata*]


Forewing: Sc and R1 fused towards apex; M1+2 only just touching Rs, with free oblique basal stem; limited cross veins near base of Rs; pterostigmatic veinlets often forked; terminal branches of CuA and CuP not transverse.


lineata Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

parvula Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

baronne Lambkin, 1988; Gayndah Beds; Esk Graben.

sp. A Lambkin, 1988; UTrias; Mt Crosby Fm; Ipswich





Family OSMYLOPSYCHOPIDAE

Martynova, 1949


Osmylopsychops Tillyard, 1923b [*spillerae*]

Forewing: Sc and R1 fused apically; all veins abundantly branched; CuA many branched; Sc, R1 and Rs separate almost to base; endtwiggling limited; costal veinlets mainly forked and connected by scattered cross veins; CuP and anals well branched.

spillerae Tillyard, 1923b; UTrias; Blackstone Fm; Ipswich.



Osmylopsychops spillerae; GSQI314; 14mm long. Line drawing (left) of Tillyard, 1923b, text-fig. 93.



Protopsychochops venosa; GSQI160a; 9.5mm long (right). Line drawing (left) of Tillyard, 1917, pl. 8, fig. 3.



Protopsychochops Tillyard, 1917 [*venosa*]

Wing broad, moderately pointed; large number of forked apical veins. Forewing: broad costal margin, with numerous costal cross veins, some forked; Sc and R separated by numerous cross veins; distally Sc and R approach close to one another a short distance before the apex; R and Rs further apart, with many cross veins; Rs 12-branched, with delicate cross veins; 2 rows of gradate veins in almost complete transverse lines across wings; another series of stepwise crossveins between the two.

venosa Tillyard, 1917; UTrias; Blackstone Fm; Ipswich.


Petropsychochops Riek, 1956 [*superba*]


Sc fused to R1; no distinct 'ven triplica'; R bent strongly away from Sc at base; Rs arising almost basally; Cu diverging strongly from the Sc-R veins; M forming pectinate series of many anterior branches occupying large area of wing.

superba Riek, 1956; UTrias; Blackstone Fm; Ipswich.


Petropsychochops superba; UQC2136; 28mm long along R.

Archeopsychochops triassica; GSQI137; 6.7mm long.







Triassopsychops superba; GSQI284a; 29mm long. Line drawing (above) of Tillyard, 1922b, text-fig. 89.



Proberotha superba; UQC1404; 7mm long (below). Line drawing (above) of Riek, 1955, fig. 18.



Archeopsychops triassica; GSQ137; line drawing of Tillyard, 1919a, text-fig. 27.

Family PSYCHOPSIDAE Handlirsch, 1906

?*Archeopsychops* Tillyard, 1919a [*triassica*]

Forewing: costal space markedly expanded towards base but tapering rapidly towards pterostigma; Sc, R1 and Rs probably forming a vena triplica.

triassica Tillyard, 1919a; UTrias; Blackstone, MtCrosby Fms; Ipswich.

Triassopsychops Tillyard, 1922b [*superba*]

Forewing: very broad; apex rounded; Sc, R1, Rs strong from base to just beyond midlength, joined distally by two strong cross veins; the 3 continuing for short distance to be joined by cross veins again; Sc and R1 continuing a further short distance to another cross vein then all dividing into terminal veinlets; costal area with recurrent veinlet near base sending a few veinlets to margin; most costal veinlets arising from Sc at 45°; M and Cu1 not fused distally; Rs with about 14 branches from Rs; M 5-branched; basal r-m short, strong; cross veins between branches of Rs numerous; Cu simple for long way, distally multibranched; A1 and A2 multi-branched.

superba Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.

Family PROBEROTHIDAE Riek, 1955


Proberotha Kruger, 1923 [*prisca*]

Forewing: costal space markedly expanded over middle portion; margin of wing almost entire; Sc and R1 fused near apex; normally 3 cross veins between R1 and Rs; M and Rs connected only by a cross vein.

superba Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

Proberothella Riek, 1955 [*elongata*]

Forewing: costal space markedly expanded over middle portion; margin of wing almost entire; Sc and R1 fused near apex; 3 cross veins re-m; A1 deeply forked.



Proberothella elongata; UQC2196; 9mm long (left). Line drawing (above) of Riek, 1955, fig. 19.

elongata Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

Order GLOSSELYTRODEA

Family JURINIDAE Zalessky, 1929

Polycytella Tillyard, 1922b [*triassica*]

Tegmen: surface ornament of raised polygonal cellules forming meshwork; Sc, R, M and Cu1 unbranched, running straight to margin; R finishing near midlength; M finishing near apex; clavus narrow.

triassica Tillyard, 1922b; UTrias; Blackstone Fm; Ipswich.




Polycytella triassica; GSQI81a; 7.5mm long.


Permoberothella Riek, 1953 [*perplexa*]

Forewing: pterostigma absent; Rs arising near base of wing; M1+2 fused with Rs proximally, then almost straight to the apex; M3+4 fused with Cu1 for short distance then parallel to M1+2; a distinct groove between MA and MP; MP 2-branched, with terminal twigging; Cu1 and Cu2 simple; 3 simple anals connected by irregular cross veins.


perplexa Riek, 1953; UPerm; Newcastle CM; Sydney.



Permoberothella perplexa; AMF42520; 7mm long. Line drawing (above) of Riek, 1953, text-fig. 53.



Cladochorista belmontensis; AMF28054; 4mm long. Line drawing (right) of Riek, 1953, text-fig. 50.



Cladochoristella bryani; UQC1746; 12mm long. Line drawing of Riek, 1955, fig. 16.



Prorhyacophila colliveri; UQC1741; 5mm long. Line drawing (above) of Riek, 1955, fig. 17.

Order TRICHOPTERA (caddisflies)

Family **CLADOCHORISTIDAE** Riek, 1953

Cladochorista Riek, 1953 [*belmontensis*]

Costal space with numerous oblique crossveins; R1 branched at apex; R2+3 branching before R4+5; sc-r before origin of Rs; M 4-branched; cubito-median Y-vein distinct; Cu1 branched at or beyond its middle; Cu2 simple, weak; anal veins looped.

belmontensis Tillyard, 1926b; UPerm; Newcastle CM; Sydney.



Cladochoristella Riek, 1955 [*bryani*]

Forewing: Sc ending in series of veinlets; 4 transverse costal veinlets; R1 forked in pterostigma, 3-branched; arms of cubito-median Y-vein unequal; R2+3 forking before R4+5; M 5-branched, with extra branch on Ms. Hindwing: Sc forked, ending before R1; Rs forked as in forewing.

bryani Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

Prorhyacophila Riek, 1955 [*colliveri*]

[=*Eocorona* Tindale, 1980; type *iani*]

Forewing: Sc forked at apex, with only single cross vein to costal margin; R1 simple; CuA deeply forked; CuP straight; distinct cubito-median Y-vein.

colliveri Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.

iani Tindale, 1980; UTrias; Mt Crosby Fm; Ipswich.



Prorhyacophila iani (holotype of *Eocorona iani*); UQC2327; 8mm long.

Family **CALAMOCERATIDAE** Ulmer, 1905

Calamoceratid pupa indet.

Abdomen with fringe of fine long hairs from segment 2 back, with dorsal hook plates on segments 2-7,

Calamoceratid pupa indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Oecetis sp.;
NMVP103184;
9mm long.

Family **LEPTOCERIDAE** Leach, 1815

Oecetis sp.

Case of fine sand grains, straight, slightly tapered; apex slightly oblique; sand grains increasing in size towards aperture.

Oecetis sp. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.


Family **STEREOCHORISTIDAE** Tillyard, 1919

Stereochorista Tillyard, 1919a [frustrata]


Wing small; CuA forked; R2+3 forking before R4+5; r-m strong

frustrata Tillyard, 1919a; UTrias; Blackstone Fm; Ipswich.


[Tillyard (1919a) assigned the genus to the Mecoptera and detailed some peculiar morphology e.g. the 3-branched 1A. Riek (1956, fig. 1) interpreted the wing the other way up from Tillyard and assigned the genus to the Trichoptera. Carpenter (1992) considered it among genera that could not be assigned to Order within the Neoptera].






Calamoceratid pupa indet.;
NMVP102632; x8. Line drawing
(above right) of Jell & Duncan,



Stereochorista frustrata; GSQI218; 7.5mm long.
Line drawing (above) of Riek, 1956, fig. 1.



Aphryganoneura anomala; AMF29033; 12mm long.



Neopetromantis australis; AMF40211; 5mm long. Line drawing (right) of Riek, 1953, fig. 27.

Order MECOPTERA (scorpionflies)

Family INCERTAE SEDIS

Aphryganoneura Tillyard, 1926b [*anomala*]
Forewing: branches of main veins parallel, separated by grooves as in Neuroptera Plannipennia, unbranched distally.

anomala Tillyard, 1926b; UPerm; Newcastle CM; Sydney.

Family AGETOPANORPIDAE Carpenter, 1930

Agetochoristella Riek, 1953 [*adscita*]
Forewing: Sc 3-branched; costal space expanded; Rs 5-branched, dichotomously, with extra fork on R5; M 5-branched, with extra fork on M2; cubito-median Y-vein distinct; Cu1, Cu2, A1 and A2 all simple.

adscita Riek, 1953; UPerm; Newcastle CM; Sydney.

Agetochoristella adscita; AMF43656; 10mm long. Line drawing (above left) of Riek, 1953, fig. 28.

Neoageta Riek, 1953 [*elongata*]
Forewing: Sc 3-branched; Rs 5-branched, with extra fork on R5; M 6-branched, extra forks on M2 and M4; Cu1, Cu2 and anal veins simple; Cu1 and M touching; cubito-median Y-vein absent.

elongata Riek, 1953; UPerm; Newcastle CM; Sydney.

Neoageta elongata; AMF43969; 12mm long. Line drawing (above left) of Riek, 1953, fig. 26.

Neopetromantis Riek, 1953 [*australis*]
Forewing: costal space not expanded; Sc 3-branched; Rs 4-branched; M 6-branched, extra forks on M2 and M4; Cu1, Cu2 and A1 simple; cubito-median Y-vein absent.


australis Riek, 1953; UPerm; Newcastle CM; Sydney.



Phipoides Riek, 1953 [*elegans*]

Forewing: costal space expanded, Sc 3-branched; Rs 5-branched, extra fork on R2; M 6-branched; Rs 5-branched, extra fork on R2; M 6-branched, extra forks on M2 and M4. Hindwing: costal space not expanded; Sc short, simple; R1 forked near apex; Rs same as in forewing; M 4- or 5-branched.

elegans Riek, 1953; UPerm; Newcastle CM; Sydney.



Phipoides elegans; AMF40144; 8mm long. Line drawing (above) of Riek, 1953, fig. 25.

Family **ARCHIPANORPIDAE** Tillyard, 1917


Archipanorpa Tillyard, 1917 [*magnifica*]

Forewing: pterostigma short, irregular; R arching up into pterostigma; Sc reaching costal margin beyond midwidth; Rs and M branching repeatedly and dichotomously; Cu1 2-branched; Cu2 simple.

magnifica Tillyard, 1917; UTrias; Blackstone Fm; Ipswich.



Archipanorpa magnifica; GSQ1106; 33mm long (right). Line drawing (above) of Tillyard, 1917, pl. 8, fig. 5.



Family **BELMONTIIDAE** Tillyard, 1919b


Belmontia Tillyard, 1919b [*mitchelli*]


Forewing: costal space narrow, not expanded near base; Sc forked near its apex, connected to R1 by a cross vein; no distinct pterostigma; R1 simple; Rs 6-branched; M 4-branched, occasionally 5-branched; cubito-median Y-vein well-developed; Cu1 normally forked near apex; Cu2 simple; anal veins simple with cross veins.

mitchelli Tillyard, 1919b; UPerm; Newcastle CM; Sydney.



Belmontia mitchelli; AMF28469; 16mm long. Line drawing (right) of Tillyard, 1919b, text-fig. 1.





Parabelmontia Tillyard, 1922a [*permiana*]
Sc forked distally; 2 r3-r4; M fused basally with RRs and M each 6-branched; Cu1 not forked distally;
permiana Tillyard, 1922a; UPerm; Newcastle CM; Sydney.




Parabelmontia permiana
(hindwing); AMF28461; 18mm long. Line drawing (above left) of Riek, 1953, text-fig. 7.


Family **CHORISTIDAE**
Ebsen-Petersen, 1915

Chorista Klug, 1836 [*australis*]
Forewing: costal area narrow, with one simple cross vein; Sc reaching margin at or just beyond midlength; Cu1 touching or fused with M for short distance; Rs 4-branched; M 5-branched, extra fork on M4.



sobrina Riek, 1952a; Palaeocene; Redbank Plains Fm: Ipswich.



Chorista sobrina Riek, 1952; UQF10628; x4. Line drawing (right) from Riek, 1952a, text-fig. 1.



Cretacochorista parva;
NMVP 103248; 7mm long. Line drawings (below) of Jell & Duncan, 1986, fig. 41C; forewing (right) and hindwing (left).



Mesochorista proavita [holotype of *Eoses triassica*]; UQF7853; 11mm long.

Cretacochorista Jell & Duncan, 1986 [*parva*]
Rs 5-branched; M 5-branched in forewing, 4-branched in hindwing.

parva Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **PERMOCHORISTIDAE** Tillyard, 1918b

Mesochorista Tillyard, 1916 [*proavita*] [= *Eoses* Tindale, 1945; type *triassica* (= *proavita*)] [= *Permochorista* Tillyard, 1918b; type *australica*]

Forewing: Sc 2-branched; Rs 4-branched; M 6-branched, extra forks on M2 and M4; CuA simple; Cu-M Y-vein variable. Hind: Sc simple; Rs as in fore; M 4-branched; CuP and A1 fused for part of length.

proavita Tillyard, 1916; Trias; Blackstone, Mt Crosby Fms; Ipswich.

australica (Tillyard, 1918b); UPerm; Newcastle CM; Sydney.

jucunda (Tillyard, 1926b); UPerm; Newcastle CM; Sydney.

dubia Riek, 1953; UPerm; Newcastle CM; Sydney.

phipa Riek, 1953; UPerm; Newcastle CM; Sydney.

Parachorista Tillyard, 1926b [*pincombeae*]

Forewing: R2+3 forming pectinate series of 4 or 3 branches; R4+5 2-branched; M 6-branched, extra forks on M2 and M4; Cu1 and anals simple; cubito-median Y-vein distinct. Hindwing: R branched as in forewing; M 4-branched; Cu1 fused to M for short distance.

pincombeae Tillyard, 1926b; UPerm; Newcastle CM; Sydney.

splendida Tillyard, 1926b; UPerm; Newcastle CM; Sydney.

bairdae (Tillyard, 1922b); UPerm; Newcastle CM; Sydney.

warnerensis Tillyard, 1926b; UPerm; Newcastle CM; Sydney.



Family **MESOPANORPODIDAE** Tillyard, 1918b


Mesopanorpodes Tillyard, 1918b [*wianamattensis*]

Forewing: Sc forked, 2-branched; Rs 4-branched; R2+3 forking before R4+5; M 4-branched; cubito-median Y-vein variable; Cu1, Cu2, A1, A2 simple; A3 forked close to its base; forking to Rs abnormal, similar to Trichoptera.


wianamattensis Tillyard, 1918b; MTrias; Ashfield Fm; Sydney.

belmontensis Riek, 1953; UPerm; Newcastle CM; Sydney.


robustus Riek, 1953; UPerm; Newcastle CM; Sydney.





Mesochorista proavita; GSQI32. Line drawing from Tillyard, 1916, pl. 2, fig. 2. [Parfrey (1996) noted the type missing since being loaned to Tillyard in 1936. It is now in AM (Fletcher, 1971)].





Mesochorista australica; AMF40141; 10mm long.




Parachorista splendida; AMF28048; 15mm long. Line drawing (left) of Riek, 1953, fig. 24.



Mesopanorpodes wianamattensis; AMF39326; 7.5mm long. Line drawing (above) of Tillyard, 1918b, text-fig. 10




Prochoristella leongatha; NMVP102512; x9. Line drawing (above) of Jell & Duncan, 1986, fig. 43C.



Prochoristella megaloprepia; AMF40123; 7mm long (left). Line drawing of Riek, 1953, text-fig. 37.



Mesopsyche superba; AMF39263; 24mm long.



Mesopsyche elongata; AMF39233; 13mm long. Line drawing (right) of Tillyard, 1917, pl. 7, fig. 1.

Prochoristella Riek, 1953 [*megaloprepia*]

Forewing: Sc long, 2-branched; R1 simple, pterostigma well-developed; Rs 4-branched; cubito-medial Y-vein variable, arms equal of CuA and M fused; CuA and CuP simple; A1, A2 simple; A3 apparently simple.

megaloprepia Riek, 1953; UPerm; Newcastle CM; Sydney.

anagaura Riek, 1953; UPerm; Newcastle CM; Sydney.

exilis Riek, 1953; UPerm; Newcastle CM; Sydney.

pusilla Riek, 1953; UPerm; Newcastle CM; Sydney.

belli (Tillyard, 1926b); UPerm; Newcastle CM; Sydney.

concinna Riek, 1953; UPerm; Newcastle CM; Sydney.

leongatha Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **MESOPSYCHIDAE** Tillyard, 1917

Mesopsyche Tillyard, 1917 [*triareolata*]


(= *Aristopsyche* Tillyard, 1919a; = *Neuropsychyche* Tillyard, 1919a; = *Triassopsyche* Tillyard, 1917a) Forewing: normal Mecoptera except for series of costal veinlets; Rs and M each normally 4-branched; distinct cubito-medial Y-vein. Hindwing: often with small end twigging on branches of Rs; Sc shorter than in forewing; no cubito-medial Y-vein.

triareolata Tillyard, 1917; UTrias; Blackstone Fm; Ipswich.

dunstani (Tillyard, 1917); UTrias; Blackstone Fm; Ipswich.

elongata (Tillyard, 1919a); UTrias; Blackstone Fm; Ipswich.

superba (Tillyard, 1919a); UTrias; Blackstone Fm; Ipswich.



Family **NANNOCHORISTIDAE** Tillyard, 1917

Nannochorista Tillyard, 1917 [*dipteroides*]

Head with heavily sclerotised caudal margin; pronotum with sclerotised shield having scalloped anterior and convex lateral margins; abdominal segments increasing in length from 1 to 3; with middle tergites twice as long as wide; segments 7-10 decreasing in length and width; anal hooks long, tapering to point, as long as 10th segment.

Nannochorista sp. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Nannochorista sp; NMVP103245; 13mm long.

Nannochoristella Riek, 1953 [*reducta*]



Forewing: Sc forked; Rs 3-branched; M 3-branched; no cubito-median Y-vein; M and Cu1 fused for short distance. Hindwing: apex a little more rounded; Rs and M as in forewing.

reducta Riek, 1953; UPerm; Newcastle CM; Sydney.


Neochoristella Riek, 1953 [*optata*]

Forewing: Sc long, probably forked; Rs 3-branched; M 4-branched; cubito-median Y-vein absent; Cu1 and M fused for some distance; Cu1 and Cu2 simple; A1, A2 simple.


optata Riek, 1953; UPerm; Newcastle CM; Sydney.



Nannochoristella reducta; AMF39968; 5mm long. Line drawing (above) of Riek, 1953, text-fig. 47.



Neochoristella optata; AMF41608; 4mm long. Line drawing (above) of Riek, 1953, text-fig. 46.




Robinjohnia tillyardi; AMF52312; x8. Line drawings of Riek, 1968, fig. 1A, forewing (upper left), 1B, hindwing (lower left), 1C, metanotum and abdomen (right).



Robinjohnia tillyardi; AMF52312; x8.



Choristopanorpa bifasciata; AMF38263; x3.



Choristopanorpa drinnani; NMVP102506; 7.5mm long. Line drawing (right) of Jell & Duncan, 1986, fig. 41A.

Robinjohnia Martynova, 1948 [*tillyardi*]

Forewing: costal space not expanded; Sc forked, short; R simple, straight, except at base; pterostigma large but not sharply defined; Rs and M 4-branched; R2+3 forking close to margin; M1+2 forking after M3+4; discoidal cell closed by cross vein; Cu1 simple, straight; Cu2 parallel to Cu1 strongly curved at margin, finishing away from Cu1; Cu1 and M fused near bases; basal stem of Cu1 transverse, joining M level with humeral cross vein. Hindwing: Sc simple, short; humeral cross vein distinct; R fused with Sc as far as humeral cross vein; Cu1 simple, distinctly curved back at margin; Cu1 very oblique, just touching M; Cu2 well separated from Cu1; Cu2 and A1 fused for short distance.

tillyardi Martynova, 1948; UPerm; Newcastle CM; Sydney.

Family **NEORTHOPHLEBIIDAE**

Handlirsch, 1939

Archebittacus Riek, 1955 [*exilis*]

Forewing: very narrow base; A3 absent or very short; M4 a distinct stem before it branches.

exilis Riek, 1955; UTrias; Mt Crosby Fm; Ipswich.



Archebittacus exilis; UQC2139; 16mm long (left). Line drawing of Riek, 1955, fig. 11.


Family **ORTHOPHLEBIIDAE** Handlirsch, 1906


Choristopanorpa Riek, 1950 [*bifasciata*]

Forewing: Sc long, reaching into pterostigma; Rs 5-branched, extra fork on R2; R2+3 forking before R4+5; M 5-branched, extra branch on M4; cubito-median Y-vein well-developed; 3 anal veins; A3 forked. Hindwing: Sc long, simple; R1 forked distally; Rs 5-branched, rarely 6-branched; M 4-branched; cubito-median Y-vein absent; CuP and A1 fused for part of its length.

bifasciata Riek, 1950; MTrias; Hawkesbury Sst; Sydney.

drinnani Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.





Choristopantorpa bifasciata, line drawing of holotype forewing, AMF30959 (above) and a typical hindwing, AMF39195 (below).

Neoparachorista Riek, 1955 [*perkinsi*]

Forewing: Sc forked distally; R1 forked in pterostigma; Rs with 5-8 branches; without cubito-median Y-vein; CuA almost straight; M 5- or 6-branched. Hindwing: same as forewing but M 4-branched; Sc shorter and simple; CuP and A1 fused for part of their length.

perkinsi Riek, 1955; UTrias; MtCrosby Fm; Ipswich.

splendida Riek, 1955; UTrias; MtCrosby Fm; Ipswich.

semiovena Riek, 1955; UTrias; MtCrosby Fm; Ipswich.



clarkae Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **PANORPIDAE** Latreille, 1804


Austropanorpa Riek, 1952a [*australis*]

Forewing: Sc ending close to or at pterostigma; R2 pectinately branched; R2 4-, 5- or 6-branched; Cu1 not fused to M, connected to it by cross vein; A1 reaching margin beyond origin of Rs.



australis Riek, 1952a; Palaeocene; Redbank Plains Fm; Ipswich.




Austropanorpa australis; UQF10636; x4. Line drawing (above) of Riek, 1952a, fig. 2.





Neoparachorista clarkae; NMVP102529; body 9mm long. Line drawing (above) of Jell & Duncan, 1986, fig. 41B.



Neoparachorista perkinsi; UQC1897; 24mm long. Line drawing (left) of Riek, 1955, fig. 6.




Austropanorpa australis; UQF15572;



Permomeropidae australis; AMF28045; 8.5mm long. Line drawing (above) of Riek, 1953, fig. 1.



Neopermopanorpa mesembria; UQC643; 7.5mm long. Line drawing (above right) of Riek, 1955, fig. 2.



Xenopanorpa didymovena; UQC1043; 11mm long. Line drawing (right) of Riek, 1955, fig. 3.


Family PERMOMEROPIDAE Riek, 1953

Permomeropidae australis Tillyard, 1926b

Forewing: costal space slightly expanded over basal half; Rs with numerous dichotomous branches; M normally 8-branched; CuA forked at least once near apex. Hindwing: costal space narrow basally; Sc not reaching pterostigma but fused to R1; pterostigma well-developed in apical fork of R1; Rs with fewer branches than forewing; M usually with 8 dichotomous branches; cubito-median Y-vein not clear; CuA forked at least once, generally twice near apex.

australis Tillyard, 1926b; UPerm; Newcastle CM; Sydney.

nanus Riek, 1953; UPerm; Newcastle CM; Sydney.



Family PERMOPANORPIDAE Tillyard, 1926

Neopermopanorpa Riek, 1955 [*mesembria*]


Forewing: similar to *Permopanorpa* but with 7-branched M; M4 with strong fork; Sc forked near apex; pterostigma well-developed; Rs 7-branched; CuA simple.

mesembria Riek, 1955; UTrias; MtCrosby Fm; Ipswich.

Xenopanorpa Riek, 1955 [*didymovena*]

Forewing: R2+3 and R4+5 both very short; R2+3 branching just before R4+5; Rs 7- or 8-branched; M 7- or 8-branched, dichotomous divisions; Sc forked near apex.

didymovena Riek, 1955; UTrias; MtCrosby Fm; Ipswich.



Family **PERMOTANYDERIDAE** Riek, 1953

Permotanyderus Riek, 1953 [*ableptus*]

Forewing: Sc long, simple, close to costal margin; pterostigma well-developed; Rs 4-branched; M 4-branched; Cu1 simple, curved over apical part; 2 anal veins.

ableptus Riek, 1953; UPerm; Newcastle CM; Sydney.

Choristotanyderus Riek, 1953 [*nanus*]

Forewing: Sc long, close to costal margin; pterostigma well-developed, well before apex of wing; Rs 4-branched; M 4-branched, cubito-median Y-vein present; Cu1 with slight bend towards apex; e anal veins.

nanus Riek, 1953; UPerm; Newcastle CM; Sydney.



Permotanyderus ableptus; AMF40675; 5mm long. Line drawing (above) of Riek, 1953, text-fig. 48.




Mesotanyderus Riek, 1955 [*jonesi*]

Forewing: long, narrow; Sc long, simple; Rs arising close to wing base; cubito-median Y-vein absent; 3 anal veins; A2 and A3 short; Sc, R and M fused at base; R strongly angled at base; a few cross veins over apical half.

jonesi Riek, 1955; UTrias; MtCrosby Fm; Ipswich.



Choristotanyderus nanus; AMF39964; 4mm long. Line drawing (left) of Riek, 1953, text-fig. 49.



Mesotanyderus jonesi; UQC1054; 7mm long (right). Line drawing (above) of Riek, 1955, fig. 15.





Family **PERMOTIPULIDAE** Tillyard, 1929

Permotipula Tillyard, 1929 [*patricia*]

Forewing: Sc short, only to midlength; pterostigma well-developed; Rs 3-branched; R2+3 simple; M 4-branched, arising close to base; cubito-median Y-vein reduced; only 2 anal veins.

patricia Tillyard, 1929; UPerm; Newcastle CM; Sydney.



Permotipula patricia; BMNH In45382 (2nd above). Wing venation from Willman, 1989a, fig. 2B (above left) and Tillyard, 1929 (above).



Xenochorista splendida; AMF40157; 11mm long.
Line drawing (above) of Riek, 1953, fig. 33.



Xenochoristella hillae; UQC2115; 10.5mm long.
Line drawing (above) of Riek, 1955, fig. 13.



Austrochoristella whitehousei; UQC2172; 5.5mm long.
Line drawing of Riek, 1955, fig. 14.

Family XENOCHORISTIDAE Riek, 1953

Xenochorista Riek, 1953 [*splendida*]

Forewing: costal space moderately expanded; Sc forked near midlength, 3-branched; Rs and M each 4-branched; cubito-median Y-vein well-developed; CuA, CuP and anals (with possible exception of A3) simple.

splendida Riek, 1953; UPerm; Newcastle CM; Sydney.

sobrina Riek, 1953; UPerm; Newcastle CM; Sydney.

Xenochoristella Riek, 1955 [*hillae*]

Forewing: costal space moderately expanded; Sc 3-branched, with distinct cross vein to margin between 1st and 2nd branches; Rs and M each 4-branched; cubito-median Y-vein with unequal arms; CuA, CuP and anals (except A3) simple.

hillae Riek, 1955; UTrias; MtCrosby Fm; Ipswich.

Family MESOPANORPODIDAE Tillyard, 1918b

Austrochoristella Willmann, 1989b

[*Prochoristella whitehousei* Riek, 1955]

M and Cu1 fused for short distance near base.

whitehousei (Riek, 1955); UTrias; MtCrosby Fm; Ipswich.


Family TRIASSOCHORISTIDAE Willmann, 1989

Triassochorista Willmann, 1989b

[*Neoparachorista nana* Riek, 1955]

Rs 5-branched; M 5-branched; M and Cu1 fused for short section at base

nana (Riek, 1955); UTrias; MtCrosby Fm; Ipswich.



Triassochorista nana; UQC2080; 12mm long. Line drawing of Riek, 1955, fig. 10.

Order SIPHONAPTERA (fleas)

Family INCERTAE SEDIS

Tarwinia Jell & Duncan, 1986 [*australis*]

Antennae 18 or 19-segmented; head not excessively compressed laterally; thoracic segments relatively long; tergal combs absent; tibiae all with stout bristles; tarsi very long, with long tarsal claws, with stout bristles under apical tarsal segment. Abdomen enlarged, with typical male genitalia.

australis Jell & Duncan, 1986; LCret;
Koonwarra FB; Gippsland.

Niwratia Jell & Duncan, 1986 [*elongata*]

Legs long, slender; body typically siphonapteran.

elongata Jell & Duncan, 1986; LCret;
Koonwarra FB; Gippsland.

Family PULICIDAE Stephens, 1829

Pulicid indet.


Wingless, laterally compressed

Pulicid indet. Jell & Duncan, 1986; LCret;
Koonwarra FB; Gippsland.

[Carpenter (1992:395) noted the 2 apterous insects from the Lower Cretaceous of Australia assigned to the Siphonaptera by Riek (1970a) but on the opinions of "specialists who have examined the fossils" he dismissed them from the order. Rasnitsyn in Rasnitsyn & Quicke, 2002:242 accepted one of these, *Tarwinia australis*, as a siphonapteran but dismissed others. Zherikhin in Rasnitsyn & Quicke, 2002:366 considered *Niwratia elongata* to be a young dragonfly nymph and the Pulicid indet. to be roach nymphs].




Tarwinia australis; NMVP26202, body 7mm long.



Pulicid indet.; NMVP103056; body 3mm long. Line drawing (above) of Jell & Duncan, 1986, fig. 45A.



Niwratia elongata; NMVP102517; body 3.5mm long.



Chironomaptera collessi; larva NMVP103262 (left), 5mm long; pupae NMVP103253 (centre), 4mm long; NMVP 103139 (right), 5mm long.



Order DIPTERA (flies)

Family CHAOBORIDAE

Edwards, 1920

Chironomaptera Ping, 1928

[*Samarura gregaria* Grabau, 1923]

Larva: Head very large, with eyes; Thorax and 7th abdominal segment with narrow elongate air sacs (tracheal chambers). Abdominal segment 10 with typical elongate culicoid ventral brush of 16 or more hairs, dense anteriorly; dorsal brush small with short hairs; siphon with pointed tip.

Pupa: abdominal segments with transverse, symmetrical, tapering, tuft-like sets of markings (presumably hairs); segment 7 shorter than segment 8; posterior paddles wide.

collessi Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.


Family CHIRONOMIDAE Macquert, 1838

Subfamily TANYPODINAE Skuse, 1889

Tanypodine indet.

Head small, with lateral surface almost completely occupied by eyes; antenna long, stout, with short pedicel, longer scape, at least 10 flagellar segments, plumose distally. Thorax large, strongly convex; metanotum heavily sclerotised; legs with long thin femora and tibia; basitarsus 0.6 of length of tibia, with rest of tarsus including long claw less than half as long as basitarsus. Abdomen twice as long as thorax, of 8 segments and terminalia, with first 6 segments of equal length, with 7th and 8th shorter. Wing with strong R; R4+5 forward of apex; R1 meeting costal margin beyond midlength of wing; base of M3+4 distal to M3+4 -CuA junction.

Tanypodine indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Tanypodine indet.; NMVP103282 (left above), body 2.1mm long; NMVP103285 (left), 5.5mm long. Line drawing (above) of Jell & Duncan, 1986, fig. 55B.

Tanypodine pupae indet.

Two types of pupae of typical form, with 8 abdominal segments plus caudal lobes; each segment with median carina and lateral longitudinal furrows; caudal lobes in one type broad, flat, with rounded outer margin and straight almost sagittal margins along posteromedial cleft; in other type they are large, flat, with outer margin inverted pear-shaped, with posterior spines, wide posterior embayment and fringe of fine hairs around whole margin.

Tanypodine pupa indet. 1 Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Tanypodine pupa indet. 2 Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Tanypodine pupae indet.; dorso-ventrally compressed NMVP103187 (above left), x8; laterally compressed MUGD3733 (above), x7; different species NMVP103134 (left), x6. Line drawing of Jell & Duncan, 1986, fig. 58A (above left), 58B (above).


Family DIXIIDAE Brauer, 1880

Dixid indet. Jell & Duncan, 1986

Head small, rounded, with large lateral eyes; antenna longer than body, with stout scape, with long hairy tapering markedly more slender flagellum. Thorax broad, with sclerotised wing attachment points.

Wing: Sc and R parallel, close from humeral vein; R1 curving distally just inside margin; Rs arising closer to the base of the wing than Sc; Legs extremely long, slender, with long femur, tibia and basitarsus. Abdomen large, with bluntly rounded posterior.

Dixid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.




Family DITOMYIIDAE

Australosymmerus Freeman, 1954


Sc lacking;

imperfecta Riek, 1954a; Palaeocene; Redbank Plains Fm; Ipswich.


[named by Riek as *Centrocnemis? imperfecta* but genus preoccupied; *Centrocnemis* replaced by *Australomyia* Freeman, 1951, replaced by *Australosymmerus*].




Dixid indet.; NMVP102499; body 4.5mm long. Line drawing of wing (above) of Jell & Duncan, 1986, fig. 50J.



Australosymmerus imperfecta; UQF14345; 5mm long. Line drawing (left) of Riek, 1954, fig. 2.



Aneura apicalis; UQF14361; 2.5mm long. Line drawing (right) of Riek, 1954, fig. 5



Protasmanina nana; UQF14347; 4mm long. Line drawing (above) of Riek, 1954, fig. 3.

Pseudalysiinia mimicans; NMVP103290; body 3.5mm long. Line drawing (below) of Jell & Duncan, 1986, fig. 50I.



Mycetophilidae gen. nov.; NMVP103286; 3.2mm long. Line drawing (right) of Jell & Duncan, 1986, fig. 50H.



Family MYCETOPHILIDAE Macquert, 1838

Aneura Marshall, 1896 [*boletinoides*]

Wing: Sc without an apical crossvein to R1; M1+2 forking towards the wing margin, with stem longer than branches, with stem longer than r-m; r-m long; only 1A distinct.

apicalis Riek, 1954a; Palaeocene; Redbank Plains Fm; Ipswich.

Protasmanina Riek, 1954a [*nana*]

Wing with Sc ending in C after origin of Rs, with crossvein to R1; Rs transverse from its origin to r-m; r-m more or less longitudinal, about equal to stem of M1+2; M3+4 arising from CuA more basad than forking of M1+2; CuP long but weak; 1A ending well before wing margin; 2A strong, longer than 1A, reaching nearer wing margin than 1A.

nana Riek, 1954a; Palaeocene; Redbank Plains Fm; Ipswich.

Pseudalysiinia Tonnoir, 1929 [*mimicans*]

Sc long, not quite reaching midlength of wing; ending at costal margin; R1 strong, simple; Rs leaving R1 well beyond apex of Sc; r-m long; M3+4 branching from CuA close to wing base; CuP close to CuA, both strongly curved.


mimicans Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

[Blagoderov (1994) considered this species may possibly represent a new genus of Mesosciophilidae].

Mycetophilidae gen. nov. Jell & Duncan, 1986

Sc short, close to R1, fading out rapidly laterally to costal margin; R1 curving forward to costal margin; R5 simple, arising from R1 near wing base, just touching R1 at point where latter turns to costal margin; r-m short, before fork of M; M1+2 leaving R near base, forking just basal to touching of R1 and Rs; M3 simple meeting Cu near wing base; Cu simple, strong, convex forward; single anal reaching posterior margin.

Mycetophilidae gen. nov. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.






Family RHAGIONIDAE Latreille, 1802

Atherimorpha White, 1915 [*vernalis*]

Stigma from apex of R1 between Sc and R2+3 to transverse line through r-m, tapering basally; apices of Sc, R1 and R2+3 equally spaced, well apart; r-m meeting discal cell about $\frac{1}{4}$ its length from base; M1 with short basal portion; wing apex between R4, R5.

festuca Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Atherimorpha festuca; NMVP102498; 10mm long. Line drawing (right) of Jell & Duncan, 1986, fig. 61B.



Family SIMULIIDAE Newman, 1834

Simuliid indet.

Adult: Head small; eye large, occupying most of lateral view of head. Thorax large, with mesopleuron markedly produced ventrally; foreleg with coxa longer than wide and trochanter smaller than coxa. Abdomen longer than head and thorax combined, heavily sclerotised only dorsally and at apex.

Larva: Head subrectangular, longer than wide, heavily sclerotised, with base supporting cephalic fan ray extending from anterolateral corner of head, with fan rays more than half length of head, with prominent anteromedial projection on head, with 7-8 rays per fan. Abdomen only slightly expanded towards apex around segments 5-7; proleg large, extending well under head, truncated, sclerotised at apex; terminal sucker well-developed, with sclerotised radial bars about $\frac{1}{3}$ diameter of sucker.

Simuliid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Simuliid indet.; adult
NMVP 103050; x12.
Line drawing (left) of
Jell & Duncan, 1986, fig.
60A.

Simuliid indet.; larva
(far right)
NMVP103178; x8.
Line drawing (right)
of Jell & Duncan,
1986, fig. 60C.



Family TIPULIDAE Leach, 1815


Protolimnophila Riek, 1954a [*superba*]

Wing like Archilimnophila but with discal cell more truncate basally, with M1 and M2 more widely separated, with M more sinuous. m-cu sloping in opposite direction; CuP almost or completely reaching wing margin.


superba Riek, 1954a; Palaeocene; Redbank Plains Fm; Ipswich.



Protolimnophila superba; UQF14344; 10mm long. Line drawing (above) of Riek, 1954a, fig. 1.




Tipulid indet.; NMVP103289; 3.5mm long.




Tipulid pupa indet.; NMVP103254; x10.



Tipulid pupa indet.; NMVP 103188; x7.



Tipulid larva indet.; NMVP 103251; x9



Limoniid indet.; NMVP103287; 5mm long. Line drawing (above) of Jell & Duncan, 1986, fig. 50A.

Tipulid? indet.

Head small, with large eyes dorsolaterally, with elongate frons; antennae geniculate above scape; thorax deep, with greatly expanded mesonotum, with small prosutural portion; legs extremely long and slender; hind femur (1.5mm long) more than half abdominal length; abdomen long,, widest near midlength, tapering posteriorly to acute termination.

Tipulid? indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Tipulid larva indet.

Head small, sclerotised, well back on body with only half protruding forward of the thorax, with distinct median longitudinal furrow; thorax and abdomen apparently soft; free pleural margins weakly convex;

Tipulid larva indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Tipulid pupae indet.

These pupae are assigned here by comparison with the pupae of living genera *Ischnotoma* and *Antocha*.

Tipulid pupa 1 indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Tipulid pupa 2 indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family LIMONIIDAE Engel, 1915

Limoniid indet

Wing long, narrow; Sc long; R1 long; Rs 4-branched; r-m from R5 to M; long distances between forks in general but particularly on Rs1

Limoniid indet. 1 Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Limoniid indet. 2 Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.


Family **CERATOPOGONIDAE** Skuse, 1890

Leptoconops Skuse, 1890 [*stygius*]


C poorly defined beyond R1; origin of R4+5 and M near base of wing; venation generally similar to that of living members of the genus.

Simuliid indet. 2 Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

[Generic assignment by Borkent (1997)]



Leptoconops sp.; NMVP 103203; 1.8mm long.



Leptoconops sp.; NMVP103203. Sketches from Borkent, 1997, fig. 2G,H.

Family **TILLYARDIPTERIDAE**

Lukashevich & Shcherbakov, 1998



Tillyardiptera Lukashevich & Shcherbakov, 1998 [*prima*]

C thick along anterior margin (except base), narrow beyond R4 apex; Sc terminating distal of R4+5 bifurcation, the latter level with that of M3+4; free R2 absent; R4+5 fork about 7 times longer than R4+5 stem; M1+2 fork as long as d cell; Sc apex at 61%; R fork at 29%; Rs fork at 54%; M fork at 45%; CuA apex at 64%; CuP apex at 53% of wing length.


prima Lukashevich & Shcherbakov, 1998; Mt Crosby Fm; UTrias; Ipswich.



Tillyardiptera prima; BMNHIn 44958; 10mm long. Line drawing (above right) from Lukashevich & Shcherbakov, 1998, fig. 2.



Archexyela crosbyi; UQC2232; 9mm long. Line drawing (above) of Riek, 1955, fig. 1. Second specimen mentioned by Riek (1955) UQC2233 (top and middle at right (part and counterpart), 9mm long. *Archexyela* sp. nov., QMF44154 (lower at right), 10mm long.



Eoichneumon duncanae; NMVP103104; 7mm long. Line drawing (above) of Jell & Duncan, 1986, fig. 65B.

Order HYMENOPTERA

Suborder SYMPHYTA



Gerstaecker, 1867

Family XYELIDAE Newman, 1835

Archexyela Riek, 1955 [*crosbyi*]

Forewing: like modern *Macroxyela* but apical cross vein from pterostigma to Rs passing to R2+3 not to stem of Rs; Sc free to base of wing; M curved back towards apex of wing.

crosbyi Riek, 1955; UTrias; MtCrosby Fm; Ipswich.



Suborder APOCRITA

Gerstaecker, 1867

Family EOICHNEUMONIDAE Jell & Duncan, 1986

Eoichneumon Jell & Duncan, 1986 [*duncanae*]

Forewing with basal m-cu joining M just after its separation from Rs; areolet 6-sided. Metasoma like Pimplinae (Ichneumonidae), with tergites short and broad; 1st tergite with pair of diverging carinae arising from base, only slightly longer than succeeding segments; ovipositor strongly exserted; basitarsus of hindleg longer than segments 2-5 combined.

duncanae Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **AULACIDAE** Shuckard, 1841**Aulacid** indet.

Antenna 14-segmented, with short scape and funicle; Mesoscutellum with deep percurrent notaulices; scutellum large, as long as scutum; metanotum half length of scutellum; propodium abruptly declivous; metasoma inserted high, well above hind coxal insertion. Wings with long radial cell ending before apex, with thickened margin from pterostigma to apex of radial cell; pterostigma narrow.

Aulacid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Aulacid indet.; NMVP103168; x10. Line drawing (above) of Jell & Duncan, 1986, fig. 65E.

Family **PROCTOTRUPIDAE** Latreille, 1802**Proctotrupid** indet.


Propodeum coarsely ornamented; apex produced into slight neck; metanotum vertical at meson; scutellum raised; hindleg with large coxa; trochanter long, thin, more than 0.5 length of coxa; femur long, thin, narrow at base; tibia as long as femur; middle leg shorter than hindleg, with small coxa; metasoma arising from high on mesosoma, consisting mainly of greatly enlarged first segment, with its sternite defined at base ventrally and over very short area dorsally, with tergite produced ventrally to enclose sternite over distal two thirds.

Proctotrupid indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.


Superfamily **BETHYLOIDEA** Haliday, 1840**Bethyloid?** indet.

Head elongate, flattened; antennae and large lateral eyes present but poorly preserved; thorax as long as abdomen; wings folded back along abdomen, not quite reaching end of body; legs incomplete, coxae and femora markedly expanded, with more slender tibiae and tarsi. Metasoma well segmented, expanded throughout but slightly more enlarged anteriorly.

Bethyloid? indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Proctotrupid indet.; NMVP103106; x10. Line drawing (above) of Jell & Duncan, 1986, fig. 65A.



Bethyloid? indet.; NMVP103320; 5mm long. Line drawing (left) of Jell & Duncan, 1986, fig. 68F.



Pemphredonine indet.; NMVP102740; 5mm long.
Line drawing of Jell & Duncan, 1986, fig. 68C.



Family **SPHECIDAE** Leach, 1815

Pemphredonine? indet.

Head with occipital carinae converging forward, not curved around foramen. Propleuron longer than wide; pronotum rounded posterolaterally, with posterolateral spiracle cover lobe sharply differentiated from central extension of pronotum; episternum large, with fore margin clearly defined at meson, with short concave anterolateral margin; prepectus evident; middle coxa small, with elongate trochanter; hind coxa larger, stouter, trochanter not clearly defined, femur only slightly widened, with a narrow dorsal flange defined by a groove or ridge, with tibia narrower than femur. Metasoma longer than mesosoma, with sutures between all 7 segments; first sternite very short, with convex apical margin; 3rd-6th sternites becoming shorter, with caudal margins becoming more concave. posterior with pair of large claspers over apical segment laterally.

Pemphredonine? indet. Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.

Family **FORMICIDAE??** Latreille, 1802

Cretacoformica Jell & Duncan, 1986

[*explicata*]

Single petiole segment; abdominal segment IV non-tubulate; pronotum subrectangular, narrower than rest of mesosoma; forewing with reduced



Cretacoformica explicata; NMVP102501; wingspan 7.5mm. Line drawing of wing venation from Jell & Duncan, 1986, fig. 69C.

venation, with enclosed costal cell, with pterostigma, with M+CuA, M, CuA and 1A not tracheate. Legs similar, various segments longer to posterior; femur expanded medially; tibia expanded distally; basitarsus of uniform width, longer than rest of tarsus combined.

explicata Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.


[Holldobler & Wilson (1990) doubted the family assignment of *Cretacoformica* and Naumann (1993) gave a very detailed description and discussion of the taxon. He concluded that it could not be satisfactorily assigned to any family of the Apocrita principally because the nature of the petiole could not be discerned due to the attitude of preservation—a difficulty expressed by Jell & Duncan (1986).]

Family **PRAEAULACIDAE** Rasnitsyn, 1972


Westratia Jell & Duncan, 1986 [*nana*]

First funicle segment longest. Hindleg short; femur distinctly shorter than metasoma; basitarsus shorter than segments 2-5 combined. Anterior of mesoscutum apparently not produced into transverse ridges. Wing with wide costal cell, prominent pterostigma, slightly reduced forewing venation similar to that of *Austronia rubrithorax* Riek, 1955. Ovipositor strongly exserted.

nana Jell & Duncan, 1986; LCret; Koonwarra FB; Gippsland.



Westratia nana; NMVP103105; x18 (below left). Line drawing (above) of Jell & Duncan, 1986, fig. 65D.



Westratia nana; NMVP103324; x18. Line drawing (above) of Jell & Duncan, 1986, fig. 65C.



Order MEGALOPTERA (alderflies)

Family SIALIDAE

Austrosialis Tillyard [*ignicollis*]

RP3+4 simple; MA 2-branched; MP1+2 2-branched;
MP1+2 forked at about same level as representative;
MP3+4 2-branched; forks of MA, MP and CuA not
differing much in depth.

sp. Lambkin, 1992; Palaeocene; Redbank
Plains Fm; Ipswich.



Austrosialis sp.; QMF21638; 11mm long. Line drawing of Lambkin, 1992, fig. 1.

ACKNOWLEDGEMENTS

I thank Andrew Ross and Phillip Crabb, Natural History Museum for providing photographs of specimens in the collection under their care, Bob Jones, Australian Museum for the loan of specimens, Robert Beattie for providing upto date information on the Belmont/Warners Bay fossil insect sites, Paul Avern and Sue Parfrey, Queensland Museum, for help with photography and Megan Lloyd, Queensland Museum, for help in obtaining difficult literature.

LITERATURE CITED

- ANSORGE, J. 1996. Insekten aus dem oberen Lias von Grimmen (Vorpommern, Norddeutschland). *Neue Paläontologische Abhandlungen* 2: 1-132, pls 1-17.
- ARCHER, M., GODTHELP, H., HAND, S.J. & MEGIRIAN, D. 1989. Fossil mammals of Riversleigh, northwestern Queensland: preliminary overview of biostratigraphy, correlation and environmental change. *Australian Zoologist* 25: 29-65.
- ARCHER, M., GODTHELP, H. & HAND, S.J. 1991. Riversleigh: the story of animals in ancient rainforests of inland Australia. (Reed: Sydney). 264p.
- BALME, B.E. 1964. The palynological record of Australian pre-Tertiary floras. Pp. 49-80. In Cranwell, L. (ed.) *Ancient Pacific floras*. (University of Hawaii Press: Honolulu).
- BLAGODEROV, V.A. 1994. Dipterans (Mesoscophilidae) from the Lower Cretaceous of Transbaykal. *Paleontological Journal* 27(1A): 123-130.
- BORKENT, A. 1997. Upper and Lower Cretaceous biting midges (Ceratopogonidae: Diptera) from Hungarian and Austrian amber and the Koonwarra Fossil Bed of Australia. *Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie)* 249: 1-10.
- CARROLL, E.J. 1962. Mesozoic fossil insects from Koonwarra, South Gippsland, Victoria. *Australian Journal of Science* 25: 264-265.
- CARPENTER, F. 1992. Superclass Hexapoda. Pp. 1-655. In Kaesler, R. (ed.) *Treatise of Invertebrate Paleontology, Part R. Arthropoda 4, Volumes 3 & 4*. (Geological Society of America & University of Kansas: Lawrence, Kansas).
- DAVIS, C. 1942. Hemiptera and Copeognatha from the Upper Permian of New South Wales. *Proceedings of the Linnean Society of New South Wales* 67: 111-112.
1943. A new species of *Permithone* (Neuroptera Planipennia) from the Upper Permian of New South Wales. *Proceedings of the Linnean Society of New South Wales* 68: 11-12.
- DETTMANN, M.E. 1986. Early Cretaceous palynoflora of subsurface strata correlative with the Koonwarra Fossil Bed, Victoria. *Memoirs of the Association of Australasian Palaeontologists* 3: 79-110.
- DODDS, B. 1949. Mid-Triassic Blattoidea from the Mount Crosby Insect Bed. *University of Queensland Papers, Department of Geology* 3: 1-11.
- DRINNAN, A.N. & CHAMBERS, C. 1986. Flora of the Lower Cretaceous Koonwarra Fossil Bed (Korumburra Group), South Gippsland, Victoria. *Memoirs of the Association of Australasian Palaeontologists* 3: 1-77.
- DUNCAN, I.J., BRIGGS, D.E.G. & ARCHER, M. 1998. Three-dimensionally mineralized insects and millipedes from the Tertiary of Riversleigh, Queensland, Australia. *Palaeontology* 41: 835-851.
- DUNCAN, I.J. & BRIGGS, D.E.G. 1996. Three-dimensionally preserved insects. *Nature* 381: 30-31.
- DUNSTAN, B. 1916. Mesozoic and Tertiary insects of Queensland and New South Wales. *Stratigraphical features. Geological Survey of Queensland Publication* 253: 1-10, pls 8-9.
1923. Mesozoic insects of Queensland. Part 1. Introduction and Coleoptera. *Geological Survey of Queensland Publication* 273: 1-74, pls 1-7.
- ENGEL, B.A. 1966. Explanatory notes on the Newcastle 1:250,000 Geological Sheet, SI 56-2. (Geological Survey of New South Wales: Sydney). 63p.
- ETHERIDGE, R. Jr 1913. Palaeontological contributions to the geology of Western Australia 4. *Geological Survey of Western Australia Bulletin* 55: 1-34.
- ETHERIDGE, R. Jr & OLLIFF, A.S. 1890. The Mesozoic and Tertiary insects of New South Wales. *Memoirs of the Geological Survey of New South Wales, Palaeontology* 7: 1-18, pls 1-2.
- EVANS, J.W. 1943a. Two interesting Upper Permian Homoptera from New South Wales. *Transactions of the Royal Society of South Australia* 67: 7-9, pl. 1.
- 1943b. Upper Permian Homoptera from New South Wales. *Records of the Australian Museum* 21: 180-198.
1947. A new fossil homopteran from Kimble's Hill, Belmont (Upper Permian). *Records of the Australian Museum* 21: 431-432.
1950. A re-examination of an Upper Permian insect, *Paraknightia magnifica* Ev. *Records of the Australian Museum* 22: 246-250.
1956. Palaeozoic and Mesozoic Hemiptera (Insecta). *Australian Journal of Zoology* 4: 165-258.
1957. Some aspects of the morphology and inter-relationships of extinct and recent Homoptera. *Transactions of the Royal Entomological Society of London* 109: 275-294.
1958. New Upper Permian Homoptera from the Belmont Beds (Homoptera: Insecta). *Records of the Australian Museum* 24: 109-114.

1961. Some Upper Triassic Hemiptera from Queensland. *Memoirs of the Queensland Museum* 14: 13-23.
1963. The systematic position of the Ipsvichiidae (Upper Triassic Hemiptera) and some new Upper Permian and Middle Triassic Hemiptera from Australia (Insecta). *Journal of the Entomological Society of Queensland* 2: 17-23, pl. 1.
- EVANS, J.W. 1971. Some Upper Triassic Hemiptera from Mount Crosby, Queensland. *Memoirs of the Queensland Museum* 16: 145-152.
- FLECK, G. & NEL, A. 2003. Revision of the Mesozoic family Aechmidae (Odonata: Anisoptera). *Zoologica* 153: 1-172.
- FLEMING, P.J.G. 1966. Notes on Triassic fossils from the Mount Crosby Formation in the Parish of Chuwar. *Queensland Government Mining Journal* 63: 119-120.
- FLETCHER, H.O. 1971. Catalogue of type specimens of fossils in the Australian Museum Sydney. *Memoirs of the Australian Museum* 13: 1-167.
- GILL, E.D. 1955. Fossil insects, centipede and spider. *Victorian Naturalist* 72: 87-92.
- HAMILTON, K.G.A. 1992. Lower Cretaceous Homoptera from the Koonwarra Fossil Bed in Australia, with a new superfamily and synopsis of Mesozoic Homoptera. *Annals of the Entomological Society of America* 85: 423-430.
- HAWLEY, S.P. & BRUNTON, J.S. 1995. The Newcastle Coalfield. Notes to accompany the 1:100,000 Newcastle Coalfield regional geology map. Department of Mineral Resources, Geological Survey of New South Wales Report GS1995/256: 1-93.
- HILL, D., PLAYFORD, G. & WOODS, J.T. 1965. Triassic fossils of Queensland. (Queensland Palaeontographical Society: Brisbane).
1968. Cretaceous fossils of Queensland. (Queensland Palaeontographical Society: Brisbane).
1970. Cretaceous fossils of Queensland. (Queensland Palaeontographical Society: Brisbane).
- HOLLOBLER, B. & WILSON, E.O. 1990. The ants. (Belknap Press: Cambridge).
- JELL, P.A. 1993. Late Triassic homopterous nymph from Dinmore, Ipswich Basin. *Memoirs of the Queensland Museum* 33: 360.
- JELL, P.A. & DUNCAN, P.M. 1986. Invertebrates, mainly insects, from the freshwater, Lower Cretaceous, Koonwarra Fossil Bed (Korumburra Group), South Gippsland, Victoria. *Memoirs of the Association of Australasian Palaeontologists* 3: 111-205.
- JELL, P.A. & LAMBKIN, K.J. 1993. Middle Triassic orthopteroid (Titanoptera) insect from the Esk Formation at Lake Wivenhoe. *Memoirs of the Queensland Museum* 33: 258.
- JELL, P.A. & ROBERTS, J. (eds) 1986. Plants and invertebrates from the Lower Cretaceous, Koonwarra Fossil Bed, South Gippsland, Victoria. *Memoirs of the Association of Australasian Palaeontologists* 3: 1-205.
- KOVALEV, V.G. 1983. New dipteran family from the Triassic of Australia and its presumed descendants (Diptera, Crosaphididae fam.n., Mycetobiidae). *Entomology Review* 62(4): 130-136.
- KUKALOVÁ, J. 1966. Protelytroptera from the Upper Permian of Australia, with a discussion of the Protocoleoptera and Paracoleoptera. *Psyche* 73: 89-111, pl. 8.
- KUKALOVÁ-PEK, J. 1988. A substitute name for the extinct genus *Stenelytron* Kukalová (Protelytroptera). *Psyche* 94: 339.
1991. Fossil history and the evolution of hexapod structures. Pp. 141-179. In CSIRO (ed.) *The insects of Australia: A textbook for students and research workers*. 2nd ed. vol. 1 (Melbourne University Press: Carlton). 542p.
- LAMBKIN, K.J. 1987. A re-examination of *Euporismites balli* Tillyard from the Palaeocene of Queensland (Neuroptera: Osmiidae: Kempyninae). *Neuroptera International* 4: 295-300.
1988. A re-examination of *Lithosmylidia* Riek from the Triassic of Queensland with notes on Mesozoic 'osmylid-like' fossil Neuroptera (Insecta : Neuroptera). *Memoirs of the Queensland Museum* 25: 445-458.
- 1992a. Re-examination of the venation of *Osmiopsychops spillerae* Tillyard from the Triassic of Queensland. *Memoirs of the Queensland Museum* 32: 183-188.
- 1992b. A record of *Austrosialis* Tillyard from the Queensland Palaeocene (Insecta: Megaloptera: Sialidae). *Queensland Naturalist* 31: 84-86.
- LITTLE, M.P., BOYD, R.L., BRUNTON, J., IVES, M., RIGBY, R. & TOBIN, C. 1996. The Newcastle Coal Measures: stratigraphy, sedimentology and sequence stratigraphy. Pp. 77-84. In *Proceedings of the 30th Newcastle Symposium on Advances in the study of the Sydney Basin* (Newcastle University: University).
- LUKASHEVICH, E.D. & SHCHERBAKOV, D.E. 1998. A new Triassic family of Diptera from Australia. Pp. 81-89. In *Proceedings of the First International Paleontological Conference*. (AMBA projects: Moscow).
- MARTYNOVA, O.M. 1948. Data on the evolution of the Mecoptera. *Trudy Paleontologicheskogo instituta akademii nauk SSSR* 14: 1-76, pls 1-3.
- McKEOWN, K.C. 1937. New fossil insect wings (Protohemiptera, Family Mesotitanidae). *Records of the Australian Museum* 20: 31-37, pls 4-7.
- NAUMANN, I.D. 1993. The supposed Cretaceous ant *Cretacoformica explicata* Jell and Duncan (Hymenoptera). *Journal of the Australian Entomological Society* 32: 355-356.
- PARFREY, S.M. 1996. Geological Survey of Queensland Catalogue of type, figured and cited

- fossils. (Queensland Department of Mines & Energy: Brisbane). 268p.
- RASNITSYN, A.P. 1992. *Strashila incredibilis*, a new enigmatic mecopteroid insect with possible siphonapteran affinities from the Upper Jurassic of Siberia. *Psyche* 99: 323-333.
- RASNITSYN, A.P. & QUICKE, L.J. (eds) 2002. History of insects. (Kluwer: Dordrecht, Holland). 517p.
- RIEK, E.F. 1950. A fossil mecopteran from the Triassic beds at Brookvale, N.S.W. Records of the Australian Museum 22: 254-256.
- 1952a. The fossil insects of the Tertiary Redbank Plains Series. Part 1: an outline of the fossil assemblage with description of the fossil insects of the Orders Mecoptera and Neuroptera. University of Queensland Papers, Department of Geology 4: 3-14, pls 1-2.
- 1952b. Fossil insects from the Tertiary sediments at Dinmore, Queensland. University of Queensland Papers, Department of Geology 4: 17-22, pl. 1.
1953. Fossil mecopteroid insects from the Upper Permian of New South Wales. Records of the Australian Museum 23: 55-87, pls 5-6.
- 1954a. The fossil Diptera of the Tertiary Redbank Plains series, Queensland. Proceedings of the Linnean Society of New South Wales 79: 58-60.
- 1954b. A re-examination of the Upper Tertiary mayflies described by Etheridge and Olliff from the Vegetable Creek tin-field. Records of the Australian Museum 23: 159-160, pl. 10.
- 1954c. Further Triassic insects from Brookvale, N.S.W. (Orders Orthoptera Saltatoria, Protorthoptera, Perlaria). Records of the Australian Museum 23: 161-168, pl. 11.
- 1954d. A second specimen of the dragonfly *Aeschnidiopsis flindersiensis* (Woodward) from the Queensland Cretaceous. Proceedings of the Linnean Society of New South Wales 79: 61-62.
1955. Fossil insects from the Triassic beds at Mt Crosby Queensland. Australian Journal of Zoology 3: 654-691, pls 1-4.
1956. A re-examination of the mecopteroid and orthopteroid fossils (Insecta) from the Triassic beds at Denmark Hill, Queensland, with descriptions of further specimens. Australian Journal of Zoology 4: 98-110, pls 1-2.
1962. Fossil insects from the Triassic at Hobart, Tasmania. Papers and Proceedings of the Royal Society of Tasmania 96: 39-40, pls 1-2.
- 1967a. Further evidence of the Panorpidae in the Australian Tertiary (Insecta: Mecoptera). Journal of the Australian Entomological Society 6: 71-72.
- 1967b. A fossil cockroach (Blattodea: Poroblattinidae) from Mt. Nicholas Coal Measures, Tasmania. Journal of the Australian Entomological Society 6: 73.
- 1968a. *Robinjohnia tillyardi* Martynova, a mecopteran from the Upper Permian of Belmont, New South Wales. Records of the Australian Museum 27: 299-302, pl. 44.
- 1968b. Undescribed fossil insects from the Upper Permian of Belmont, New South Wales (with an appendix listing the described species). Records of the Australian Museum 27: 303-310, pl. 45.
- 1968c. On the occurrence of fossil insects in the Mesozoic rocks of Western Australia. Records of the Australian Museum 27: 311-312, pl. 46.
- 1970a. Lower Cretaceous fleas. *Nature* 227: 746-747.
- 1970b. Origin of the Australian insect fauna. Pp. 593-598. In Theron (ed.) Proceedings and Papers of the Second Gondwana Symposium, South Africa, 1970.
- 1970c. Fossil history. Pp. 168-186. In CSIRO (ed.) The insects of Australia: A textbook for students and research workers. (Melbourne University Press: Carlton). 1029p.
1971. The presumed heads of Homoptera (Insecta) in the Australian Upper Permian. *Palaeontology* 14: 211-221.
1973. A Carboniferous insect from Tasmania. *Nature* 244: 455-456.
1976. Neosecoptera, a new insect suborder based on specimen discovered in the Late Carboniferous of Tasmania. *Alcheringa* 1: 227-234.
- ROZEFELDS, A.C. 1985. A fossil zygopteran nymph (Insecta: Odonata) from the Late Triassic Aberdare Conglomerate, southeast Queensland. Proceedings of the Royal Society of Queensland 96: 25-32.
- 1988a. Insect leaf mines from the Eocene Anglesea locality, Victoria, Australia. *Alcheringa* 12: 1-6.
- 1988b. Lepidoptera mines in *Pachypteris* (Coryspermaceae: Pteridospermo-phyta) from the Upper Jurassic/Lower Cretaceous Battle Camp Formation, north Queensland. Proceedings of the Royal Society of Queensland 99: 77-81.
- ROZEFELDS, A.C. & SOBBE, I. 1987. Problematic insect leaf mines from the Upper Triassic Ipswich Coal Measures of southeastern Queensland. *Alcheringa* 11: 51-57.
- SHAROV, A.G. 1968. Phylogeny of the Orthopteroidea. *Trudy Paleontologicheskogo Instituta Akademiyi Nauk SSSR* 118: 1-218.
- SMITHERS, C.N. 1972. The classification and phylogeny of the Psocoptera. *Memoirs of the Australian Museum* 14: 1-107.
- SUKATCHEVA, I.D. 1982. Istoricheskoe razvitiye otryada rucheinikov [Historical development of the caddisflies]. *Trudy Paleontologicheskogo Instituta Akademiyi Nauk SSSR* 197: 1-107.
- TALENT, J.A., DUNCAN, P.M. & HANDBY, P.L. 1966. Early Cretaceous feathers from Victoria. *Emu* 66: 81-86, pls 1-3.
- TILLYARD, R.J. 1917. Mesozoic insects of Queensland. No. 1. Planipennia, Trichoptera, and the new order Protomecoptera. Proceedings of the

- Linnean Society of New South Wales 42: 175-200, pls 7-9.
- 1918a. Mesozoic insects of Queensland. No. 2. The fossil dragonfly *Aeschnidiopsis* (*Aeschna*) *flindersiensis* Woodward, from the Rolling Downs (Cretaceous) Series. Proceedings of the Linnean Society of New South Wales 42: 676-692, pls 42-43.
- 1918b. Permian and Triassic insects from New South Wales, in the collection of Mr. John Mitchell. Proceedings of the Linnean Society of New South Wales 42: 720-756.
- 1918c. A fossil insect wing from the roof of the coal-seam in the Sydney Harbour Colliery. Proceedings of the Linnean Society of New South Wales 43: 260-264.
- 1918d. Mesozoic insects of Queensland. No. 3. Odonata and Protodonata. Proceedings of the Linnean Society of New South Wales 43: 417-436, pls 44-45.
- 1918e. Mesozoic insects of Queensland. No. 4. Hemiptera Heteroptera: the family Dunstaniidae with a note on the origin of the Heteroptera. Proceedings of the Linnean Society of New South Wales 43: 568-592, pl. 59.
- 1919a. Mesozoic insects of Queensland. No. 5. Mecoptera, the new order Paratrachoptera, and additions to Planipennia. Proceedings of the Linnean Society of New South Wales 44: 194-212, pl. 67.
- 1919b. A fossil insect wing belonging to the new order Paramecoptera, ancestral to the Trichoptera and Lepidoptera, from the upper Coal-Measures of Newcastle, N.S.W. Proceedings of the Linnean Society of New South Wales 44: 231-256, pls 12-13.
- 1919c. Mesozoic insects of Queensland. No. 6. Blattoidea. Proceedings of the Linnean Society of New South Wales 44: 358-382.
- 1919d. Mesozoic insects of Queensland. No. 7. Hemiptera Homoptera; with a note on the phylogeny of the suborder. Proceedings of the Linnean Society of New South Wales 44: 857-896, pls 7-9.
- 1921a. Mesozoic insects of Queensland. No. 8. Hemiptera Homoptera (Contd.). Proceedings of the Linnean Society of New South Wales 46: 270-284, pls 16-21.
- 1921b. Two fossil insect wings in the collection of Mr. John Mitchell, from the Upper Permian of Newcastle, N.S.W., belonging to the Order Hemiptera. Proceedings of the Linnean Society of New South Wales 46: 413-422, pl. 35.
- 1922a. Some new Permian insects from Belmont, N.S.W. in the collection of Mr John Mitchell. Proceedings of the Linnean Society of New South Wales 47: 279-292, pls 33-34.
- 1922b. Mesozoic insects of Queensland. No. 9. Orthoptera, and additions to the Protorthoptera, Odonata, Hemiptera and Planipennia. Proceedings of the Linnean Society of New South Wales 47: 447-470, pls 51-53.
- 1922c. An insect wing in a crystal of selenite (order Orthoptera). Records of the Geological Survey of New South Wales 10: 205-207, pl. 15.
- 1923a. On a Tertiary fossil insect wing from Queensland (Homoptera Fulgoroidea), with description of a new genus and species. Proceedings of the Royal Society of Queensland 35: 16-20, pl. 1.
- 1923b. Mesozoic insects of Queensland. No. 10. Summary of the Upper Triassic insect fauna of Ipswich, Q. (With an appendix describing new Hemiptera and Planipennia). Proceedings of the Linnean Society of New South Wales 48: 481-498, pl. 43.
1924. Upper Permian Coleoptera and a new order from the Belmont Beds, New South Wales. Proceedings of the Linnean Society of New South Wales 49: 429-435, pls 45-46.
1925. A new fossil insect wing from Triassic beds near Deewhy, N.S.W.. Proceedings of the Linnean Society of New South Wales 50: 374-377, pl. 36.
- 1926a. Upper Permian insects of New South Wales 1. Introduction and the Order Hemiptera. Proceedings of the Linnean Society of New South Wales 51: 1-30, pl. 1.
- 1926b. Upper Permian insects of New South Wales. 2. The orders Mecoptera, Paramecoptera and Neuroptera. Proceedings of the Linnean Society of New South Wales 51: 265-282, pls 15-16.
1929. Permian Diptera from Warner's Bay, N.S.W. Nature 123: 778-779.
- 1935a. Upper Permian insects of New South Wales. 3. The order Copeognatha. Proceedings of the Linnean Society of New South Wales 60: 265-279.
- 1935b. Upper Permian insects of New South Wales. 4. The order Odonata. Proceedings of the Linnean Society of New South Wales 60: 374-384, pl. 12.
- 1935c. Upper Permian insects of New South Wales. 5. The order Perlaria or stoneflies. Proceedings of the Linnean Society of New South Wales 60: 385-391, pl. 12.
1937. A small collection of fossil cockroach remains from the Triassic Beds of Mount Crosby, Queensland. Proceedings of the Royal Society of Queensland 48: 35-40.
1916. Mesozoic and Tertiary insects of Queensland and New South Wales. Description of the fossil insects. Geological Survey of Queensland Publication 253: 11-60, pls 1-7.
- TINDALE, N.B. 1945. Triassic insects of Queensland. 1. *Eoses*, a probable lepidopterous insect from the Triassic beds of Mt. Crosby, Queensland. Proceedings of the Royal Society of Queensland 56: 37-46, pl. 5.
1980. Origin of the Lepidoptera, with description of a new mid-Triassic species and notes on the

- origin of the butterfly stem. *Journal of the Lepidopterists' Society* 34: 263-285.
- VISHNIAKOVA, V.N. 1981. New Palaeozoic and Mesozoic lophioneurids (Thripida, Lophioneuridae). *Trudy Paleontologicheskogo instituta, Akademiya Nauk SSSR* 183: 43-63.
- WALDMAN, M. 1971. Fish from the freshwater Lower Cretaceous of Victoria, Australia, with comments on the palaeo-environment. *Special Papers in Palaeontology* 9: 1-124, pls 1-18.
- WHITWORTH, H.F. 1954. A faked fossil. *Museum Journal* 53: 319-320.
- WILLMAN, R. 1989a. Rediscovered: *Permotipula patricia*, the oldest known fly. *Naturwissenschaften* 76: 375-377.
- 1989b. Evolution und Phylo- genetisches System der Mecoptera (Insecta: Holometabola). *Abhandlungen der Sencken- bergischen Naturforschenden Gesellschaft* 544: 1-153.
- WOODWARD, H. 1884. On the wing of a neuropterous insect from the Cretaceous limestone of Flinders River, north Queensland, Australia. *Geological Magazine*, 3rd decade 1: 337-339, pl. 1.
- WOOTON, R.J. 1963. Actinoscytinidae (Hemiptera-Heteroptera) from the upper Triassic of Queensland. *Annals and Magazine of Natural History*, series 13, 6: 249-255.

INDEX

Entries for Orders are in upper case and bold face type. Entries for other suprageneric taxa (mainly families) are in upper case. All species entries are in lower case and are followed by the generic assignment. Generic and specific entries in italics indicate the taxon has gone into synonymy.

<i>aberrans</i> Triassoscytinopsis	59
<i>ableptus</i> Permotanyderus	97
<i>abnormis</i> Ademosynoides	70
<i>abrupta</i> Austrocypha	31
<i>abrupta</i> Mesonirvana	56
<i>Actinoscytina</i>	65
<i>acutipennis</i> Ipsicia	45
<i>Adelidium</i>	79
<i>Ademosyne</i>	70
<i>Ademosynoides</i>	70
<i>adscita</i> Agetochoristella	88
<i>adunca</i> Ademosyne	70
<i>Aeroplana</i>	30
AEROPLANIDAE	30
AESCHNIDIIDAE	15
<i>Aeschnidiopsis</i>	15
<i>affine</i> Mesogereon	46
<i>affine</i> Permosyllidium	50
<i>affinis</i> Mesoscytina	38
<i>affinis</i> Paradunstania	42
<i>affinis</i> Permosyne	71
<i>Agetochoristella</i>	88
AGETOPANORPIDAE	88
<i>alata</i> Crosbella	55
<i>Alotrifidus</i>	36
<i>alternata</i> Ademosynoides	70
<i>amplicata</i> Triassocotis	39
<i>analis</i> Triassothea	54
<i>andersoni</i> Clatrotitan	29
<i>Aneura</i>	102
<i>angaura</i> Prochoristella	92
<i>angusta</i> Eochiliocyclus	44
<i>angusta</i> Lophioneura	31
<i>angustata</i> Ademosynoides	70
<i>angustata</i> Stenovicia	62
<i>angustius</i> Elateridium	76
<i>anomala</i> Aphryganoneura	88
<i>anomala</i> Crosaphis	35
<i>anomala</i> Stenoperlidium	17
<i>anomala</i> Triassoscelis	59
<i>Anomaloscytina</i>	48
<i>Anthocorid</i> indet.	68
<i>antinodalis</i> Mesophlebia	14
<i>Antitaxinura</i>	17
<i>Apheloodes</i>	73
<i>Apheloscyta</i>	60

APHIDIDAE	35
<i>Aphryganoneura</i>	88
<i>apicalis</i> Aneura	102
APOCRITA	106
<i>Archebittacus</i>	94
ARCHEOSMYLIDAE	81
<i>Archeosmylus</i>	81
<i>Archepsychops</i>	84
ARCHESCYTINIDAE	35
<i>Archexyla</i>	106
<i>Archipanorpa</i>	89
ARCHIPANORPIDAE	89
<i>Aristopsyche</i>	92
<i>Atalophlebia</i>	11
<i>Atherimorpha</i>	103
<i>Aulacid</i> indet.	107
<i>australe</i> Eurymelidium	38
<i>australe</i> Megapsocidium	33
<i>australe</i> Protosyllidium	52
<i>australis</i> Mesochorista	91
<i>australicus</i> Triassocixius	41
<i>australiensis</i> Ademosyne	70
<i>australiensis</i> Stenoscytina	58
<i>australiensis</i> Triassagrion	15
<i>australis</i> Austropanorpa	95
<i>australis</i> Etheridgea	72
<i>australis</i> Mesoscytina	38
<i>australis</i> Neopetromantis	88
<i>australis</i> Permojassus	58
<i>australis</i> Permomerope	96
<i>australis</i> Tarwinia	99
<i>australis</i> Triassocotis	39
<i>Australoblattula</i>	19
<i>Australosymmerus</i>	101
<i>Australurus</i>	11
<i>Austroblattula</i>	5
<i>Austrochoristella</i>	98
<i>Austrocypha</i>	31
<i>Austrodictya</i>	7
<i>Austroelytron</i>	23
<i>Austroidelia</i>	18
<i>Austrolestidion</i>	7
<i>Austromylacrites</i>	20
<i>Austropanorpa</i>	95
<i>Austroprosbole</i>	48
<i>Austropsocidium</i>	33
<i>Austroscarites</i>	39
<i>Austroscytina</i>	35
<i>Austrosialis</i> sp.	110
BAETIDAE	11
<i>bairdi</i> Parachorista	91
<i>balli</i> Euporismites	82
<i>baretti</i> Austrocypha	31
<i>baronne</i> Lithosmylidia	82
<i>Beaconiella</i>	64
<i>Bekkerscytina</i>	35

belli Prochoristella	92	clarkae Neoparachorista	95
Belmontocarta	43	clathrata Mesothoris	80
belmontensis Actinoscytina	65	Clatrotitan	3, 9, 29
belmontensis Cladochorista	86	Clavopsyllidium	50
belmontensis Mesopanorpodes	91	Cleon	11
belmontensis Orthoscytina	57	COENAGRIONIDAE	16
belmontensis Permithone	81	Coenagrionid indet.	16
belmontensis Permofulgor	25	colliveri Prorhyacophila	86
belmontensis Permoglyphis	57	COLEOPTERA	70
belmontensis Permopsychops	81	collessi Chironomaptera	100
belmontensis Permosyne	71	colorata Hylicella	45
belmontensis Tychticoloides	37	compressum Mesogereon	46
Belmontia	89	concinna Prochoristella	92
BELMONTIIDAE	89	congener Ademosyne	70
Belpsylla	50	conjuncta Lophioneura	31
Bethyloid indet.	107	conservativum Dermelytron	24
bifasciata Choristopanorpa	94	corbouldi Austrodictya	7
bifida Psocoscytina	53	cordatum Adelidium	79
bilineatus Grammositus	71	costalis Archeosmylus	81
bipunctatum Elaterium	76	crassifemur Proparagryllacris	26
biserialis Permorpisma	82	Cretacochorista	90
BLATTARIA	19	Cretacoformica	108
BLATTIDAE	20	Crosaphis	35
blatteloides Samaroblatta	19	Crosbella	55
Blattotermes	22	crosbyi Archexyela	106
brachyclada Mesocixioides	41	crosbyi Xenopterum	28
brevis Ademosyne	70	cubitus Triassoaphis	35
bryani Cladochoristella	86	culleni Atalophlebia	11
bryani Eoscarteroides	44	CURCULIONIDAE	72
bryani Scolypopites	63	curvata Ademosyne	70
burrettae Psychroptilus	1, 13	cylindrica Simmondsia	78
Calamoceratid pupa indet.	87	DASCILLIDAE	73
cameroni Ademosyne	70	davisia Psyllidiana	54
Cantharid indet.	72	delicatum Eopsyllidium	36
capulus Permophilus	25	delicatus Zoropsocus	32
Centrocnemis	101	denmeadi Triassoblatta	19
cephalota Promirara	11	DERMELYTRIDAE	24
CERAMBYCIDAE	72	Dermelytron	24
CERATOPOGONIDAE	105	DERMESTIDAE	74
CERCOPIDAE	36	didymovena Xenopanorpa	96
Chanoselytron	24	DIPTERA	100
CHAOBORIDAE	100	distincta Permagra	60
Chiliocycla	38	DITOMYIDAE	101
CHILIOCYCLIDAE	38	Dixiid indet.	101
chinai Triassodoecus	67	drinnani Choristopanorpa	94
Chironomaptera	100	dubia Mesochorista	91
CHIRONOMIDAE	100	dubia Permobrachus	57
Chorista	90	dubius Permojassus	58
CHORISTIDAE	90	Dulcimanna	12
Choristopanorpa	94	duncanae Eoichneumon	106
Choristotanyderus	97	duncani Eodinotoperla	17
Cicada?	6	Duncanovelia	68
CICADELLIDAE	38	dunstani Ligavena	40
CIXIIDAE	41	dunstani Mesopsyche	92
Cladochorista	86	dunstani Mesorhynchophora	73
Cladochoristella	86	Dunstanina	42

DUNSTANIIDAE	42	frustrata Stereochorista	87
Dunstanioipsis	42	FULGORIDAE	64
duaringae Austrolestidion	7	Gelastocorid indet.	65
DYSMORPHOPTILIDAE	43	giganteus Mesogryllacris	26
Dysmorphoptiloides	43	giganteus Mesotitan	9
Dytiscid indet.	75	gingiva Chanoselytron	24
Edgariokia	32	globulus Permocapitus	10
ELATERIDAE	76	GLOSSELYTRODEA	85
Elateridium	76	gracilipes Ligavena	40
Elaterium	76	grandis Mesodiphthera	56
ELCANIDAE	26	grandis Mesothoris	80
Elcanopsis	26	grandis Metrorhynchites	79
elegans Ipsviciopsis	45	grandis Triassocoris	69
elegans Phipoides	89	granulatum Phyllelytron	23
Elliptoscarta	55	Grammositus	71
elongata Crosbella	55	granulata Lobites	71
elongata Dysmorphoptiloides	43	granulata Tillyardiopsis	73
elongata Mesacridites	18	granulata Tryoniopsis	74
elongata Mesopsyche	92	GRYLLACRIDIDAE	27
elongata Neoageta	88	Gryllacridoid indet.	28
elongata Niwratia	99	GRIPOPTERYGIDAE	17
elongata Proberothella	84	Gyrinid indet.	78
elongatum Stenopsocidium	34	HAGLIDAE	26
Elytrathrix	24	Helodid indet.	76
emnavillensis Cleon	11	HEMIPTERA	35
enervatum Labidelytron	25	Heterochterus	65
Eodinotoperla	17	Heterojassus	66
Eochiliocyclus	44	Heteronella	66
Eocorona	8, 86	HETEROPTERA	64
Eoichneumon	106	Heteroscytina	66
EOICHNEUMONIDAE	106	Hexascytina	66
Eopsyllidium	36	hillae Xenochoristella	98
Eoscarterella	43	hirsuta Elytrathrix	24
EOSCARTERELLIDAE	43	hirtus Permophilus	25
Eoscarteroides	44	Homaloscytina	56
Eoscytina	36	Homopterulum	40
Eoses	8, 91	Hydraenid indet.	78
EPHEMEROPTERA	11	Hydrophyllid indet.	78
epiphleboides Triassolestes	16	HYDROPHYLIDAE	77
Etheridgea	72	Hylicella	45
Eupincombea	47	HYLICELLIDAE	44
Euporismites	82	HYMENOPTERA	106
EURYMELIDAE	44	iani Prorhyacophila	86
Eurymelidium	38	IDELIIDAE	18
EUSTHENIIDAE	17	imperfecta Australosymmerus	101
exilis Archebittacus	94	imperfecta Austroscytina	35
exilis Prochoristella	92	imperfecta Prohagla	27
explicata Cretacoformica	108	incerta Palaeoviccia	60
extensa Duncanovelia	68	incompleta Anomaloscytina	48
femoratus Austroscarites	39	incompleta Eoscytina	36
fennahi Beaconiella	64	incompleta Triassoscytina	59
festuca Atherimorpha	103	indistincta Orthoscytina	57
Fletcheriana	46	indistinctus Permofulgor	25
flindersiensis Aeschnidiopsis	15	insignita Triassoblatta	19
folium Phyllelytron	23	insolita Permopsylloides	51
FORMICIDAE??	108	intercalata Samaroblatta	19

intermedia Ademosyne	70	major Ademosyne	70
intermedia Reeveana	74	major Reeveana	74
intermedia Triassoblatta	19	MALACODERMIDAE	79
interruptus Alotrifidus	36	marginata Mesojassula	49
ipsviciensis Australoblattula	19	marksei Heteronella	66
ipsviciensis Mesojassus	44	maryae Steinhartia	29
Ipsvicia	45	MASTOTERMITIDAE	22
IPSVICIIDAE	45	maxima Lophiocypha	31
Ipsviciopsis	45	MECOPTERA	88
irregularis Orthoscytina	57	media Eoscarterella	43
ISOPTERA	22	media Psocopsyllidium	53
JASCOPIIDAE	40	megaloprepia Prochoristella	92
jelli Homopterulum	40	MEGALOPTERA	110
jonesi Ipsvicia	45	Megapsocidium	33
jonesi Mesotanyderus	97	MEGASECOPTERA	13
jonesi Samaroblatta	19	melinum Phyllelytron	23
jonesi Triassoblatta	19	membranaceus Heterojassus	66
jucunda Mesochorista	91	Mesacridites	18
Jurinidae	85	mesembria Neopermopanorpa	96
kimblensis Stenoglyphis	58	MESOBLATTINIDAE	19
Knezouria	64	mesocampta Apheloscyta	60
knighti Permocephalus	10	Mesochorista	91
Labidelytron	25	Mesocicadella	63
lata Ademosyne	70	Mesocixioides	41
latipennis Permothea	51	Mesocixius	41
latus Austromylacrites	20	Mesodiphthera	56
Leiodes	74	Mesogereon	46
leongatha Prochoristella	92	MESOGEREONIDAE	46
LEPTOCERIDAE	87	Mesogryllacris	26
Leptoconops	105	Mesojassula	49
LEPTOPHLEBIIDAE	11	Mesojassus	44
leptoptera Triassolocusta	27	MESOMANTIDIIDAE	14
Ligavena	40	Mesomantidion	14
LIGAVENIDAE	40	Mesonirvana	56
ligula Xenelytron	25	Mesonotoperla	17
ligulatus Platycrossos	77	Mesopanorpodes	91
Limoniid indet.	104	MESOPANORPODIDAE	91, 98
lineata Lithosmylidia	82	Mesophlebia	14
Lithosmylidia	82	MESOPHLEBIIDAE	14
Lithosmylidia sp. A	82	Mesopsyche	92
Lobites	71	MESOPSYCHIDAE	92
locustoides Mesorthopteron	18	Mesorhynchophora	73
LOCUSTOPSEIDAE	27	MESORTHOPTERIDAE	18
Lophiocypha	31	Mesorthopteron	18
Lophioneura	3, 31	Mesoscytina	38
LOPHIONEURIDAE	31	Mesostigmodera	72
lowei Cicada?	6	Mesotanyderus	97
maculata Austroprosbole	48	Mesothoris	80
maculata Ipsvicia	45	Mesothymbris	39
magna Ipsviciopsis	45	Mesotitan	4, 9, 10
magna Psyllidella	53	MESOTITANIDAE	29
magnifica Ademosynoides	70	MESOVELIIDAE	68
magnifica Archipanorpa	89	metapteryx Anomaloscytina	48
magnifica Paraknightia	47	Methana sp.	20
magnopunctata Willcoxia	72	Metrorhynchites	79
magnus Permobrachus	57	Microscytinella	67

<i>migdisovae</i> Eoscytina	36	<i>oliffi</i> Ademosyne	70
<i>mimicans</i> Pseudalysiinia	102	<i>optata</i> Neochoristella	93
<i>minor</i> Ademosynoides	70	<i>orthoclada</i> Mesocixioides	41
<i>minor</i> Permophilus	25	ORTHOPHLEBIIDAE	94
<i>minor</i> Reeveana	74	ORTHOPTERA	26
<i>minuta</i> Protopsyllops	52	<i>Orthoscytina</i>	57
<i>minuta</i> Tricrosbia	64	OSMYLIDAE	82
<i>minutum</i> Clavopsyllidium	50	OSYMLOPSYCHOPIDAE	83
<i>minutus</i> Polysitum	77	<i>Osmylopsychops</i>	83
<i>mirabilis</i> Aeroplane	30	<i>ovalis</i> Elliptoscarta	55
<i>mirabilis</i> Pincombea	47	<i>ovalis</i> Triassocoris	69
<i>mitchelli</i> Belmontia	89	PALAEONTINIDAE	46
<i>mitchelli</i> Notoblattites	21	<i>Palaeoviccia</i>	60
<i>mitchelli</i> Orthoscytina	57	PANORPIDAE	95
<i>mitchelli</i> Permopsyllidium	50	<i>Parabelmontia</i>	90
<i>mitchelli</i> Permoscarta	3,61	<i>Parachorista</i>	91
<i>mitchelli</i> Permosyne	71	<i>paradoxa</i> Platyscytinella	67
<i>mitchelli</i> Protocoleus	23	<i>Paradunstania</i>	42
<i>Mitchelloneura</i>	3,49	<i>Paraknightia</i>	47
<i>Mordellid</i> indet.	79	PARAKNIGHTIIDAE	47
<i>multiseriata</i> Perissophlebia	16	<i>paranotalis</i> Triassoscytinopsis	59
<i>multivenata</i> Beaconiella	64	<i>parva</i> Ademosyne	70
MYCETOPHILIDAE	102	<i>parva</i> Cretacochochista	90
<i>Mycetophilidae</i> gen. et sp. nov.	102	<i>parva</i> Dysmorphoptiloides	43
<i>myersi</i> Triassocoris	69	<i>parvula</i> Lithosmylidia	82
MYLACRIDAE	20	<i>patricia</i> Permotipulus	97
<i>nana</i> Protasmanina	102	<i>pectinata</i> Tettigoides	28
<i>nana</i> Triassochorista	98	<i>pectinatus</i> Archeosmylus	81
<i>nana</i> Westratia	109	<i>Pemphredonine</i> indet.	108
<i>Nannochorista</i> sp.	93	<i>Peraphlebia</i>	15
<i>Nannochoristella</i>	93	<i>perfecta</i> Belmontocarta	43
NANNOCHORISTIDAE	93	<i>perfecta</i> Trifidella	37
<i>nanus</i> Permomerope	96	<i>Perissophlebia</i>	16
<i>nanus</i> Choristotanyderus	97	<i>perkinsi</i> Mesothymbris	39
<i>Neoageta</i>	88	<i>perkinsi</i> Neoparachorista	95
<i>Neochoristella</i>	93	PERLARIA	17
<i>Neohagla</i>	26	PERMAESCHNIDAE	15
<i>Neoparachorista</i>	95	<i>Permagra</i>	60
<i>Neoparachorista nana</i>	98	<i>permiana</i> Lophiocypha	31
<i>Neopermopanorpa</i>	96	<i>permiana</i> Mitchelloneura	3,49
<i>Neopetromantis</i>	88	<i>permiana</i> Parabelmontia	90
NEORTHOPHLEBIIDAE	94	<i>permianum</i> Stenoperlidium	17
<i>neoxenus</i> Blattotermes	22	<i>permianus</i> Zygopsocus	34
<i>neoxenus</i> Permithone	81	<i>Permithone</i>	81
<i>Neuropsyche</i>	92	PERMITHONIDAE	81
NEUROPTERA	81	<i>Permoberothella</i>	85
<i>neuropunctatum</i> Mesogereon	46	<i>Permobrachus</i>	57
<i>Niwratia</i>	99	<i>Permocapitus</i>	10
<i>Notoblattites</i>	21	<i>Permocentrus</i>	61
<i>obliqua</i> Apheloodes	73	<i>Permocephalus</i>	10
<i>obliqua</i> Orthoscytina	57	PERMOCHORISTIDAE	91
<i>obscura</i> Permoviccia	61	<i>Permodiphthera</i>	49
<i>obscura</i> Protopincombea	48	<i>Permofulgor</i>	25
<i>obtusa</i> Ademosynoides	70	PERMOFULGORIDAE	25
<i>Oecetis</i> sp.	87	<i>Permoglyphis</i>	57
<i>oliarcoides</i> Permithone	81	<i>Permojassus</i>	58

Permomerope	96	Prochoristella	92
PERMOMEROPIDAE	96	<i>Prochoristella whitehousei</i>	98
PERMOPANORPIDAE	96	Proctotrupid indet.	107
PERMOPHILIDAE	24	PROGONOCIMICIDAE	65
Permophilus	25	progressivum Psychelytron	24
Permopsychops	81	Prohagla	27
Permopsyllidiops	51	Promirara	11
Permopsyllidium	50	PROPARGYLLACRIDIDAE	26
Permopsylloides	51	Propargyllacris	26
Permorapisma	82	Prorhyacophila	8, 86
Permoscarta	3, 61	PROSBOLIDAE	48
<i>Permosmylus</i>	81	prosboloides Ligavena	40
Permosyne	71	Protasmanina	102
PERMOTANYDERIDAE	97	PROTELYTOPTERA	23
Permotanyderus	97	PROTOCOLEIDAE	23
Permothea	51	Protocoleus	23
Permotheella	52	Protolimnophila	103
PERMOTIPULIDAE	97	PROTOMYRMELEONTIDAE	15
Permotipulus	97	Protopsychopsis	83
Permovicia	61	Protopsyllops	52
perplexa Austroidelia	18	Protopincombea	48
perplexa Permoberothella	85	PROTOPSYLLIDIIDAE	50
petalon Phyllelytron	23	Protopsyllidium	52
petrica Etheridgea	72	PROTORTHOPTERA	18
Petropsychops	83	Pselaphid indet.	79
PHASMATODEA	30	Pseudalysiinia	102
phipa Mesochorista	91	PSOCIDIIDAE	33
Phipoides	89	Psocopsyllidium	53
Phyllelytron	23	PSOCOPTERA	33
pigmentatum Dermelytron	24	Psocoscytina	53
Pincombea	47	Psychelytron	24
pincombeae Parachorista	91	PSYCHOPSIDAE	84
pincombeae Permosyne	71	PSYCHROPTILIDAE	13
pincombeae Permithone	81	Psychroptilus	1, 13
pincombei Austropsocidium	33	Psyllid indet.	62
pincombei Orthoscytina	57	Psyllidella	53
pincombei Permophilus	25	Psyllidiana	54
PINCOMBEIIDAE	47	pulchra Dunstania	42
plana Homaloscytina	56	pulchra Stanleyana	62
plana Leiodes	74	Pulcid indet.	99
Platycrossos	77	punctata Ademosyne	70
Platyscytinella	67	punctata Mesocicadella	63
plecioides Triassopsylla	54	punctata Tryoniopsis	74
plexus Australurus	11	punctatus Polysitum	77
Polycytella	85	punctomarginum Elaterium	76
Polysitum	77	pusilla Prochoristella	92
Polytaxineura	15	pygmaea Leiodes	74
postica Eupincombea	47	pygmaeus Triassomantis	28
PRAEAULACIDAE	109	quadripartita Mesothoris	80
prima Tillyardiptera	105	quadrivittata Shepperdia	77
primitiva Bekkerscytina	35	queenslandicum Mesomantidion	14
proavita Mesochorista	8, 91	quinquemedi	57
proavitus Triassojuassus	40	radians Microscytinella	67
Proberotha	84	ramacostata Ademosyne	70
Proberothella	84	reducta Hylicella	45
PROBEROTHIDAE	84	reducta Nannochoristella	93

reductus <i>Xenogryllacris</i>	27	STEREOCHORISTIDAE	87
Reeveana	74	stigmata <i>Triassophlebia</i>	14
reticulata <i>Belpsylla</i>	50	stigmaticum <i>Austropsocidium</i>	33
reticulata <i>Samaroblatta</i>	19	stigmatus <i>Archeosmylus</i>	81
RHAGIONIDAE	103	strenulata <i>Triassoscytinopsis</i>	59
RINCANIIDAE	63	striatella <i>Ademosynoides</i>	70
Robinjohnia	94	stricta <i>Triassocotis</i>	39
robusta <i>Permodiphthera</i>	49	subcostalis <i>Notoblattites</i>	21
robustus <i>Mesopanorpodes</i>	91	subcostalis <i>Orthoscytina</i>	57
rotundata <i>Triscytina</i>	67	subcostalis <i>Triassoscarta</i>	37
rugulosa <i>Ademosyne</i>	70	subpyriformis <i>Simmondsia</i>	78
rugosa <i>Aphelodes</i>	73	subtumidus <i>Platycrossos</i>	77
Samaroblatta	19	subulatum <i>Elateridium</i>	76
Samarura sp.	16	superba <i>Mesopsyche</i>	92
<i>Scarites</i>	39	superba <i>Petropsychops</i>	83
scolopoides <i>Chiliocycla</i>	38	superba <i>Proberotha</i>	84
Scolypopites	63	superba <i>Prohagla</i>	27
scullyi <i>Clatrotitan</i>	29	superba <i>Protolimnophila</i>	103
sculptor <i>Dulcimanna</i>	12	superba <i>Triassopsychops</i>	84
scutulum <i>Triassocoris</i>	69	superbum <i>Mesogereon</i>	46
SCYTINOPTERIDAE	55	sydneiensis <i>Elcanopsis</i>	26
scytinopteroides <i>Permotheella</i>	52	sydneiensis <i>Metrorhynchites</i>	79
semiovena <i>Neoparachorista</i>	95	SYMPHYTA	106
shepperdi <i>Mesogereon</i>	46	Tanypodine indet.	100
Shepperdia	77	Tanypodine pupa indet.	101
SIALIDAE	110	Tarwinia	99
Simmondsia	78	tasmanica <i>Triassoblatta</i>	19
Simuliid indet.	103	TENEBRIONIDAE	79
Simuliid indet. 2	105	tenuiquadrata <i>Mesothoris</i>	80
sinuata <i>Mesonotoperla</i>	17	termioneura <i>Mesocixioides</i>	41
sinuata <i>Neohagla</i>	26	Terpandrus <i>horridus</i>	8
sinuatum <i>Protopsyllidium</i>	52	tetraneura <i>Orthoscytina</i>	57
SIPHONURIDAE	11	tetrastichia <i>Peraphlebia</i>	15
Siphonuridae gen. et sp. nov.	12	Tettigoides	28
SIPHONAPTERA	99	TETTIGONIDAE	28
sobrina <i>Chorista</i>	90	thysanella <i>Lophiocypha</i>	31
sobrina <i>Xenochorista</i>	98	THYSANOPTERA	31
SPHECIDAE	108	tillyardi <i>Austroelytron</i>	23
spilleri <i>Osmyllopsychops</i>	83	tillyardi <i>Clatrotitan</i>	29
splendida <i>Neoparachorista</i>	95	tillyardi <i>Heteroscytina</i>	66
splendida <i>Parachorista</i>	91	tillyardi <i>Robinjohnia</i>	94
splendida <i>Xenochorista</i>	98	Tillyardiopsis	73
Stanleyana	62	Tillyardipectera	105
stanleyi <i>Lophiocypha</i>	31	TILLYARDIPTERIDAE	105
stanleyi <i>Permopsyllidiops</i>	51	timmsii <i>Heterochterus</i>	65
stanleyi <i>Polytaxineura</i>	15	TIPULIDAE	103
stanleyi <i>Zoropsocus</i>	32	Tipulid indet.	104
Staphylinid indet.	79	Tipulid larva indet.	104
Steinhardtia	5, 29	Tipulid pupa indet.	104
STENELTRIDAE	25	TITANOPTERA	29
Stenoglyphis	58	transecta <i>Hexascytina</i>	66
Stenoperlidium	17	transversum <i>Elateridium</i>	76
Stenopsocidium	34	triareolata <i>Mesopsyche</i>	92
Stenoscytina	58	triassica <i>Archeopsychops</i>	84
Stenovicia	62	triassica <i>Dunstaniopsis</i>	42
Stereochorista	87	triassica <i>Fletcheriana</i>	46

<i>triassica</i> Mesochorista	91	tuberculata Lobites	71
<i>triassica</i> Polycytella	85	tuberculata Tillyardiopsis	73
<i>triassica</i> Samaroblatta	19	tumidus Platycrossos	77
<i>triassicum</i> Stenoperlidium	17	Tychticoloides	37
<i>triassicus</i> Mesocixius	41	typica Mesostigmodera	72
Triassagrion	15	typica Triassoblatta	19
Triassoaphis	35	Ulomites	80
Triassoblatta	19	una Undacypha	32
Triassochochista	98	Undacypha	32
TRIASOCHORISTIDAE	98	unicus Knezouria	64
Triassocixius	41	ustulata Lophioneura	3, 31
Triassocoris	69	variotuberculata Tillyardiopsis	73
Triassocotis	39	Veliid indet.	68
Triassodoecus	67	venosa Mesocicadella	63
Triassojoassus	40	venosa Permithone	81
Triassoolestes	16	venosa Protopsychopsis	83
TRIASOLESTIDAE	16	vittamargina Ademosyne	70
Triassolocusta	27	wadei Tripsyllidium	55
TRIASOMANTEIDAE	28	warnensis Parachorista	91
Triassomantis	28	westraliensis Mesothoris	5, 80
Triassophlebia	14	Westratia	109
<i>Triassopsyche</i>	92	whitehousei Austrochoristella	98
Triassopsychops	84	wianamattense Elateridium	76
Triassopsylla	54	wianamattensis Ademosyne	70
Triassoscarta	37	wianamattensis Mesopanorpodes	91
Triassoscelis	59	wianamattensis Notoblattites	21
Triassoscytina	59	willcoxi Ulomites	80
Triassoscytinopsis	59	Willcoxia	72
Triassothea	54	woodwardi Mesothymbris	39
TRICHOPTERA	86	Xenelytron	25
Tricrosbia	64	Xenochorista	98
TRIDACTYLIDAE	28	Xenochoristella	98
Tridactylid indet.	28	XENOCHORISTIDAE	98
Trifidella	37	Xenogryllacris	27
Tripsyllidium	55	Xenopanorpa	96
Triscytina	67	Xenopterum	28
triserialis Permorapisma	82	XYELIDAE	106
trivenulatus Permocentrus	61	Zoropsocus	32
trivittata Lobites	71	ZYGOPSOCIDAE	34
Tryoniopsis	74	Zygopsocus	34

NOTE ADDED IN PROOF

Since this volume went to press I have become aware that Rasnitsyn & Aristov (2004) have erected *Jarmilacladus variabilis* gen. et sp. nov. (Hypoperlida: Anthracopeltidae) [holotype BMNHIn45933] and *Belmophenopterum pectinatum* gen. et sp. nov. (Grylloblattida: Sylvaphlebiidae) [holotype BMNHIn45837] from the Upper Permian Newcastle Coal Measures near Belmont, south of Newcastle.

Sue Parfrey has kindly brought to my attention a fossil insect in a drill core through the Upper Permian Baralaba Coal Measures in the southeastern Bowen Basin, central Queensland. It is a stonefly nymph but a more detailed description will be provided in another paper. The significance of this specimen is as the first

insect from the considerable thickness and extent of the Permian sediments in the Bowen Basin. Further Permian insects may be expected from the Basin.

Mesochorista australica, the type of *Permochorista* (page 91), was considered by Riek (1953) to be the senior synonym of 8 other species referred to *Permochorista* by Tillyard, namely:- *mittelli* in 1918b, *sinuata* and *affinis* in 1922a and *collinsi*, *pincombei*, *angustipennis*, *osbornei* and *inaequalis* in 1926b.

FURTHER LITERATURE CITED

- RASNITSYN, A.P. & ARISTOV, D.S. 2004. Two new insects from the Upper Permian (Tatarian) of Belmont, New South Wales, Australia (Insecta: Hypoperlida: Anthracopeltidae=Permarraphidae; Grylloblattida: Sylvaphlebiidae). *Paleontological Journal* 38(supplement 2): 158-163.



Jell, P. A. 2004. "The fossil insects of Australia." *Memoirs of the Queensland Museum* 50, 1–124.

View This Item Online: <https://www.biodiversitylibrary.org/item/233107>

Permalink: <https://www.biodiversitylibrary.org/partpdf/303982>

Holding Institution

Queensland Museum

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.