

PAPILIO DEIPHOBUS. *Var.*

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatæ.

Alæ (sedentis) erectæ sursumque conniventes, (Volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

PAPILIO alis caudatis; subtus basi rubro maculatis; posticis maculis septem rubris subannularibus.


Lin. Syst. Nat. p. 746.

Var. Alis inferioribus macula media palmata alba, marginibus interioribus ochraceis.

Pap. Achates.

Cram. t. 243.

In India præcipue conspicitur Papilio Deiphobus, cujus veram magnitudinem cernere est in tabula.



704



R. W. Godwin Del. et Sculp.

DEIPHOBUS. *Var.*

GENERIC CHARACTER.

Antennæ thickening towards the end, and commonly terminating in a clavated tip.

Wings (when at rest) meeting upwards; (Flight diurnal.)

SPECIFIC CHARACTER, &c.

Black Butterfly with tailed wings; beneath spotted at the base with red; the lower pair with seven subannular red spots.

Var. With the lower wings marked by a pal-mated, white middle spot, and with the inner margins ochre-coloured.

The *Papilio Deiphobus* is principally found in India, and is represented on the plate in its natural size.



Shaw, George. 1806. "Deiphobus, Papilio deiphobus [Pl. 704]." *The Naturalist's Miscellany* 17(CCII), <https://doi.org/10.5962/p.311141>.

View This Item Online: <https://www.biodiversitylibrary.org/item/281954>

DOI: <https://doi.org/10.5962/p.311141>

Permalink: <https://www.biodiversitylibrary.org/partpdf/311141>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.