

INDEX

Synonyms are printed in *italics*; new names in **bold-face** type

- Abelia Bodinieri*, 334
 — *Cavaleriei*, 335
 — *deutziaefolia*, 334
 — *Mairei*, 334
 — *myrtilloides*, 334
 — *parvifolia*, 334
 — *Schumannii*, 334
 — *verticillata*, 334
Adina racemosa, 319
Aeschynanthus Dunnii, 315
Agrostis canina, 88
Akebia pentaphylla, 358
 — *quinata* × *trifoliata*, 358
 — *trifoliata pentaphylla*, 358
Allium canadense, 90
Allocarya alternifolia, 193
 — *calycosa*, 193
 — *corallicarpa*, 193
 — *divaricata*, 192
 — *figurata*, 193
 — *hirta*, 193
 — *media*, 192
 — *Scouleri hirta*, 193
Alomia tenuifolia, 445
Amelanchier § **Malacomeles**, 449
 — § *Nagelia*, 449
 — **paniculata**, 449
 — *stolonifera*, 92
Amorpha fruticosa, 93
Ampelopsis brevipedunculata Maximowiczii, 470
Amsinckia angustifolia, 203
 — *Douglasiana*, 197, 202, 203
 — *intermedia*, 197, 203
 — *Lemmonii*, 202
 — *lycopsoides*, 199
 — *nigricans*, 198
 — *parviflora*, 203
 — **Scouleri**, 202
 — *spectabilis*, 198
Amylothea sp., 206
 An endemic *Sophora* from Rumania, 76 pl. 123, 124, fig.
Anchusa Sellowiana, 194
 ANDERSON, EDGAR, An endemic *Sophora* from Rumania, 76, pl. 123, 124, fig.
 — and REHDER, ALFRED, New hybrids from the Arnold Arboretum, 358
 — and SAX, KARL, Chromosome numbers in the Hamamelidaceae and their phylogenetic significance, 210, figs.
Anemone quinquefolia, 92
Antennaria canadensis, 96
 — *neglecta*, 96
 — *plantaginifolia*, 96
Antirhea bifida, 166
 — *frangulacea*, 166
Apios americana, 93
 — *tuberosa*, 93
Aquilegia vulgaris, 92
Aralia hispida, 94
Arctostaphylos novoleontis, 450
Argusia, 161
Arguzia, 161
 — *cimmerica*, 164
 — *Messerschmidia*, 164
 — *repens*, 164
 — *rosmarinifolia*, 164
Aristida dichotoma, 88
 — *gracilis*, 88
Aristolochia Clematitis, 91
 — *Kaempferi*, 91
 Arnold Arboretum, Bibliography of the published writings of the staff and students of the, 1934–1935, 455
 Arnold Arboretum during the fiscal year ended June 30, 1935, The, 455
 Arnold Arboretum Expedition, 1930–1932, Lorantheae collected in the Solomon Islands by L. J. Brass and S. F. Kajewski, on the, 206, pl. 129
 Arnold Arboretum, New hybrids from the, 358
 Arnold Arboretum, 1934–35, Staff of the, 469
Aronia arbutifolia, 92

- Aster acuminatus*, 96
 — *linariifolius leucactis*, 96
 — *multiflorus*, 96
Atriplex Chapinii, 443
Avicennia marina, 70
 — — *resinifera*, 70
 BAILEY, I. W. and KERR, THOMAS, The visible structure of the secondary wall and its significance in physical and chemical investigations of tracheary cells and fibers, 273, pl. 140–149
 BAKHUIZEN VAN DEN BRINK, R. C., Notes on some of the Ebenaceae and Verbenaceae of the Solomon Islands collected on the Arnold Arboretum Expedition, 1930–1932, 68, pl. 120–122
Benthamia lycopsoides, 200
Benzoin aestivale, 92
Berberis Thunbergii, 92
Boltonia asteroides, 96
 Boraginaceae, X, Studies in; The Boraginaceae of northeastern South America, 1
 Boraginaceae, XI, Studies in the, 145
Borellia asper, 11
Brandisia racemosa, 315
 BRASS and S. F. KAJEWSKI, Lorantheae collected in the Solomon Islands by L. J. . . . on the Arnold Arboretum Expedition, 1930–1932, 206, pl. 129
Buchenavia capitata, 45
Buddleia plectranthoidea, 311
Callicarpa pedunculata, 70
 — *pentandra furfuracea*, 70
 — — *genuina*, 70
Callitriche heterophylla, 93
 — *palustris*, 93
Canthium Cavaleriei, 324
 — *Dunnianum*, 323
 — *Henryi*, 328
 — *Labordei*, 323
Carex angustior, 89
 — *brevior*, 90
 — *canescens disjuncta*, 89
 — *caryophyllea*, 89
 — *communis*, 89
 — *Crawfordii*, 89
 — *crinita gynandra*, 90
 — *Goodenowii*, 89
Carex laxiflora gracillima, 89
 — — *leptonervia*, 89
 — *longirostris*, 90
 — *lupulina*, 90
 — *panicea*, 89
 — *pennsylvanica lucorum*, 89
 — *stellulata angustata*, 89
 — *Straminea*, 89
 — *tenera*, 89
 — *varia*, 89
Carya mexicana polyneura, 448
 — *ovalis*, 91
Caryopteris Esquirolii, 313
 — *fluminis*, 312
Cavaleriella Dunniana, 333
Celastrus orbiculatus, 94
Cephalanthus Cavaleriei, 319
 — *Navillei*, 319
Cephalosporium wilt of elms, New facts concerning, 453
Cerdana alliadora, 8
 Chromosome numbers in the Hamamelidaceae and their phylogenetic significance, 210, figs.
 Chromosome structure and behavior in mitosis and meiosis, 423, pl. 161–164
 Chromosome structure in the meiotic chromosomes of *Rhoeo discolor* Hance, 216, pl. 130, 131
Chrysanthemum segetum, 96
Clematis paniculata, 92
Clerodendron amicum, 71
 — *Buchanani*, 70
 — *confusum*, 70
 — *inerme*, 71
Coldenia conspicua, 183
 — *decumbens*, 185
 — *Nuttallii*, 185
Colquhounia elegans pauciflora, 312
 — *Seguini*, 311
Colutea media, 93
Conyza scorpioides, 185
Cordia, 3
 — *aberrans*, 173
 — *alliadora*, 8
 — *amplifolia*, 174
 — *Anderssoni*, 175
 — *Aubletii*, 41
 — *axillaris gymnocarpa*, 179

- Cordia bahiensis*, 43
 — *bicolor*, 23
 — *bifurcata*, 34
 — **Braceliniae**, 177
 — *calophylla*, 18
 — *campestris*, 178
 — *carnosa*, 23
 — *cordifolia*, 11
 — *coriacea*, 23, 27
 — *corymbosa*, 34
 — *curassavica*, 45
 — *dichotoma*, 23
 — *echitioides*, 44
 — *exaltata*, 29
 — — **melanoneura**, 30
 — **fallax**, 10
 — *flavescens*, 44
 — *formicarum*, 13
 — **fulva**, 20
 — *galapagensis*, 176
 — *grandiflora*, 32
 — *graveolens*, 36
 — *guazumaefolia*, 179
 — *guianensis*, 10, 36
 — *gujanensis*, 36
 — *Haenkeana*, 179
 — *hebecarpa*, 21
 — *heterophylla*, 21
 — *hirsuta*, 13
 — **hirta**, 29
 — *hispidissima*, 13
 — *Hookerianum*, 174
 — *ierensis*, 25
 — *insignis*, 179
 — *interrupta*, 36
 — **laevifrons**, 16
 — **laevior**, 180
 — *leucophlyctis*, 176
 — *linearis*, 174
 — *Lockartii*, 23
 — *lucida*, 39
 — *lutea*, 44
 — *macrostachya*, 36
 — *martinicensis*, 45
 — *melanoneura*, 30
 — *Meziana*, 174
 — *miranda*, 13
 — *monosperma*, 34
 — *mucronata*, 30, 173
 — *Cordia multispicata*, 43
 — *muneco*, 11
 — *Myxa*, 45
 — **naidophila**, 32
 — *Neowediana*, 177
 — *Neowidiana*, 177
 — *nervosa*, 18
 — *nodosa*, 13
 — — *angustifolia*, 14
 — — *hispidissima*, 13
 — *oxyphylla*, 36
 — *panicularis*, 26
 — **polycephala**, 33
 — *pubescens*, 21
 — *revoluta*, 174
 — **ripicola**, 180
 — *rufa*, 32
 — **Sagotii**, 27
 — *sarmentosa*, 44
 — *scabrida*, 30
 — *scabrifolia*, 22
 — *scandens*, 44
 — *Schomburgkii*, 39
 — *Scouleri*, 175
 — *sebestena*, 9
 — *sericalyx*, 25
 — — *latifolia*, 25
 — **setigera**, 176
 — *speciosa*, 9
 — *Sprucei*, 17
 — *taguahyensis*, 174
 — *tetrandra*, 11
 — *tetraphylla*, 44
 — *tobagensis*, 39
 — — *Broadwayi*, 39
 — *tomentosa*, 41
 — *toqueve*, 21
 — *trichostyla*, 23
 — **ucayaliensis**, 181
 — *Ulei ucayaliensis*, 181
 — *ulmifolia*, 34
 — *umbraculifera*, 11
 — *umbrosa*, 14
 — *volubilis*, 14
 — **Weddellii**, 173
Cornus Esquirolii, 319
 — *stolonifera*, 94
Corydalis bulbosa, 92
Cotoneaster § *Malacomeles*, 449

- Cotoneaster § *Naegelia*, 449
 Crataegus from Missouri, Two new species of, 353, figs.
 Crataegus *Danielsii*, 355, fig. 2
 — — *glabra*, 357
 — — *hannibalensis*, 353, fig. 1
 — — *rotundifolia*, 92
 CREAGER, D. B., New facts concerning Cephalosporium wilt of elms, 453
 Crepis capillaris, 97
 Cristaria insularis, 445
 CROWELL, IVAN H., The hosts, life history and control of Gymnosporangium clavipes C. and P., 367, pl. 155-160
 Cryptantha affinis, 171
 — albida, 172
 — alyssoides, 169
 — ambigua, 171, 172
 — — robustior, 172
 — aprica, 169, 170
 — barbigera, 171
 — Buchtienii, 169
 — campylotricha, 170
 — candelabrum, 170
 — Candolleana, 170
 — capituliflora compacta, 170
 — circumscissa, 195
 — clandestina, 195
 — Clevelandii, 171
 — — florosa, 172
 — confusa, 171
 — decipiens corollata, 172
 — diffusa, 170, 171
 — diplotricha, 170, 171, 172
 — echinella, 172
 — fallax, 170
 — Famatinae, 171
 — Fendleri, 171, 172
 — flaccida, 171, 172
 — globulifera, 170, 171
 — glomerata, 170, 172
 — *glomerata*, 195
 — glomerulifera, 169
 — grandiflora anulata, 172
 — granulosa, 172
 — Hansenii, 173
 — — pulchella, 173
 — Hendersonii, 172
 — hispida, 172
 Cryptantha hispidissima, 172
 — hispidula, 172
 — — Elmeri, 172
 — Hossei, 171
 — intermedia, 173
 — Kingii, 170
 — *Lappula*, 195
 — leiocarpa eremocaryoides, 171
 — limensis, 172
 — linearis, 170
 — Lyallii, 171
 — macrocalyx, 169
 — *microcarpa*, 196
 — microstachys, 171
 — mirabunda, 172
 — modesta, 170
 — muricata Jonesii, 173
 — nevadensis, 171
 — parvula, 171
 — phacelioides, 169
 — Philippiana, 172
 — Rattanii, 172
 — Romanii, 172
 — simulans, 172, 173
 — Stuebelii, 173
 — talquina, 169
 — Torreyana, 171, 172
 — utahensis, 172
 — Vidalii, 170
 — Watsoni, 171
 Cynanchum yunnanense, 325
 Cyperus esculentus, 89
 Cyphomattia lanata, 194
 Cypripedium acaule, 90
 Cystacanthus *yangtsekiangensis*, 315
 Dactylophora *salomonica*, 206, pl. 129 a-b
 DANSER, B. H., Loranthaceae collected in the Solomon Islands by L. J. Brass and S. F. Kajewski, on the Arnold Arboretum Expedition, 1930-1932, 206, pl. 129
 DARKER, GRANT D., Hypodermella Hiratsukae, a new species of Hypodermataceae from Japan, 364, pl. 154
 Datura Metel, 95
 — Stramonium, 95
 Dendrophthoe falcata, 209
 Deutzia *funnebris*, 315

- Dichapetalum scandens*, 44
 — *vestitum scandens*, 44
Diervilla Lonicera, 95
Diospyros ellipticifolia, 68
 — *ferrea*, 68
 — — *salomonensis*, 68
 — *insularis*, 69, pl. 120, 121
 — *maritima*, 69
 — *samoensis*, 69, pl. 122
Dipelta yunnanensis, 333
Diplotaxis muralis, 92
 Dispersal of viable basidiospores of the
 Gymnosporangium rusts, The, 411,
 figs.
Echinochloa crusgalli longiseta, 88
 — *muricata*, 88
Echium vulgare, 95
 Effect of temperature on nuclear differ-
 entiation in microspore development,
 The, 301, pl. 150, fig.
Ehretia Esquirolii, 320
 Elaioplasts in Iris: a morphological
 study, 225, pl. 132-137
 Elms, New facts concerning Cephalo-
 sporium wilt of, 453
Elsholtzia Cavaleriei, 311
 — *Dielsii*, 313
 — *fruticosa*, 312
 — *Labordei*, 312
 — *lampradena*, 313
 — *Lychnitis*, 313
 — *ochroleuca*, 313
 — *polystachya*, 312
 — *rugulosa*, 312
 — *Souliei*, 313
 — *tristis*, 313
Emmenopterys Henryi, 318
Epifagus virginiana, 95
Epilobium hirsutum, 94
Eragrostis cilianensis, 88
 — *megastachya*, 88
 — *pectinacea*, 89
 — *peruviana*, 443
Eritrichium Rengifoanum, 169
Erythronium americanum, 90
Eurya Degeneri, 351
 — — *grandifolia*, 352
 — — *stenophylla*, 352
 — *sandwicensis*, 348
Eurya sandwicensis β , 352
 — — *grandifolia*, 352
 — — *sessilifolia*, 348
 — — *prostrata*, 350
Evonymus obovatus, 93
Faradaya amicorum, 71
 — — *salomonensis*, 71
 FAULL, ANNA F., Elaioplasts in Iris: a
 morphological study, 225, pl. 132-137
Festuca ovina, 89
 — *rubra commutata*, 89
Firensia, 44
 — *hirsuta*, 13
 — *lutea*, 45
 Flora of San Felix Island, The, 440,
 pl. 165
Galium asprellum, 95
Gardenia florida, 321
 — *jasminoides*, 321
 — *Schlechteri*, 321
Gillenia trifoliata, 92
Glyceria septentrionalis, 89
Gmelina moluccana, 72
 — *salomonensis*, 72, pl. 122
Gymnosporangium clavipes C. and P.,
 The hosts, life history and control of,
 367, pl. 155-160
Gymnosporangium globosum Farl. and
 their relative susceptibility, The hosts
 of, 98, pl. 125-128, figs.
Gymnosporangium rusts, The dispersal
 of viable basidiospores of the, 411,
 figs.
Hackelia caerulea, 194
 — *diffusa caerulea*, 194
 — *grisea*, 194
 — *Nelsonii*, 194
 — *patens*, 194
 Hamamelidaceae and their phylogenetic
 significance, Chromosome numbers in
 the, 210, figs.
Handeliiodendron, a new genus of Sapin-
 daceae, 65, pl. 119, fig.
Handeliiodendron, 65
 — *Bodinieri*, 66, pl. 119, fig.
Havilandia opaca, 190
 — *papuana*, 191
 — *robusta*, 191
Hedyotis Bodinieri, 316

- Hedyotis Esquirolii*, 316
 — *macrostemon*, 316
 — *Mairei*, 329
 — *yunnanensis*, 332
Helenium nudiflorum, 96
Heliophytum indicum, 59
 — *passerinoides*, 61
Heliopsis helianthoides, 96
Heliotropium, 57
 — *angiospermum*, 186
 — *curassavicum*, 57
 — *filiforme*, 62
 — *fruticosum*, 61
 — *Fumana*, 61
 — *gnaphalodes*, 56, 165
 — *helophilum*, 63
 — *humile*, 186
 — *indicum*, 59
 — *inundatum*, 60
 — *japonicum*, 164
 — *lagoense*, 58
 — *lineare*, 167
 — *messerschmidioides*, 166, 167, 168
 — *micranthos*, 167
 — *pannifolium*, 168
 — *procumbens*, 60
 — *salicoides*, 62
 — *scandens*, 150
 — *ternatum*, 61
 — — *Fumana*, 62
 — *tiaridioides*, 186
 — — *genuina*, 186
 — — *schizocarpum*, 186
 — *transalpinum*, 186
 — — *tiaridioides*, 470
 — *trinitense*, 58
 — *tuberculosum*, 167
 — *viridiflorum*, 153
 — *zeylanicum*, 167, 168
 — *sp.*, 61
Hibiscus Trionum, 94
Hieracium florentinum, 97
 — *vulgatum*, 97
 Hosts of *Gymnosporangium globosum* Farl. and their relative susceptibility, The, 98, pl. 125–128, figs.
 Hosts, life history and control of *Gymnosporangium clavipes* C. and P., The, 367, pl. 155–160
Houstonia caerulea, 95
 HUNTER, LILLIAN M., A preliminary note on life history studies of European species of *Milesia*, 143
Huodendron, a new genus of *Styracaceae*, 341, pl. 151, 152, fig.
Huodendron, 341
 — *biaristatum*, 344
 — — *parviflorum*, 346
 — *tibeticum*, 342
Hypochaeris radicata, 96
Hypodermella Hiratsukae, a new species of *Hypodermataceae* from Japan, 364, pl. 154
Hypodermella Hiratsukae, 364
Incarvillea Delavayi, 315
 Iris: Elaioplasts in, a morphological study, 225, pl. 132–137
Ixora Henryi, 322
 JOHNSTON, IVAN M., Studies in *Boraginaceae*, X; The *Boraginaceae* of northeastern South America, 1
 — Studies in the *Boraginaceae*, XI, 145
 — The flora of San Felix Island, 440, pl. 165
Juniperus communis depressa, 88
 Kajewski, Lorantheae collected in the Solomon Islands by L. J. Brass and S. F. . . . on the Arnold Arboretum Expedition, 1930–1932, 206, pl. 129
 KERR, THOMAS, and BAILEY, I. W., The visible structure of the secondary wall and its significance in physical and chemical investigations of tracheary cells and fibers, 273, pl. 140–149
 KOBUSKI, CLARENCE E., Studies in *Theaceae*, I. *Eurya* subgen. *Ternstroemiopsis*, 347, pl. 153
Lactuca sagittifolia, 97
 — *scariola*, 97
 — *spicata integrifolia*, 97
Lagenaria leucantha, 96
Lantana corymbosa, 33
Lappula coerulescens, 194
 — *decumbens*, 194
 — *echinata*, 194
 — *grisea*, 194
 — *subdecumbens coerulescens*, 194
Lasianthus Biermanni, 323

- Lasianthus Dunniana*, 323
 — *Esquirolii*, 323
 — *Hartii*, 323
 — *Hookeri*, 323
 — *Labordei*, 323
 — *sp.*, 324
Lasiarrhenum pinetorum, 187
Lathyrus pannonicus versicolor, 93
Lechea intermedia, 94
 — *tenuifolia*, 94
Lepidocordia, 45
 — *punctata*, 45
Leptodermis Chaneti, 328
 — *Esquirolii*, 326
 — *Mairei*, 327
 — *motsouensis*, 327
 — *oblonga*, 328
 — *pilosa glabrescens*, 327
 — *Potanini*, 326
 — — *glauca*, 327
 — — *tomentosa*, 327
 — *tongchouanensis*, 327
Lespedeza hirta, 93
Leucosceptrum Bodinieri, 311
 — *plectranthoideum*, 311
 — *sinense*, 311
 Lévillé from eastern Asia, Notes on
 the ligneous plants described by, 311
Liatris scariosa, 96
Ligustrum Thea, 317
 — *vulgare*, 95
Liriodendron Tulipifera, 92
Lithocardium Anderssonii, 175
 — *Aubletii*, 41
 — *bicolor*, 23
 — *calophyllum*, 18
 — *cordifolium*, 11
 — *exaltatum*, 29
 — *flavescens*, 44
 — *galapagosenum*, 176
 — *grandiflorum*, 32
 — *guazumifolia*, 179
 — *hebecarpum*, 21
 — *heterophyllum*, 21
 — *hispidissimum*, 13
 — *Hookerianum*, 174
 — *leucophlyctis*, 176
 — *Lockartii*, 23
 — *Mezianum*, 174
 — *Lithocardium mirandum*, 13
 — *multispicatum*, 43
 — *muneco*, 11
 — *Neowiedianum*, 177
 — *nervosum*, 18
 — *nodosum*, 13
 — *oxyphyllum*, 36
 — *paniculare*, 26
 — *pubescens*, 21
 — *revoluta*, 174
 — *scabridum*, 30
 — *Schomburgkii*, 39
 — *Scouleri*, 175
 — *sericalyx*, 25
 — *Sprucei*, 17
 — *tetrandrum*, 11
 — *tetraphyllum*, 45
 — *tomentosum*, 41
 — *toqueve*, 21
 — *umbraculiferum*, 11
Lithospermum calycinum, 203
 — *clandestinum*, 195
 — *lycopsoides*, 202
 — *minutum*, 191
 — **Muelleri**, 187
 — *viridiflorum*, 153
Litsea Muelleri, 449
Lonicera acrophila, 337
 — *alpigena*, 95
 — *buxifolia*, 336
 — *dioica*, 95
 — *Esquirolii*, 339
 — *Fauriei*, 339
 — *fragilis*, 337
 — *Guilloni*, 338
 — *gynopogon*, 338
 — *Henryi setuligera*, 338
 — *japonica*, 339
 — *Koehneana*, 338
 — *lanceolata*, 337
 — *ligustrina*, 335
 — *macrantha*, 338
 — *Mairei*, 339
 — *Menelii*, 315
 — *missionis*, 336
 — *Pampaninii*, 338
 — *pileata*, 336
 — *Rocheri*, 335
 — *tangutica*, 335

- Lonicera yunnanensis*, 339
 Loranthaceae collected in the Solomon Islands by L. J. Brass and S. F. Kojewski, on the Arnold Arboretum Expedition, 1930-1932, 206, pl. 129
Loranthus falcatus, 209
Luculia gratissima, 317
Luzula nemorosa, 90
Lycapsus tenuifolius, 445
Lycopersicon esculentum, 95
Lycopus rubellus, 95
Maba buxifolia, 68
 — *elliptica*, 68
 MacLACHLAN, J. D., The dispersal of viable basidiospores of the Gymnosporangium rusts, 411, figs.
 — The hosts of Gymnosporangium globosum Farl. and their relative susceptibility, 98, pl. 125-128, figs.
Macromeria barbiger, 189
 — *leontis*, 188
Mallotonia, 161
 — *gnaphalodes*, 165
Malus baccata mandshurica, 92
 — *hupehensis*, 92
 — *theifera*, 92
Malva parviflora, 94
 — *verticillata crispa*, 94
 McKELVEY, SUSAN DELANO, Notes on *Yucca*, 268, pl. 138, 139
Medicago hispida, 93
Menodora Muellerae, 451
Messerschmidia, 161
 — *angustifolia*, 168
 — *argentea*, 164
 — *Argunia*, 164
 — *Arguzia*, 164
 — *cancellata*, 168
 — *floribunda*, 168
 — *fruticosa*, 168
 — *gnaphalodes*, 165
 — *hispida*, 168
 — *incana*, 56, 64
 — *montana*, 153
 — *rosmarinifolia*, 164
 — *sibirica*, 164
 — *syringifolia*, 48
Messerschmidia, 161
 — *Argusia*, 164
 — *Mexico*, Some new trees and shrubs from, 448
Micromeria biflora, 312
Milesia, A preliminary note on life history studies of European species of, 143
Mirabilis Jalapa, 91
Montjolya guianensis, 36
 — *tomentosa*, 41
Moritzia dasyantha Sellowiana, 194
 — *Sellowiana*, 194
Mussaenda Bodinieri, 320
 — *Cavaleriei*, 318
 — *Esquirolii*, 319
 — *Mairei*, 318
 — *pubescens*, 320
 — *Wilsonii*, 319
Myosotis Scouleri, 192
Nägelia, 449
Nagelia, 449
Napaea dioica, 94
Neonauclea Navillei, 319
 New facts concerning *Cephalosporium* wilt of elms, 453
 New hybrids from the Arnold Arboretum, 358
 Notes on some of the Ebenaceae and Verbenaceae of the Solomon Islands collected on the Arnold Arboretum Expedition, 1930-1932, 68, pl. 120-122
 Notes on the ligneous plants described by Lévillé from Eastern Asia, 311
 Notes on *Yucca*, 268, pl. 138, 139
Notothixos leiophyllus, 209
Ocotea commutata, 44
Oldenlandia Bodinieri, 316
 — *Esquirolii*, 316
 — *macrostemon*, 316
 — *Mairei*, 329
 — *yunnanensis*, 332
Omphalodes erecta, 204
Ophiorrhiza cantoniensis, 316
 — *Cavaleriei*, 316
 — *japonica*, 316
 — *Labordei*, 316
 — *Seguini*, 316
 — *violaceo-flammea*, 317
Oxalis europaea villicaulis, 93

- Oxalis stricta*, 93
Paederia Bodinieri, 325
 — *Cavaleriei*, 326
 — *chinensis*, 324
 — *Dunniana*, 324
 — *Esquirolii*, 324
 — *foetida*, 324
 — *Mairei*, 324
 — *scandens*, 324
 — *tomentosa*, 324, 326
 — — *Mairei*, 324
 — — *purpureo-coerulea*, 325
 — *Wallichii*, 325
 PALMER, ERNEST J., Supplement to the spontaneous flora of the Arnold Arboretum, 81
 — Two new species of *Crataegus* from Missouri, 353, figs.
Panicum commutatum Ashei, 88
 — *depauperatum* psilophyllum, 88
 — *dichotomiflorum*, 88
 — *lanuginosum* Lindheimeri, 88
 — — *septentrionale*, 88
 — *latifolium*, 88
 — *Lindheimeri*, 88
 — *linearifolium*, 88
 — *oligosanthes* Scribnerianum, 88
 — *philadelphicum*, 88
 — *Scribnerianum*, 88
Parietaria debilis, 443
 — *felicianae*, 443
Pertya Bodinieri, 340
Petraeovitex multiflora salomonensis, 73
Phlogacanthus pubinervis, 315
Phlox paniculata, 95
Physalis heterophylla nyctaginea, 95
Plagiobothrys calandrinoides, 193
 — *hirtus*, 193
 — — *corallicarpus*, 193
 — — *figuratus*, 193
 — *medius*, 192
 — *Scouleri*, 192
 — — *corallicarpus*, 193
 — — *hirtus*, 193
Plantago major, 95
Plectranthus coetsa, 314
 — *Mairei*, 314
 — *ternifolius*, 313
Pogostemon glaber, 313
Polygala sanguinea, 93
Polygonum Carey, 91
 — *lapathifolium*, 91
 — *punctatum leptostachyum*, 91
 — *Sieboldii*, 91
Polystichum acrostichoides, 87
Populus tremuloides, 91
Potamogeton epihydrus Nuttallii, 88
Potentilla canadensis villosissima, 93
 Preliminary note on life history studies of European species of *Milesia*, A, 143
Premna Esquirolii, 329
 — *integrifolia*, 73
 — *Valbrayii*, 331
Prismatomeris brevipes, 328
 — **Henryi**, 328
Prunus Cerasus, 93
 — **Juddii**, 358
 — *pumila* susquehanae, 93
 — *Sargentii* × *yedoensis*, 358
 — *yedoensis* × *Sargentii*, 358
Psychotria elliptica, 322
 — *Esquirolii*, 322
 — *Henryi*, 322
 — *Prainii*, 322
 — *rubra*, 322
Pyrola americana, 94
Quercus ilicifolia × *velutina*, 91
 — *imbricaria* × *velutina*, 91
 — *Leana*, 91
 — *prinoides*, 91
 — *Rehderi*, 91
 — *velutina* × *ilicifolia*, 91
 — — × *imbricaria*, 91
Ranunculus bulbosus, 92
 REHDER, ALFRED, *Handeli* *dendron*, a new genus of Sapindaceae, 65, pl. 119, fig.
 — *Huodendron*, a new genus of Styracaceae, 341, pl. 151, 152, fig.
 — Notes on the ligneous plants described by Léveillé from Eastern Asia, 311
 — Some new trees and shrubs from Mexico, 448
 — and ANDERSON, EDGAR, New hybrids from the Arnold Arboretum, 358
Rhoeo discolor Hance, Chromosome structure in the meiotic chromosomes of, 216, pl. 130, 131

- Ribes sativum*, 92
 — *vulgare*, 92
Rochelia patens, 194
 — *sp.*, 168
Rorippa sylvestris, 92
Rubus allegheniensis, 93
 — *flagellaris*, 93
 — *Idaeus strigosus*, 92
 — *Jeckylanus*, 93
 — *parvifolius*, 93
 — *Randii*, 93
Rumex crispus × *obtusifolius*, 91
Saccellium brasiliense, 181
 — *lanceolatum*, 182
 — *Oliverii*, 182
Sagina decumbens, 91
Sagittaria latifolia gracilis, 88
Salix alba calva, 90
 — *fragilis*, 90
 — *humilis*, 90
 — *lucida*, 90
 — *pedicellaris*, 90
 — *pyrifolia*, 91
Sambucus Argyi, 328
 — *javanica Argyi*, 328
 San Felix Island, The flora of, 440, pl. 165
Saponaria officinalis, 91
 SAX, HALLY JOLIVETTE and SAX, KARL, Chromosome structure and behavior in mitosis and meiosis, 423, pl. 161–164
 SAX, KARL, Chromosome structure in the meiotic chromosomes of *Rhoeo discolor* Hance, 216, pl. 130, 131
 — The effect of temperature on nuclear differentiation in microspore development, 301, pl. 150, fig.
 — and ANDERSON, EDGAR, Chromosome numbers in the Hamamelidaceae and their phylogenetic significance, 210, figs.
 — and SAX, HALLY JOLIVETTE, Chromosome structure and behavior in mitosis and meiosis, 423, pl. 161–164
Schleidenia filiformis, 62
 — *Fumana*, 61
 — *lagoensis*, 58
 — *subracemosa*, 62
Scilla sibirica, 90
Scirpus rubrotinctus, 89
Sebestena linearis, 174
 — *revoluta*, 174
Senecio aureus, 96
 — *viscosus*, 96
Sida hermaphrodita, 94
Sideroxylon Bodinieri, 66
Silene antirrhina, 91
 — *Armeria*, 91
Sisyrinchium angustifolium, 90
Sogerianthe versicolor, 207, pl. 129
 c-e
Solanum aculeatissimum, 314
 — *Bodinieri*, 314
 — *Cavaleriei*, 314
Solidago odora, 96
 — *ulmifolia*, 96
 Solomon Islands, Loranaceae collected in the . . . by L. J. Brass and S. F. Kajewski, on the Arnold Arboretum Expedition, 1930–1932, 206, pl. 129
 Solomon Islands, Notes on some of the Ebenaceae and Verbenaceae of the . . . collected on the Arnold Arboretum Expedition, 1930–1932, 68, pl. 120–122
 Some new trees and shrubs from Mexico, 448
Sonchus arvensis glabrescens, 97
 — *asper*, 97
 — *oleracea*, 97
Sophora from Rumania, An endemic, 76, pl. 123, 124, fig.
Sophora alopecuroides, 78
 — — *tomentosa*, 78, pl. 124, fig. A
 — *Prodanii*, 76, pl. 123, 124, fig. B
Spergula arvensis, 91
 Spontaneous flora of the Arnold Arboretum, Supplement to the, 81
Strobilanthes deutziaefolia, 334
 — *hypericifolia*, 334
 — *yangtsekiangensis*, 315
Strobilanthes deutziaefolius, 334
 — *hypericifolius*, 334
 Studies in Boraginaceae, X; The Boraginaceae of northeastern South America, 1
 Studies in the Boraginaceae, XI, 145

- Studies in Theaceae I. *Eurya* subgen.
 Ternstroemiopsis, 347, pl. 153
Styrax biaristatus, 344
 — *parviflora*, 346
 — *tibeticus*, 342
Suaeda divaricata microphylla, 444
 — *nesophila*, 444
 Supplement to the spontaneous flora of
 the Arnold Arboretum, 81
Syringa diversifolia, 362
 — *oblata* Giraldii $\times$ *pinnatifolia*, 362
 — *vulgaris*, 95
Tarenna incerta, 321
 — *mollissima*, 320
 — *pallida*, 321
 — *zeylanica*, 321
Tecoma Mairei, 315
Ternstroemiopsis sandwicensis, 348
Tetrandra glabra, 150
 — *Wallichii*, 150
 — *Zollingeri*, 150
Teucrium Esquirolii, 313
Teysmannioidendron Ahernianum, 74
Thamnoseric laceratus, 445
 — *lobata*, 445
Thaumatocaryon dasyanthum Sellowianum, 194
 — *Sellowianum*, 194
Thelypteris spinulosa intermedia, 87
Thymus Cavaleriei, 312
Toquera tomentosa, 21
Tournefortia, 46, 145
 — § *Arguzia*, 161
 — § *Eutournefortia*, 145
 — § *Mallota*, 161
 — § *Mallotonia*, 161
 — § *Messerschmidia*, 161
 — § *Oskampia*, 146
 — § *Pittonia*, 146
 — § *Pittoniae*, 145
 — § *Tetrandra*, 146
 — *acclinis*, 147
 — *acuminata*, 159
 — *alba*, 52
 — *angulosa*, 166
 — *angustifolia*, 166
 — *arborea*, 164
 — *arborescens*, 159
 — *argentea*, 164
Tournefortia Arguzia, 164
 — — *angustior*, 164
 — — *cynanchoides*, 164
 — — *latifolia*, 164
 — — *rosmarinifolia*, 164
 — *Aubletii*, 52
 — *bicolor*, 51
 — — *calycosa*, 52
 — *bifida*, 166
 — *bifida*, 160
 — *Bojeri*, 160, 166
 — *Bojeri*, 159
 — *Boniana*, 154
 — *brasiliensis*, 185
 — *Candollii*, 154
 — *coriacea*, 52
 — *cuspidata*, 54
 — *cymosa*, 56, 166
 — *Edgeworthii*, 167
 — *foetidissima*, 56
 — *frangulaefolia*, 147
 — *fruticosa*, 167
 — *Gaudichaudii*, 154
 — *glabra*, 52, 150
 — *glabrifolia*, 147
 — *gnaphalodes*, 56
 — *gnaphalodes*, 165
 — *Hartwegiana*, 167
 — *Heyneana*, 152, 166
 — *Heyneana*, 154
 — *hirsuta*, 147
 — *hirsutissima*, 56
 — *Hookeri*, 151
 — — *subtropica*, 151
 — *Horsfieldii*, 147
 — *Hostmanni*, 48
 — *incana*, 56
 — *incana*, 64
 — *khasiana*, 154
 — *laevigata*, 51
 — — *latifolia*, 51
 — *linearis*, 167
 — *luzonica*, 156
 — — *sublucens*, 157
 — *macrophylla*, 147
 — *melanochaeta*, 55
 — *Messerschmidia*, 167
 — *Meyeri*, 56
 — *Meyeri*, 64

- Tournefortia micranthos*, 167
 — *Miquelii*, 51
 — *Mocquerysi*, 158
 — *mollis*, 167
 — *mollis*, 157
 — *montana*, 153
 — **Muelleri**, 157
 — *mutabilis*, 167
 — *obscura*, 54
 — *orientalis*, 147
 — *ovata*, 153
 — *paniculata* **spigeliaeflora**, 49
 — *peruviana*, 48
 — *puberula*, 158
 — — **Kirkii**, 158
 — *reticosa*, 152
 — *rosmarinifolia*, 164
 — *Roxburghii*, 153
 — *Royleana*, 167
 — *Sampsoni*, 153
 — *sarmentosa*, 147
 — — *hirsuta*, 147
 — — *magnifolia*, 147
 — *Schomburgkii*, 52
 — *setifera*, 54
 — *sibirica*, 164
 — — *angustior*, 164
 — *spigeliaeflora*, 49
 — *stenoraca*, 167
 — *subulata*, 167
 — *surinamensis*, 48
 — *syringaefolia*, 48
 — *syringaefolia*, 51
 — *tetragona*, 150
 — *tetrandra*, 150
 — — *angustifolia*, 151
 — — *glabra*, 150
 — — *hirsuta*, 147
 — — *longiflora*, 150
 — — **Walkerae**, 151
 — *tuberculosa*, 167
 — *Ulei*, 51
 — *Urvilleana*, 147
 — *velutina*, 159
 — *viridiflora*, 153
 — — *Griffithii*, 154
 — *volubilis*, 47
 — *Walkerae*, 151
 — *Wallichii*, 150

Tournefortia Wightii, 153
 — *zeylanica*, 168
Tradescantia virginiana, 90
Trientalis americana, 95
 Two new species of *Crataegus* from Missouri, 353, figs.
Typha latifolia, 88
Uncaria scandens, 319
Vaccinium corymbosum, 94
Vangueria tomentosa, 167
Varronia canescens, 175
 — *corymbosa*, 34
 — *dichotoma*, 34
 — *grandiflora*, 32
 — — *glabrata*, 33
 — — *Sprucei*, 33
 — *guasumaeifolia*, 179
 — *guianensis*, 36
 — *lantanoides*, 32
 — *leucophlyctis*, 176
 — *linearis*, 174
 — *macrostachia*, 470
 — *macrostachya*, 36
 — *monosperma*, 34
 — *polycephala*, 33
 — *revoluta*, 174
 — *scaberrima*, 176
 — *Scouleri*, 175
 — *spicata*, 43
 — *tomentosa*, 41
 — *ulmifolia*, 34
Vernonia scorpioides, 185
Veronica peregrina, 95
Viburnum ajugifolium, 331
 — *barbigerum*, 332
 — *bitchiuense* × *Carlesii*, 360
 — *Bodinieri*, 332
 — *botryoideum*, 328
 — *Cavaleriei*, 329
 — *Chaffanjonii*, 330
 — *congestum*, 329
 — *corylifolium*, 332
 — *crassifolium*, 330
 — *cylindricum crassifolium*, 330
 — *Dunnianum*, 332
 — *erosum punctatum*, 333
 — — *Taquetii*, 333
 — *erubescens*, 328
 — *foetidum*, 331

- Viburnum foetidum* rectangulatum, 331
— **Juddii**, 360
— *Mairei*, 329
— *Martini*, 329
— oliganthum, 328
— *pinfaense*, 330, 331
— Schneiderianum, 331
— sempervirens, 331
— setigerum, 332
— *Stapfianum*, 329
— sympodiale, 329
— *Taquetii*, 333
— ternatum, 330
— *theiferum*, 332
— *Touchanense*, 331
— trilobum, 95
Viola fimbriatula × papilionacea, 94
— latiuscula, 94
— odorata, 94
— pedata lineariloba, 94
— sagittata, 94
— sororia, 94
Visible structure of the secondary wall
and its significance in physical and
chemical investigations of tracheary
cells and fibers. The, 273, pl. 140–149
Vitex Aherniana, 74
— cofassus, 74
Webera Cavaleriei, 321
— *Henryi*, 321
— *pallida*, 321
Wendlandia Cavaleriei, 318
— *Dunniana*, 318
— *Feddei*, 318
— ligustrina, 317
— longidens, 318
— salicifolia, 317
— uvariifolia *Dunniana*, 318
Yucca, Notes on, 268, pl. 138, 139
Yucca arizonica, 270
— *brevifolia*, 270
— *brevifolia Jaegeriana*, 269, pl. 139
— *puberula*, 270
— **Thornberi**, 268, pl. 138
— *Treleasei*, 270



1935. "Index [vol. 16]." *Journal of the Arnold Arboretum* 16(4), 471–483.
<https://doi.org/10.5962/p.324788>.

View This Item Online: <https://www.biodiversitylibrary.org/item/33592>

DOI: <https://doi.org/10.5962/p.324788>

Permalink: <https://www.biodiversitylibrary.org/partpdf/324788>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Arnold Arboretum of Harvard University

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.