

hyperborea atlantica near Westbourne, Manitoba; this bird was the third in a flock of immature snow geese. It is an adult female, weight 6 lbs. 13 oz.; T. 1.30.04; W. 17.1; spread 22.1; T. 2.5; C. 1.22; Tail 6.2. This species

at which the land drops abruptly to the lower level of the peninsula which is occupied by a black spruce-sphagnum bog extending southward to the Hamilton River and northward to the C. River. The C. River is a small stream



Scott, W. B. 1951. "Lake Erie ciscoes free of *Triaenophorus crassus*." *The Canadian field-naturalist* 65(2), 80–80. <https://doi.org/10.5962/p.341314>.

View This Item Online: <https://www.biodiversitylibrary.org/item/90126>

DOI: <https://doi.org/10.5962/p.341314>

Permalink: <https://www.biodiversitylibrary.org/partpdf/341314>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Ottawa Field-Naturalists' Club

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.