

# QUEENSLAND'S RAINFOREST CANOPIES - A MITEY CORNUCOPIA

David Evans WALTER

Department of Entomology and Centre for Tropical Pest Management, University of Queensland, Brisbane, Qld, 4072

Recent research, relying on the rain of arthropods that falls after the application of a chemical fog or spray, has clearly demonstrated that insects are abundant and diverse in the canopies of Australian rainforests (Bassett 1991; Kitching *et al.* 1993). Although estimates of insect species number may have been too enthusiastic (Monteith 1990), actual arthropod diversity and abundance are grossly underestimated by these chemical knockdown techniques because of their strong bias against small arthropods, especially mites (Walter *et al.* 1994). Like the soil beneath the canopy, rainforest trees themselves are infused with a terrestrial plankton, composed primarily of minute, scurrying, eight-legged arthropods.

To demonstrate this, I obtained estimates of the number of mites inhabiting the leaves of rainforest trees. Extendable pole-pruners were used to clip leaves from the canopies of a scentless rosewood and a native gardenia adjacent to Mick's Tower, O'Reilly's Rainforest Retreat, Lamington National Park (one of the sites reported in Kitching *et al.* 1993) on 10.iii.1993, and to clip the large, compound leaves of a brown tamarind at the CSIRO forestry research tower at Curtain Fig near Atherton on 24.iv.1993. Leaves were placed in plastic bags and chilled until mites could be counted and collected using a fibre optic-illuminated stereomicroscope. I also counted the number of leaves per branch on a sample of branches, and the number of branches per tree on each of these, relatively small, rainforest trees. The mean number of leaves per branch (rounded down) times the number of branches gave conservative estimates of the number of leaves per tree (Table 1).

**Table 1.** Estimates of the number of mites inhabiting leaves of some rainforest trees in Queensland. Only estimates of mites living on leaves are given.

Site	Tree Species	Height	Leaves/Tree	Mites/Leaf	Mites/Tree
Curtain Fig	<i>Diploglottis</i>	26 m	4,000	95.4 ± 12.2	381,600
	<i>diphyllostegia</i>			(n = 15)	
O'Reilly's	<i>Synoum</i>	12 m	16,500	28.3 ± 2.8	466,950
	<i>glandulosum</i>			(n = 45)	
O'Reilly's	<i>Randia</i>	10 m	34,000	10.92 ± 1.3	371,280
	<i>benthamiana</i>			(n = 50)	



Rather than being relatively rare (Kitching *et al.* 1993) or absent (Basset 1991), as implied by previous studies, Australian rainforest canopies are covered in mites, numerous species and uncountable numbers (Table 1). Many thousands of other mites live on stems and bark, in flowers and epiphytes, and as commensals and parasites on many of the insects and all of the vertebrates that also inhabit these trees. This richness of animals must have consequences for the canopy system, especially those resulting from the feeding of the predatory and scavenging-fungivore mites that made up the vast majority of the mites discovered.

### References

- BASSET, Y. 1991. The taxonomic composition of the arthropod fauna associated with an Australian rainforest tree. *Australian Journal of Zoology* **39**: 171-190.
- KITCHING, R.L., BERGELSON, J.M., LOWMAN, M.D., McINTYRE, S. and CARRUTHERS, G. 1993. The biodiversity of arthropods from Australian rainforest canopies: General introduction, methods, sites and ordinal results. *Australian Journal of Ecology* **18**: 181-191.
- MONTEITH, G.B. 1990. Rainforest insects: Biodiversity, bioguesstimation, or just hand-waving? *Myrmecia* **26**: 93-95.
- WALTER, D.E., O'DOWD, D.J. and BARNES, V. 1994. The forgotten arthropods: foliar mites in the forest canopy. *Memoirs of the Queensland Museum*



Walter, David Evans. 1993. "Queenslands rainforest canopies - a mitey cornucopia." *Australian Entomological Magazine* 20(4), 115–116.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/313083>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/347495>

**Holding Institution**

Museums Victoria

**Sponsored by**

Atlas of Living Australia

**Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Entomological Society of Queensland

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.