Case 3149

Proposed conservation of 31 species-group names originally published as junior primary homonyms in *Buprestis* Linnaeus, 1758 (Insecta, Coleoptera)

C.L. Bellamy

Natural History Museum of Los Angeles County, 900 Exposition Boulevard, Los Angeles, California 90007, U.S.A. (e-mail: cbellamy@nhm.org)

Abstract. The purpose of this application is the conservation of 31 specific names which have been in use for buprestid beetles for very many years but which, when originally published in combination with *Buprestis*, were junior primary homonyms. The species are now placed in many different genera. In none of these cases have the species denoted by the homonyms been considered congeneric since the 19th-century, if at all, and this case is submitted in accord with Article 23.9.5 of the *Code*.

Keywords. Nomenclature; taxonomy; Coleoptera; BUPRESTIDAE; *Buprestis*; buprestids; jewel beetles.

1. According to Article 23.9.5 of the Code, when two or more primary homonyms are in use 'but the names apply to taxa not considered congeneric after 1899, an author must not automatically replace the junior homonym; the case should be referred to the Commission and meanwhile prevailing usage of both names is to be maintained'. Therefore, I refer a number of primary homonymy situations from the very well-known genus Buprestis Linnaeus, 1758 (type genus of the jewel beetle family BUPRESTIDAE Leach, 1815), and propose the conservation of the existing usage of 31 specific names which have not appeared in homonymous combinations since the 19th-century and in some cases have never been considered congeneric with their senior homonyms. In the 18th century, and in the early part of the 19th, Buprestis was a 'catch-all' genus (by 1800 more than 250 species had been described in combination with that generic name), but for more than a hundred years taxa first published in Buprestis have been divided amongst many genera and the original primary homonymies have been overlooked or ignored. In some cases the senior homonyms have not been in use because they have older synonyms. To now abandon the longstanding usage of numerous specific names would be taxonomically extremely confusing, and it is fortunate that the new Code dictates that this must no longer be done unless the Commission so rules.

2. The cases of primary homonymy of specific names originally published in combination with *Buprestis* are given below in alphabetical order. The junior name(s) is/are given first.

(1) *B. arcuata* Laporte & Gory, 1837 (p. 159). Preoccupied by *B. arcuata* Say, 1825 (p. 251). *B. arcuata* Laporte & Gory was transferred to *Cinyra* Laporte & Gory, 1837 by Gemminger & Harold (1869, p. 1381) and later into *Spectralia* Casey, 1909; it is now known as *S. arcuata*. *B. arcuata* Say was placed in *Agrilus* Curtis, 1825 by Say (1839, p. 162) and is now known as *A. arcuatus*.

(2) *B. aurata* Fabricius, 1787 (p. 178). Preoccupied by *B. aurata* Pallas, 1776 (p. 719). *B. aurata* Pallas was transferred to *Erythyrea* Serville in Dejean, 1833 by Marseul (1865, p. 190) and is now known as *E. aurata*. The Fabricius taxon was transferred into *Chrysochroa* Laporte & Carcel, 1833 by Laporte & Gory (1835, p. 16), and later into *Chrysaspina* Théry, 1926 (a replacement name for the junior homonym *Chrysaspis* Saunders, 1869); it is now known as *Chrysaspina aurata*. Another species was called *B. aurata* by Thunberg (1787, p. 52), but since Saunders (1871, p. 50) it has been treated as a synonym of *Aristosoma suturale* (Thunberg, 1789) and I do not propose the conservation of its name.

(3) *B. bella* Gory, 1840 (p. 116). Preoccupied by *B. bella* Guérin-Méneville, 1830 (p. 66). *B. bella* Guérin-Méneville was moved into *Anthaxia* Eschscholtz, 1829 by Mannerheim (1837, p. 86) and is currently treated as a junior synonym of *Curis maulica* (Molina, 1782). *B. bella* Gory was transferred to *Poecilonota* Eschscholtz, 1829 by Lacordaire (1857, p. 37) and is now placed in *Scintillatrix* Obenburger, 1956.

(4) *B. bilineata* Latreille, 1813 (p. 60). Preoccupied by *B. bilineata* Weber, 1801 (p. 74). *B. bilineata* Latreille was transferred to *Psiloptera* Solier, 1833 by Dejean (1833, p. 76) and is currently placed in *Pseudolampetis* Obenberger, 1926. *B. bilineata* Weber was placed in *Agrilus* Curtis, 1825 by Say (1839, p. 162) and is now known as *A. bilineatus*.

(5) *B. cayennensis* Herbst, 1801 (p. 56). Preoccupied by *B. cayennensis* Gmelin, 1790 (p. 1931). The name *B. cayennensis* has been used for three different species. *B. cayennensis* Gmelin was placed in *Actenodes* Dejean, 1833 by Gemminger & Harold (1869, p. 1421) and is known as *A. cayennensis*. *B. cayennensis* Herbst was first transferred to *Chrysobothris* Eschscholtz, 1829 by Dejean (1833, p. 79) and later to *Colobogaster* Solier, 1833; it is now known as *Colobogaster cayennensis*. *B. transferred* to *Chrysobothris* Eschscholtz, 1829 by Dejean (1833, p. 79) and later to *Colobogaster* Solier, 1833; it is now known as *Colobogaster cayennensis*. *B. cayennensis*. *Chevelse*. *Cheve*

(6) *B. coerulea* Olivier, 1790 (p. 21). Preoccupied by *B. coerulea* Thunberg, 1789 (p. 91). The name *B. coerulea* has been used for three different species. *B. coerulea* Olivier was transferred to *Psiloptera* Solier, 1833 by Redtenbacher (1843, p. 505) and later to *Lampetis* Dejean, 1833; it is now known as *L. coerulea*. *B. coerulea* Thunberg, 1789 was transferred to *Meliboeus* Deyrolle, 1864 by Saunders (1871, p. 106) and is now known as *M. coeruleus*. *B. coerulea* Rossi, 1790 (p. 407) has been treated as a synonym of *Agrilus cyanescens* Ratzeburg, 1837 since Saunders (1871, p. 106) and I do not propose its conservation.

(7) *B. cuprifera* Laporte & Gory, 1836 (p. 59). Preoccupied by *B. cuprifera* Kirby, 1818 (p. 457). *B. cuprifera* Laporte & Gory has been placed in *Polybothris* Spinola, 1837 since Kerremans (1892). The Kirby taxon was transferred to *Plagiope* Saunders, 1868 by Saunders (1869, p. 13); it is now placed in *Torresita* Gemminger & Harold, 1869, a replacement name for the preoccupied *Plagiope*. A subsequent use of *B. cuprifera* by Laporte & Gory (1837, p. 119) is currently treated as a synonym of *Melobasis propinqua* (Laporte & Gory, 1837, p. 120) and I do not propose its conservation.

(8) *B. cyanea* Rossi, 1790 (p. 20). Preoccupied by *B. cyanea* Fabricius, 1775 (p. 223). The name *B. cyanea* has been used for three different species. *B. cyanea*

Rossi has been placed in *Agrilus* Curtis, 1825 since that genus was established. *B. cyanea* Fabricius is the type species of *Phaenops* Dejean, 1833. *B. cyanea* Olivier, 1790 (p. 91) has been treated as a junior synonym of *Agrilus viridis* (Linnaeus, 1758) since Dejean (1833, p. 83) and I do not propose its conservation.

(9) *B. cyanipes* Say, 1823 (p. 164). Preoccupied by *B. cyanipes* Fabricius, 1787 (p. 178). *B. cyanipes* Say has been placed in *Poecilonota* Eschscholtz, 1829 since LeConte (1860, p. 254) and is a well-known North American species. *B. cyanipes* Fabricius is a Jamaican species placed in *Polycesta* Serville in Dejean, 1833 since Kerremans (1892, p. 161).

(10) *B. depressa* Fabricius, 1775 (p. 219). Preoccupied by *B. depressa* Linnaeus, 1771 (p. 533). The name *B. depressa* has been used four times, with two of the junior homonyms replaced earlier. *B. depressa* Fabricius has been placed in *Pelecopselaphus* Solier, 1833 since Solier (1833, p. 287). *B. depressa* Linnaeus, 1771 has been in *Polycesta* Serville in Dejean, 1833 since Lacordaire (1857, p. 63). Since Laporte & Gory (1838, p. 3) *B. depressa* Fabricius, 1775 has been known by the replacement name *Strigopteroides aegyptiaca* (Gmelin, 1790, p. 1932). *B. depressa* Olivier, 1790 (p. 39) is known by the replacement name *Polycesta olivieri* Waterhouse, 1904 (p. 255).

(11) *B. drummondi* Kirby, 1837 (p. 157). Preoccupied by *B. drummondi* Laporte & Gory, 1836 (p. 37). *B. drummondi* Kirby was transferred to *Melanophila* Eschscholtz, 1829 by Gemminger & Harold (1869, p. 1385); it is currently known as *Phaenops drummondi*. *B. drummondi* Laporte & Gory was transferred to *Psiloptera* Solier, 1833 by Gemminger & Harold (1869, p. 1366) and is currently placed in *Lampetis* Dejean, 1833.

(12) *B. excellens* Klug, 1855 (p. 644). Preoccupied by *B. excellens* Klug, 1825 (p. 421). *B. excellens* Klug, 1855 has been placed in *Acmaeodera* Eschscholtz, 1829 since Lacordaire (1857, p. 68), while the species named in 1825 has been placed in *Conognatha* Eschscholtz, 1829 since Dejean (1833, p. 76).

(13) *B. fasciata* Villers, 1789 (p. 339). Preoccupied by *B. fasciata* Fabricius, 1787 (p. 177). The name *B. fasciata* has been used for three species. *B. fasciata* Villers has been placed in *Coraebus* Gory & Laporte, 1839 since Saunders (1871, p. 104). *B. fasciata* Fabricius was placed in *Ancylochira* Eschscholtz, 1829 by Dejean (1833, p. 78) and is now known as *Cypriacus fasciatus*. *B. fasciata* Voet, 1806 (p. 96) has been treated as a junior synonym of *Dismorpha linearis* (Linnaeus, 1758) since Saunders (1871, p. 112) and I do not propose its conservation.

(14) *B. femorata* Olivier, 1790 (p. 47). Preoccupied by *B. femorata* Villers, 1789 (p. 338). *B. femorata* Olivier has been placed in *Chrysobothris* Eschecholtz, 1829 since Dejean (1833, p. 80) and is a widespread and economically important species in North America. The Villers taxon has long been synonymized under *Anthaxia hungarica* (Scopoli, 1772).

(15) *B. flavofasciata* Herbst, 1801 (p. 306). Preoccupied by *B. flavofasciata* Piller & Mitterpacher, 1783 (p. 84). Herbst's taxon has been treated as a subspecies of *Acmaeodera trifasciata* (Thunberg, 1789, p. 95) since Gemminger & Harold (1869, p. 1412). *B. flavofasciata* Piller & Mittenpacher is the type species of *Acmaeodorella* (*Carininota*) Volkovitsh, 1979.

(16) *B. foveicollis* Gory, 1840 (p. 95). Preoccupied by *B. foveicollis* d'Urville in Boisduval, 1835 (p. 73). The southern African *B. foveicollis* Gory was transferred to

Damarsila Thomson, 1878 by Thomson (1878, p. 33) and is now placed in *Lampetis* Dejean, 1833. *B. foveicollis* d'Urville was designated by Deyrolle (1864, p. 46) as the type species of his genus *Cyphogastra* and is still placed in that genus.

(17) *B. geminatus* Say, 1823 (p. 163). Preoccupied by *B. geminatus* Illiger, 1803 (p. 244). Say (1839, p. 162) transferred his *B. geminatus* of 1823 to *Agrilus* Curtis, 1825; it is still placed in that genus and is a widespread eastern North American species. Since Dejean (1833, p. 81) *B. geminatus* Illiger has been treated as a junior synonym of *Sphenoptera rauca* (Fabricius, 1787, p. 177).

(18) *B. gibbicollis* Say, 1823 (p. 161). Preoccupied by *B. gibbicollis* Illiger, 1803 (p. 239). *B. gibbicollis* Say was transferred to *Ptosima* Serville in Dejean, 1833 by LeConte (1860, p. 221) and is widespread and well-known in eastern North America. Illiger's taxon from Iberia was transferred to *Coraebus* Gory & Laporte, 1839 by Gistel (1856, p. 135) and is now placed in *Meliboeus* Deyrolle, 1864.

(19) *B. haemorrhoidalis* Olivier, 1790 (p. 38). Preoccupied by *B. haemorrhoidalis* Herbst, 1780 (p. 97). Olivier's species was transferred to *Stigmodera* Eschscholtz, 1829 by Laporte & Gory (1838, p. 55) and is currently placed in the Neotropical genus *Conognatha* Eschscholtz, 1829. *B. haemorrhoidalis* Herbst is a widespread Palaearctic species still classified in *Buprestis*.

(20) *B. interrupta* Laporte & Gory, 1837 (p. 81). Preoccupied by *B. interrupta* Olivier, 1790 (p. 26). Gemminger & Harold (1869, p. 1368) transferred *B. interrupta* Laporte & Gory to *Psiloptera* Solier, 1833; it is now regarded as a subspecies of *Lampetis rugosa* (Palisot de Beauvois, 1807, p. 44). Olivier's taxon has been known as *Sternocera interrupta* since Dejean (1883, p. 74).

(21) *B. maculipennis* Gory, 1841 (p. 118). Preoccupied by *B. maculipennis* Laporte & Gory, 1837 (p. 111). Gory's taxon of 1841 remains in *Buprestis*, while the earlier homonym has been used as *Halecia maculipennis* (Laporte & Gory, 1837) since Gemminger & Harold (1869, p. 1361).

(22) *B. mucronata* Laporte & Gory, 1836 (p. 62). Preoccupied by *B. mucronata* Klug, 1825 (p. 426). Gemminger & Harold (1869, p. 1364) placed *B. mucronata* Laporte & Gory in *Polybothris* Dejean, 1833 and on p. 1442 transferred Klug's species to *Agrilus* Curtis, 1825; these placements have been maintained ever since.

(23) *B. nobilis* Fabricius, 1787 (p. 180). Preoccupied by *B. nobilis* Linnaeus, 1758 (p. 410). *B. nobilis* Fabricius has been placed in *Chrysobothris* Eschscholtz, 1829 since Saunders (1871, p. 95), while Dejean (1833, p. 80) designated *B. nobilis* Linnaeus as the type species of his new genus *Actenodes*, in which it remains today.

(24) *B. picta* Thunberg, 1827 (p. 47) and *B. picta* Waterhouse, 1882 (p. 15). Both preoccupied by *B. picta* Pallas, 1773 (p. 719). The name *B. picta* has been used for three species. The oldest homonym was transferred to *Melanophila* Eschscholtz, 1829 by Gemminger & Harold (1869, p. 1385) and later to *Trachypteris* Kirby, 1837, of which it is the type species; it is currently known as *T. picta* (Pallas, 1773). *B. picta* Thunberg was transferred to *Julodis* Solier, 1833 by Saunders (1871, p. 3) and since Kerremans (1902, p. 13) has been known as *Neojudolis picta* (Thunberg, 1827). *B. picta* Waterhouse, 1882 is still placed in *Buprestis*.

(25) *B. pumila* Klug, 1829 (p. 37). Preoccupied by *B. pumila* Illiger, 1803 (p. 275). *B. pumila* Klug has been placed in *Anthaxia* Eschscholtz, 1829 at least since Gemminger & Harold (1869, p. 1390), while *B. pumila* Illiger has been in *Trachys* Fabricius, 1801 at least since Lacordaire (1857, p. 88). (26) *B. salicis* Lewis, 1893 (p. 337). Preoccupied by *B. salicis* Fabricius, 1776 (p. 237). The placement by Lewis of his taxon in *Buprestis* is inexplicable; it was transferred to *Trachys* Fabricius, 1801 by Kerremans (1903, p. 310) and is now treated as a subspecies of *T. minutus* (Linnaeus, 1758). *B. salicis* Fabricius has been placed in *Anthaxia* Eschscholtz, 1829 since Dejean (1833, p. 80).

(27) *B. sulcata* Fischer von Waldheim, 1824 (p. 197). Preoccupied by *B. sulcata* Thunberg, 1789 (p. 90). The Fischer von Waldheim taxon has been placed in *Sphenoptera* Dejean, 1833 since Gemminger & Harold (1869, p. 1416). *B. sulcata* Thunberg, 1789 was transferred to their new genus *Chrysodema* by Laporte & Gory (1835, p. 18) and is currently placed in *Iridotaenia* Deyrolle, 1864. A subsequent use of the name *B. sulcata* by Thunberg (1827, p. 44) has been treated as a junior synonym of *Pelecopselaphus depressus* (Fabricius, 1775, p. 219) since Saunders (1871, p. 19).

(28) *B. variolosa* Fabricius, 1801 (p. 109. Preoccupied by *B. variolosa* Paykull, 1799 (p. 219). The Fabricius taxon was transferred to *Psiloptera* Solier, 1833 by Gemminger & Harold (1869, p. 1369) and is currently placed in *Lampetis* Dejean, 1833. *B. variolosa* Paykull was transferred to his new genus *Lampra* by Dejean (1833, p. 78); it was designated as the type species of *Poecilonota* Eschecholtz, 1829 in Opinion 1825 (1996). Another species called *B. variolosa* by Laporte & Gory (1836, p. 30) is now known by the replacement name *Lampetis coeruleitarsis* (Saunders, 1871, p. 23).

(29) *B. ventralis* Waterhouse, 1882 (p. 14). Preoccupied by *B. ventralis* Laporte & Gory, 1837 (p. 158). *B. ventralis* Waterhouse remains in *Buprestis*, while the Laporte & Gory taxon was transferred to *Cinyra* Laporte & Gory, 1837 by Lacordaire (1857, p. 46) and to *Halecia* Laporte & Gory, 1837, in which it is now placed, by Saunders (1871, p. 20)

(30) *B. vetusta* Boisduval, 1835 (p. 85). Preoccupied by *B. vetusta* Ménétries, 1832 (p. 152). The Boisduval taxon is the type species by monotypy of *Nascio* Laporte & Gory, 1837. *B. vetusta* Ménétries was transferred to *Acmaeodera* Eschscholtz, 1829 by Gemminger & Harold (1869, p. 1414) and is currently known as *Acmaeoderella vetusta*.

3. As stated at the outset, the junior primary homonyms listed above have been in use for considerably more than 100 years and to displace them now would upset the established nomenclature in numerous genera without serving any purpose. In some cases there are available synonyms which could be introduced as substitute names, but in others entirely new names would have to be formulated; in neither situation would the changes offer anything but loss of information and confusion. This application for the conservation of prevailing usage is submitted in the interests of stability and in accord with Article 23.9.5 of the *Code*.

4. The International Commission on Zoological Nomenclature is accordingly asked to use its plenary power to rule that the following specific names are not to be treated as invalid by reason of having been originally published as junior primary homonyms in combination with *Buprestis* Linnaeus, 1758, and to place these names on the Official List of Specific Names in Zoology:

- (1) arcuata Laporte & Gory, 1837;
- (2) aurata Fabricius, 1787;
- (3) *bella* Gory, 1840;
- (4) bilineata Latreille, 1813;

- (5) cayennensis Herbst, 1801;
- (6) coerulea Olivier, 1790;
- (7) cuprifera Laporte & Gory, 1836;
- (8) cyanea Rossi, 1790;
- (9) cyanipes Say, 1823;
- (10) depressa Fabricius, 1775;
- (11) drummondi Kirby, 1837;
- (12) excellens Klug, 1855;
- (13) fasciata Villers, 1789;
- (14) femorata Olivier, 1790;
- (15) flavofasciata Herbst, 1801;
- (16) foveicollis Gory, 1840;
- (17) geminatus Say, 1823;
- (18) gibbicollis Say, 1823;
- (19) haemorrhoidalis Olivier, 1790;
- (20) interrupta Laporte & Gory, 1837;
- (21) maculipennis Gory, 1841;
- (22) mucronata Laporte & Gory, 1836;
- (23) nobilis Fabricius, 1787;
- (24) (a) *picta* Thunberg, 1827;
 (b) *picta* Waterhouse, 1882;
- (25) *pumila* Klug, 1829;
- (26) salicis Lewis, 1893;
- (27) sulcata Fischer von Waldheim, 1824;
- (28) variolosa Fabricius, 1801;
- (29) ventralis Waterhouse, 1882;
- (30) vetusta Boisduval, 1835.

Acknowlegements

A draft of this manuscript was reviewed by R.L. Westcott, Oregon Department of Agriculture, Salem, Oregon, U.S.A., G.H. Nelson, Blue Springs, Missouri, U.S.A. and S. Bílý, National Museum, Prague, Czech Republic.

References

- **Boisduval, J.B.A.D. de.** 1835. Voyage de découvertes de l'Astrolabe exécuteé ... pendant les années 1826–1827–1828–1829, sous les commandement de M.J. Dumont d'Urville. Faune entomologique de l'Océan Pacifique, avec l'illustration des insectes nouveaux recueillis pendant le voyage. Deuxième partie. Coléoptères et autres ordres. vii, 716 pp. Tastu, Paris.
- Boisduval, J.B.A.D. de & Lacordaire, J.T. 1835. Faune entomologique des environs de Paris ..., vol. 1. 17, cii, 696 pp. Méquignon-Marvis, Paris.
- **Dejean, P.F.M.A.** 1833. *Catalogue des Coléoptères de la collection de M. le Comte Dejean*, fasc. 1. 96 pp. Méquignon-Marvis, Paris.
- Deyrolle, H. 1864. Description des Buprestides de la Malaisie recueillés par M. Wallace. Annales de la Société Entomologique de Belgique, 8: 1–312.
- Fabricius, J.C. 1775. Systema entomologiae ... 832 pp. Kortii, Flensburg.
- Fabricius, J.C. 1776. Genera insectorum. xiv, 310 pp. Bartsch, Chilonii.
- Fabricius, J.C. 1787. Mantissa insectorum ..., vol. 1, 348 pp; vol. 2, 382 pp. Proft, Hafniae.
- Fabricius, J.C. 1801. Systema Eleutheratorum . . ., vol. 2. 687 pp. Bibliopolii Academici Novi, Kiliae.

- Fischer von Waldheim, G. 1824. Entomographia Imperii Russici. Genera Insectorum systematica exposita et analysi iconographica instructa. Coleoptera. Volume 2 (1823–1824). xx, 262 pp., 40 pls.
- Gemminger, M. & Harold, E. von. 1869. Catalogus coleopterorum hucusque descriptorum synonymicus et systematicus, vol. 5 (pp. 1347–1608). Monachii, London.
- Gistel, J.N.F.X. 1856. Pleroma zu den Mysterien der europäischen Insectenwelt. Mit einem systematischen Verzeichniss der Schmetterlinge und Käfer Europas. 250 pp. Schorner, Straubing.
- Gmelin, J.F. 1790. Caroli a Linné Systema Naturae, Ed. 13, vol. 1 (4, Insecta, pp. 1517–2224). Lipsiae.
- Gory, H.L. 1840–1841. Histoire naturelle et iconographie des insectes Coléoptères. Monographie des buprestides, Supplement. Duméril, Paris.
- Guérin-Méneville, F.E. 1830. Crustacées, Arachnides et Insectes. In: Duperrey, L.J., Voyage autour du monde sur la Coquille. Zoologie, vol. 2, pt. 2, div. 1. 319 pp., 22 pls.
- Harold, E. von. 1869. Berichtigungen und Zusätze zum Catalogus Coleopterorum synonymicus et systematicus. Coleopterologische Hefte 5, pp. 112–119, 122–125.
- Herbst, J.F.W. 1780. Beschreibung aller Prachtkäfer ... Schriften der Berlinerischen Gesellschaft Naturforschender Freunde, 1: 85–100.
- Herbst, J.F.W. 1801. Natursystem aller bekännten in-und ausländischen Insecten, als eine Fortsetzung der Büssonschen Naturgeschichte. Der Käfer, vol. 9. 344 pp.
- Holm, E. 1978. Monograph of the genus Acmaeodera Eschscholtz (Coleoptera: Buprestidae) of Africa south of the Sahara. Department of Agricultural Technical Services, Republic of South Africa, Entomology Memoir no. 47. 210 pp.
- Illiger, J.C.W. 1803. Verzeichniss der in Portugal einheimischen Käfer. Erste Lieferung. Magazin für Insectenkunde, 2: 186–258.
- Kerremans, C. 1892. Catalogue synonymique des Buprestides decrits de 1758 à 1890. Mémoires de la Société Entomologique de Belgique, 1: 1–304.
- Kerremans, C. 1902. Coleoptera, Fam. Buprestidae. In: Wytsman, P. (Ed.), Genera Insectorum, Fasc. 12a (pp. 1–48). Verteneuil & Desmet, Bruxelles.
- Kerremans, C. 1903. Coleoptera, Fam. Buprestidae [contd.]. In: Wytsman, P. (Ed.), Genera Insectorum, Fasc. 12b, 12c and 12d (pp. 49–338). Verteneuil & Desmet, Bruxelles.
- Kirby, W. 1818. A description of several new species of Insects collected in New Holland by Robert Brown. *Transactions of the Linnean Society of London*, **12**: 454–482.
- Kirby, W. 1837. The Insects. In: Richardson, J. (Ed.), Fauna Boreali-Americana, part 4. 325 pp. Longman, London.
- Klug, J.C.F. 1825. Entomologiae Brasilianae specimen alterum, sistens insectorum coleopterorum nondum descriptorum centuriam. *Nova Acta Physico-Medica Academiae Caesareae-Leopoldino Carolinae, Naturae Curiosorum*, **12**(2): 421–476.
- Klug, J.C.F. 1829. Symbolae physicae seu icones et descriptiones insectorum quae ex itinere per Africam borealem et Asiam occidentalem Friderici Gulielmi Hemprich et Christiani Godofredi Ehrenberg ..., fol. 5. Mittler, Berlin.
- Klug, J.C.F. 1855. Diagnosen der neuen ... Coleopteren ... von Mossambique. Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 1855: 643–660.
- Lacordaire, J.T. 1857. *Histoire naturelle des insectes*. Genera des Coléoptères ..., vol. 4. 554 pp. Roret, Paris.
- Laporte, F.L. de L. (Comte de Castelnau) & Gory, H.L. 1835–1838. Histoire naturelle et iconographie des insectes coléoptères, vol. 1. Duméril, Paris.
- Latreille, P.A. 1813. Insects de l'Amérique équinoxiale recueillis pendant le voyage de MM. de Humboldt et Bonpland. Pp. 127–252 *in* Humboldt, F.H.A. von & Bonpland, A.J.A., *Voyage aux régions equinoxiales du Nouveau Continent, fait en 1799–1804*, vol. 2, part 2. Paris.
- LeConte, J.L. 1860. Revision of the Buprestidae of the United States. Transactions of the American Philosophical Society, (n.s.)11: 187–258.
- Lewis, G. 1893. On the Buprestidae of Japan. Journal of the Linnean Society of London (Zoology), 24: 327–338.
- Linnaeus, C. 1758. Systema Naturae, Ed. 10, vol. 1. 824 pp. Salvii, Holmiae.

Linnaeus, C. 1771. Mantissa Plantarum. 588 pp. Holmiae.

- Mannerheim, C.G. von. 1837. Enumération des Buprestides, et description de quelques nouvelles espèces de cette tribu de la famille des Sternoxes, de la collection de M. Le Comte Mannerheim. Bulletin de la Société Impériale des Naturalistes de Moscou, 8: 1–126.
- Marseul, S.A. de. 1865. Monographie des buprestides, famille des sternoxes de Latreille. L'Abeille. *Mémoires d'Entomologie*, **2**: 1–289.
- Molina, J.I. 1782. Saggio sulla storia naturale del Chili. 367 pp. Bologna.
- Olivier, G.A. 1790. Entomologie, ou histoire naturelle des insectes Coléoptères, genera 9-34. Vol. 2(32). 485 pp., 63 pls. Baudouin, Paris.
- Palisot de Beauvois, A.M.F.S. 1807. Insectes recueillis en Afrique et en Amérique, vol. 16. 276 pp. Levrault, Paris.
- Pallas, P.S. 1776. Reise durch verschiedene Provinzen des Russischen Reiches in den Jahren 1768–1774. Pisani.
- Paykull, G. von. 1799. Fauna Suecica, vol. 2. Insecta (Col.). 234 pp. Edman, Upsaliae.
- Piller, M. & Mitterpacher, L. 1783. Iter per Poseganum, Sclavoniae provinciam mensibus Junio et Julio 1782 susceptum. 147 pp. Budae.
- Redtenbacher, L. 1843. Abbildungen und Beschreibungen neuer und seltner Thiere und Pflanzen, in Syrien und im westlichen Taurus gesammelt von Th. Kotschy. Coleoptera. 254 pp., 2 pls. Schweizerbart, Stuttgart.
- Rossi, P. 1790. Fauna Etrusca, sistens Insecta, quae in provinciis Florentina et Pisana prasertim collegit P. Rossius, vol. 1. xii, 348 pp. Thomae Masi & Sociorum, Liburni.
- Saunders, E. 1869. Descriptions of nine new species of Buprestidae. *Transactions of the Royal Entomological Society of London*, 17: 1–8.
- Saunders, E. 1870. Catalogue of the species contained in the genus Buprestis of Linnaeus, previous to its subdivision by Eschscholtz in 1829, referring each to its present genus. 37 pp. van Voorst, London.
- Saunders, E. 1871. Catalogus Buprestidarum synonymicus et systematicus. 171 pp. Janson, London.
- Say, T. 1823. Descriptions of coleopterous insects collected in the late expedition to the Rocky Mountains performed by order of Mr. Calhoun, Secretary of War, under command of Major Long. *Journal of the Academy of Natural Sciences of Philadelphia*, 3: 9–54, 73–104, 139–216.
- Say, T. 1825. Descriptions of new American species of *Buprestis*, *Trachys* and *Elater*. Annals of the Lyceum of New York, 1: 249–269.
- Say, T. 1839. Description of new North American insects, and observations on some already described. *Transactions of the American Philosophical Society*, (2)6: 155–190.
- Solier, A.J.J. 1833. Essai sur les Buprestides. Annales de la Société Entomologique de France,
 2: 261–316.
- Thomson, J. 1878. Typi Buprestidarum Musaei Thomsoniani. 103 pp. Deyrolle, Paris.
- **Thunberg, C.P.** 1787. *Museum Naturalium Academiae Upsalensis*. Dissertation (pp. 43–58). Uppsala.
- Thunberg, C.P. 1789. Novas Insectorum species. 5. Dissertation (pp. 85-106). Uppsala.
- **Thunberg, C.P.** 1827. Coleoptera Capensia, antennis filiformibus, enumerata et nova descripta. Nova Acta Regiae Societatis Scientiarum Upsaliensis, 9: 1–52.
- Villers, C.J. de. 1789. Caroli Linnaei Entomologia. 24, 765 pp. Piestre et Delamolliére, Lugduni.
- Voet, J.E. 1806. Catalogus systematicus Coleopterorum, vol. 2. Bakhuysen, Haag.
- Waterhouse, C.O. 1882. In: Godman, F.D. & Salvin, O. (Eds.), Biologia Centrali-Americana, vol. 3, part 1. Insecta, Coleoptera, Buprestidae. xvi, 690 pp., 27 pls. London.
- Waterhouse, C.O. 1904. Observations on Coleoptera of the family Buprestidae, with descriptions of new species. Annals and Magazine of Natural History, (7)14: 245-267.
- Weber, F. 1801. Observationes entomologicae, continentes novorum quae condidit generum characteres, et nuper detectarum specierum descriptiones. 12, 116pp. Kiliae.

Comments on this case are invited for publication (subject to editing) in the *Bulletin*; they should be sent to the Executive Secretary, I.C.Z.N., c/o The Natural History Museum, London, SW7 5BD, U.K. (e-mail: iczn@nhm.ac.uk).



Bellamy, C. L. 2001. "Case 3149. Proposed conservation of 31 species group names originally published as junior primary homonyms in Buprestis Linnaeus, 1758 (Insecta, Coleoptera)." *The Bulletin of zoological nomenclature* 58, 24–31.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/105441</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/41794</u>

Holding Institution Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: International Commission on Zoological Nomenclature License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.