

XXXI.—*Note on Tarentola americana, Gray.*

By G. A. BOULENGER.

ON one of my last visits to the Paris Museum I was enabled, by the courtesy of Prof. Vaillant, to examine the type specimen of the Gecko named *Tarentola americana* by Gray and *Platydictylus Milbertii* by Duméril and Bibron. This specimen, sent by Milbert from New York to the Paris Museum, probably came from some West-Indian Island,—which we cannot say, as the lizard does not appear to have been rediscovered since. It bears most affinity to the Cuban *T. cubana*, Gundl. & Peters, the type of which I examined in the Berlin Museum (*cf.* Cat. Liz. iii. p. 490), and the West-African *T. Delalandii*, D. & B., but represents unquestionably a valid species. As in these species, the supraorbital bone is absent. The tubercles are larger than in *T. Delalandii*, scarcely keeled, in 12 series on the back; the enlarged caudal tubercles are smooth and nearly flat. The ear-opening is entirely surrounded by a fringe of converging conical tubercles. The symphysial (mental) shield is about once and a half as long as it is broad in the middle.

The number of well-defined species of *Tarentola* has now risen to nine, which may be distinguished by means of the following key:—

- I. Supraocular region bony; 13 to 20 scales across the head, from eye to eye.
 - A. Tail rounded on the sides, elliptical in section.
 - Dorsal tubercles strongly keeled 1. *T. mauritanica*, L.
 - Dorsal tubercles smooth or feebly keeled 2. *T. ephippiata*, O'Sh.
 - B. Tail flat inferiorly, with sharpish lateral edge.
 - Anterior border of ear denticulate 3. *T. annularis*, Geoffr.
 - Anterior border of ear not denticulate 4. *T. senegalensis*, Blgr.
- II. Supraorbital bone present, but small; 10 to 12 scales from eye to eye. 5. *T. neglecta*, Strauch.
- III. No supraorbital bone; 15 to 23 scales from eye to eye.
 - A. Anterior border of ear not denticulate; dorsal tubercles smooth or feebly keeled; symphysial shield at least twice as long as broad in the middle.
 - Symphysial shield not thrice as long as broad in the middle. 6. *T. Delalandii*, D. & B.
 - Symphysial shield thrice as long as broad in the middle 7. *T. gigas*, Bocage.
 - B. Anterior border of ear denticulate.
 - Symphysial shield about once and a half as long as broad in the middle; dorsal tubercles very feebly keeled. 8. *T. americana*, Gray.
 - Symphysial shield about twice as long as broad in the middle; dorsal tubercles strongly keeled 9. *T. cubana*, Gundl. & Peters.



Boulenger, George Albert. 1893. "XXXI.—Note on Tarentola americana, Gray." *The Annals and magazine of natural history; zoology, botany, and geology* 12, 204–204. <https://doi.org/10.1080/00222939308677603>.

View This Item Online: <https://www.biodiversitylibrary.org/item/78509>

DOI: <https://doi.org/10.1080/00222939308677603>

Permalink: <https://www.biodiversitylibrary.org/partpdf/62280>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.