


March 5, 1872.

John Gould, Esq., F.R.S., in the Chair.

An extract was read from a letter addressed to the Secretary by Mr. Walter T. Scott, C.M.Z.S., dated Vale of Herbert, Cardwell, Queensland, December 4, 1871. Mr. Scott wrote as follows of the supposed "Native Tiger" of Queensland, concerning which Mr. Sclater had previously communicated the evidence given by Mr. Sheridan (see P. Z. S. 1871, p. 629):—

"As to the Tiger, I am inclined to think there really is some large carnivorous animal as yet undescribed in this neighbourhood. A Mr. Hull, Licensed Surveyor, was lately at work with a party of five men, surveying on the Murray and Mackay rivers, north of Cardwell. They were lying in their tents one night between eight and nine o'clock, when they were all startled by a loud roar close to the tents. They seized their guns and carefully reconnoitred; but the animal had departed. In the morning they found the tracks of the unknown visitor, of which Mr. Hull took the measurements and a rough sketch. I send you part of a leaf of Mr. Hull's field-book,



Footprint of "Native Tiger," reduced one half.

containing the original sketch—and also his drawing of the track, of the natural size. Mr. Hull assures me that the drawing was a very faithful one, the soft ground having taken the impression with all its details. I have also examined some of the men who were with Mr. Hull. They all tell the same story, and say they heard the animal three nights in succession.

I think that I have already mentioned to you that a bullock-driver of ours, as long ago as 1864, came in one day with a story that he had seen a Tiger; but as he was a notorious liar we did not believe a word of it at the time. Yet it is possible he may really have seen the same animal, which must I think, from its claws, be allied to the Tasmanian Thylacine (*Thylacinus cynocephalus*)."

---



The following papers were read:—

1. On the Occurrence of *Falco barbarus* and *Cypselus pallidus* on the Continent of Europe. By HOWARD SAUNDERS, F.Z.S.

[Received March 4, 1872.]

In the excellent account of *Falco barbarus* given by Mr. O. Salvin in 'The Ibis,' 1859, p. 184 *et seq.*, he recommends that a look-out should be kept for it in Spain; and I have now great pleasure in exhibiting an example of this miniature Peregrine obtained near Granada, Spain, in January 1871. It appears to be a bird of the year, and proved to be a female on dissection. As Messrs. Salvin and Brodrick observe in their 'Falconry in the British Isles,' p. 101, "although smaller by nearly one fourth than the true Peregrine, it has the organs of destruction, such as the beak, feet, and talons, fully as large." Indeed in the present specimen the middle toe is *very nearly as long* as that of a magnificent adult female Peregrine, and *rather longer* than that of an adult male, her mate, shot near Seville, and rivalling in size the largest northern specimens. As Mr. Salvin remarks, the small stature, powerful feet and claws, and ruddy under plumage of *Falco barbarus* are its best characteristics.

In 'The Ibis,' 1870, p. 445, Capt. G. E. Shelley described *Cypselus pallidus* as new, from a specimen he had obtained in Egypt, where it would appear to take the place of *Cypselus apus*. He subsequently identified with this species specimens brought by Major Irby from Tangiers; and that gentleman further remarked that he had seen it in Spain. I am not aware that he has hitherto been successful in obtaining specimens in the Peninsula, and have therefore great pleasure in exhibiting a solitary specimen obtained at Granada on the 28th May, 1870, and sent to me along with a number of the common species, from which it may be distinguished by its lighter colour, white throat, and lighter forehead. From the date, it was probably breeding.

2. Notes on an Ostrich lately living in the Society's Collection. By A. H. GARROD, B.A., Prosector to the Society, and FRANK DARWIN, B.A.

[Received March 5, 1872.]

A male Ostrich (*Struthio camelus*) has been in the Society's Gardens since April 1869, and was quite healthy until last October, when its appetite began to fail, and it did not take kindly to its food from that time until its death on the 6th ult. In September last the keeper noticed on several occasions that after running about as it was accustomed to do in play, it turned giddy and apparently



tripped, but never quite fell. For the last four months it had lost flesh gradually. Whenever any fresh food was offered it, it would take a little and then refuse any more, and would do thus, however many new things were presented to it.

It had suffered from diarrhœa more or less ever since October, the excrement having a yellowish-green colour.

Latterly it had been nearly continually in the sitting position, and would stand very unwillingly. It also frequently rubbed its head and eyes with its foot, as if something was irritating it there.

In the post mortem examination very little structural disease was found; and the cause of death is more probably connected with the contents of the stomach rather than with any other agency.

There was more than half a gallon of stones in the stomach: most of them were about the size of cob-nuts or peas; and they fully dilated the organ and pulled it down abnormally. Mixed up with these stones were numerous copper coins and pieces of coins in a much worn state. There were two pennies and fifteen halfpence; and very few showed the least trace of the stamp they had previously borne, and those only by an oblique light, the difference in density of the metal, produced by the stamping, having caused them to wear unevenly. Most of them were slightly curved, being meniscoid in form. They were all highly polished and not in the least corroded. Many were in pairs, with a layer of softish green matter, about  $\frac{1}{20}$  of an inch thick, interposed. The chips of coins were very numerous and of all sizes below that of the coins themselves. No silver was found, and nothing else except a glove-button and a nut, the latter being at the bottom of the œsophagus.

All the contents of the stomach were of a green colour; and two small boluses of hay which it contained were tinged deeply with green.

Four more coins, deeply corroded and greenish black, were found in one of the intestinal cæca, together with a few stones. There were also a few stones in the other cæcum; and the mucous membrane of both cæca was congested and unhealthy in appearance, which was not the case in the stomach to any extent.

There were no symptoms of jaundice, which frequently accompanies copper poisoning. The liver appeared healthy, except that scattered about were a few dense white lumps about the size of peas, mostly near the surface: it weighed 3 lb. 9 oz. No gall-bladder was present.

The spleen was very small, and altogether weighed just under 2 oz. There was very little healthy tissue preserved, it mostly consisting of spheroidal dense masses of matter which were about the size of chestnuts, and by protruding beyond the general surface produced an appearance of knobs. These masses, on cutting through the capsule, separated entirely, and were then seen to be rough and altogether very like urinary calculi; they were of a fawn-colour. The organ was situated nearly in the middle line, just above the kidneys.

The heart weighed 1 lb. 7 oz., and gave origin to two carotid arteries, one from each main branch, which ran to the head, a distance of about 3 feet 6 inches, side by side, in front of the cervical vertebræ, in the groove formed by the anteriorly projecting processes



of those bones ; and they never showed any tendency to unite or cross one another. They were thickly covered by the anterior cervical muscles, and sent off symmetrical branches\*.

Superficially on each side of the neck ran a vein with the pneumogastric nerve ; but that on the left side was not bigger than a crow-quill, while that on the right had a diameter at the lower part of the neck of two thirds of an inch. This condition is constant in many birds.

This right (practically the only) jugular vein, after coursing about half or a little more up the neck, sent two branches to the head, the second running in the middle line, just behind the trachea and in front of the œsophagus, the first being a direct continuation of the main trunk.

The intestinal canal was 34 feet long ; and the two cæca, each 2 feet long and arranged like a spirally twisted cone, were situated 11 feet from the pylorus, which is very different from their situation in most birds, as has been noticed by Owen.

The diaphragm was well marked. It formed a partition which divided the thoracic cavity into two parts, one posterior and small containing the lungs, and the other anterior and large containing the heart and liver. It was a fibrous membrane, concave forwards, with a muscular attachment at either side to the ribs and intercostal tissues, which it joined in about the middle of their course. This muscular part was formed of transverse fibres in the middle and upper part of the chest, while the lower ones slanted slightly upwards as they coursed towards the median line. They were about 2 inches long, and formed a thin layer. The pleural cavity was closed above and below by the fibrous diaphragm becoming blended with the first and last ribs.

The anterior thoracic cavity, which contained the pericardium-coated heart in its upper part, entirely independent of the pleural cavity, was divided into two by a dense fibrous membrane which sprang from two vertebral crura, much as the human diaphragm, and extended above the line to join the sternum, along the border which articulated with the ribs, leaving the heart entirely in front of it ; its concavity was directed downwards and forwards ; and it was separated from the diaphragm proper by very large air-cells. The œsophagus also ran in the interval ; but the aorta was included in the pleural cavity, being clearly seen through the membrane of the diaphragm, along the median line, before its removal.

The liver was completely separated from the abdominal cavity by a fibrous membrane, so that when the included viscera had been removed it was not at all brought into view. The mesentery was very dense and strong, the vessels, especially the veins, being of large size.


\* The presence of two carotids in this bird, while there is only one in *Rhea*, would require that they should be far separated in Nitzsch's classification of birds according to the number of these vessels—the Ostrich being in his first class, with a carotid from each main aortic branch, and the *Rhea* in the fourth class, with only the left developed. See Nitzsch's 'Pterylography' (English edition), App. p. 171.



Some further points in the anatomy of this bird are not without interest.

There are three parietal abdominal muscles as usual, the muscular fibres of the external and internal being nearly parallel and transverse, while those of the intermediate one are longitudinal. They each send down a dense fascial attachment to the pubic bone; and a semilunar free margin between the ilium and the superior pubic crest appears closely allied to Poupart's ligament, the anterior crural vessels and nerves going underneath it to enter the leg. It may be here mentioned that the main vein of the thigh is the internal saphenous; but the main artery is the one that goes through the sciatic notch, therefore the sciatic. These come into relation with one another in the loop for the biceps tendon at the knee.

Exactly in the middle of the anterior border of the pubic portion of the innominate bone there is a small thin plate of osseous tissue which is connected with the pubis by strong fibrous bands, and which is continued anteriorly and superiorly by cartilage for some distance, when it becomes continuous with the tendons of the parietal abdominal muscles, being most connected with the external oblique.



Portions of the external surface of the left pubis and ischium of the Ostrich. The small osseous plate (*a*) attached to the pubis is represented partly surrounded by cartilage.

In dry skeletons a slight thickening of the anterior border of the pubic bone indicates the attachment of this ossification in most; but in one of the three skeletons in the British Museum this bone is ankylosed on one side, and Mr. Gerrard has specimens in which both are still attached. A diagram of the Ostrich's pelvis in Mr. Houghton's paper also shows this bone ankylosed, though no mention is made of it in his paper.

It would be extremely interesting to make out the homology of this small but perfectly independent ossification. Its relation to the



muscles of the abdominal wall would favour the idea of its corresponding to the marsupial bone of the Kangaroo and its allies; and if that is the case, the whole of the anterior prolongation of the Ostrich's pubis would correspond to the small ridge of bone on either side of the superior margin of the symphysis pubis in the Mammalia.

The *obturator internus* also arises from the superficial surface of this bone and its cartilage, as well as from the adjacent surface of the ischium and from the pubis, extending so far forward that the muscles of the opposite sides are only separated from one another by an inch or so at the symphysis pubis.

Mr. Macalister, in his description of the myology of this bird, has omitted a few of the muscles, some of which from the head will be described, together with those of the leg, which Mr. Frank Darwin has allowed me to introduce in this communication, from his notes and the dissection of that limb in this individual specimen.

*Pterygoid*.—From the inferior surface of the posterior part of the palate-bone, and from the process of bone which connects it with the main portion of it, this is fibrous—also from the whole of the inferior surface of the pterygoid bone, extending inwards almost to the basisphenoidal rostrum.

The fibres are all directed backwards, and are inserted in two ways:—the outer, and some of those from the palatine longitudinal process, into the anterior surface of the transverse ridge at the angle of the mandible, which posteriorly receives the insertion of the digastric muscle; the inner, and others from the palate-bone, into a fibrous band which runs from the side of the median Eustachian aperture and its cartilaginous continuation to the prominent ridge behind and internal to the condyloid articular surface for the mandible, thus forming an arch under which run the arteries and veins to the head.

This second portion of the muscle acts partly as an opener of the Eustachian aperture, partly as a retractor of the slightly movable pterygoid and palatine bones.

*Quadrato-mandibular*.—From the whole of the longitudinal ridge which forms the superior internal portion of the quadrate bone, and from the surface of the bone external to it. The fibres are directed outwards and downwards to be inserted into the inner surface of the mandible, in front of the articulation, not extending to the inferior margin, nor forwards further than the optic foramen.

*Quadrato-cranial*.—From the back of the orbit, below and behind the origin of the recti muscles and the exit of the nerve, from a surface bounded above by a semicircular line, and extending down in the space between the orbit and the quadrate bone. The fibres are directed outwards to the corresponding, internal surface of the quadrate bone, a slight ridge separating the superior ones from those of the quadrato-mandibular.

*Gastrocnemius* consists of two enormous masses of muscle blending together at their origins round the proximal end of the tibia, and separating lower down into the gastrocnemius anticus, which laps



round the anterior half, and the gastrocnemius posticus, which surrounds the back part of the tibial section of the limb. Gastrocnemius anticus arises partly from the tibia, partly by blending with gastrocnemius posticus; the latter arises from the distal extremity of the femur, the tendon of the quadriceps extensor and patella, and from the tibia. At the tibio-tarsal joint the gastrocnemii form a sheath fitting into the trochlea of the tibia for the passage of the flexor tendons of the toes; this is effected by the tendons becoming very much thickened and semicartilaginous (especially gastrocnemius posticus), and uniting with each other at their edges, the anterior element of the sheath being formed by gastrocnemius anticus, the posterior by gastrocnemius posticus. Just above the joint, gastrocnemius anticus sends off a slip which passes down in a special sheath along the outer surface of the contiguous heads of tibia and tarso-metatarsal bone, and is inserted into the tendon of flexor perforatus. Gastrocnemius anticus is inserted into the posterior surface of the tarso-metatarsal bone just below the tibio-tarsal joint. Gastrocnemius posticus is inserted into the external and internal lips of the posterior border of the tarso-metatarsal bone, forming a sheath for the passage of the flexor tendons; it subsequently forms, with a "sesamoid" cartilage presently to be described, a pulley for the same tendons at the tarso-phalangeal joint, and ends by blending with the fascia covering the sole of the foot. Mr. Macalister\* describes the gastrocnemius as ending in one tendon only, which he says forms a sheath for the deeper tendons on the back of the metatarsus.

The flexors of the toes are flexor magnus (perforatus), flexor perforans, flexor externi digiti, flexor interosseus, flexor profundus.

*Flexor magnus* arises (1) by a tendon from the upper part of the external surface of the outer condyle of the femur, the tendon winding over the knee, and then ending in the muscle; just before it does so, it receives the insertion of the rectus femoris (Cuvier and Meckel), (the pectineus of Owen); (2) from posterior surface of distal end of femur; (3) proximal end of tibia. The muscle ends in a broad tendon, which passes through the gastrocnemial sheath at the tibio-tarsal joint, and is here pierced by the tendon of flexor externi digiti. It passes down the tarso-metatarsal bone in the sheath formed by gastrocnemius posticus, receiving a tendinous slip, already described, from gastrocnemius anticus. At the tarso-phalangeal joint it passes through a sheath formed anteriorly by a "sesamoid" cartilage, posteriorly by the tendon of gastrocnemius posticus. This cartilage is ligamentously attached to the proximal end of the first phalanx of the internal digit, and to the synovial membrane of the tarso-phalangeal joint; it is deeply grooved posteriorly for the flexor tendons, and has two shallow grooves on its anterior surface, which fit on to the condyles of the tarso-metatarsal bone when the digits are extended; there is a smaller "sesamoid" cartilage for the external digit. A small muscle arises from the larger cartilage and by a few

\* Alexander Macalister, L.R.C.S.I., Demonstrator of Anatomy, Royal College of Surgeons, "On the Anatomy of the Ostrich (*Struthio camelus*)," Proceedings of the Royal Irish Academy, vol. ix. part 1 (1865).



fibres from the smaller one, and is inserted into the anterior surface of the flexor profundus tendon. The flexor magnus is pierced, as it passes through the sesamoid sheath or pulley, by the tendons of flexor profundus and flexor perforans, and ends by dividing into two slips, which are inserted into the proximal end of the second phalanx of the internal digit.

Mr. Macalister gives as the origin of the flexor magnus the deep pit above the condyles of the femur, the tendon of the rectus femoris, the external lateral ligaments, and the back of the fibula. In the specimen which I dissected, the tendon of the rectus femoris was much smaller than the tendinous head of the flexor profundus, and was inserted into it, which is an arrangement differing very slightly from its usual insertion in birds, which is, I believe, into the fleshy part of flexor magnus (Owen, Anat. Vert. vol. ii. p. 107).

The Rev. S. Haughton\* describes the rectus as becoming provided with a second muscular belly (p. 53), which does not at all describe its condition in my specimen. He considers this "digastric rectus femoris muscle" to be "the key to the explanation of the complicated muscular apparatus of the Ostrich's leg" (p. 50). Speaking of it, he says, "it acts before the extensor muscles come into full play; it binds down the two patellæ, braces up the heel-joint, and gives the signal for the m. gastrocnemio-solæus and other associated muscles to contract, and thus produces what may be regarded as one of the most striking phenomena in nature, viz. that the delicate bones and ligaments of a bird's leg, acted on by muscles equal to those influencing a horse's hind leg, shall remain uninjured under the sudden action of forces the slightest error in the application of which would break to pieces the machinery on which they act." This arrangement of the rectus, which Mr. Haughton considers so important, is only a well-developed form of what is found in most birds, and which Prof. Owen says is used in perching, by flexing the toes when the knee is bent (*loc. cit.*).

Mr. Macalister does not mention the muscle from the flexor profundus tendon to the sesamoid cartilage; but says that the flexor magnus sends a slip to it. The function of this muscle must be to keep the pulley-like sesamoid cartilage firmly in its place when the toes are extended preparatory to their flexion in the spring of the bird.

*Flexor digiti interni* arises from the outside of the tubercle of the tibia, and from the tendon of the quadriceps extensor; the tendon passes through the gastrocnemial sheath and the sesamoid pulley. It pierces the flexor magnus, receiving a tendinous slip from it, and then spreads out and binds the tendon of flexor profundus to the grooved under surface of the second phalanx of the internal digit, and is inserted by two slips into the proximal end of the third phalanx.

*Flexor digiti externi* arises from the posterior surface of the distal end of the femur, and from the tendinous head of flexor magnus. It passes through the gastrocnemial sheath, pierces the tendon of

\* The Rev. S. Haughton, M.D., "Notes on Animal Mechanics.—No. 3. On the Muscular Mechanism of the leg of the Ostrich," Proc. Roy. Acad. Dub. vol. ix. part 1 (1865).



flexor magnus, passes through the lesser sesamoid sheaths, and is inserted into each of the three proximal phalanges of the external toe.

*Flexor profundus* arises by two heads—one from the posterior surface of distal end of femur, the other from the posterior surface of the upper half of the tibia and part of the fibula. The tendon of the external muscle passes at the tibio-tarsal joint through a canal in the tendon of gastrocnemius anticus. The inner tendon does not enter the gastrocnemial sheath till below the joint; it passes down the inner surface of the contiguous ends of the tibia and the tarso-metatarsal bone, bound down to them by a special aponeurotic sheath, and joins the outer tendon near the tarso-phalangeal joint. The common tendon passes over the sesamoid pulley, piercing the flexor magnus tendon; it is here much thickened and hardened, and fits into the grooved and thickened tendon of flexor perforans. It is inserted into the fourth phalanx of the internal toe, sending off a short strong slip to the third phalanx, and an elastic slip to the second phalanx, as well as a small but long slip to flex the fifth phalanx of the external digit.

Mr. Macalister describes the tendon of the flexor profundus as being inserted only into the last phalanges of both digits; the insertion, as it was in my specimen, accords, I believe, with the usual condition of this tendon in birds.

The *flexor interosseus* is a delicate and weak muscle, which consists of numerous very short oblique fibres arising from the posterior surface of the tarso-metatarsal bone, and inserted into an aponeurosis stretching the whole length of the muscle; this aponeurosis ends in a tendon which is inserted into the outer surface of the first phalanx of the external digit. Its action appears to be to abduct and flex the toe.

A small muscle which Mr. Macalister does not describe, but merely mentions as probably representing the dorsal interosseus, arises from a small triangular space on the anterior surface of the distal end of the tarso-metatarsal bone, and is inserted into the capsular ligament of the tarso-phalangeal joint. There were some differences between the specimen which I dissected and that described by Mr. Macalister, in the precise origin of some of the muscles, which I have not thought to be worthy of note.

The *extensor communis digitorum* presents no peculiarities; the very small *extensor unguis* mentioned by Mr. Macalister was present.

### 3. Note on a Collection of small Mammalia made by Mr. Monteiro in Angola. By Dr. W. PETERS, F.M.Z.S.

[Received February 13, 1872.]

Mr. Sclater has sent to me for determination a collection of small Mammals made by Mr. J. J. Monteiro, C.M.Z.S., during his last travels in Benguela and Cambembe. The collection contains ex-



amples of eight species, of which, in the interests of science, I think it advisable to record the names and exact localities.

1. *Pterocyon stramineus*, Geoffroy. Benguela.
2. *Epomophorus macrocephalus*, Ogilby. Benguela.
3. *Epomophorus gambianus*, Ogilby. Benguela.
4. *Phyllorhina commersonii*, Geoffroy. Benguela.
5. *Nyctinomus limbatus*, Ptrs. Benguela.
6. *Mus barbarus*, Linné. Cambembe.
7. *Mus musculus*, Linné. Cambembe.
8. *Georychus hottentottus*, Lesson. Cambembe.

#### 4. Synopsis of the *Lamellirostres* of the Argentine Republic. By Dr. HERMANN BURMEISTER, F.M.Z.S.

[Received January 22, 1872.]

The tribe of natatorial birds named *Lamellirostres*, from the peculiar structure of the sides of the bill, which are furnished at the edges with small perpendicular corneous lamellæ, is very numerous represented in the vast territory of the Argentine Republic. As the list of species named in my travels through that country (vol. ii. p. 512 *et seq.*) is not entirely complete, and as I am now, after a residence of ten years in Buenos Ayres, better acquainted with the geographical distribution of many of the species than at the time of my first publication, I have thought it would be useful to prepare a second synopsis, giving additional remarks on most of the species, and correcting some of my former views from new observations.

#### I. PHÆNICOPTERINÆ.

##### 1. PHÆNICOPTERUS.

1. PHÆNICOPTERUS IGNIPALLIATUS, Is. Geoff. Mag. de Zool. 1832, *Aves*, pl. 2; Burm. Reise, ii. p. 512.

This Flamingo is common on the lagunes in the southern parts of the pampas of the Republic. I have obtained it at Mendoza and at Buenos Ayres. As I have already stated in my 'Reise,' the colour of the base of the bill is not rose-colour but white, and the legs of young birds are yellowish, with bluish-red articulations, but greenish grey in the old birds, with darker reddish articulations and toes.

2. PHÆNICOPTERUS ANDINUS, Philippi, Reise durch die Wüste Atacama, p. 164, pls. 4 & 5.

This well-marked species is found also in the north-western extremity of the Republic, on the lagunes in the eastern valleys between the Cordilleras and the adjacent mountains, and has been observed there by one of my countrymen, Hr. Schickendantz.



## II. ANATINÆ.

## 2. CYGNUS.

## 3. CYGNUS NIGRICOLLIS, auct. ; Burm. Reise, ii. p. 512.

A common bird in the southern parts of the Republic near the river Paraná and the great lagunes of the interior. It is often brought into the market at Buenos Ayres, and is occasionally to be seen in flocks consisting of from four to five individuals, flying at a great height over the town towards the river.

## 4. CYGNUS COSCOROBA, Gray ; Burm. Reise, ii. p. 512.

This bird is more numerous than the preceding species, and is generally seen in large flocks near Santa Fé on the river Salado, and in the south of Buenos Ayres on the same river. It is very common in Patagonia on the rivers Colorado and Negro, and is also to be seen in Buenos Ayres, flying high over the town in flocks which I have never observed to exceed six individuals in number. On the shore these flocks generally unite and form large bodies.

## 3. SARCIDIORNIS.

## 5. SARCIDIORNIS REGIA, Eyton ; Burm. Reise, ii. p. 513.

This large and beautiful Duck I have found only in the northern province of Tucuman, and never in the southern parts of the Republic near Paraná or Buenos Ayres. Azara says the same, and that the bird is common in Paraguay.

## 4. BERNICLA.

## 6. BERNICLA MELANOPTERA, Gay ; Philippi, Wiegmann. Arch. 1863, i. p. 185.

*Chloëphaga melanoptera*, Eyt. Monogr. p. 93 ; Burm. Reise, ii. p. 513.

This bird is found in the valleys of the Cordilleras from 8000 to 10,000 feet above the level of the sea, not descending into the plains. It is nowhere common, at least in the southern districts of the Republic.

I only found this bird while on my way from Copacavana to Copiapo on the river Blanco. They were always in pairs ; the sexes are alike in colour. The name given to this well-known bird by the inhabitants is "Pinquen."

There are three other species of *Bernicla* in the Argentine Republic, all living in the most southern parts of that country, from Bahia Blanca to the Straits of Magellan, where I have never been. On the approach of winter two of these species remove nearer to Buenos Ayres, where they are found in the vicinity of Tondil and the Laguna de los Padres, but exceptionally. They are more common at Bahia Blanca and El Carmen, on the Rio Negro, from which locality I have lately received both species. The third is an



entirely antarctic bird, as its name imports; and its range extends from Terra del Fuego to the inlet of Santa Cruz, where, I have been informed by a friend, this bird is found every year.

My distinguished countryman, Prof. R. A. Philippi, of Santiago in Chile, has published (in conjunction with L. Landbeck) in Wiegmann's 'Archiv' (1863, i. p. 187 *et seq.*) a valuable synopsis of the Chilian species of *Bernicla*, to which I refer for further information.

7. *BERNICLA ANTARCTICA*, Gmel.; Burm. Reise, ii. p. 514; Phil. Wiegmann. Arch. 1863, i. p. 199.

This species is never seen in the northern districts of Patagonia, but only in the south near the Straits of Magellan, going in the winter to the inlet of Santa Cruz. My former notice (in my Journey) that *B. antarctica* is found on the Sierra Tinta is an error caused by my having taken the following species (which was known to me at that time only by the description of my son) for *B. antarctica*.

8. *BERNICLA DISPAR*, Philippi, Wiegmann. Arch. 1863, i. p. 190. 2.

*Bernicla magellanica*, Cassin, Gilliss' Exped. ii. p. 201, pl. 24.

To this species must be referred my notice of a Goose living in the south of Buenos Ayres on the Sierra Tinta, near Tandil, and the adjacent country. The full description of Professors Philippi and Landbeck renders it unnecessary for me to add any thing further. I can only now extend the habitat of the species more to the south, having received both sexes from El Carmen on the river Negro, where this Goose is common in the summer, proceeding to the northern districts in the winter.

9. *BERNICLA POLIOCEPHALA*, Gray.

*Bernicla inornata*, Des Murs, in Gay, Faun. Chil. i. p. 444.

*Bernicla chiloënsis*, Phil. & Landb. Wiegmann. Archiv, 1863, i. p. 195.

This beautiful Goose, the smallest of the species of our country, was described by Des Murs as the female of the *Anser inornatus* of King, which is, according to the very probable opinion of Philippi and Landbeck, the young of *Bernicla magellanica*. Therefore they describe the species under the new name of *Bernicla chiloënsis*. But this name is by no means suitable, because the bird lives also on the continent, and its range extends over the whole of Patagonia, where it is one of the most common Geese. Both sexes are nearly alike in colour and form, but the colours of the males are more brilliant, and the females have numerous black semicircles on the reddish brown colour of the neck, breast, and commencement of the back. The full description of this bird given in Wiegmann's 'Archiv' renders unnecessary any further remarks from me. I received the bird from El Carmen, Bahia Blanca, in the vicinity of Dolores, where it is not uncommon in the winter.



## 5. CAIRINA.

10. CAIRINA MOSCHATA auct. ; Burm. Reise, ii. p. 514.

This large Duck I have seen only in the eastern and north-eastern parts of the country, near the river Paraná, and occasionally from Santa Fé and Tucuman ; but I have never observed it in the market at Buenos Ayres.

## 6. DENDROCYGNA.

11. DENDROCYGNA FULVA auct. ; Burm. Reise, ii. p. 514.

12. DENDROCYGNA VIDUATA auct. ; Burm.
- l. c.*
- p. 515.

The range of both these birds extends from Buenos Ayres to Tucuman ; and they are nowhere rare. They are often brought to the market of Buenos Ayres, where I have frequently purchased specimens of them.

The second species is generally seen in the evening ; and during the night its presence may be known by its peculiar cry while flying over the town in flocks, bird answering to bird, as has been already observed by Azara. This species is very common in Paraguay.

## 7. DAFILA.

13. DAFILA BAHAMENSIS (Linn.) ; Burm. Reise, ii. p. 515.

In the same parts of this country, but not in the west. Often brought to the market in Buenos Ayres.

14. DAFILA SPINICAUDA (Vieill.) ; Burm. Reise, ii. p. 515 ;
- 
- Slater, Proc. Zool. Soc. 1870, p. 666, pl. 38.

This Duck is found in all the southern provinces from Buenos Ayres to Mendoza, and is nowhere rare. As the eastern and western specimens are somewhat different in colour, I thought of separating the species into two, giving to the western form the name of *oxyura*, after Meyen (*cf.* Reise, ii. p. 515) ; but as I have not any western individuals for examination, I will accept the opinion of Messrs. Slater and Salvin (Proc. Zool. Soc. 1869, p. 157) that these two may be the same. The specimens from Buenos Ayres are somewhat more brilliant in colour, the edges of the tectrices and secondary remiges being yellow, and not white. This Duck was first described by Azara, who obtained specimens at Buenos Ayres (Apunt. iii. p. 421), and then by Meyen, who brought it from Chili, where the bird is very common.

## 8. QUERQUEDULA.

14. QUERQUEDULA FLAVIROSTRIS (Vieill.) ; Burm. Reise, ii. p. 516.

15. QUERQUEDULA MACULIROSTRIS (Licht.) ; Burm.
- ibid.*

16. QUERQUEDULA TORQUATA (Vieill.) ; Azara, Apunt. iii. p. 452 (male) ;
- ibid.*
- no. 352, "
- Anas leucophrys*
- , Vieill." (female).

This beautiful Duck is rare, and was brought to me by a friend



some years ago, in both sexes, from the river Paraná, near Las Conchas, where the bird lives on the numerous branches of the river between the flat islands of its delta. I have never seen specimens in the market of Buenos Ayres, where the two former species of *Querquedula* are very common, and sold every day during winter (from May to September). Azara has described the sexes of *Q. torquata* under different names.

17. *QUERQUEDULA BRASILIENSIS* (Briss.); Burm. Reise, ii. p. 517.

*Ipecutiri*, Azara, Apunt. iii. p. 445.

Also a very common bird in the interior and northern provinces. Very numerous near Paraná, Santa Fé, and Tucuman, and likewise in Paraguay and the Brazils, where this Duck is the most common of all species. Rare in the vicinity of Buenos Ayres, and very seldom brought into the market.

#### 9. PTEROCYANEA.

18. *PTEROCYANEA CYANOPTERA* (Vieill.); Burm. Reise, ii. p. 514.

Common in all the southern provinces from Buenos Ayres to Mendoza; rarer in the north to Paraná and Santa Fé.

#### 10. SPATULA.

19. *SPATULA PLATALEA* (Vieill.); Burm. *l. c.* p. 517.

Only in the eastern districts of the Republic, from Buenos Ayres on the river to Paraguay, and not rare.

#### 11. MARECA.

20. *MARECA CHILOËNSIS*, King; Burm. Reise, ii. p. 517.

On both sides of the Cordilleras, and likewise common on the plains of Patagonia and up to Buenos Ayres, where this Duck may be found every day in the market during the winter. Not abundant in the northern parts of the country, and never seen on the river near Paraná.

#### 12. METOPIANA.

21. *METOPIANA PEPOSACA* (Vieill.); Burm. Reise, ii. p. 518; Sclater, Proc. Zool. Soc. 1870, p. 666, pl. 37.

Likewise on both sides of the Cordilleras, and common throughout the whole of Patagonia to Buenos Ayres and Paraná, where I have observed this Duck in large flocks on the lagunes near the river.

#### 13. ERISMATURA.

22. *ERISMATURA FERRUGINEA*, Eyton; Gay, Fauna Chilena, i. p. 458.

*Anas cyanorhyncha*, Licht. MS.



This peculiar Duck is also found on both sides of the Cordilleras, and may be seen occasionally in the vicinity of Buenos Ayres, where it has been sometimes killed by my hunter on the Laguna Matanza at the south of the town. As the description given in Gay's work is very short and only applicable to the summer dress, I think it desirable to give a full description of this bird.

The old male in the summer dress is of a dark reddish-brown colour on the back, and somewhat paler, mixed with grey, on the underside. The head and neck are black, or dark blackish grey, mixed in the younger plumage with reddish margins to the feathers. The wings are dark blackish brown with reddish spots; and the tail is clear yellowish grey; the bill is blue-grey; and the legs are black, the iris dark red.

The bill, somewhat shorter than the skull, is very high at the base, and the plumes on the front are somewhat produced; the flat apical half is nearly of the same breadth, and the hook on the tip very small. The small nostrils are placed in the middle, where the flat apical half begins. The much narrower under mandible is whitish, the upper mandible bluish grey, the hook reddish brown.

The plumage is of a silky appearance, very soft and very compact; all the feathers are broad and rounded, and those on the back very large. The wings are short, and do not reach the tail; and the first of the pointed remiges is the longest. The most peculiar organ is the tail, which consists of eighteen small very rigid feathers, successively longer from the outer to the middle, the first on each side being shorter than the half of the middle. This strong rigid tail is therefore cuneated, and passes directly out of the plumage without any tail-coverts, as is the case with many other Ducks, the plumage being transversely cut off at the commencement of the tail on the upper and under parts. The feet are large; and the outer toe is twice as long as the tarsus, which is much compressed; the hinder toe is small and furnished with a small membrane; the whole foot is black.

My specimens measure 14 inches long; the bill  $1\frac{1}{2}$  inch, the closed wings (which have a small tubercle on the hand-joint) 5 inches, the tail 3 inches, the tarsus  $1\frac{1}{4}$  inch, and the outer phalanx  $2\frac{1}{2}$  inches.

The young male and the female are of entirely different colours, but have the same style of plumage. The whole of the upperside of the body is a blackish brown, each feather having a thin yellowish grey or whitish margin, which gives to the back an undulating appearance. The underside is whitish grey, or yellowish grey, with faint blackish-grey undulations, as every feather is dark grey, but with a large white or yellowish edging, which covers the whole of the grey part so that only here and there the grey ground is to be seen.

The wings and tail are blackish grey, the latter being somewhat yellowish. A small whitish or yellowish stripe commences at the sides of the bill, and passes under the eye to the neck, terminating over the ear. This white stripe is lighter in the female; the whole of the throat and sides of the neck are of the same colour, so that



the plumage on the fore side of the neck forms a darker ring under the white throat, between it and the whitish breast.

All this whitish colour is more yellowish in the winter dress of the males, and nearly of a reddish colour on the breast.

This bird is very ready to dive, and disappears under water immediately on observing the sportsman, not coming again to the surface until some distance off. Owing to the shortness of its wings its power of flight is very limited; so it adopts the expedient of diving in order to escape its foes.

All the before-mentioned species of *Anatidæ* are represented in the Public Museum of Buenos Ayres.

Azara describes two other species, which I have seen, but which have not yet reached my hands.

These are:—

23. *ANAS BICOLOR*, Vieill.; Azara, Apunt. iii. p. 443. no. 436.

*Anas collaris*, Merrem, Ersch. & Gruber, Encycl. tom. xxxv. p. 25.

This beautiful Duck I occasionally saw on the lagunes near my quinta, when I lived at Paraná in the year 1859; but I never had my gun, and so could not obtain it. It is not common, and lives in society with the other species there.

24. *ANAS MELANOCEPHALA*, Vieill.; Azara, Apunt. iii. p. 447. no. 438.

*Anas atricapilla*, Merrem, *ibid.* p. 26; Gill, U. S. N. Astr. Exp. ii. p. 202, pl. 25.

Of this Duck I have only seen one specimen, which was in the possession of a private collector, who killed it on a lagune in the interior; it seems to be very rare in our province. The Prince Max. of Wied mentions it as having been obtained from Rio Grande do Sul in Brazil.

I will close this paper with some remarks on the number of the tail-feathers of the Ducks, which varies in different species. Possessing a large number of specimens of *Erismatura*, I was induced to give this genus a careful examination.

*Erismatura ferruginea* has eighteen rigid feathers, the largest number observed by myself in this group of birds; and a like number was found by Azara in *Cairina moschata*.

The long-tailed group of *Dafila* (*D. bahamensis* and *D. spinicauda*) have sixteen tail-feathers, and Azara mentions that he found the same number in *Anas bicolor*, as also in *Spatula platalea*.

In *Sarcidiornis regia*, *Metopiana peposaca*, *Mareca chiloënsis*, *Dafila viduata*, *D. fulva*, *Querquedula brasiliensis*, and *Q. maculirostris* I have found fourteen tail-feathers.

Most of the smaller species have twelve tail-feathers: such are *Pterocyanea cyanoptera* and the other *Querquedula*. Azara also gives the same number for *Anas melanocephala*.



## 5. On some Persian, Himalayan, and other Reptiles.

By J. ANDERSON, M.D.

[Received January 15, 1872.]

## CYCLEMYS OLDHAMI, Gray.

I have received a living specimen of this species and a perfect shell from Samagooting in the Naga Hills in Assam, to the east of the Brahmaputra. The first measures  $8\frac{1}{2}$  inches in length, and the second is 9 inches long.

In both specimens the pectoro-abdominal sternal suture is unanchylosed, conferring, in the living example, distinct mobility on the lobes of the sternum, but of a more limited character than in *Cuora*. In the dried shell the mobility between the pectoral and abdominal plates is at once demonstrated when the sternum is moistened in water, while before doing so there is but little evident motion.

In the living specimen the portion of the abdominal plates overlapping the line of the pectoro-abdominal joint has become more or less fractured, so to speak, evidently by the motion of the two halves of the sternum on each other; and in the other specimen (the shell) the fracturing is complete, and the suture or joint is carried through the plates from side to side. Anterior to the outer third of the suture on each side is the separated portion of the abdominal plates, broken up in an area, which Theobald has compared to a curious fossa. Behind the middle third of the joint is a separated piece of each pectoral. The more perfectly fractured character of the portions of the abdominal plates that overlap the suture in the larger, as compared with their imperfect fracturing in the smaller specimen, would seem to favour Theobald's observation that the suture of the lobes of the sternum becomes more developed with age. If Dr. Gray\*, who combats this observation of Theobald's on the ground that it is opposed to the experience of most zoologists and the specimens in museums, had been familiar with the animal in life, or had examined the moistened sternum of a museum specimen, it is questionable whether he would have committed himself to dispute the correctness of the observations of a zoologist who spoke from personal knowledge of the living animal, and who did not confine himself to a crude knowledge gained from museum specimens. In connexion with Theobald's observation it is curious to remark that Dr. Gray did not observe any mark of the transverse suture on the sternum in a specimen which he doubtfully regarded as a younger stage of *C. ovata*, as compared with another example of the same species, which he regarded as more aged than the former, on which the cross suture of the sternum was much eroded on the edge, doubtless in the same way as I have described in the present species; so that Dr. Gray's own facts substantiate Theobald's conclusion which he disputes.

\* Suppl. Cat. Shield Rept. B. M. 1870, p. 23.



The colour of the body of this species is pale yellowish, the limbs, tail, head, and neck being brownish, with a tinge of olive, the head being unspotted.

*CUORA AMBOINENSIS*, Daud.

I have received a male of this species from Samagooting in the Naga Hills, Assam, measuring 7" 8" in length.

VARANIDÆ.

*HYDROSAURUS SALVATOR*, Linn.

This fine Lizard is not uncommon in Assam.

LACERTIDÆ.

*LACERTA STRIGATA*, Eichw.

Scales oblong, granular, placed obliquely but keeled longitudinally. Ventral plates in six rows, the two central lines of scales each about half the size of the one external to it. Small scales along the external margin of the outer row, which has twenty-five scales from the axilla to the groin. Preanal plate large, semicircular, its hinder margin slightly convex; surrounded by two rows of small scales and a rudimentary third. Nineteen to twenty-two femoral pores. Verticils of tail regular, of moderate length and breadth, strongly keeled. Nostril in the lower hinder angle of the nasal, with the suture of the rostral and first labial touching the middle of its lower margin; two small shields behind it of nearly equal size, the lowest one triangular, with its apex directed downwards and backwards, the shield above it pentagonal. Frontal of moderate size, hexagonal. Postfrontals each about the same size as the frontal, heptagonal, forming a moderately broad suture with each other, but a very narrow one with the anterior loreal and parietal. Vertical broader anteriorly than posteriorly, rounded in front, obscurely pointed; lateral margins concave; posterior margins oblique, straight, or slightly concave. Parietals larger than the postfrontals, triangular, with the lateral margins convex. Postparietals smaller than parietals, pentagonal, forming a very small suture with the vertical, and one of nearly equal breadth with the parietals and preoccipitals. One or two small shields behind the postparietals, one before the other. Preoccipitals as large as the postfrontals, pentagonal, forming a broad suture with each other, the postparietals and exoccipitals and sutures of nearly equal width with the vertical and occipital, which are partially wedged in between them. Occipital rather smaller than the parietal, its anterior margins meeting at an obtuse angle, lateral margins convergent, posterior end truncated. A small truncated conical postoccipital. Exoccipitals larger than any of the other head-plates, hexagonal, external posterior angle rounded. Two temporals equalling the length of the exoccipitals, and forming a suture with them externally, the foremost one being longest and largest. An elongated supraaural placed obliquely



across the ear below the hindermost temporal. An area of moderately large and small scales between the temporals and upper labials. Two loreals, the posterior one the larger. Seven upper labials, the one below the eye much larger than the others. Seven lower labials. Ear vertically oval. Four pairs of large shields behind the mental, increasing in size from before backwards, the posterior pair about four times the size of the anterior one. The fold, from ear to ear, not prominent. The anterior margin of the neck-fold with ten large scales from side to side.

Dark greenish olive-brown above, with five fine bluish-green narrow longitudinal lines from the head to the base of the tail—one from the ear along the side, another from above the ear over the former, and the third from the postoccipital along the middle of the back: the area between these spotted with blackish. Under surface rich green, yellowish about the anal region.

Snout to vent  $2\frac{4}{12}$ "; vent to top of tail  $4\frac{9}{12}$ "; anterior extremity  $\frac{9}{12}$ ", posterior extremity  $1\frac{4}{12}$ "; fourth toe  $\frac{11}{24}$ "; snout to postoccipital (hinder margin)  $\frac{13}{24}$ ".

*Hab.* Shiraz, Persia. Two specimens.

Duméril and Bibron regard this species as a variety of *L. viridis*; and in this they are followed by Prof. Filippi, who records it from Tiflis and Senkoran.

EREMIAS CÆRULEO-OCCELLATA, D. & Bib. Erpét. Générale, vol. v. pp. 295, 296.

Head much pointed; tail not enlarged at the base, long and slender. Frontal obliquely quadrangular, with an angle in the middle line before and behind, entire or longitudinally divided in two, in contact with two of the nasals, anterior loreal, and the postfrontals. Postfrontals hexagonal, forming a small suture together. Vertical hastate, lateral margins concave, convergent. Preoccipitals triangular, with the small wedge-like occipital indenting their line of union posteriorly. Exoccipitals large, subquadrangular, narrower before than behind. A line of six small linear granuloid scales along their external margin. Temporal region granular. Two large supraciliaries, together forming an oval surface. Two or four small scales before them, with two rows of small granules along their external border. Anterior loreal small, quadrangular; posterior large, subquadrangular. Five linear scales along the ridge of the eye, the anterior one very much longer than the others. Two small scales above and behind the posterior angle of the eye. Ten to eleven upper labials, exclusive of the inferior orbital plate, which has two small rounded scales behind it; seven to eight lower labials; five pairs of large plates below them, the first and last pairs the smallest. A fold across the throat from ear to ear. The prevertical fold with three rather large scales in its middle. Dorsal scales minute, smooth, ovally rhomboidal, arranged more or less in transverse rows, with a minute granule between each scale posteriorly. Scales of tail oblong, arranged in verticils. Scales on front of thighs and under surface of tibial portion of leg very large, hexagonal. Either



two moderately-sized preanal plates surrounded by smaller ones, or all the plates of one size and not enlarged. Femoral pores thirteen or fourteen. Ventrals square, smooth, arranged in transverse rows of fourteen to eighteen scales.

Yellowish olive-brown above, with a series of yellowish or bluish spots, with black margins along each side of the body, spotted with black on the sides of the head, neck, and body, and on the dorsal surface. Under surface yellowish white. Tail uniform dark olive-green above and below in its two posterior thirds.

Tip of snout to vent  $2\frac{4}{12}$ " ; vent to tip of tail  $3\frac{6}{12}$ " ; snout to occiput  $\frac{7}{12}$ " ; anterior limb  $\frac{10}{12}$ " ; posterior limb  $1\frac{1}{12}$ " ; fourth toe from base of fifth  $\frac{7}{12}$ ".

*Hab.* Yarkand.

I have followed Duméril and Bibron in regarding this as distinct from Pallas's *Lacerta velox* and Eichwald's *L. argulus*, for the reasons stated by the former authors. The specimens before me, although they were obtained in Yarkand, agree in every particular with the description of the types which came from the Crimea. Filippi records *E. variabilis*, Fitz., as a common species throughout Persia.

#### OPHIOPS ELEGANS, Ménétriés.

Frontal quadrangular, its anterior margins slightly convex or straight, meeting at an obtuse angle with the two upper shields of the rostral, which separate it from the rostral. Postfrontals either transversely triangular, with the base directed outwards, or more or less pentagonal; in the former case they may or may not form a suture behind the rostral, while in the latter instance they invariably meet in the middle line, and the suture is usually broad. Vertical moderately elongate, rounded and rather pointed anteriorly, its margins meeting at an obtuse angle; lateral margins concave, convergent; posterior extremity either slightly pointed or transversely truncated. Preoccipitals small, pentagonal, forming a broad suture together with a small lozenge-shaped occipital behind them, with a tumid disk in its centre. A small triangular postoccipital. Two large pentagonal exoccipitals with two linear temporals along the outer margins, the posterior one the smallest; temporal region below them granular. Three supraciliaries, the anterior one very small, two posterior ones very large, together forming an oval, with a flattened external margin, along which there is a longitudinal row of fine granules. Two or three small almost granular plates behind the posterior one. Two loreals, one before the other, and contributing to form the "canthus rostralis;" the anterior one small. Eight upper labials, the fifth forming the lower support to the eye, with three small labials behind it; one small shield above the last, separating it from the supraaural plate, which is either oval or oblong. Eight lower labials with six large shields below, three on each side, the fourth pair from the mental the largest. The throat in some (twelve) specimens shows a distinct fold from ear to ear, while it cannot be detected in others. Scales from occiput to



nearly in a line with the axilla almost granular and small-keeled. Those on the remainder of the body rhomboidal, strongly keeled, imbricate, with their tips directed backwards and upwards. The scales on the middle line of the back larger than the others, and the scales generally on the lumbar and mesial regions larger than any of the rest, and very strongly keeled. Scales on tail arranged in verticils, very strongly keeled above; those on the anterior fifth of the under surface smooth, on the remaining four fifths keeled. Eight rows of ventral scales, with a lateral line of small scales intermediate between them and the dorsal scales. The two rows in the middle of the belly are smaller than the row on either side of them. A large transversely elongated preanal shield, with a smaller one before it, surrounded by some still smaller shields.

*Colour* olive, brownish- or even reddish-bronze above, with two whitish longitudinal lines along each side, the lower one proceeding from the snout along the lower margin of the eye over the shoulder to the groin; the other and higher, through the "canthus rostralis," over the eye and along the side of the back. Below the former, the sides of mouth (labials), neck, and sides of body are spotted with blackish. The area between these two white lines is either reddish brown, or reddish and black-spotted; and there is a band of black spots along their upper margins. A short narrow dark-brown line from the occiput on to the neck. A few black spots on the side of base of tail. Limbs dark brown or black-spotted above, most markedly on the hinder limbs. Under surface yellowish.

Snout to vent  $2''$ ; vent to tip of tail  $3\frac{9}{12}''$ ; snout to occiput  $\frac{11}{2}''$ ; anterior extremity  $\frac{8}{24}''$ ; posterior limb  $1\frac{3}{12}''$ ; fourth toe from base of fifth  $\frac{7}{12}''$ .

*Hab.* Shiraz, Persia.

This species is evidently very variable, both in colouring and in some of the details, chiefly affecting its posterior frontals, which are sometimes quite separate from each other, while in other examples they are contiguous, and form a broad suture. The specimen figured by Duméril and Bibron shows the latter peculiarity. It is evidently, from its peculiar coloration, a form inhabiting a dry country; and it is probably an inhabitant of arid hill-sides, where its colour will hardly be distinguishable from the soil. Ménétrés's specimens were from Bahun; it has also been obtained at Smyrna, and Eichwald includes it in his fauna.

Four species of this genus have been recognized in India:—*O. jerdoni*, Blyth; *O. theobaldi*, Jerdon; *O. beddomii*, Jerdon; *O. microlepis*, Blanford.

*EUMECES LADACENSIS*, Gthr. Rept. of Brit. Ind. p. 88.


This species has a transparent eyelid, no supranasals, four supraciliary shields, thirty-eight rows of scales round the body, and fifty-six to fifty-eight transverse series between the fore and hind limbs. The opening of the ear is denticulated, and the subcaudals are broad, and there are two enlarged preanal scales. Günther states that



the fore limb, when laid forward, reaches to the snout; but in the specimen before me it reaches only to the anterior angle of the eye. The tail of my individual is forked at its middle, the two prongs being of nearly equal length, the longest measuring  $1\frac{4}{12}$ " and the undivided portion of the tail  $\frac{9}{12}$ ", total length of the organ being  $2\frac{1}{12}$ ". The body is  $1\frac{11}{12}$ " long, the head  $\frac{5}{12}$ ", the fore limb  $\frac{8}{12}$ ", and the posterior limb  $\frac{10}{12}$ " in length.

*Hab.* Ladak.

Fig. 1.



*Anguis orientalis.*

ANGUIS ORIENTALIS, n. sp. (Fig. 1.)

Rostral triangular. Two supranasals on each side enclosing an azygos quadrangular shield in contact with the rostral in front, and the internasal and postfrontals behind. Internasal triangular, pointed in front, with the fronto-nasals on each side of it. Fronto-nasal pentagonal or elongately oval and pointed at each end, lying between the posterior supranasal, internasal, and frontal, in contact behind with two shields, one of the side of the head, and one of a longitudinal line of eight small shields between the supraciliaries and the eye. Frontal rather large, triangular, broadly truncated in front, with a wavy posterior margin. Three postfrontals in a transverse row, more or less quadrangular. Six rather large supraciliaries; two small scales behind the eye. Vertical very large, broader in front than behind, lateral margins divergent, and then passing inwards at an obtuse angle to the posterior margin, which is transversely truncated. Two pairs of quadrangular parietals behind the supraciliaries, the external shield the largest, the inner one touching by its internal margins the vertical and parietal. The latter is a long wedge-like pentagonal shield truncated anteriorly, but with its lateral margins convergent to a point behind. A pair of large oblong obliquely placed exoccipitals, with a triangular posteriorly pointed small shield between their posterior margins. Two rows of temporals, three on either side, but each external to the parietals and exoccipitals. A single row of small shields from the fourth lower labial round anteriorly to the corresponding labial of the opposite side. Two rows of small shields from and below the fourth to the last lower labial. A large azygos chin-shield behind the infralabial line of scales, with four large pairs posterior to it,



the shields of the first pair forming a broad suture, the others separated by a number of small scales. Thirty rows of scales around the body, one inch behind the head. Two longitudinal rows of hexagonal scales, with their long diameter set nearly transversely to the body; the rows immediately external to them and on the sides slightly smaller and rhomboidal, and distributed in oblique rows. Ventral scales hexagonal, larger than those on the vertebral line. Vent with four pairs of anal plates, the central pair the largest. The tail contracts rather suddenly a little beyond its middle; and the remaining portion is slight, turned up, and covered above and on its sides with large brown scales. Scales on under surface of tail larger than on ventral surface of body, with rounded rather pointed margins.

Brown above, a dark brown band along the vertebral line of scales, and another one of the same hue along the fourth line of scales external to the former, the four longitudinal lines below shaded with dark brown, which fades away on the sides. General hue of under surface and lower half of sides olive-yellowish. The scales of the side and under surface have each a dark brown centre, with a broad brownish-yellowish or greenish-yellowish margin; and the predominance of the two last-mentioned colours on the sides and under surface determines the general tints of these parts, the dark centres, however, of the scales being quite distinct.

Length: snout to vent  $4\frac{9}{12}$ ", vent to tip of  $\frac{9}{12}$ ", snout to occiput  $\frac{9}{12}$ ".

*Hab.* Rehst, on the Caspian Sea.

This species is distinguished from the *A. fragilis*, Linn., by the greater number of shields on its head and scales round its body.

#### SEPIDÆ.

##### GONGYLUS OCELLATUS, Wagler.

Head much pointed; rostral rounded in front, concave posteriorly. Two supranasals, separated by a minute prefrontal. Post-nasal pentagonal, in contact with the first and second labials. Two large loreals, the anterior one about thrice as large as the posterior shield, behind which there are three scale-like preoculars, one behind the smallest loreal, and two in a line above it. Frontal large and broad, in contact with the prefrontal, supranasal, vertical, anterior loreal, and anterior supraocular, which rests on the two uppermost preoculars. Vertical large and elongate, exceeding the distance between its anterior extremity and the tip of the rostral. A small azygos shield wedged into its posterior extremity. One pair of occipitals about the same size as the vertical. Mental not quite so large as the shield behind it. Eyelid with a longitudinally elongate transparent disk. Ear round, of moderate size, not denticulated. Scales in thirty-two rows round the body.

Upper surface (in spirits) of body and tail brown, every alternate line of scales marked by a transverse black band, with a longitudinally elongated bluish-white spot in the centre of each scale.



The bands are lost on the head, where their place is taken by black spots with whitish centres. Some of the bands run into each other, so that the transverse arrangement is not perfect. Labials with black margins, and sides of neck and body black, reticulately spotted on a yellow ground.

Under surface immaculate.

*Hab.* Bushire, Persia.


This Lizard has been figured no less than three times—first by its discoverer, Olivier, afterwards by Geoffroy, and lastly by Gervais in the ‘Dictionnaire Universelle d’Histoire Naturelle;’ and it is doubtful whether Geoffroy’s figure excels the first, which certainly the last-mentioned does not.

#### GECKOTIDÆ.

##### HEMIDACTYLUS PERSICUS, n. sp. (Fig. 2.)

Back covered with numerous white, rather large, trihedral tubercles, with blackish-brown ones intermixed; nearly all the tubercles about half the size of the opening of the ear, which is longitudinally crescentic, the concavity being directed forwards. There are no tu-

Fig. 2.



*Hemidactylus persicus.*

bercles on the side of the neck; and those on the nape are less than half the size of those on the loins, where they have a tendency, as in the body generally, to be arranged in longitudinal rows, fourteen such lines occurring before the loins; all the tubercles are minutely striated in a radiate manner from their heads. A patch of large rounded granules between the nostril and eye, and another behind the nostril. The ventral scales are small, and forty-five to fifty rows



occur in the middle of the belly; those on the throat are very minute. Ten upper, and eight lower labials. A pair of large chin-shields behind the azygos lower labial, and forming a pretty broad suture behind the latter, succeeded by another pair, the shields of which are widely apart, their anterior extremities only touching the first pair of labials. Two or three lines of largish scales below the remaining lower labials. Tail with regular transverse rows on its upper surface of rather pointed trihedral tubercles, seven in each row. A small preanal region of enlarged scales in the female.

Colour pale yellowish brown, with six faint brownish transverse narrow dorsal bands, the tubercles in these areas being almost black; a darkish brown streak from the nostrils through the eye, above the ear, with a whitish line above it. Lips whitish.

From snout to vent  $2\frac{5}{8}$ ". Tail imperfect.

*Hab.* Persia.

This species appears to be closely allied to *H. trihedrus*, from which it may be distinguished, however, by its smaller ventral scales and tubercles, and by its peculiar coloration.

*PENTADACTYLUS KHASIENSIS*, Jerdon; Proc. As. Soc. Bengal, 1870, p. 75.

*Gymnodactylus khasiensis*, Anders. Proc. Zool. Soc. 1871, p. 162.

I have received seventeen specimens of this species from Cherra Punji in excellent condition. The former examples, which made me first acquainted with the species, had become hardened by too strong spirit, and had the close hood so contracted that they had the facies of a *Gymnodactylus*. The recent specimens, however, clearly show that Dr. Jerdon was quite correct in referring them to *Pentadactylus*. The largest specimen measures from the snout to the vent  $3'' 2'''$ , vent to tip of tail  $3'' 8'''$ . It appears to be a common species at Cherra Punji.

*EUBLEPHARIS MACULARIUS*, Blyth, Journ. As. Soc. Beng. vol. xxiii. pp. 737, 738.

*Eublepharis macularius*, Theobald, Cat. Rept. As. Soc. Mus. 1868, p. 32; Jerdon, Proc. As. Soc. 1870, p. 75; H. T. Blanford, Journ. As. Soc. vol. xxxix. p. 338.

Body covered with moderate-sized, conical, backwardly pointed tubercles, separated from each other by numerous small flat rounded granules, an arrangement that prevails all over the body as far forwards as on a line with the posterior angles of the eye; before this, the granules narrow in size; and anterior to the front angle of the eye the upper surface of the snout is covered with pointed tubercles, smaller than those of the rest of the body and arranged in a tessellated manner. Nostril oval, in a single plate over the first labial, with a large shield at its anterior superior margin between its plate and the rostral. Ear large and crescentic, the straight anterior margin with a few pointed tubercles. Tubercles on the loins and thighs larger than on the rest of the body. Three shields of different sizes be-



hind the rostral; ten upper and thirteen lower labials. Belly covered with slightly imbricate rhombic scales, increasing in size from before backwards. Tail thick, conical, and verticillated, covered below with quadrangular, rhombic, or almost rounded, slightly imbricate, moderate-sized scales; the upper surface with smaller granular scales, with a median lateral row of rounded or pointed enlarged tubercles. The verticils are defined below by a transverse line of fine granules. Preanal pores in an angular line of thirteen. Eighty-six small cylindrical conically pointed teeth in the upper jaw, and eighty in the lower jaw.

Olive-brown above, covered with numerous deep-black spots, most frequent about the occiput and nape. A black spot from the nostril to the eye. Under surface yellowish.

Length: snout to vent 4" 10"', vent to tip of tail 3" 3"', head 1" 2"', fore limb 4" 6"', hind limb 2".

*Hab.* Salt range, Punjab; and Amritzur.

This species is closely allied to *E. hardwickii*, from which it is distinguished by the tubercles being separated from each other by flat rounded granules instead of smaller tubercles, and by the arrangement of the tubercles on the head, which in *E. hardwickii* present a tessellated appearance all over the head from snout to occiput; while in *E. macularius* the tubercles are arranged on the head as on the body, as far forwards as the posterior angle of the eye, and anterior to that they gradually assume the paved arrangement. The coloration, too, of these species is markedly distinct; and *E. macularius* is essentially a north-western species, while *E. hardwickii* is confined (so far as is known) to the eastern side of India.

Blyth's type, now before me, is a half-grown individual, and is so bleached that it shows no trace of colouring beyond a uniform whitish hue. His description of the species is almost exclusively directed to the colouring and to a speculation as to the probable colour of the young. He mentions that Mr. Theobald informed him that the species attains to more than double the size ( $3\frac{1}{8}$  inches) of the specimen from which he drew up his notes, and that it is remarkable, when alive, for the beauty of its prevailing rosy caraneous hue.

Dr. Jerdon's \* *E. fasciatus*, from Hurriana, appears to me to belong to this species; for he describes the tubercles as "*larger and finer*" than in this species, and less close and narrower than in *E. hardwickii*, which is exactly the character which distinguishes *E. macularius* from the last-mentioned species. The coloration of his single very young and imperfect specimen was exactly that of the young of *E. macularius* as described by Blyth.


Although the difference of colour between the young and adult is very marked, as will be gathered from my description and the account given of the coloration by Blyth and Jerdon (who both describe the young as beautifully banded on the body, with a nuchal and three dorsal white bands), still a careful comparison of

\* Proc. As. Soc. Bengal, 1870, p. 75.



my adult with the young type does not reveal any structural character that would entitle me to separate them.

Fig. 3.

*Cyrtodactylus yarkandensis.*

**CYRTODACTYLUS YARKANDENSIS, n. sp. (Fig. 3.)**

Upper surface uniformly smoothly granular, some of the granules much larger than the others, especially on the hinder part of the body, none tubercular. Two pointed tubercles on the under surface of the tail at the side of the vent posteriorly; three large tubercles on the side of the tail at the base, the external large and pointed. (Tail absent in both specimens.) Ventral scales small, and those on the throat granular. Nostril over the suture of the first labial and rostral defined in front by the rostral, below by the first labial, and behind and above by a semicircle of two or three shields, the upper one of which is the largest. Ten to twelve upper labials, the first sometimes confluent with the rostral; nine lower labials. Two pairs of chin-shields, the anterior the largest, with the rostral wedged between its plates. (Females, no femoral pores.) Ear small, obliquely oval from above downwards and backwards. Toes moderately long, slender.

Bluish grey, with seven broad blackish waved bands, with a well-marked black posterior margin.

Length: snout to vent 2", vent to tip of tail — — — ?, snout to occiput  $\frac{6}{12}$ ", length of fore limb  $\frac{9}{12}$ ", of hind limb  $\frac{11}{12}$ ", fourth toe from base of fifth  $\frac{4}{12}$ ", fifth toe  $\frac{3}{12}$ ".

This species is from Yarkand; but I can give no information regarding the elevation at which it was found.

**AGAMIDÆ.**

**CALOTES VERSICOLOR, Daud.**

I have received seven fine adults of this species from Nasrick with a much more elevated crest than occurs in specimens found in the eastern side of India. They are all coloured in the same way: viz. the general colour is light, rather olive-yellow, the head and the anterior half of the trunk being suffused with pink, which is most intense on the crest and throat; the hinder half of the body is speckled with blackish, and the upper surface of the tail is marked with broad dark-coloured bars. The specimens are much larger



than any I have met with on this side of India, constituting a well-marked western race.

*CALOTES MYSTACEUS*, D. & B.

I have received twelve examples of this species from Cherra Punji.

*CALOTES MARIA*, Gray.

This is not an uncommon species in the Khasya Hills. The body of the largest specimen in the museum from that locality measures from snout to vent 5 inches; the tail is unfortunately broken; but in another individual measuring in body 4.8 inches, the tail is  $14\frac{1}{2}$  inches long. The elbow, knee, and heel are marked with white; and a white line runs along the outside of the fourth toe. The back in the largest specimen is ornamented by eleven transverse white bands intermixed with reddish; they extend only a short way on to the sides. They are continued on to the tail, where they ultimately resolve themselves into pairs of whitish spots, distributed at regular intervals.

*Hab.* Cherra Punji, Khasi Hills.

*CALOTES JERDONI*, Günther, Proc. Zool. Soc. 1870, p. 779, pl. xlv. fig. A.

I have received three specimens of this handsome species, obtained at Cherra Punji. A male and a female have a pale red band along the back and on to the base of the tail, and two rows of white spots at regular intervals on the sides between the fore and hind limbs. The other specimen is uniformly coloured green.

*CHARASIA DORSALIS*, Gray.

This species occurs as far east as the Rajmahal Hills; and Blandford has found it sixty miles west of Raipur, and, he believes, also on the Godavery.

In the young the scales of the under surface are all markedly keeled; but this character is soon all but lost, although traces of it may be detected in adult examples. The cross-rowed character of the scales of the tail is not so marked as on the body, but it can be distinctly detected, although they are imbricate in both localities.


*STELLIO PERSICUS*, n. sp. (Fig. 4, p. 383.)

The middle line of the back from between the shoulders covered with eight to nine longitudinal rows of very small, rounded, keeled scales of different sizes, those along the vertebral line very small. All the surface external to this, including the sides, very finely granular, with interspersed large granules, chiefly in the axillar region, no scattered spines on the sides. There is a square area on the middle of the sides, covered with enlarged keeled scales of different sizes, arranged more or less in irregularly transverse series. The largest scales are roundly oval, with a rather strong tubercular-like



keel in the centres, whilst the smaller ones, which are the most numerous, are also rounded, but more conical or tubercular than keeled. All the surfaces of the neck very finely tubercular, with numerous little patches of rounded, enlarged, and, in some cases, spiny tubercles, which occur also above and further back than the shoulder.

Fig. 4.

*Stellio persicus.*

Large strongly keeled scales on the upper surface of both limbs. A few large spined scales along the posterior upper margin of the thigh. Toes long and tapering (5'' 5'''), clawed, slightly compressed, with transverse plates on the under surface, with a strong spine on the angle formed by the lateral and anterior margin. Throat more coarsely granular than the neck; the rest of the under surface covered with very small rhomboidal, slightly imbricate, smooth scales, in transverse rows, as many as fifty in a row, across the middle of the belly. No femoral or preanal pores. Tail slightly swollen at the base, surrounded with regular verticils of strongly keeled scales. Head rather broad; snout short and moderately pointed. Loreal region concave. Nostril round, in a triangular nasal below the canthus rostralis, nearer the end of the snout than the eye, and separated from the upper labials by three longitudinal rows of scales. Three scales between the nasal and rostral. A line of large, more or less linear, keeled, tubercular scales from below the anterior angle of the eye, under the eye to over the ear, where they form a group of large tubercular-like plates. Area in front of and below the ear very finely granular, with large spiny shields at the anterior and inferior margins of the ear. Upper surface of the head covered with numerous small shields, a group of prominent ones in the middle of the frontal region, and another and more extended series on the occiput. All the scales and shields on the snout and sides of the head have numerous dark-brown dots on their margins. Upper labials 12, low and long; 13 lower labials, which are higher and shorter than the upper ones. Several rows of small shields parallel to the lower labials. Mental shield with a small azygos shield behind it, and an enlarged shield on each of its sides, and in contact with the front labial, with a line of five to six enlarged scales behind each. Skin of neck loose, thrown into numerous folds; a strong fold from below the ear to the shoulder, and another below it again; small folds on the nape; two transverse folds on the under surface



of the neck, enclosing short longitudinal folds. The prominent portions of these folds bear patches of more or less spiny tubercles.

Olive above, marbled on the sides and back by about eight narrow, transverse, black bands, which scarcely meet in the vertebral line, and enclosing pale whitish round spots. Tail pale yellowish olive above, with narrow dark olive-brown bands that do not extend to the under surface. Throat yellow, finely marbled with narrow reticulate dark-olive lines. Under surface of belly, limbs, and tail yellowish.

Length: snout to vent  $1\frac{1}{4}2''$ , vent to tip of tail  $2\frac{5}{8}7''$ , snout to occiput  $6''$ , fore limb  $1''1''$ , hind limb  $1''8''$ .

*Hab.* Teheran, Persia.

A closely allied species to this is the *S. himalayensis*, Steind., from which the present specimen is distinguished by the abrupt separation of its enlarged dorsal scales from the lateral granules, which in *S. himalayensis*, now before me, pass gradually one into the other—and by the presence of the tubercular keeled area on its sides, which does not exist in the allied species. There are other points of difference; but these are among the most prominent, combined with the difference in coloration.

#### AGAMA AGILIS, Olivier.

I can add nothing to the characteristic description given of this Lizard by Duméril and Bibron, except as far as regards a point in coloration. In the four specimens before me, of all ages, not only the throat but the belly has longitudinal dark lines, which, however, are more indistinct on the latter than on the former. In one adult, the greater part of the throat, and the sides, and under surface of the neck and the sides of the belly are deep purplish black. All have a deep spot of a similar colour on the side of the neck before the shoulder.

*Hab.* Shiraz.

Length: snout to vent  $3''6''$ , vent to tip of tail  $4''8''$ , head  $11''$ , fore limb  $1''11''$ , hind limb  $2''6''$ .

This Lizard is recorded by Blyth from the Salt range of the Punjab, but not included by Günther in his 'Reptiles of British India.' Filippi describes a new species, *A. lessonæ*, from Ispahan.

#### TRAPELUS RUDERATUS, Olivier.

Scales of different sizes, more or less feebly keeled, with numerous large, rather erect, spiny, tubercular scales scattered over the body, either singly or in groups, more numerous in the males than in the females; some of them, instead of being spinose, are only thickened posteriorly; they pass on to the root of the tail, but not beyond it. Tail a little less than twice the length of the body and head, covered with rhomboidal keeled scales, dilated at its base in the male, and tapering. A fold across the neck, another from the lower angle of the jaw, over the shoulder. A central group of large tubercular furrowed shields on the frontal region, surrounded by slightly smaller ones of the same character; a group of similar shields on the occiput,



and enlarged tubercular, almost spiny, scales on the parotoid region, and numerous spiny tubercles on the nape of the neck. On the sides of the neck the ordinary scales are small and almost granular. A group of large, flattened, furrowed shields on the temporal region, extending forwards to the posterior angle of the eye. Ear oval, its upper margin guarded by 3 to 4 dependent spines, with some tubercles on either side of them; 28 to 32 upper labials; 26 to 29 lower shields, excluding the rostral and mental. Ventral scales smooth, but terminating in a very minute but spinose extremity, about 16 oblique rows between the fore limbs. A transverse row of 14 preanal pores, separated from the margin of the cloaca by about a similar number of pores in a broken line of much narrower and more pointed scales, and with another line of 8 larger pores anterior to the former. No pores in the female. No trace of a crest on the back or neck. Limbs rather slender, digits strong, and armed with long sharp claws, the dorsal ridge of which is black, and the sides horn-yellow. The palms, soles, and under surface of fingers and toes are strongly keeled, each scale terminating in an apical spine.

The colour varies considerably, doubtless either due to the animal having the power to change the hues of its skin, or, it may be, depending on sexual causes. An adult gravid female is a rich reddish buff tinged with greenish on the head, especially on the parotoids, with six transverse bands of oblong pink spots, situated in groups of large spiny tubercles. A very obscure greenish-buff line along the vertebral ridge, terminating on the base of the tail in a bright yellow line, which runs along its middle to its posterior third, with a series of large reddish-brown spots along each side of it, becoming fainter posteriorly. A narrow, wavy yellow line along the back of the thigh and along the side of the base of the tail, broadly but interruptedly margined with blackish. Under surface uniform yellow. The male has the same tint as the female, only more yellow; while the other is slightly darker, with an olive hue. In the former, the red spots are the same, but they are separated on the vertebral line by a stripe of elongated bright yellow spots, which is prolonged on to the base of the tail. The yellow line on the back of the thigh is present, but it is crossed by an intensely black spot. In the latter specimen the vertebral spots have the same character and distribution as in the former, but the transverse bands of red spots are darker red; the line on the back of the thigh is very markedly white, with its dark margins very brightly developed. The upper dark margin is continued on to the tibial portion of the leg, which, along with the outside of the foot, is more or less banded with dark olive-brown. The two lateral spots on the side of the base of the tail are reddish or even yellowish, with dark margins. In both the males the axillæ are metallic pink. In one male the whole of the ventral scales, nearly as far forwards as the axilla, are of a darker and duller yellow than the general colour of the underparts, but they are not perceptibly tumid.



Length: snout to vent 2" 4"', 2" 2"', 2" 7"'; vent to tip of tail 3" 7"', 3" 10"', 3" 7"'; head 9"', 7"', 9"'; fore limb 1" 4"', 1" 3"', 1" 5"'; hind limb 1" 10"', 1" 9"', 1" 11"/>.

Dentition, upper-jaw, m.  $13 + 13 = 26$ , can.  $1 + 1 = 2$ , inc.  $2 + 2 = 4$ .

,, lower-jaw, m.  $12 + 12 = 24$ , can.  $1 + 1 = 2$ , inc.  $1 + 1 = 2$ .

The external incisor on either side of the upper jaw is much larger than the middle pair, but considerably shorter than the canines, with which they can hardly be classed, although Duméril and Bibron speak of four canines and two incisors in the upper jaw.

*Hab.* Teheran, Persia.

Duméril and Bibron state that the tail of the *Agama mutabilis* is three times the length of the body, which is certainly not the case with the *Trapelus* figured by Olivier, to which the above-mentioned specimens appear to belong.

#### PHRYNOCEPHALUS OLIVIERI, D. & B.

The five specimens of the peculiar-looking Lizard which I refer to this species, agree with the original description in all its important characters. Duméril and Bibron state, however, "le dessus des membres est recouvert d'écailles carénées," a character which is not applicable to the limbs of my specimens, inasmuch as only one individual shows a few faintly keeled scales on the tibial portion of its leg. I observe, however, that those specimens which have lost the hard epithelial covering of the scales and become shrivelled have an appearance that might be taken for keeling. It seems probable that Duméril and Bibron's specimens may have been in this condition; for in describing the tail, they state that it is "semée de petites verrues," a description applicable to my specimens which have lost their skin, but not to the perfect ones, which have the base of the tail with a few scattered spiny tubercles. These authors describe the scales of the tail as faintly keeled, a character which the specimens before me have as well.

The number of the labials is very variable; in one I count 30 upper and 30 lower plates, whilst in others the numbers mentioned by Duméril and Bibron, 27 and 26, prevail. The extent of the variation in the upper lip is from 27 to 30, and in the lower from 22 to 30. From an examination of a large series of Lizards, both of this and the Geckoid types, I am convinced that the number of labials is not a reliable specific character, although it may be useful as a generic one in a few instances.

There is another point in connexion with this Lizard worthy of notice, and not referred to by original describers, viz. that there is a rounded eminence on each side of the neck, of larger granules than those surrounding it. It is persistent in all my specimens. Those naturalists have also pointed out that what at first sight might strike an observer as an enlargement and flattening of the root of the tail is a character which depends not so much on the tail itself, but on an enlargement of the body before the vent—which, however, to me appears to swell somewhat behind the vent, and then to contract



rather suddenly to the tail proper, which tapers to a very fine point. The tail in all the specimens is nearly the length of the head longer than the body, and appears from its character to be more or less prehensile, although the Museum collector informs me that he found them on a grassy hill-side.

Snout to vent 1" 10''; vent to tip of tail 2" 5''; snout to occiput 6''; neck (above) 3''; anterior extremity 1''; posterior extremity 1" 7''.

*Hab.* Shiraz, Persia.

Prof. Filippi (*l. c.*) records that he obtained many examples of this species.

*PHRYNOCEPHALUS CAUDIVOLVULUS*, Pallas; *Gthr. Rept. of India*, p. 161.

Scales granular on body and tail; those on the lower halves of the limbs slightly larger. Ventral scales scarcely larger than dorsal ones. Nasals separated by an azygos scale and a longitudinal pair on each side of it. Two large scales before the anterior angle of the eye, followed by a row of seven smaller superciliaries. A large rounded scale behind the posterior angle of the orbit, removed from the superciliaries. Three or four large median plates above the nasals, the vertex being occupied by plates larger than those of the supraorbital region, but smaller than those on the occiput. All of those scales or plates are smooth. A rather large plate on the vertex between the posterior angles of the eye, with a tumid centre. Upper labials 27 to 29, the one nearest the corner of the mouth the largest. Lower labials 25. Twelve to thirteen teeth in the upper jaw, the one posterior to the first two incisors elongated like a canine. Twelve teeth in the lower jaw, the one posterior to the incisor of its side canine-like. Ear hidden; a fold below the neck and over the shoulder. No anal or femoral pores. Tail slightly dilated at its base, and flattened and rather thick towards its extremity; not prehensile.

Darker or lighter olive-grey above, either with some of the granules paler than the others or all uniform. Dark forms reticulated or spotted with blackish; light forms feebly spotted yellowish and blackish, some with rather large reddish spots, with black borders on either side of the mesial line, most distinctly marked on lumbar region and base of tail. Lips more or less barred or spotted with blackish brown and pale yellowish. Under surface either uniform yellowish greyish white with a slight median blackish band, or with a broad black longitudinal area involving the whole of the throat, the centre of the chest, and expanding on the belly, but not extending as far back as the groin. Under surface of the tail yellowish or greyish white, its sides spotted with brownish almost to the degree of being annulated, the posterior fourth deep black, most distinctly seen on the under surface.

The peculiar coloration of this Lizard would lead me to believe that the black coloration of the belly is either sexual or seasonal.




Vent to snout 2"; vent to tip of tail 2" 6"; snout to occiput 6"; neck 2"; anterior extremity 1"; posterior limb 1" 7".

*Hab.* Yarkand.

I have received ten specimens of this species from the foregoing locality; and the number would indicate that it is a common species in the high region of Yarkand and Tibet. Mr. Theobald obtained this Lizard on the shores of the Chomorreri Lake. He describes the female as smaller than the male, and the species as monogamous in its habits, a pair occupying a burrow a few inches deep in the sandy soil, the opening of which is often concealed by a stone or tuft of grass.

Fig. 5.



*Phrynocephalus persicus.*

**PHRYNOCEPHALUS PERSICUS**, Filippi, Viaggio in Persia, 1862, pp. 343, 344. (Fig. 5.)

Nostril plates anterior, but separated by five rows of scales. Body covered with small imbricate smooth scales, their tips directed backwards and slightly upwards; sides granular or nearly so; interspersed oval, rather pointed tubercles, either single or in groups of three or five; when more than two or three, one of the tubercles is larger than the others; they are largest on the back and root of the tail, but are not continued on to the tail beyond its basal fourth. Tail with rather small smooth imbricate scales. Two rather flattened tubercular scales above the anterior angle of the eye. Upper eyelid with a fringe of twelve rather oblong scales; lower eyelid with ten larger pointed scales. A group of large tubercles above the region of the hidden ear, and continued forwards to below the middle of the eye. Twenty-four upper and twenty-eight lower labials. Nape with a short obscure crest of five small dark brown rounded spines. A fold across the neck to behind the angle of the jaw (more or less distinct on the nape), whence another is prolonged over the shoulder, marked by two groups of moderate-sized pointed tubercles. Scales on the upper surface of the hinder extremity, below the knee, slightly keeled. Scales of ventral aspect smooth. Rest of tail flattened and laterally much dilated, covered with spiny tubercular scales, rapidly contracting at the anterior



fourth and subconic. Scales of toes serrated; claws pale yellowish, long and pointed. Uniform olive-grey in spirit, almost pale yellow on the hands and feet. Sides of body and upper surface of extremities and of the tail with a few scattered minute blackish spots. A very obscure yellowish spot with a dark blue margin on each side of the neck. Sides of the tail with rather large brownish spots; under surface whitish. The chin and throat obscurely black-spotted. Labials with black spots.

Length from snout to vent 2"; vent to tip of tail 2" 1"; snout to occiput 5"; neck (above) 2"; anterior extremity 11"; posterior extremity 1" 6".


*Hab.* Awada, seven days north of Shiraz, Persia.

Prof. Filippi, who apparently describes the colours from a pale specimen, gives the ground-colour as an earthy, somewhat reddish grey; and the under surface he mentions as a dirty white with somewhat of a rose colour, the throat with vermicular lines forming an azure-grey marbling. He describes the side of the neck with a large indigo ash-coloured spot surrounded by a delicate rusty-coloured band, which the action of alcohol causes quickly to disappear. Some brown angulated spots on the sides of the back are also described by its discoverer, and two similar ones at the base of the tail; after these occur other spots more numerous and more rounded; a few other transverse brown spots on the legs.

Prof. Filippi says that the above markings are quite constant, and that the species is profusely spread over the desert countries of Sultanieh and Teheran.

It is closely allied to *P. helioscopus*, Kaup, from which it is distinguished by the shorter and more rounded head, the long fringed scales of its lower eyelid, the five rows of scales between the nasals, the greater size of the tubercles over the auricular region, their anterior prolongation, along with the more dilated base of its tail.

Fig. 6.



*Phrynocephalus maculatus.*

**PHRYNOCEPHALUS MACULATUS, n. s.** (Fig. 6.)

Ear hidden. Scales small, smooth, directed backwards and upwards. A few lines of keeled scales on the lower arms and legs;



scales on the first fourth of the tail smooth, arranged in transverse rows, more or less keeled on the remaining three fourths, forming three longitudinal ridges on the under surface. Ventrals of moderate size, smooth. Head covered on the mesial line with flat, non-tuberculated scales larger than the others; a large oval one occupying the centre of the occiput, with a tumid centre. Nasals separated by three scales one above the other, and a lateral scale on either side of the inferior one. Upper labials 31, increasing in size towards the angle of the mouth, where the largest occurs. The two corresponding to the rostral are more transversely elongated than the others, which are immediately behind, and which are square, rather rounded at their free extremities. Lower labials 26; a longitudinal line of five enlarged scales behind the mental on each side, separated from the labials by a line of smaller scales. Upper eyelid with a fringe of ten oblong scales; lower eyelid margined with ten long, pointed scales. Upper jaw with 10 teeth on either side, the anterior two separated by a considerable interval; 20 in the lower. The teeth gradually decrease in size from behind forwards, showing no indication of differentiation into incisors or canines. The nails are yellowish, long, and pointed. The tail is long and prehensile, exceeding the length of the body by half. Pale yellowish brown; the head pencilled with dark brown and pale yellow; three short, dark-brown longitudinal lines on the nape. Labials minutely punctulated with brownish. The back and sides and upper surface of the limbs with moderate-sized dark-brown spots, and small brown dots of the same colour, with numerous yellow spots either involving one or three scales. Four brown spots, in transverse series, between the thighs. The base of the tail with two transverse brown bands, with a very faint one between them; the rest of the tail with blackish spots along its sides, almost forming rings, but interrupted on the upper surface by small yellow spots nearly constituting a longitudinal line. The last fourth entirely black, separated by a yellow band from a black ring anteriorly. The under surface on the anterior half is rich orange-yellow. Ventral surface generally yellowish white.

Vent to snout 1" 8"; vent to tip of tail 2" 11"; snout to occiput 5"; neck 3"; anterior extremity 1" 1"; posterior limb 1" 9".

*Hab.* Awada, Shiraz, Persia.

This species is distinguished from *P. caudivolvulus* by its smooth scales and peculiar coloration, and by the number of its teeth.

**PHRYNOCEPHALUS FORSYTHII, n. s. (Fig. 7.)**

Scales small, granular, ovally rhomboidal or quadrangular, smooth, arranged more or less in transverse rows, with a very few enlarged white scales occurring at intervals. No tubercles. Two enlarged scales on each side of the occiput, with a rather large tumid one in the mesial line before them, with those on the vertex only slightly larger than the ones external to them. Superciliary scales small. Scales of the lower eyelid forming a moderately pronounced fringe of ten scales with rounded points. Ear hidden. Nostrils anterior,



separated from each other by three to five longitudinal lines of scales, and widely so from the labial-like ventral by rows of granular-like scales. Twenty-six to twenty-eight upper labials, forming a toothed line. Twenty-six lower labials. A fold across the lower aspect of the neck, continued above the shoulder, and thrown into a number of folds at the side of the neck. Scales of limbs imbricate, almost

Fig. 7.

*Phrynocephalus forsythii.*

granular, partially keeled. Ventral scales of moderate size, some of them internal to the fore limb, showing a tendency to keeling. No femoral pores. Scales of dorsal surface of tail slightly larger than those on the body, and somewhat imbricate, and arranged transversely. Scales on under surface like those on abdomen, but increasing in size posteriorly, especially on the middle line. Base of tail dilated, with a group of spiny tubercles on the side of the most prominent portion. Tail long and tapering, but not prehensile. A tendency to the formation of a slight crest, by the skin falling into a longitudinal fold in the middle line of the back. Twelve teeth in the upper jaw on each side, with only one incisor, the tooth succeeding it being rather long and canine-like; twelve in the lower jaw, with the same characters and arrangement as in the upper. The last three teeth in both jaws are sensibly larger than the ones preceding them.

Snout to vent 1" 16''' ; vent to tip of tail 2" 6''' ; snout to occiput 5''' ; neck 2''' ; anterior limb 10''' ; posterior limb 1" 3'''.

Brownish yellow above, with five pairs of dark-brown spots on either side of the mesial line of the back, and a few more obscure ones on the base of the tail. Sides and upper surface of the body generally faintly spotted with brown and yellowish. Cheeks and labials punctulated with dark brown, varying in intensity in different



specimens. Sides and under surface of the tail with brown spots, the under surface of the posterior fourth blackish. Under surface of body yellowish, showing in some a faint blackish line down the centre of the chest and belly.

*Hab.* Yarkand. (Five specimens.)

#### OLIGODONTIDÆ.

*SIMOTES RUSSELLII*, Schleg.

Katmandoo, Nepaul. (Two specimens.)


#### COLUBRIDÆ.

*ABLABES FUSCUS*, Blyth.

This has all the characters of this species, but its postfrontal is divided. The anterior pair of chin-shields are twice the size of the posterior pair.

Katmandoo, Nepaul.

Fig. 8.



*Cyclophis persicus*.

#### *CYCLOPHIS PERSICUS*, n. s. (Fig. 8.)

Body slightly compressed, tail moderately long. Head distinct from the neck, rather broad across the occipital region. Eye of moderate size. Rostral broader than high. Anterior frontals rounded in front, broader than long. Ventral rather broad, with its lateral margins nearly parallel, its posterior margins slightly rounded. Occipitals large, divergent posteriorly. Nasal much elongated, abruptly truncated in front, and pointed posteriorly, with the nostril in its middle. One anterior and one posterior ocular, the former extending on to the surface of the head. Seven upper labials, the third and fourth entering the orbit, the seventh and eighth the largest. Temporals  $1 + 1 + 1 = 3$ , the last the longest. Two pairs of chin-shields; the first pair the largest, oblong, rather broad, and in contact with four labials; the posterior pair less than half the length of the former. Fifteen rows of smooth rhomboidal scales, with an apical groove. Ventrals 144, slightly keeled, and extending up the sides. Anal bifid. Subcaudals 77.

Pale olive-brownish, buff above, greenish yellow below. A large black spot on the surface of the head encircling the greater part of the postfrontals and the other shields behind them as far as the



posterior half of the occipitals, giving off a fine line to the nostril and a black ring round the eye. A buff band across the last half of the occipitals, followed by a broad black collar, which passes down on the sides, but does not meet below. Length of body 10" 3"', tail 3" 2"/>.

*Hab.* Bushire, Persia.

COMPSOSOMA HODGSONI, Gthr.

Katmandoo, Nepaul. (Three specimens.)

PTYAS MUCOSUS, Linn.

Katmandoo, Nepaul, and Cashmere. (Eight specimens.)

ZAMENIS PERSICUS, Jan, Iconograph. Gén. des Ophid. 23<sup>e</sup> livr. pl. 11. fig. 1.

*Zamenis ladacensis*, Anders. Journ. As. Soc. Beng. vol. xl. pp. 16, 17.

Since describing *L. ladacensis* I have received the part of Jan's 'Iconographie' containing the figure of *Z. persicus*, with which the former appears to be identical.

I have before me three specimens from Shiraz agreeing with *Z. persicus* in all their structural details, and only differing from Jan's figure in the absence of the black band between the eyes. The largest measures, body 30" 3"', tail 10" 7"/>.

ZAMENIS CAUDOLINEATUS, Gthr. *l. c.*

*Periops caudolineatus*, Gthr., Filippi, Viaggio in Persia, p. 355.

*Zamenis caudolineatus*, Gthr., Jan, Iconograph. Gén. des Ophid. 23<sup>e</sup> livr. pl. iii.

I have received three specimens of a snake from Shiraz and Iswhan, Persia, which I identify with this species, which has been figured by Prof. Jan. The largest individual measures 43" 3"', of which the tail forms 10" 8"". There are 21 rows of scales in all the specimens, each scale being terminated by two apical pores.

ZAMENIS CLIFFORDII, Schlegel.

Iswhan, fifteen days' journey north of Shiraz.

TROPIDONOTUS STOLATUS, Linn.

Katmandoo, Nepaul.

TROPIDONOTUS NATRIX, Linn.

Snout moderately pointed. Rostral twice as broad as high, hexagonal. Anterior frontals tapering, but transversely truncated anteriorly. Frontals bent well down on the side of the head, larger than the anterior frontals. Vertical moderately broad, with its lateral margin convergent, straight or more or less concave, most so in the young. Occipitals pointed posteriorly and divergent. Loreal quadrangular; one preocular, reaching to the upper surface of the head,



but not in contact with the vertical; three postoculars (one specimen out of six has four postoculars on one side); one large anterior temporal in contact with the two lower postoculars. A pair of posterior temporals touching the upper posterior angle of the last or seventh labial. Upper labials 7; sometimes 8, by the division of the second (this occurs in two specimens out of six). In the normal number, 7, the third and fourth enter the orbit, and in the latter case the fourth and fifth. The fifth labial is by far the largest. Nineteen rows of strongly keeled scales round the body. Scales elongate and leaf-like, and slightly notched at their free extremity. Ventrals vary from 177 to 180; subcaudals from 71 to 77.

Adults either uniform dark olive-black above, with indications of two longitudinal pale-coloured dorsal bands by some of the scales at regular intervals being more or less tinged with light brownish, or general colour dark olive-brown, with two longitudinal pale brown bands the breadth of two lines of scales. The bands and sides with rather obscure black spots, a scale's length, at regular intervals of from two to three rows, the ones on the sides alternating with those on the bands. Under surface yellowish anteriorly, with large bluish-black spots, which become larger and more numerous from before backwards, till at last the yellow is entirely replaced by the spots, the hinder part of the belly and the whole of the tail being deep black. In the young the confluence of the spots is not so perfect, and the ground-colour on the under part of the body is bluish grey. A transverse pale yellowish brown band from the angle of the mouth to the posterior extremities of the occipitals, very indistinct on the centre of the neck; a large black spot behind each, the small lateral spots being as it were a continuation of them. Upper labials yellowish, with black margins, those of the fourth, fifth, and sixth being very intense.

Largest specimen measures from snout to vent 38" 3", tail 10"; total 48" 3".

*Hab.* Rehst, on the Caspian Sea.

The Caspian specimens belong to a melanoid form of this species, probably the var. *atra* of Nordmann.

#### TROPIDONOTUS HYDRUS, Pallas.

I have received five specimens of this snake—one from Rehst, on the Caspian Sea, another from Shiraz, Persia, and three young specimens from Teheran. The first is not adult. It is dark-olive brown, covered at regular distances with black spots tending to form transverse bands. The last seventh of the belly and the whole under-surface of the tail are quite black. Its prefrontals are pointed; it has three anterior oculars and four postoculars; eight upper labials, the fourth only entering the orbit; and nineteen rows of strongly keeled scales. The Shiraz specimen is a light yellowish olive, with black spots, those on the back being placed obliquely; those on the side alternating with the former, and only distinguished by the black skin and edges of the bases of the scales. Two or three of the scales between each of these dark areas are more or less margined with



bright yellow, almost forming a series of yellow transverse bands on the side and between each of those above; one or two of the scales are longitudinally margined with a like colour. The under surface of the posterior five sixths of the body is much finely spotted or punctulated with black, and the last sixth is almost wholly black, as is the under surface of the tail. This specimen has nineteen rows of scales, pointed prefrontals, only two anterior oculars and three postoculars; but it is evident, from the way in which the superciliary shield is prolonged down behind the eye, that it is confluent with the fourth postocular; eight upper labials, the fourth entering the orbit.


This species is recorded by Prof. Filippi in his work on Persia.

#### DENDROPHIIDÆ.

##### GONYOSOMA DORSALE, n. sp. (Fig. 9.)

Rostral rather prolonged on to the top of throat. Head elongately oval and rather pointed. Anterior frontal quadrangular, rounded in front, nearly as large as the posterior frontals. Loreal subquadrangular. Two anterior oculars, the upper one large, and reaching the vertical; the inferior ocular evidently a separated portion of the fourth and fifth labials. The fourth labial also divided below the posterior half of the loreal. Fifth and sixth labials entering the orbit, the sixth prolonged backwards before the seventh labial, which is the largest. Two posterior oculars; vertical rather elongate. Lateral margins rather deeply concave.

Fig. 9.



*Gonyosoma dorsale.*

Occipitals rather large, and nearly oblong, abruptly transversely truncated behind. Temporals  $2+3+4=9$ . A line of elongated temporals along the occipitals, and another similar line along the labials. The two anterior shields of each of these lines in contact, the remainder of the lines enclosing three other shorter temporals. Scales smooth, long, lanceolate, with an apical groove, in nineteen rows. Ventrals keeled, prolonged up the side, 227; caudals 107. Anal bifid.

Pale yellowish green above, with a bright pink longitudinal band from the posterior extremity of the vertical on to the base of the tail, where it disappears. Under surface bright yellow. Area round the



eye yellowish, with a small black speck below the eye; a faint blackish green one behind the eye, and another at the angle of the mouth, continued more or less into each other. Length of body 23"; tail 8" 2".

*Hab.* Shiraz, Persia.

*CHERSYDRUS GRANULATUS*, Schneider; Gthr. *l. c.* p. 336.

A specimen of this snake, 36 inches in length, from the sands of the Puri beach. The smaller scales that cover the sides have thin keels, each distinctly terminating in a minute recurved spine. In the Indian Museum, Calcutta, there is another specimen of this snake, from the mouth of the Hughli, so that its distribution is from the southern to the northern extremity of the western side of the Bay of Bengal; and it is probable that it will be found to have a corresponding distribution on the eastern side of the same sea.

*HYDROPHIS CHLORIS*, Daud.; Gthr. *l. c.* p. 70.

The specimen referred to this species measures  $27\frac{5}{8}$  inches. The length of the long thin neck is more than one third of the total. There is one postocular; the third and fourth labials enter the orbit, the former being widely separated from the nasal by the preocular and rather large second labial. Two rather large, nearly equal-sized temporals. Two pairs of chin-shields, in contact with each other. Thirty-three rows of scales round the neck, very fully keeled, with a minute, nearly apical tubercle. Ventrals 495, distant on the neck, where they are about twice as large as the neighbouring scales; but they are relatively smaller on the thick part of the trunk. Four anal shields, the external scales being large. The dorsal half of the trunk is greenish olive, and the ventral half yellowish. Fifty-three blackish bands encircle the body, their dorsal halves being very black, broad above and narrowing to the middle line, and the ventral halves very faint on the thick portion of the body, but quite distinct along the ventral line, where they are connected with each other by an obscure black band. On the neck the transverse rings are very black, and so broad that the yellow interspaces are reduced to pairs of round spots, the head and under surface of the neck being quite black. Length  $27\frac{5}{8}$  inches.

*HYDROPHIS GRACILIS*, Shaw.

I have received a specimen from Puri, with all the characters of this species as diagnosed by Günther. It has twenty-one rows of bitubercular scales round the neck; 232 ventrals, those on the neck nearly twice as large as the adjoining scales, while those on the compound limb are divided into two halves, placed opposite each other, each half bearing two tubercles as the ordinary scales.

Length  $33\frac{2}{8}$  inches, of which the tail is  $2\frac{7}{8}$ .

*HYDROPHIS LINDSAYI*, Gray; Gthr. *l. c.* p. 371.

With the exception of having only twenty-six rows of scales round the neck, and 455 ventrals, very few of which are divided, a



*Hydrophis* which I have received from Puri differs in no other respect from *H. lindsayi*. It measures 27 inches in length.

*HYDROPHIS STRICTICOLLIS*, Gthr.

Head small, neck long and slender; scales not imbricate; thirty-seven round the neck, having either one or more tubercles in a straight line, tending to form an obscure keel; the central tubercle is the largest, the others very indistinct. On the posterior five sixths of the trunk there is one central tubercle to each scale. The ventrals are 385 in number, and on the anterior two thirds of the trunk are twice as broad as the adjoining scales. They show a distinct tendency to division, and are marked by a central groove, which nearly divides them; each half is marked by a central tubercle and a very obscure one behind it. There are six small anal scales, the middle one of the three of each side being the largest. Five upper labials, with a scale-like shield behind the last. The second upper labial is the largest, and touches the preocular. One postocular, with a single large temporal behind, which is as high as broad, rests on the scale-like posterior labial. Two pairs of chin-shields in contact with each other; fifty-one blackish rings round the trunk, broader than the groundwork between them, not confluent on the under surface, but running together on the back from behind the anterior third of the trunk. Tail with eleven vertical blackish bars, confluent on the under surface.

Length, snout to vent 36" 6"; vent to tip of tail 3".

*Hab.* Sandheads, mouth of the Hughli river; Bay of Bengal.

*HYDROPHIS SPIRALIS*, Shaw; Gthr. *l. c.* p. 366.

I have received what appear to be two young specimens of this species from Pooree, Cuttack, measuring respectively 22 and 16½ inches. The former has twenty-nine rows of scales round the neck. The scales are imbricate on the trunk, with a central tubercle. There are 332 ventrals, generally twice the size of the scales adjoining them; and almost all are undivided. The rostral, although it is well prolonged upwards, is broader than high. The third and fourth labials enter the orbit. There is only one postocular; and of the three temporals, the first is much the largest. Two pairs of chin-shields, the posterior being only slightly smaller than the anterior pair, and both are in contact. There is a black line from the rostral to the eye; but the rostral and the upper labial margin are yellowish white, and the whole under surface of the chin and throat are of the same colour. The upper surface of the head, from the frontals to behind the occipitals, is blackish, but more or less spotted with dirty yellowish, a spot occurring in the centre of the vertical and of each occipital. A faint, short, blackish line from the angle of the mouth either backwards or slightly upwards; fifty-nine black rings round the trunk, interrupted on the sides in the greater part of its extent, but perfect on the neck. These perfect or imperfect rings are broadest on the ventral surface, and on the neck they are connected with each other by a black line running along the ventrals.



Where the dorsal and ventral segments are not connected, the latter form rather broad, almost cone-shaped markings; the upper surface corresponding to the dorsal segments is dark olive, owing to the basal half of each yellow scale being black. The lower half of the length of the snake is rather bright yellow; after the twelfth ring from the head there is a round black spot on the dorsal surface between the rings, one or two blanks occurring only here and there. The latter half of the tail is black.

In the second specimen the shields of the head and scales are as those of the species; but I only count twenty-six rows of the latter round the neck, and 283 ventrals, which on the neck are rather more than twice as large as the neighbouring scales. The ventrals are undivided, and distinct throughout the whole length of the trunk. There is the black line from the rostral along the upper lip, the lower half of which is yellowish white, along with the chin, throat, and greater part of frontals; it is olive over the eye and temporals. From the posterior border of the frontals backwards to the hinder edge of the occiput, and including the two internal temporal shields, is black, with a faint yellowish spot in the centre of the vertical and each occipital. The trunk is encircled with forty perfect black rings, broadest on the back and ventral surface; on the latter region the rings are all connected with each other by a black longitudinal line twice the breadth of the ventral scales. One round, dorsal, black spot occurs between the second and third rings, but it is partially connected with the former. Between the sixth and seventh rings from the tail, another similar but separate spot occurs; and the coloration of the part of the body where it is placed represents exactly the coloration of the type of the species. The dorsal half of the body is coloured as in the former specimen; but its ventral half is not so light in colour—a circumstance which may be due to its greater youth.

*HYDROPHIS CANTORIS*, Gthr. *l. c.* p. 374.

I have received a young *Hydrophis* (also from Puri, and measuring  $20\frac{2}{8}$  inches in length) that fully agrees with Günther's diagnosis of the above species. The elongated neck is nearly one half of the total length of the snake; the rostral shield is much longer than broad, and produced backwards, its hinder margin being in a line with the posterior upper angle of the first labial. The third labial is in contact with the nasal. The third labial enters the orbit, while the fourth is subocular; the postocular and two large temporals on each side of the elongated occipitals. Two pairs of chin-shields in contact with each other. Twenty-six rows of scales round the neck; 443 ventrals, those on the neck twice as large as the adjoining scales, and those on the compressed part of the body divided into two, or distinct, each with two central tubercles. Six small anal shields. Head and belly entirely black; the body surrounded with fifty-three black transverse bands, nearly all of which are confluent with the black of the under surface, and many of them with each other on the back, so that the greenish yellow of the ground-colour forms large



oval lateral spots. The first fourteen black bands of the neck are not confluent above, but are separated from each other by light greenish-yellow bands half the width. On the posterior half of the thick part of the body the black bands extend only halfway down the side. The tail has eight black vertical bands; and the tip is black. The coloration is intermediate in this specimen to what Günther describes it in the adult and young.

*HYDROPHIS NIGRA*, n. sp.

Neck but moderately slender, less than two thirds the length of the body. Head broader than neck, but long and oblong, with nearly straight sides, the preorbital breadth equalling the temporal diameter. Snout moderately long, broad and rounded, and rather spatulate. Rostral much broader than high (only feebly notched on its inferior surface), its posterior extremity being in a line with the rostro-labial suture. Nasals broader posteriorly than they are long. The third labial is not in contact with the nasal, and it enters the orbit, from which, however, it is almost excluded by the fourth labial. One postocular. The fifth and sixth labials are divided transversely. Two temporals, of which the anterior is the largest. Two pairs of chin-shields, the anterior in contact, the shields of the posterior pair separated by an azygos scale. Thirty-two rows of scales round the body, slightly imbricate, and elongately hexagonal, smooth. Ventrals 248; the first twenty-five or so sometimes as large as the adjoining scales, which are rather small. They diminish in size as they are traced backwards, but nearly all remain distinct and undivided. Three pairs of anal shields, of which the outer are the largest. Uniform intense black, without any true markings. Length 19 inches, of which the tail is 2 inches; length from vent to occiput  $\frac{7}{1\frac{1}{2}}$ ; breadth at angle of mouth  $\frac{4}{1\frac{1}{2}}$ ; breadth before eyes  $\frac{4}{1\frac{1}{2}}$ ; breadth in a line with nasal suture  $\frac{3}{1\frac{1}{2}}$ ; snout to eye  $\frac{3}{1\frac{1}{2}}$ ; eye to angle of mouth  $2\frac{1}{2}$  lines; angle of jaws to tip of snout 8 lines.

*Hab.* Puri, Cuttack coast, India.

*HYDROPHIS CRASSICOLLIS*, Anders. Journ. As. Soc. Bengal, 1871, p. 19.

I have received another specimen of this snake from Puri. It measures  $57\frac{3}{8}$  inches, and is slightly lighter-coloured than my first specimen, and with the transverse bands much narrower. The upper surface of the head is coloured as in the type; but the chin and labials have a clear gamboge-yellow tint, and from the angle of the mouth there is a distinct similarly coloured yellow line passing backwards to the yellow longitudinal area of the side. The under-surface, except where the transverse black bands occur, is a clear, warm, yellowish white. I count 326 ventrals, and twenty-six rows of scales round the neck.

*HYDROPHIS STEWARTII*, n. sp.

The neck moderately long and slender, and the head rather short



and not much broader than the neck; the remainder of the body much compressed. Rostral considerably broader than high. The nasals as broad posteriorly as their common suture is long. The third and fourth labials enter the orbit, the former not being in contact with the nasal. Three temporals, the anterior being the largest. Two postoculars. Vertical much pointed behind. Occipitals long and narrow. Two almost quadrangular chin-shields in contact with each other. Thirty-three rows of scales round the neck. The scales hexagonal, not imbricate, with a feeble central tubercle. Ventrals 387, smooth, the first forty on the neck being about four times as large as the adjoining scales, those behind them becoming small and narrow as they are traced backwards, and hardly discernible on the last 6 inches of the trunk. Two pairs of small scale-like anals. Tail broad and markedly dilating from its root. Lips yellowish; upper surface of head, upper half of neck, and dorsal two fifths of compressed portion of body greenish olive; undersurface of head and all the remaining portion of the neck and body salmon-coloured. Fifty-seven very obscure darker olive, almost black rhomboidal bars on the dark dorsal area, and not extending on to the light-coloured sides. The tail pale greenish olive, mottled and tipped with black.

Length 38" 3", of which the tail constitutes 2" 8"; girth round neck 2 inches, behind head 2" 5"; greatest depth of body (5 inches before tail) 1" 8"; greatest thickness at last-mentioned point 6"; thickness at upper margin of lower third in same locality 3"; thickness at ventral margin 3" 2"; snout to occiput 11"; breadth across gape 7".

*Hab.* Puri, Cuttack.

*HYDROPHIS VIPERINA*, Schmidt; Gthr. *l. c.* p. 378.

I have received a specimen of this snake, also from Puri, with thirty-two rows of scales round the neck and with 268 ventrals. The first thirty-five or forty ventrals are six times as large as the adjoining scales. Beyond these the ventrals decrease in size, and ultimately are not much larger than ordinary scales. The third labial does not reach the orbit, below which are the fourth and fifth lip-shields. The latter labial and the sixth and seventh are transversely divided, as is also the first large temporal on the right side. The nasals are broader posteriorly than they are long; and the dimensions of the vertical in these directions hold a similar relation to each other. Two posterior temporals. Scales keeled. The upper surface of the head and of the body generally is dark olive, without any trace of spots or markings of any kind; and the sides, upper lips, and under surface are yellowish. Length  $29\frac{5}{8}$  inches, of which the tail is 3 inches; snout to occiput  $\frac{6}{8}$  inch; breadth across gape  $\frac{5}{8}$  inch; breadth in a line with posterior margin of nasals  $\frac{4}{8}$  inch.

I have received another specimen, which appears to be the young of this species. It measures  $13\frac{1}{2}$  inches, of which the tail forms



$1\frac{5}{8}$  inch. It has, however, thirty-two rows of scales round the neck, 240 ventrals, and three præanal shields on one side and two on the other. It has twenty-six black rhombic spots on the back, continued as vertical bars on to the sides, and confluent on the ventral shields, which are wholly black and six times as large as the neighbouring scales, which are smooth. It has on either side a minute detached præocular.

I have received from Muscat, Arab'a, another specimen of this species, measuring 27 inches, of which the tail forms  $2\frac{5}{8}$ , with 238 ventrals and twenty-six rows of keeled scales round the neck.

Two other specimens from Puri have been sent to this Museum since the above was written; and I am inclined to regard both as forms of *H. viperina*. They are young, and measure respectively  $13\frac{1}{8}$  inches, of which the tail forms  $\frac{1}{2}$  inch, and  $11\frac{6}{8}$ , of which the tail is  $1\frac{3}{8}$  inch. The first specimen has twenty-eight rows of scales round the neck and 267 ventral shields, while the second has twenty-nine cervical rows and 255 ventrals. Both have three pairs of anal shields; but as the condition of these shields in the above-mentioned young specimen would appear to indicate that these structures are variable in their number, too much importance cannot be attached to their numerical increase in the specimens under consideration as compared with the type. The third labial does not enter the orbit. On one side in one specimen there are two præoculars. The first temporal, on both sides in one specimen and on one side in another, is considerably higher than broad; while on the remaining side of the last and on both sides of the former a narrow temporal, longer than high, is separated, as it were, from the great anterior temporal.

These three young specimens were caught at one haul of a net, which would appear to indicate that the young of this species swim in shoals. When adult Hydrophides are caught in the net, there are, as a rule, always two or three of a species.

As one of these young snakes killed a chicken that it was made to bite, the species would appear to be endowed with an active poison.

*HYDROPHIS JERDONI*, Gray; Gthr. *l. c.* p. 362.

I have received a specimen from Puri measuring  $38\frac{6}{8}$  inches in length, the tail measuring 4 inches.

*ENHYDRINA VALACADYEN*, Boie.

One specimen, from Puri, with a divided postocular on one side.

#### CROTALIDÆ.

*TRIMERESURUS MONTICOLA*, Gthr.

Katmandoo, Nepaul.

*HALYS HIMALAYANUS*, Gthr.

I have received a Pit-viper from Goduk which may be provisionally referred to the above species. It has, however, its anterior frontals in a straight transverse line, as in *H. pallasii*, and only the

PROC. ZOOL. SOC.—1872, No. XXVI.



second large temporal confluent with the sixth labial. In these characters it is intermediate between the two species. I have another specimen before me with the divergent anterior frontals of *H. himalayanus*; but it has only the sixth labial confluent with the middle temporal. In the first-mentioned specimen the anterior are nearly as large as the posterior frontals; but in the second they are considerably smaller than the hinder frontals, which are more or less pointed anteriorly. If this individual had had its small anterior frontals in a straight transverse line, with only one temporal entering the lip, it might with equal propriety have been referred to one or other of the two species. The bearing of these variations is apparent. The colouring is as described in the 'Reptiles of British India.'

#### BATRACHIA SALIENTA.

##### *RANA ESCULENTA*, Linn.

Vomerine teeth in a transverse row between the inner nostrils. These specimens have the membrane reaching to the extremity of the outer margin of the first, second, and third toes and to the end of the inner margin of the fifth. The third and fifth toes are nearly equal, the last being, if any thing, longer than the first-mentioned; but to all practical purposes they may be regarded as equal. The fourth is one third longer than the third and fifth. The internal tubercle is oblong and laterally compressed, and of moderate size; the external tubercle round and obscure. The lateral glandular fold is never pale-coloured; and the dorsal white line occurs in four out of twelve specimens. The dark band along the canthus rostralis and over the tympanum can be faintly detected in a few. There are no true dorsal glandular folds; but the back is covered with rounded, not prominent, glandular spots.

General colour dark olive, profusely or sparsely covered with black spots; the limbs either banded or black-spotted. Some specimens have the under surface of the hind limbs with one or two black spots; while others have a much larger number, and the spots of the side extending on to the belly, while the whole of the under surface is more or less very obscurely apparently reticulated with blackish, but so faintly that it is hardly noticeable.

These specimens are larger than the European examples of the species, seven out of the twelve exceeding 3 inches, the largest measuring 4" 1" from the snout to the vent, and the hind limb 6" 7".

*Hab.* Shiraz, Persia.

##### *RANA CYANOPHLYCTIS*, Schneid.

*Hab.* Katmandoo, Nepaul.

##### *RANA GRACILIS*, Wiegman.

*Hab.* Katmandoo, Nepaul.

##### *BUFO VIRIDIS*, Laur.

I have received seven specimens of this species from Shiraz,



Persia. The largest is a very characteristically and highly marked specimen, measuring 3" 11" in length, and the hind limb 4" 3". The coloration is exactly that of Laurenti's figure\*. The others are young, and their colours are comparatively dull, the spots fewer and even proportionally smaller than in the adult.

The Museum collector also gathered, in the Himalayas, on his way to Ladak, seven specimens of a toad agreeing in every way with the young specimens from Shiraz. This species, however, had been previously obtained at Simla by Dr. Stoliczka.

*HYLORANA ERYTHRÆA*, Schl.; Gthr. *l. c.* p. 425.

I have received a young specimen of this frog from the Garo Hills, to the east of the Brahmaputra. The area between the two glandular lines is deep black; and there is a narrow black band along the dorsal margin of the uppermost glandular fold from above the eye to the side of the vent. The posterior surface of the fore and hind limbs marbled with deep brownish, a light line from the vent along the back of the thigh. In all its other characters it agrees with the characters given by Günther.

Length of body 1"  $\frac{1}{2}$ "; vent to heel 11  $\frac{1}{3}$ ".

This species appears to extend as far west as Eastern Bengal, the fauna of which is decidedly much more Malayan than Indian properly so called.

*POLYPEDATES ANNECTENS*, Jerdon, Proc. As. Soc. Beng. 1870, p. 83, 84; Anders. Proc. Zool. Soc. 1871, p. 210.

I have received twelve specimens of this handsome species from the southern slopes of the Shillong plateau of the Khasia hills.

The largest specimen measures as follows:—

Snout to vent 1" 7"; vent to tip of fourth toe 2" 6  $\frac{1}{4}$ "; vent to knee 8  $\frac{3}{4}$ "; knee to heel 8  $\frac{1}{2}$ "; heel to tip of fourth toe 1" 1"; length of fourth toe 8". These measurements of the hind limb show the difference of only  $\frac{1}{4}$  of a line between the dimensions of it and of the specimen formerly described by me.

*RHACOPHORUS MACULATUS*, Anders.

*Rhacophorus reinwardtii*, Jerdon, Proc. As. Soc. Beng. 1870, p. 84.

*Rhacophorus maculatus*, Anders. Journ. As. Soc. Bengal, vol. xl. p. 27.

This Museum has received seven other specimens of this frog from Shillong, on the Khasia hills. They in no way differ from the original specimens from which the species was described.

*HYLA ARBOREA*, Linn.

The specimens from the undermentioned locality have the dark lateral streak narrow and margined by a broader yellowish-white line. A dark transverse band over the vent, margined above with a pale band similar to the former. A pale dark streak from the knee

\* Syn. Rept. p. 111, pl i. fig. 1.



along the outer side of the leg to the fifth toe, margined posteriorly with pale yellowish. No spots. The membrane of the fingers very rudimentary; the toes two thirds webbed.

*Hab.* Rehst, Caspian Sea.

EPICRIUM GLUTINOSUM, Linn.

There is a median longitudinal fold from the symphysis of the lower jaw backwards as far as the length of the gape behind the angle of the mouth, abruptly defined posteriorly by a transverse fold separate and distinct from the body-folds but only reaching the sides. About halfway between it and the angle of the mouth there is another permanent fold that nearly encircles the body, but is interrupted behind the occiput. Anterior to this fold there is a short indistinct transverse fold on the throat, slightly posterior to the angle of the mouth.

The upper and under parts are not black, as described by Günther, but a rich slaty blue, the lateral line, however, being yellow. Whenever the specimens are removed from spirit they become dark brown, almost black.

I have received this species from Goalpara, Assam, and from Shillong, in the Khasia hills.

6. Catalogue of the Birds found in Ceylon; with some Remarks on their Habits and Local Distribution, and Descriptions of two New Species peculiar to the Island. By E. W. H. HOLDSWORTH, F.L.S., F.Z.S., &c.


[Received February 21, 1872.]

(Plates XVII.-XX.)

In the following Catalogue I have endeavoured to give a correct account of what is known at the present time of the birds resident in and visiting Ceylon. I have included no species about which there appears to be any doubt, except in a few cases; and in those cases I have mentioned the grounds on which their reported occurrence seems likely to be true.

The latest original list of Ceylon birds is that comprised in Mr. E. L. Layard's valuable and generally trustworthy "Notes on the Ornithology of Ceylon" published in the 'Annals and Magazine of Natural History' for 1853-54. Since that time there has been hardly any one in Ceylon who has given systematic attention to the avifauna of the island; and in the preparation of this Catalogue the considerable collection of birds made by myself and Mr. Layard's "Notes" have been the principal materials I have had at my command. I have been able, however, to make use of the extensive knowledge of eastern birds possessed by Lord Walden, the President of this Society, and his large collection of specimens, both of which





J. Smit lith

M&N Hanhart imp

1 ERYTHROSTERNA HYPERYTHRA ♂, 2 ♂ juv









J. Smit lith.

M&N Hanhart imp

BRACHYPTERYX PALLISERI.









J. Smith lith.

M. & N. Hanhart imp.

1. ARRENCIA BLIGHI ♂, 2 ♀ juv.









J. Smit lith.

M & N Hanhart imp

1. ZOSTEROPS PALPEBROSUS.  
2. .... CEYLONENSIS.







have been most kindly placed at my service; and they have been of great assistance to me in determining questions of species and nomenclature. I am glad also to acknowledge the help I have received from Mr. J. H. Gurney, Mr. Edmund Harting, Mr. Howard Saunders, and others, in the different groups of birds to which those gentlemen have given their special attention.

Mr. Layard's industry during his eight years' residence in Ceylon resulted in more than doubling the number of birds known in the island; and he did not leave much for his successors to find out in the low country, where he principally worked. His list included 315 species; but I have found it necessary to omit a few which appear to have been wrongly identified or to have been recorded by the late Dr. Kelaart on doubtful evidence. I have added 25 species which have been met with during the last few years, and among them two birds hitherto undescribed, making altogether 325 species which apparently have a good claim to be regarded as belonging to the avifauna of Ceylon.

The number of birds not hitherto recognized out of Ceylon is remarkably large considering its small extent of country (equal to only five sixths of the size of Ireland). A few species at one time thought to be peculiar to the island have since been recognized, or are believed to be found, in India or Malacca; but, so far as is known, the following 37 species are exclusively confined to Ceylon, and are pretty evenly divided between the low country and the hills, most of them, however, being found only in the southern half of the island. They form one ninth of the known Ceylon species.

#### BIRDS PECULIAR TO CEYLON.

<i>Athene castaneonota.</i>	<i>Geocichla layardi.</i>
<i>Tockus gingalensis.</i>	<i>Merula kinnisi.</i>
<i>Palæornis calthropæ.</i>	<i>Oreocincla spiloptera.</i>
<i>Loriculus indicus.</i>	<i>Alcippe nigrifrons.</i>
<i>Chrysocolaptes stricklandi.</i>	<i>Drymocapthus fuscocapillus.</i>
<i>Brachypternus ceylonus.</i>	<i>Pomatorhinus melanurus.</i>
<i>Megalaima zeylanica.</i>	<i>Layardia rufescens.</i>
— <i>flavifrons.</i>	<i>Garrulax cinereifrons.</i>
<i>Xantholæma rubricapilla.</i>	<i>Rubigula melanicteræ.</i>
<i>Centropus chlororhynchos.</i>	<i>Drymoipus validus.</i>
<i>Phœnicophaës pyrrhocephalus.</i>	<i>Zosterops ceylonensis.</i>
<i>Prionochilus vincens*.</i>	<i>Cissa ornata.</i>
<i>Eumyias sordida.</i>	<i>Temenuchus senex.</i>
<i>Erythrosterina hyperythra.</i>	<i>Eulabes ptilogenys.</i>
<i>Buchanga minor.</i>	<i>Munia kelaarti.</i>
— <i>leucopygialis.</i>	<i>Palumbus torringtoniæ.</i>
<i>Dissemurus lophorhinus.</i>	<i>Gallus stanleyi.</i>
<i>Brachypteryx palliseri.</i>	<i>Galloperdix bicalcarata.</i>
<i>Arrenga blighi.</i>	

\* Recently discovered by Mr. Legge, R.A. A notice of this species is given in a postscript to this Catalogue.



Excepting *Phaenicophaës* and *Prionochilus*, which are quite Malay forms, all these peculiar Ceylon species belong to genera found in India. Most of these genera range from India more or less to the countries east of it; and the nearest allies of *Cissa ornata* are almost confined to Eastern Asia.

I have not included *Batrachostomus moniliger* or *Kelaartia penicillata* among the species peculiar to Ceylon, as they are believed to be found also in South India; and I have likewise omitted *Malacocercus striatus*, as I much doubt its distinctness from *M. malabaricus*.

*Geographical distribution.*—In this Catalogue I have given approximately the geographical range of most of the species found in Ceylon, from which it will be observed that all those not peculiar to the island are, with very few exceptions, known in India; the majority of them extend to Burmah, many of them to some of the Malay islands and China, and a few to Australia. *Goisachius melanolophus* is a remarkable example of a common Malaccan species having four times been found in Ceylon, and, strangely enough, only on the west side of the island, although it has not yet been observed on the adjoining Indian continent. A converse example exists in *Hirundo hyperythra*, of which one or two specimens have been brought from Malacca, that species being otherwise considered quite peculiar to, as it is abundant in, Ceylon.

The Ceylon birds which range to the westward of India belong to species of generally wide distribution, and consist principally of Raptorial, Grallatorial, and Natatorial forms; the exceptions being examples of *Hirundo*, *Cypselus*, *Halcyon*, *Ceryle*, *Cuculus*, *Cisticola*, and *Pyrrhuloxia*.

Of the species which extend to Australia those belonging to *Calobates*, *Streptopelia*, and *Terekia* are of very wide distribution; the Ceylon species of *Haliaeetus*, *Excalfactoria*, *Charadrius*, *Ægialitis*, and *Mycteria* have a considerable range east and south-east of India; and *Attagen minor* and *Sterna gracilis* seem alone to be, so far as is known, especially Australian.

*Indian families absent from Ceylon.*—The *Vulturidæ*, *Eurylaimidæ*, *Pteroclidæ*, *Otididæ*, *Glareolidæ*, *Gruidæ*, and *Mergidæ*, all families included in the Indian avifauna, have no recognized representatives in Ceylon. Of the *Vulturidæ*, one species breeds so far south as the Neilgherries; but Ceylon agrees with the Indian archipelago and the countries south of continental Asia in having no Vulture. The *Eurylaimidæ* have their stronghold on the eastern side of the Bay of Bengal and in the Malay islands; and a representative of this family may yet be found in Ceylon. It is also not improbable that stragglers of the common South-Indian species of *Pteroclidæ* and *Otididæ* may one day be met with in the north of the island. *Glareola* may likewise be looked for; but the *Gruidæ* and *Mergidæ* are not likely to range so far south.

*Position and Character of the Island.*—Without entering into the question of whether Ceylon was originally a continuous portion of India or formed part of a lost Malay continent, as believed by the late Sir J. Emerson Tennent, it may be desirable to point out the



principal features of the country as it now stands. Its position (between  $6^{\circ}$  and  $10^{\circ}$  N. lat.) is almost equatorial. Practically it is an island about 35 miles (at its least distance on the extreme north) from India, increasing to nearly 60 miles at the connecting sandbank of Adam's Bridge, and to about 150 miles between Colombo and Cape Comorin. It possesses the character of a true oceanic island in having deep water (no bottom at 150 fathoms) within a very few miles of the land all round the coast, excepting only between Adam's Bridge and Point Pedro, the parts of the island nearest to India. The water shoals abruptly on the south side of Adam's Bridge, and has only a depth of a few fathoms north of it until it passes the line between the north point of Ceylon and the nearest part of India, whence it gradually deepens into the Bay of Bengal. Adam's Bridge, the narrow connecting-link between Ceylon and India, and said to be of comparative recent formation, consists of sandstone covered with loose sand, which is alternately beaten up on and removed from the north and south sides by the sea and wind during the successive north-east and south-west monsoons. It terminates on the Indian side in the island of Ramisseram, between which and the continent is the well-known Paumben Channel. On the Ceylon side the bridge ends in the island of Mannar, which is separated from the mainland by a considerable expanse of shallow water or mud banks, according to the state of the tide, with a narrow winding channel deep enough for the passage of small native vessels. The bridge itself has also several narrow openings or "scours" at different parts, so that, although Ceylon is virtually connected with India by means of Adam's Bridge, it may be regarded as practically distinct, and, as might be expected, it has species peculiar to itself in all the great divisions of the animal kingdom. Its length is 271 miles and its greatest breadth 137 miles.

For ornithological purposes Ceylon may be divided into two parts—the northern and southern halves, the northern portion being, with the exception of a few isolated hills, entirely low country; this is continued throughout the maritime districts of the south; and the whole coast is surrounded by a narrow belt of sandy beach. The low country generally is extensively laid out with paddy-fields; but there are large tracts in the northern half of the island which are still in the normal condition of forest, or, from the poverty of the soil or the scarcity of rain, are only occasionally cultivated, and support a scattered growth of bushy jungle rarely attaining the character of forest. This last was the nature of the country round Aripo, where I spent a good deal of time and obtained a great number of the commoner birds. At the north and on the north-east side there are large lagoons or backwaters, the resort of countless Waders; and there and on the inland lakes or tanks (as they are generally called) Ducks and Terns of various kinds are abundant in winter, and many other birds at all seasons. The avifauna of the northern half of the island is quite Indian in its character. The east and south-east parts also contain a good deal of wild country; they are thinly populated, and are visited the least by Europeans. One district is the home of the few remaining Veddahs, the supposed aborigines of Ceylon, who,



although almost savages compared with the rest of the natives, are said to retain the honourable distinction of high caste. The extreme south and south-west are generally well cultivated; and paddy-fields and cocoa-nut plantations are general in that part of the country.

The mountain-districts lie almost in the centre of the southern half of the island; and in this half, at various elevations ranging from sea-level to 8000 feet, are found by far the greater number of the peculiar Ceylon birds. A conspicuous feature of the Ceylon hills is the luxuriant vegetation which clothes them from their foot to the tops of the highest ranges; and although masses of rock may be seen here and there projecting from the mountain-sides, even these are largely covered with ferns and creeping plants. The mountain region may be divided ornithologically into the lower and upper hills. The country up to between 1500 and 1600 feet, the elevation of Kandy, is only partly cultivated; and its diversified character provides suitable habitats for a great variety of birds. This is particularly the case in the neighbourhood of Kandy, where there is some really wild jungle, in which some of the rarer hill species as well as low-country birds are found at certain seasons.

From the elevation of Kandy to about 5000 feet are the coffee-districts; and where this cultivation is general the number of birds is small, and they are found mostly at the higher and lower boundaries of the estates. If, however, the soil be unsuited for coffee and the jungle remain uncleared, birds are numerous, and many of the peculiar kinds, *Athene castaneonota*, *Palæornis calthropæ*, &c., may be met with. These lower hills are the great resort for the passerine immigrants; and birds of prey abound there. From 5000 to 8200 feet (the highest point in the island) constitutes what I shall have frequent occasion to speak of as the upper hills. They are almost entirely covered with tree jungle, with a dense undergrowth of "nilloo" (*Strobilanthes*), small straggling bamboo, tree ferns, and a variety of other plants. These hills are the great stronghold of the Sambur Deer; and Elephants and Leopards mount to their summits. Nuwara Eliya, the sanatorium of the island and a place where I have collected largely, is at an elevation of 6000 feet, and lies in a narrow plain, the houses being mostly scattered along the sides at the foot of the surrounding jungle-covered hills. The birds found in this locality and the neighbouring district are not numerous in species; but they are mostly of kinds peculiar to the island, and include *Chrysocolaptes stricklandi*, *Brachypteryx palliseri*, *Cissa ornata*, *Zosterops ceylonensis*, and several others, whose range does not generally extend far below the upper hills.

*Migratory Birds.*—The migration of birds within and into Ceylon is a subject about which there is still a great deal to be learnt; but, owing to the absence of observers, there is little reason to expect much trustworthy information will be gained for some time. The migrations take place at the changes of the monsoons. The S.W. monsoon blows steadily and for the most part strongly from April to October on the west side of the island. In October there is a lull for a few days between the two winds. It is the season for cyclones in the



Bay of Bengal; and then very often there is stormy weather on the Ceylon coast. At the first decided indication of the N.E. monsoon setting in, the true migratory birds begin to appear; they are generally first seen in the north and north-west of the island, and gradually extend over the western side and to the hills. At the same time there is a large influx of resident species to these parts of the country, which during this N.E. monsoon are less exposed to the violence of the wind than in the other season. There is no positive evidence whence these birds come; but I think there can be little doubt that it is from the eastern side and some parts of the central districts. Many of them are certainly found there during the S.W. monsoon; but no continuous observations have been made on the eastern side, and there is little known of what resident Ceylon species are to be met with there at any particular season. It being a great game country, Europeans who visit the eastern jungles devote their time more to sporting than to natural history. I may give an instance showing that there must be a good deal yet to be done in certain parts of the country. In February 1871 I obtained at Nuwara Eliya two specimens of a Flycatcher (*Erythrostera hyperythra*) of which the type specimen in the Berlin Museum, obtained in 1866, was the only one known; it came from the Ceylon hills; but that species is certainly not found in the hill districts during a great part of the year, and yet it has not been observed elsewhere\*. Towards the close of February the N.E. monsoon comes to an end, and is followed by five or six weeks of fine calm weather before the usually stormy burst of the S.W. monsoon. The migratory birds now take their departure, and many species resident in the island leave its western side. A Tern, however, in immature plumage and believed to be *Sterna gracilis* has only been observed on the Ceylon coast in summer; but as the Ceylon summer is at the same time as the Australian winter, the fact of this Tern being found at Colombo in July is an additional reason for believing it to be that Australian species. I have also only seen Frigate-birds during the summer; but Mr. Layard has recorded their occurrence in February.

With respect to the breeding-season for Ceylon birds it is difficult to fix any definite rule. The climate in the low country is always hot and damp, and birds of some species or other are nesting throughout the year. In many cases the breeding-time appears to depend on the monsoons; but I believe it often varies with the same species in different parts of the island. On the upper hills, where there is the nearest approach to a cold season of any part of Ceylon, and where the midday tropical heat is succeeded by cold nights and, in January and February, by severe frost, the breeding-season follows the rule in temperate climates and usually begins about April; in other parts of the country either nesting or moulting appears to be always going on.

From what I have said of the character of the country it will be evident that Ceylon possesses, in its swamps, jungles, forests, rivers, and coasts, the conditions suitable for the existence of a great variety

\* See No. 127, footnote.



of birds; the waters on the coast and inland swarm with fishes; the country is alive with insects and reptiles; and vegetation is most luxuriant in its growth; food of all kinds abounds; and there is no winter in the low country. It is no wonder, therefore, that species and individuals are numerous; but although I have, I believe, been able to add something to the good work done by Mr. Layard, the subject is yet far from being exhausted, and much remains to be done in examining the eastern side of the island generally, in discriminating many of the wading birds, and in working out the Terns and other birds found on the coast.

In this Catalogue I have followed Jerdon's arrangement of the species as given in his 'Birds of India,' and have adopted the names he uses, except in a few cases where older titles may be more properly employed.

1. *FALCO PEREGRINUS*, Gmelin.

Europe, Asia.

2. *FALCO PEREGRINATOR*, Sund.

Ceylon, India, W. Asia.

3. *HYPOTRIORCHIS CHICQUERA*, Daud.

Ceylon, India.

These species are recorded by Layard as found in Ceylon; but they have not come under my notice.

3 *bis*. *HYPOTRIORCHIS SEVERUS*, Horsf.

In a collection of birds sent home by Mr. S. Bligh, and consisting entirely of hill species shot by himself in one of the coffee-districts, is an undoubted specimen of the Indian Hobby, an unexpected addition to the list of Ceylon birds. It was killed whilst hawking after dragonflies.

Ceylon, N. India, Malacca, Java, Philippines.

4. *TINNUNCULUS ALAUDARIUS*, Gmel.

The Kestrel is widely distributed in Ceylon; I have seen it, however, most frequently in the northern part of the island, and a pair of these birds for many weeks frequented a small clump of cocoa-nut palms near my house at Aripo. I have also observed it at Nuwara Eliya during winter; and it is often met with in the coffee-districts. Although probably a migrant, it certainly spends several months in Ceylon.

Europe, Asia.

5. *ASTUR TRIVIRGATUS*, Temm.

This is a hill species, and not very uncommon. I have examined specimens of the bird in Ceylon, and have now before me a very good one killed by Mr. Forbes Laurie.

Mountains in Ceylon and India.



6. *MICRONISUS BADIUS*, Gmelin.

Said by Layard to be very common and widely distributed in Ceylon. I have identified several specimens of it.

Ceylon, India, Burmah, Malaya, Hainan.

7. *ACCIPITER VIRGATUS*, Temm.

I have a specimen of this Sparrow-Hawk from the lower hills. Layard does not mention this species; but it may possibly have been the one recorded by Kelaart as *A. nisus*, which I have no reason to think has been found in Ceylon.

Bill dark bluish; irides yellow; feet yellow.

Ceylon, India, Burmah, Malaya, Formosa.

8. *AQUILA PENNATA*, Gmelin.

Recorded by Layard.

Ceylon, India, W. Asia, N. Africa, S. Europe.

9. *NEOPUS MALAIENSIS*, Reinw.

Tolerably numerous in the hill country, and well known in the coffee-districts. I have seen several skins in different states of plumage, which were obtained from the hills around Kandy.

Ceylon, India, Burmah, Malaya.

10. *NISAËTUS BONELLI*, Temm.

Recorded by Layard as having been obtained by the late Dr. Templeton, R.A.

Ceylon, India.

11. *LIMNAËTUS CRISTATELLUS*, Temm.

This noble bird, mentioned by Layard under the name of *Spi-zaëtus limnaëtus*, Horsf., is well known in the hill country, and not unfrequently visits the poultry-yards of the coffee-planters. I have seen it on many occasions at Nuwara Eliya, and listened to its squealing cry as it soared in wide circles over the plain. In the beginning of 1871 I procured a fine living specimen, and shipped it at Colombo for the Society's Gardens; but it died soon after the vessel sailed. The feet and claws in this species are very powerful.

Mountainous parts of Ceylon and India.

12. *LIMNAËTUS NIPALENSIS*, Hodgson.

Recorded by Layard as having been procured by the late Dr. Kelaart on the hills, at an elevation of 4000 feet. It is remarkable that this species should be found in Ceylon, as in India it is only known from the northern hills; but Mr. Blyth tells me that he identified Dr. Kelaart's specimen, and has no doubt of its being the true *L. nipalensis*, Hodgs. No other example of this bird has been recognized in Ceylon.

Ceylon, N. Indian hills, Formosa, Japan.



13. *SPILORNIS BACHA*, Daudin.

Generally distributed over the island, frequenting trees on the margin of tanks and marshy places in the low country, and near open grass-land among forest-jungle on the hills. One specimen, which I shot near Aripo, disgorged a Tree-snake (*Passerita*) more than 3 feet long and nearly uninjured. Another, obtained at Nuwara Eliya, fell to the shot as if mortally wounded, although only slightly injured in one wing; it soon recovered, and became sufficiently tame to feed from my hand. I was fortunately enabled to bring the bird with me to England; and it is now alive in the Society's Gardens.

*S. spilogaster*, Blyth, from Ceylon, is now recognized as the immature condition of *S. bacha*; and there is no doubt that the *Hæmatornis cheela*, recorded by Layard as common in Ceylon, may also be referred to the same species.

Bill dusky; irides golden yellow; cere, legs, and feet dull yellow. Ceylon, S. India, Andamans, Malaya.

14. *PANDION HALIAËTUS*, Linn.

Rare in Ceylon, and I have only seen it on one occasion; it was perched on a buoy in Galle Harbour; and I was able to watch it from a short distance for a considerable time. Lord Walden has two specimens of it from Ceylon.

Europe, Asia, Africa.

15. *POLIOAËTUS ICHTHYAËTUS*, Horsf.

This Eagle I have only seen in the north of the island, where it is not uncommon near the coast. I shot an immature specimen at Aripo in November 1866. The irides were brown, but Jerdon states (App. B. of Ind. iii. p. 869) that in the adult bird they are pale yellow.

Bill black; "irides pale yellow;" feet yellowish white. Ceylon, India, Burmah, Malaya.

16. *HALIAËTUS (CUNCUMA) LEUCOGASTER*, Gmelin.

This is the common Sea-Eagle of Ceylon, and is probably found all round the island, although I do not remember having observed it at the extreme south. It may occasionally be seen soaring over Colombo Harbour and the adjoining lake; but further north, in the neighbourhood of Aripo and Mannar, several pairs of these noble birds may generally be found, each generally in its own district, and rarely wandering far away. In the strait separating the island of Mannar, at the east end of Adam's Bridge, from the mainland of Ceylon, the narrow channel for the passage of boats in the midst of the expanse of shallow water around is marked here and there with stakes; and on these may generally be seen perched one or two pairs of this Eagle, and sometimes a pair of *Polioaëtus ichthyaëtus*. As the receding tide lays bare the extensive banks of soft mud on each side the Eagles keep a sharp look-out for the crabs, which are abundant just at the edge of the water, and, pouncing on their prey, sail



off to some favourite tree, where the hard shell of the crab is broken up and the animal devoured. One of these stations, further down the coast, was on the cross-trees of a government flagstaff at Aripo; and the ground below was always littered with crab-shells and fish-bones, the remains of many a meal provided from the refuse of the fishermen's nets, which were hauled in on the beach close by. Sea-snakes (*Hydrophis*) are said to be a favourite food of this species; and these reptiles are abundant on the Pearl-Oyster banks nine or ten miles off the Aripo coast; but I have never observed the Eagles so far from the land.

A curious instance came to my knowledge of this bird having apparently thriven on most unnatural food. My friend, Dr. Boake, the late Principal of Queen's College, Colombo, once pointed out to me an example of this Eagle of full size, but in immature plumage. It had been recently brought to him by a native, who said he had reared the bird in his own hut. In answer to an inquiry as to what he had fed the bird on, he said "rice and curry." This is the universal food of the natives; and dogs and cats appear to thrive as well upon it; but that a Sea-Eagle should have been reared on such food seemed incredible. However, the matter was soon tested by a supply of rice and curry being given to the bird; and the statement of the native was quickly confirmed by the rapid disappearance of the whole of the food. The next day some fish was given, and the Eagle, once having tasted it, could never afterwards be induced to touch rice and curry.

In a male example of this Sea-Eagle which I shot at Aripo I found the liver of an enormous size, covering the whole of the pectoral and a great part of the abdominal regions.

Bill dusky blue; cere yellow; irides brown; feet yellowish white.

Ceylon, India, Burmah, Malaya, Australia.

#### 16 *bis*. BUTEO DESERTORUM, Daudin.

Lord Walden has received a single specimen of this Buzzard from Ceylon.

Ceylon, India, Persia, S. Europe, Africa.

#### 17. CIRCUS SWAINSONII, A. Smith.

Common in the Aripo district throughout the year; and I have frequently seen it at Nuwara Eliya in July and August. The pale rump of the brown birds attracts attention as they hunt backwards and forwards over the open country.

Bill black; irides yellow; feet yellow.

Asia, Africa.

#### 18. CIRCUS CINERACEUS, Montagu.

I have only identified this species on one occasion; it was killed near Colombo. Although probably not uncommon in Ceylon, it is certainly not so numerous there as the last species.

Bill black; irides yellow; feet yellow.

Europe, Asia, Africa.



19. *CIRCUS MELANOLEUCOS*, Forster.

This species was first described from, and has since been identified as a visitor to Ceylon; but I have never met with it. Layard procured it on the west coast.

Ceylon, India, Tientsin.

20. *CIRCUS ÆRUGINOSUS*, Linn.

This Harrier is probably only an occasional visitor to Ceylon. I observed a pair of these birds near Aripo in January 1870; and after several ineffectual attempts to get near them, I at last succeeded in shooting the female, a handsome specimen, with grey wings and tail. Layard does not appear to have met with this species; but it is included doubtfully in the list of birds in Tennent's 'Natural History of Ceylon.'

Bill black; irides and cere yellow; feet deep yellow.

Europe, Asia.

21. *HALIASTUR INDUS*, Bodd.

Common on the coast, especially on the northern half of the island. Specimens in various states of plumage were obtained at Aripo. I have also seen it at Colombo and Trincomalie.

Ceylon, India, Borneo.

22. *MILVUS GOVINDA*, Sykes.

This bird has very much the same habits and distribution in Ceylon as the last species. Neither of them, however, frequents the towns so much as they both do in India. In early morning at Aripo I have seen a flock of fifty or sixty Pariah Kites, in company with about a dozen of the other species, eagerly clutching at and feeding on the winged Termites which were rising in a cloud from an ant-hill not far from my house. The Crows were busily engaged on the same work, but kept at a respectful distance, apparently not liking to join in the general scramble going on among their more powerful neighbours, the Kites.

Ceylon, India, Burmah, Malaya, Andamans, China, Formosa, Hainan.

*Note.*—There are two, perhaps three, closely allied species of Kite found in India, the smallest of which, Mr. Gurney tells me, is identical with *M. affinis* of Australia; and there is some doubt as to which is best entitled to the specific name of *govinda*. As it is not quite clear to which of these the Ceylon birds belong, the above geographical range may not be strictly correct.

23. *PERNIS PTILORHYNCHUS*, Temm.

Given by Jerdon as *P. cristata*, Cuvier. I had an opportunity of seeing this bird alive in Ceylon; and Mr. Forbes Laurie has recently shown me a good specimen which he shot on the hills. Lord Walden has also received examples of it from Ceylon. Mr. Laurie's specimen agrees pretty closely in dimensions with those given by



Jerdon ; but Mr. Gurney tells me that the birds from Ceylon are usually larger than those from India. Although this bird is well known from Ceylon, it appears not to have been hitherto recorded from that island.

Ceylon, India, Burmah to Malaya.

24. *BAZA LOPHOTES*, Cuv.

Not very numerous, but has been found both in the low country and on the hills. I have seen specimens from the Kandy district.

Ceylon, India.

25. *ELANUS MELANOPTERUS*, Daud.

I have only seen specimens of this handsome bird from the hills, where locally it is not uncommon. Layard obtained it in the low country.

Ceylon, India, part of Africa, S. Europe.

26. *STRIX INDICA*, Blyth.

Formerly included in *S. javanica*, De Wurmb., which Jerdon has recently (Ibis, 1871) stated to be more nearly allied to *S. candida*, Tickell. *S. indica* is very local in Ceylon, and is entirely confined to the north of the island. Layard gave the fort of Jaffna as the only locality for it ; but I have since obtained it at Aripo, where a pair of these Owls were resident. They frequented a government storehouse in my compound, each regularly perching in a dark corner under the roof, at opposite ends of the long building, and apparently living in harmony with the hundreds of Bats which hung from the roof and walls around. I have never observed these birds out of doors until some time after sunset.

Bill horny yellow ; irides black ; feet yellowish brown.

Ceylon, India.

27. *SYRNIUM INDRANEE*, Sykes.

This bird is found in the low country in the northern half of the island and on the lower hills ; but although well known to and dreaded by the natives as a bird of ill omen, it does not appear to be anywhere numerous. Doubts have been expressed as to whether the so-called "Devil-bird" is really an Owl ; but I have frequently questioned the native hunters about the bird, which is so notorious in Ceylon for its horrible cries ; and they have described it in such terms as to leave no doubt in my mind about its being an Owl, and probably of this species.

I have only seen specimens of it from the Kandy district ; but it has been found in several parts of the island, and I once had an opportunity of hearing the bird under very favourable circumstances near Aripo. I was lying out in wild jungle about eight miles from my house, and five from the nearest native village, watching for Bears. It was bright moonlight ; the Nightjars had long ceased their churring notes, and there was an almost unnatural stillness around—the midnight silence of the jungle, only occasionally broken



by the distant roar of the surf. For several hours I had been watching the small drinking-hole in front of me; and it was now time for the Bears to come if they meant to visit the pool at all that night. I was eagerly scrutinizing the openings among the bushes, when piercing cries and convulsive screams suddenly issued from a small patch of bushy jungle about thirty yards on the left of my hiding-place. My hunter at first thought a Leopard was there, and told me to keep quiet; but the cries increased, and became so horribly agonizing, that it was difficult to believe murder was not being committed; so, jumping up with my double rifle in my hand, I ran cautiously down to the patch of jungle, my trusty servant following with a second gun. Before I reached the place all was as silent as before, and the idea of the Devil-bird flashed across my mind. This was afterwards confirmed by the hunter, who, however, did not apparently care to talk much about it. A careful examination of the sandy ground among and around the bushes when daylight appeared resulted in no evidence of any tracks of Leopards or recent traces of other quadrupeds. I have no doubt, therefore, that it was this dreaded Owl which had disturbed our night watch; and although my sport was spoilt for the night, I did not regret having heard for once the really appalling cries of this ill-omened bird. The dimensions of a Ceylon specimen are:—Length 20 inches, wing 13, tarsus 2.

Ceylon, S. India, Malacca, Formosa (*Swinhoe*).

## 28. *HUHUA PECTORALIS*, Jerdon.

Some three or four years ago, whilst I was in Ceylon, Mr. Samuel Bligh brought to me for identification some specimens of a Horned Owl, which appeared to us, after examination, to be identical with *Huhua nipalensis*, Hodgson, except in being smaller, but agreeing in that respect with the measurements of a bird from S. India described by Jerdon as *H. pectoralis*. Considerable confusion has existed between *H. nipalensis*, Hodgs., from Nepal, *H. pectoralis*, Jerdon, from S. India, and *H. orientalis*, Horsfield, from Java; and the subject is referred to by Jerdon (B. of Ind. vol. i. p. 132) as a matter on which "materials are wanting to form a just conclusion." Jerdon has since (*Ibis*, 1871, p. 346) stated his opinion that the Nepal species will stand, and has united the other two under *H. orientalis*—but, I understand, in the absence of a specimen from S. India for direct comparison.

A comparison of one of the Ceylon birds with specimens of true *H. orientalis* and *H. nipalensis* in the British Museum has satisfactorily shown, however, that they are three very distinct species, and that the Ceylon bird is very probably the same as *H. pectoralis* from S. India. In this conclusion I am supported by Mr. Gurney and Lord Walden.

*H. pectoralis* may be described as like *H. nipalensis*, but very much smaller, both of them wanting the *closely* barred plumage of *H. orientalis*. It is, I think, evident from Jerdon's measurements of *H. nipalensis* that they were taken from a Malabar specimen of



what was supposed to be that species; but he says in his description, the brown bars of the under parts "in some tending to coalesce and form a pectoral band." In his figure of *H. pectoralis* in the 'Madras Journal' great prominence is given to this band; but in the Ceylon bird it is not very distinct; and, as Mr. Gurney has pointed out to me, this difference appears to be owing to the light intervals between the dark transverse bars on the pectoral feathers being not so light in the plate as in the Ceylon bird\*. The following are the comparative measurements of the three species:—

	Length. in.	Wing. in.	Tarsus. in.
<i>H. nipalensis</i> , from Nepal (B.M.) ..	28-29	18·5	2·5
<i>H. pectoralis</i> , from Ceylon .....	22	16	2
<i>H. orientalis</i> , from Java (B.M.) ....	20	12	1·5

*H. pectoralis* is not uncommon on the lower Ceylon hills, and has probably been mistaken, without much critical examination, for the common *Ketupa ceylonensis*.

Bill yellow; irides brown; feet dull yellow.

Ceylon, S. India.

## 29. KETUPA CEYLONENSIS, Gmel.

Generally distributed over Ceylon, but perhaps more common in the low country than on the hills. I have frequently met with them near Aripo. Large trees overhanging a tank are a favourite resort of these birds, and I have often found them in the early morning perched day after day on the same branch. They are frequently captured and kept alive by the natives.

Bill dusky yellow; irides yellow; feet dirty yellow.

Ceylon, India, Burmah, China; Palestine (*Tristram*).

## 30. EPHIALTES BAKKAMUNA, Forster.

Some difficulty exists in determining how many species of small Tufted Owl are found in Ceylon, partly on account of the confusion there has been among the species or races found in India, and variously named by different naturalists, and partly because there is some doubt about the correctness of Dr. Kelaart's identification of the species he records. There is, I think, no question, however, that the very common and widely distributed species is that given by Jerdon as *Ephialtes lempigi*, Horsf., but described from Ceylon in 1781 by Forster as *Strix bakkamuna*, an unfortunate name, as it is evidently meant for "bakha muna"—lit. "Fish-Owl," and the Singhalese name for *Ketupa ceylonensis*. Forster's plate, however, shows that his bird was the common *Ephialtes*.

*E. bakkamuna* is very common in most parts of the low country, and is also found about Kandy and on the lower hills. It was a

\* Since the above was written Mr. Bligh has sent home a specimen of this Owl for the Norwich Museum. It is generally rather darker, and probably more mature than the one in my possession; and the pectoral band is very distinct, leaving no doubt of the validity of Jerdon's species.



constant evening visitor to the trees surrounding my house at Aripo ; and its single call of *whock*, repeated at short and regular intervals, was frequently to be heard far into the night. It is a bird of rapid flight. A young bird of this species was completely tamed by Mr. Bligh in Ceylon ; it would fly to his finger, and delighted in being stroked and played with ; and this tameness continued undiminished after the bird had become adult. I have often had this amusing little pet in my hands.

The dimensions of a Ceylon specimen, a female, are :—Length 8 in., wing 6·5, tarsus 1·4, tail 3.

Bill dusky ; irides yellow ; feet greyish.

### 31. EPHIALTES SUNIA, Hodgson.

Kelaart mentions, under this name, “ a very small reddish-yellow Eared Owl, occasionally seen in the very highest parts of the island.” I have some recollection of seeing a specimen from the hills, which I believe was the bird he referred to, and I think the species may be included in the Ceylon list. It is probably the rufous phase of *E. pennatus*, Hodgs.

I have no evidence of any other species of *Ephialtes* in Ceylon than the two I have here given.

### 32. ATHENE CASTANEONOTA, Blyth.

Peculiar to Ceylon. It is probably confined to the southern half of the island, and has been killed on the hills and in the low country, but is by no means common. I obtained two specimens that were killed in the neighbourhood of Kandy. It is admitted as distinct from *A. castanoptera* from Malaya.

Bill yellow ; irides — ? ; feet greenish brown.  
Ceylon.

### 33. NINOX HIRSUTA, Temm.

Rare in Ceylon ; I have only seen one specimen, which was obtained in the central district. Layard also only met with it once in the course of eight years.

Bill green ; irides yellow ; feet dingy yellow.  
Ceylon, India, Borneo, China, Japan.

### 34. HIRUNDO RUSTICA, Linn.

Referred to by Layard under *H. gutturalis*, Scop., the Indian representative of the European species ; but the grounds for separating them appear to be of the slightest description, and I shall adopt the now general opinion that they are the same. This bird is a winter visitor to Ceylon, and generally distributed, but especially abundant in the low country. Most of these birds are young ones, without the long tail-feathers.

Asia, Africa, Europe.

### 35. HIRUNDO DOMICOLA, Jerdon.

Confined to the upper hills in Ceylon. It is a very familiar bird,



commonly nesting in the verandas at Nuwara Eliya and in the district.

Ceylon, Neilgherries, Malaya.

36. ? *HIRUNDO DAURICA*, Linn.

Layard records having obtained a single specimen at Point Pedro. It is more probable, however, that this bird was *H. erythropygia*, Sykes, the S. Indian species, which had not at that time been distinguished from *H. daurica*, a northern bird and having a wide range to the eastward. They both have the under plumage streaked.

Ceylon, S. India.

37. *HIRUNDO HYPERYTHRA*, Layard.

This Swallow was discovered by Layard in 1849, and until recently was considered peculiar to Ceylon; but I have seen a specimen lately received by Lord Walden from Malacca, and it has been otherwise recorded from that country. It is abundant in the central and, at times, in the western and southern districts of the island, both in the low country and on the lower hills; but I have not observed it at Nuwara Eliya or in the north. Its distinguishing character consists in the whole of the underparts being deep chestnut.

Bill black; irides brown; feet black.

Ceylon, Malacca.

38. *ACANTHYLIS GIGANTEA*, Temm.

This bird is said to be well known at Nuwara Eliya; and Layard mentions hearing of the native report that it breeds in hollow rhododendron trees; but there is probably some mistake, as I could hardly have failed to notice the bird under such circumstances. I have only seen it from the coffee-districts; and although specimens have undoubtedly been obtained at Nuwara Eliya, I expect it will be found to be only an occasional visitor there.

Ceylon, S. India, parts of Malaya.

39. *CYPSELUS MELBA*, Linn.

Probably a winter visitor to Ceylon. It is found in some abundance on the hills at that season, but is rather local in its distribution. I have seen it at Nuwara Eliya in the cold season; and it remains there several months, particularly frequenting some precipitous cliffs overlooking the plain on which the little town is built. In the afternoon fifty or sixty of these birds might any day be seen on the wing dashing past the hill-sides in pursuit of insects, or sweeping in wider circles at a considerable elevation.

Hill-regions in Ceylon, India, W. Asia, Africa, Europe.

40. *CYPSELUS AFFINIS*, Gray.

Layard speaks of this bird as migratory, and breeding in April in large numbers about the rocks at Damboul. I have also found it nesting, but in August, under the rocks overhanging the entrance to the famous temple at Damboul; and as it breeds in Ceylon during



the summer months, I have no doubt it is a resident species. It has been met with in other parts of the island, but is local. I have not observed it on the upper hills.

Ceylon, India.

41. *CYPSELUS BATASSIENSIS*, Gray.

Very common in the low country and particularly abundant in the north, where the palmyra is the common palm, on which it builds its nest. I have not observed this bird at Nuwara Eliya, but have known the following species sometimes mistaken for it there.

Bill black; irides brown; feet brownish.

Ceylon, India, Assam, Burmah.

42. *COLLOCALIA FUCIPHAGA*, Thunberg (1772).

There are several localities in Ceylon in which this little Swift has been known to breed; and Layard has given a good description of one which he visited. These breeding-stations are at various elevations, from close to the sea to the highest hills. Kelaart states that he has "heard from very good authority that some years ago baskets of the edible nests were obtained from a cave on the Pedrotallagalla hill. Very nutritious soup was made of them for the invalids who at that time resided at Nuwara Eliya." This hill is the highest (8200) in Ceylon and overlooks the still all-important sanatorium; but I could not ascertain the situation of the cave. The birds, however, are numerous at Nuwara Eliya in winter; and I have obtained specimens there, as well as in the low country between Colombo and Kandy.

Bill black; irides dark brown; feet purple-brown.

Ceylon, N. and S. India, Assam, Malay islands.

43. *DENDROCHELIDON CORONATUS*, Tickell.

Layard says this species is generally distributed; but I have only seen it between Colombo and Kandy and in the south. I shot a pair of these birds near Colombo at the end of May, and have seen it in abundance in Kandy itself in March. I think there is little doubt that it is a resident, although perhaps migrating from one part of the island to another. I have neither seen nor heard of it at Aripo or Nuwara Eliya.

Bill black; irides dark brown; feet bluish black.

Ceylon, S. and Central India, Pegu.

44. *BATRACHOSTOMUS MONILIGER*, Blyth.

I have only seen one skin of this curious bird. It was procured by Mr. H. Nevill close to Amblangodde Lake, a few miles north of Galle. I was shown the spot where it was killed—a small piece of recently cleared land nearly surrounded by rather low jungle. The species has been but rarely met with, and, so far as is known, is confined to the south-west of the island, in the country lying between Adam's Peak and Galle.

Mr. Blyth tells me that Jerdon's description of this species was



taken from a Ceylon specimen, but that the one from S. India is probably the same.

Ceylon, S. India?

45. *CAPRIMULGUS KELAARTI*, Blyth.

This species, first discovered in Ceylon, is entirely confined to the hills, ranging from about 2000 feet upwards, and common in the Nuwara Eliya district. It is very noisy during March and April, at the commencement of the breeding-season, appearing with great regularity a few minutes after sunset from its accustomed hiding-place in the thick jungle. I have reason to think this Nightjar leaves the upper hills during the cold season and descends to a more temperate climate.

Bill dusky; irides dark brown; feet fleshy brown.

Ceylon, Neilgherries.

46. *CAPRIMULGUS ATRIPENNIS*, Jerdon.

Tolerably common near Colombo and in the south; I have also obtained it in the interior, about twenty miles from Trincomalie. It is, I believe, a low-country bird; and I have not met with it above the foot of the hills.

The specific distinctness of this Nightjar was hardly ascertained when Layard published his "Notes on the Ornithology of Ceylon;" and when he speaks of so rare and remarkably coloured a species as *C. mahrattensis*, Sykes, being "abundant in the vicinity of Colombo and throughout the southern province," and that bird has not been met with in Ceylon by any subsequent collector, it is not unreasonable to conclude that the species intended is that which I have since identified as *C. atripennis* from the same localities.

Bill dusky; irides brown; feet pale brown.

Ceylon, S. India.

47. *CAPRIMULGUS ASIATICUS*, Latham.

Common in the low country, especially in the northern half of the island, where it is resident. I have found it breeding in September at Aripo, its two eggs being deposited on a bare sandy spot under the shelter of a bush. At Colombo it is numerous in the cinnamon-gardens during at least part of the year, hiding during the day at the foot of the bushes; but I have no recollection of seeing this bird in the south of the island. Ceylon specimens are very grey compared with those from India, a good series of the latter which I have examined having all a conspicuous rufous tinge on the upper surface. This is only observable in Ceylon birds in young specimens. There is another point in connexion with this species to which I wish to direct attention; and it may be desirable to extend it to other species of *Caprimulgidae*. Jerdon and other authors have been accustomed to group the species of *Caprimulgi* in accordance with the number of tail-feathers which have a white terminal spot, this spot being supposed to be found only in the male. I need hardly say that it is only too common for collectors to omit any notice of the sex of the birds



they shoot ; and this frequent omission has no doubt led to the overlooking of the fact that in *C. asiaticus* the female has the white tail-spot as well as the male, although about one fourth shorter. This is not a peculiarity of birds from Ceylon, as in a series of Indian specimens of unknown sex in Lord Walden's collection I was able to separate them at once into two groups agreeing on this point precisely with the known sexes of the birds of my own collecting. I first noticed the presence of the spot in the female in a specimen I shot at Aripo in 1866, and I made a note of it ; but the skin was destroyed by rats ; I have since obtained two more examples showing precisely the same character, so that the first could not have been an accidental variation. I have not been able to ascertain whether the same character is found in the two other Ceylon species, as all my specimens of them are males.

Bill dusky, tip darker ; irides very dark ; feet brownish flesh.  
Ceylon, India, Burmah.

48. *HARPACTES FASCIATUS*, Forster.

Only found among wild tree-jungle in the southern half of the island. I have seen it about twelve miles from Colombo, in a wild uncultivated district in the low country, and also at Nuwara Eliya, in February ; but it is not very commonly met with, and is perhaps somewhat local in its distribution. In its manners it resembles the Flycatchers, and has generally a peculiar fluttering mode of flight.

Bill dark blue ; cere smalt ; irides brown ; feet lavender.  
Ceylon, South and Central India.

49. *MEROPS VIRIDIS*, Linn.

Exceedingly abundant in the northern part of Ceylon, where it is a resident. It is also found sometimes at Colombo and on other parts of the coast. Whilst living at Aripo I had constant opportunities of observing these birds closely, as the railings of my veranda were a favourite perching-place for them, and they would allow me to approach within a few feet without showing any alarm. Forty or fifty of these beautiful birds generally roosted in a small bushy tree only a few yards from the house. This species seems to prefer a low station when looking out for its prey, frequently perching on a small stick only a few inches from the ground. The Ceylon birds generally have the blue throat which is found in the variety described as *M. torquatus* by Hodgson.

Bill black ; irides blood-red ; feet lead-colour.  
Ceylon, India to China.

50. *MEROPS PHILIPPINUS*, Linn.

A migratory bird ; generally distributed over the low country, but, like the preceding species, very numerous in the north. I have first observed it at Aripo at the end of September ; and it remains there till the change of the monsoon in April. It is a noisy bird, with a lofty, dashing flight, successfully pursuing the dragonflies, and then



sailing back on outstretched wings to its favourite station on the dead branch of some neighbouring tree, where the insect is killed and swallowed. In the early mornings of March, when there has been but little wind stirring, and the sea was as smooth as glass, I have frequently observed these Bee-eaters hunting for insects close to the surface, and a quarter of a mile from the shore. I have noticed this bird frequently at Colombo, but only in small parties. At Aripo I have often seen sixty or seventy on the same tree; in fact, during its stay in Ceylon, it is more numerous there than the resident species.

Bill black; irides blood-red; feet lead-colour.

Ceylon, India, Burmah, Malaya.

#### 51. MEROPS QUINTICOLOR, Vieill.

This is a hill-species, and a resident in Ceylon. I have shot it in August at the foot of the hills in the south, and I have frequently seen it on the lower hills in the neighbourhood of Kandy; but it is nowhere so numerous as either of the preceding species, and is generally seen singly or in pairs. I have not observed it on the upper hills. Of two Ceylon specimens, with the chestnut border to the black throat-band, one has the tail entirely green, and the other with the central feathers blue.

Bill black; irides blood-red; feet lead-colour.

Ceylon, India, Burmah.

#### 52. CORACIAS INDICA, Linn.

This bird, although undoubtedly locally common in the north of Ceylon, has never come under my notice in the jungly district of Aripo; nor have I seen it in the south of the island. In the country between Colombo and Kandy, however, I have frequently met with it; and its often noticed habit of perching on the top of a bare pole or the stem of a dead tree is also characteristic of the bird in Ceylon. Its flight is regular and crow-like; but when perched its head is sunk on its shoulders, giving the bird a remarkably clumsy appearance, as is also the case with the Bee-eaters when not on the wing.

Bill blackish; irides dark brown; feet fleshy yellow.

Ceylon and the greater part of India.

#### 53. EURYSTOMUS ORIENTALIS, Linn.

Layard met with three examples of this bird; but it has never come under my notice.

Ceylon, northern half of India, Burmah, Malaya, China.

#### 54. PELARGOPSIS GURIAL, Pearson.

The synonymy of this bird has been much confused. It is mentioned by Layard under the name of *Halcyon capensis*, Linn., and is described by Jerdon under the heading of *H. leucocephala*, Gmel. Mr. R. B. Sharpe (P. Z. S. 1870), however, has worked out the



question of identity, and has restored to it Pearson's original name of *gural* (1841), by which it should be known.

I have seen skins of this species, but have not met with the bird alive. Layard speaks of its frequent occurrence on the east side of Ceylon, and also of its being found about Caltura, on the west coast. The latter locality is, I have heard, a good one for this bird; and I have reason to think it is also sometimes met with on the lower hills.

Bill red; irides brown; feet dull red.  
Ceylon, India eastward.

55. *HALCYON SMYRNENSIS*, Linn.

*H. fusca*, Bodd., is now admitted as identical with the old Linnean *H. smyrnensis*. In Ceylon this Kingfisher is abundant in the low country wherever there is water, frequenting alike the neighbourhood of paddy-fields and the banks of rivers. It is perhaps less numerous in the north than elsewhere, but it was not uncommon at Aripo. Ceylonese specimens are generally more brightly coloured than those from other countries.

Bill deep red; irides brown; feet vermilion.  
Ceylon, India to China, Asia Minor.

56. *HALCYON PILEATA*, Bodd.

Recorded by Layard, under *H. atricapilla*, Gmel., as having been killed by him in the north of the island. It appears to be an eastern species, and rare both in India and Ceylon.

Ceylon, India, Burmah, Malaya, China.

57. *CEYX TRIDACTYLUS*, Pall.

Widely distributed in Ceylon, but nowhere common, and only to be procured with difficulty. I have never seen the bird alive, but at various times obtained three specimens, which were killed in the central district.

Bill coral-red; irides brown; feet red.  
Ceylon, India, Malaya.

58. *ALCEDO BENGALENSIS*, Gmel.

Common everywhere in Ceylon. It is always to be found at Nuwara Eliya, as well as in all parts of the low country.

Bill reddish, with the upper part dusky; irides brown; feet coral-red.

Ceylon, India to China, Malaya.

59. *CERYLE RUDIS*, Linn.

This is a common species, frequenting rivers more than tanks or paddy-fields. I have frequently met with it at Aripo; and it is particularly abundant on the southern rivers.

Bill black; irides brown; feet brown.  
Ceylon, India to S. China, Malaya, W. Asia, S. Europe, Africa.



60. *HYDROCISSA CORONATA*, Bodd.

Confined to wild forest jungle in the central and northern parts of the island. I have seen it occasionally a few miles from Aripo; and whilst travelling through an extensive tract of forest on the road between Kandy and Trincomalie, small parties of these birds were frequently observed on the tops of the trees, or slowly sailing across the road from one part of the forest to the other. In the early mornings their harsh cries mingled discordantly with the howlings of Monkeys (*Presbytes*), the call of the Jungle-fowl, and the more musical notes of the Long-tailed Robin (*Kittacincla*), almost the only sounds to be heard in this primitive jungle, far from the borders of cultivation, and only disturbed by occasional travellers or the bell of the light-stepping postal runner.

Ceylon, S. India.

61. *TOCKUS GINGALENSIS*, Shaw.

Considerable confusion has existed between this species and *T. griseus*, Latham; and it is desirable to mention that the species properly known as *T. gingalensis* is only found in Ceylon. Under the above heading Jerdon has inadvertently spoken of both in his 'Birds of India,' but he has since corrected the mistake (*Ibis*, 1872, p. 5).

*Tockus gingalensis* is, according to Layard, not uncommon in certain districts; and Lord Walden has received several specimens of it. It keeps, I believe, mostly to the forests; and I have only once obtained it at Aripo, where its harsh cry betrayed its presence on a low tree close to my house. The colour and shape of the bill in this bird vary a good deal with age.

Bill yellowish, more or less marked with black; irides reddish brown; feet slate grey.

Ceylon.

*Tockus griseus*, Lath., is said by Jerdon to be also found in Ceylon; but I cannot hear of any well authenticated specimens.

62. *PALÆORNIS ALEXANDRI*, Linn.

Bill red; irides buff; feet slate.

Ceylon, India, N. Burmah.

63. *PALÆORNIS TORQUATUS*, Bodd.

Bill red; irides buff; feet slate.

Ceylon, India, W. Asia, Tropical Africa?

These two species are exceedingly abundant in the north of Ceylon; but I have not seen them on the hills or in the south.

64. *PALÆORNIS ROSA*, Bodd.

I have only met with this species in the southern parts of Ceylon, where it is very destructive to the grain crops; but it is also found at times on the lower hills generally. I have seen a flock of fifty of these birds fly down one after another to a field of paddy; and each biting off a ear of the green corn, return to a neighbouring tree to



devour the plunder; and this has been repeated again and again. The above three species are constantly caged by the natives; and few native dwellings are without one or other of these favourite pets.

Bill yellow above, black below; irides buff; feet greyish.

Ceylon, India.

*Note.*—There is some doubt about the further range of this species, a closely allied form, with yellow under wing-coverts, having probably been confounded with it.

#### 65. *PALÆORNIS CALTHROPÆ*, Layard.

Peculiar to Ceylon. It was first obtained by Layard at Kandy, where it is frequently numerous; and it is said to be generally distributed over the hills. Although recorded by Kelaart from Nuwara Eliya, I suspect this beautiful bird is only a rare visitor to that cool region, as I have never seen a Parrot of any kind at that elevation, and I have always been on the look-out for this species in particular. The colouring in the sexes is alike, except that in the female the green on the side of the head is less distinct, and the bill is black instead of red.

Bill, ♂ red, ♀ black; irides buff; feet slate.

Ceylon.

*Note.*—As some confusion has existed with regard to the correct spelling of the specific name of this species, I may mention, on the direct authority of Mr. Layard, that it was given from "Calthrop," a family name.

#### 66. *LORICULUS INDICUS*, Gmel.

Peculiar to Ceylon. The history of this species has been fully discussed in a paper by Lord Walden (*Ibis*, 1867, p. 467), from which it appears that, although Edwards first figured and described the bird, it should stand as *L. indicus*, Gmel., according to the rules of zoological nomenclature. The name is unfortunate, as it is certain that the species is not found out of Ceylon; but it was not known by the earlier writers from what part of the Dutch settlements the bird described by Edwards was obtained.

This little bird is common in many parts of the southern half of Ceylon, and particularly quite in the south. It frequents cultivated ground and large native gardens; and I have sometimes seen it on the cocoanut-trees busily biting off and apparently eating the chip-like flowers. I believe it is confined to the low country. It is often caged by the natives, and, like allied species, sleeps suspended from the top of its cage by its strong curved claws. There is little difference in the colouring of the sexes; but individuals vary a good deal in the extent and brilliancy of the golden gloss on the back.

Bill reddish orange above, orange below; irides white; feet dull yellow.

Ceylon.

#### 67. *PICUS MAHRATTENSIS*, Latham.

Common in the Aripo district, and, so far as I know, only found



in the north of Ceylon. It appears to frequent low jungle, and I have rarely seen it except on dead wood near the ground and old fences. It is a resident species.

Bill slate; irides dull red; feet lead-colour.

Ceylon, India.

*Picus macei*, Vieill., has been recorded by Dr. Kelaart as being found in Ceylon; but I think its occurrence is very doubtful, in which opinion I am strongly confirmed by Mr. Blyth.

#### 68. YUNGIPICUS GYMNOPHTHALMOS, Blyth.

This little Woodpecker was discovered in Ceylon by Layard, and it is said to have been since found in S. India. It frequents the upper branches of large trees, and, although generally running over the bark in true Woodpecker fashion, may sometimes be observed perched across the smaller twigs. I have only obtained it in the neighbourhood of Colombo; but it is also found in the south.

Bill greenish slate; irides pale buff; orbital skin purple; feet greenish slate.

Ceylon, S. India?

#### 69. CHRYSOCOLAPTES FESTIVUS, Bodd.

By the kindness of Lord Walden, I am enabled to include this handsome Woodpecker in my list of Ceylon birds. The two specimens, male and female, in his collection, are labelled "November 1865, Cocarry." The name is probably that of a small native village in the north-west of the island, not far from the Aripo district, as I have reason to know that the birds collected at that date for Lord Walden were procured not many miles from where I was afterwards residing. Future collectors in Ceylon, who are not familiar with this species (described by Jerdon under the name of *C. goensis*, Gmel.), may recognize it by its black back and golden wings, the underparts being coloured much as in *C. stricklandi*, Layard.

Ceylon, parts of South and Central India.

#### 70. CHRYSOCOLAPTES STRICKLANDI, Layard.

Peculiar to Ceylon, and confined to the hills. It is abundant at Nuwara Eliya and in all tree-jungle in that district, ranging from the forest-clad Pedrotalagalla (8200 feet), the highest point in the island and overlooking the Nuwara Eliya plain, through the coffee-districts, to the Kandy country. The female has the whole top of the head and crest black, spotted with white; and a young bird of that sex had the lower part of the back black, faintly barred with white, with crimson feathers appearing among the others: the bill in this bird was only two thirds the length of that in the adult.

Layard states that the irides of this species are red-brown; but I think he must have been mistaken, as in four specimens I obtained at Nuwara Eliya, and which I myself prepared, the irides were buff, those of the young bird being rather paler than the others.

Bill greenish white; irides buff; feet greenish slate.

Ceylon hills.



71. *CHRY SOPHLE GMA CHLOROPHANES*, Vieill.

I have only procured this species at the foot of the hills in the south; but it has been also obtained in other places much nearer Colombo. When not feeding, it is in the habit of stationing itself on the highest branch of a dead tree, and there repeating its peculiar note, which has little of the harsh sound so generally characteristic of the Woodpeckers.

Bill slate, with the base yellow; irides dull red; feet dull green. Ceylon, S. India.

72. *MICROPTERNUS GULARIS*, Jerdon.

Two specimens of this Woodpecker were procured by me a few miles from Colombo. Although decidedly a scarce species, and I shot these two birds in January and July, they were both killed in native gardens not a quarter of a mile apart. Layard met with it in the south; and I have seen one or two skins from the central district.

Ceylon specimens have the lower parts rather darker than those from India. Layard gives this bird under the name of *M. phaiiceps*, Blyth.

Bill lead-grey; irides red-brown; feet slaty brown. Ceylon, S. India.

73. *BRACHYPTERNUS AURANTIUS*, Linn.

Recorded by Layard as very abundant in the Jaffna peninsula in the north of the island. I occasionally saw at Aripo what may have been this species, and heard its remarkable cry, but failed to procure a specimen.

73 *bis*. *BRACHYPTERNUS PUNCTICOLLIS*, Malh.

A specimen of this bird has been quite recently received by me from the western side of the island. A further examination of the Golden-backed Woodpeckers found in Ceylon therefore appears desirable, as the species generally met there is more likely to be *B. puncticollis*, common in Southern India, than *B. aurantius*, which has a more northerly range. *B. puncticollis* may be recognized by its white-dotted throat and under neck.

Ceylon, S. India.

74. *BRACHYPTERNUS CEYLONUS*, Forster.

Peculiar to Ceylon; not uncommon near Colombo, but very numerous in the south. Dr. Kelaart\* says it is "found in great abun-

\* The results of my own collecting at Nuwara Eliya and in the neighbouring jungles during almost every month in the year oblige me frequently to receive with suspicion the notices by the late Dr. Kelaart of the occurrence of birds, and of their abundance, in that district. The subjects to which Dr. Kelaart gave his special attention were mammals and reptiles, and in these he did good work; but ornithology was a very subordinate study with him, and he rarely, if ever, used a gun.



dance at Nuwara Eliya ;” but I have never seen it on the hills, and I have no doubt that *Chrysocolaptes stricklandi*, another red-backed Woodpecker already noticed, was mistaken by Dr. Kelaart for this species. This bird especially frequents the cocoanut-trees, and is a conspicuous object as it works its way by rapid jerks up the slender trunks of these palms. The natives in the south call it the “Toddy-bird,” and say it visits the palms for the sake of the toddy, which is largely collected in that and some other parts of the island. The insects feeding on the toddy are no doubt the real attraction. Its principal food is ants, as is the case with all the low-country Woodpeckers, their stomachs being always found more or less crammed with these ubiquitous and troublesome insects. Both sexes have the red occipital crest ; but the male has the top of the head sprinkled with the same colour, whilst the female has that part spotted with white.

Bill slate ; irides red ; feet pale greenish.

Ceylon.

#### 75. MEGALAIMA ZEYLANICA, Gmel.

Peculiar to Ceylon. This bird is closely allied to *M. caniceps*, Franklin, and is noticed under that name by Layard ; but it is a smaller bird, with the anterior portion of its plumage much browner, and the lighter markings reduced in size and distinctness. It is common in the low country, except in the north. I have never seen or heard it in the Aripo district ; and it does not ascend above the lower hills. The flight of this bird is straight, but rather heavy. It feeds on berries, and may be often seen clinging to the smaller twigs on the outside of a tree whilst eating the fruit which grows at their extremities.

Bill dull orange ; irides brown ; orbits yellow ; feet yellow.

Ceylon.

#### 76. MEGALAIMA FLAVIFRONS, Cuvier.

Peculiar to Ceylon. It is not confined to the hills, as stated by Layard, but is exceedingly abundant even close to Colombo, and ranges from near the coast to an elevation of 5000 feet. It is the only Barbet I have seen so high ; and I have not observed it there except during the N.E. monsoon, a time at which there is a great influx of migratory birds and of low-country species to the hills. I have not seen it in the north ; and it is not so numerous as the last species in the extreme south of the island. At the village of Heneratgodde, about 17 miles from Colombo, in a district abounding with native gardens, cocoanut-topes, and paddy-fields, and where I have collected a great variety of birds, the air used to resound with the loud notes of this and the preceding species of Barbet, a partial silence only occurring for an hour or two during the extreme heat of the day. *M. flavifrons* is a more sprightly bird than *M. zeylanica*, and can be readily distinguished from it when on the wing.

Bill horny yellow ; irides red-brown ; feet dark grey.

Ceylon.



77. *XANTHOLÆMA INDICA*, Lath.

Layard speaks of this bird under the name of *Megalaima philippensis*. It is confined to the north. I have only met with it at Aripo, where it is found throughout the year. Perched on some dead branch near the top of a tree, with its throat swelling and its head bowing at the utterance of each note, this handsome little Barbet repeats its monotonous cry of *poohp, poohp, poohp* for half an hour at a time, with only occasional intervals of a minute or so. Whilst thus engaged it changes the direction of its head with every note; and to this I think is mainly due the often noticed variation in the sound; but the range of direction is a full semicircle; and after often listening to the bird from different positions, I have no doubt that the voice is also dropped a little when the head is turned quite on one side. In Ceylon, as in India, this bird is known by the name of "coppersmith;" and that title is also applied about Colombo to the following species.

Bill black; irides red-brown; feet pink.

Ceylon, India, Burmah, Malaya.

78. *XANTHOLÆMA RUBRICAPILLA*, Gmel.

Peculiar to Ceylon, and common in the low country in the southern half of the island. I have also frequently seen it at Trincomalie; and Layard has procured it at Jaffna; but I have never met with it in the Aripo district. It is very common about Colombo. The note of this bird is very much like that of *X. indica*, but is not nearly so loud, and is repeated quickly four or five times without a pause; then resting for three or four seconds, the bird goes on as before. The call of *X. indica* sounds like distinct heavy blows of a hammer on a copper vessel heard in the distance; that of *X. rubricapilla* like a series of light taps on the same metal.

Bill greyish black; irides red-brown; feet pink.

Ceylon.

In a young bird I obtained in July near Colombo the bright colours about the head and neck were not developed, except a small patch of orange below the eye and a tinge of yellow on the forehead. The bill was dark grey; irides pale brown, and feet dusky flesh.

79. *CUCULUS CANORUS*, Linn.

Recorded by Layard as found in Ceylon. He obtained one example of it near Colombo; but I have not met with it.

Asia, Africa, Europe.

80. *CUCULUS SONNERATII*, Latham.

Kelaart procured several specimens; and I have seen it from near Colombo and the lower hills.

Ceylon, S. India.

81. *CUCULUS MICROPTERUS*, Gould.

This Cuckoo was recorded by Dr. Kelaart as a mountain species;



but the only two examples I met with were obtained in half-cultivated land in the low country near Colombo. To this species may probably also be referred a bird closely resembling *C. canorus* which I watched for some time in an English garden at Colombo a few days after my arrival in Ceylon.

Bill yellowish, dusky above; irides pale yellow; feet yellow.

Ceylon, India, Burmah, Malaya, China.

Layard has described a Ceylon Cuckoo under the name of *C. bartletti*; but there is some doubt about what the bird is. Jerdon places it under *C. poliocephalus*, Lath., which, however, has not been recognized in the island; it may be *C. sonneratii*.

## 82. *HIEROCOCCYX VARIUS*, Vahl.

This bird is probably a migrant from India. Layard procured it near Colombo; but I have only met with it on the hills at Nuwara Eliya, in the beginning of the year.

Bill greenish yellow, dusky above; irides yellow; feet yellow.

Ceylon, India, Burmah, Malaya.

## 83. *POLYPHASIA PASSERINA*, Vahl.

Referred to by Layard as *Cuculus tenuirostris*, Gray, and by Jerdon (B. of India, vol. i. p. 333) as *P. nigra*, apud Blyth. Jerdon has lately, however (Ibis, 1872, p. 14), gone more into the nomenclature of the species, and placed it under the above heading. It is migratory to Ceylon, but appears much later than most of the other visitants. Layard gives February for its arrival about Jaffna; but I have first seen them at Aripo in the beginning of January, and then they all at once became abundant, frequenting low bushes in the jungle, and ranging in colour from dark grey to completely rufous on the upper parts. No two specimens were exactly alike; but all were of some shade of grey beneath, and more or less barred. The rufous-bellied species is an eastern bird, and unknown in Ceylon.

Bill black above, red-brown below; irides hazel; feet dull yellow. Ceylon, India.

## 84. *SURNICULUS DICRUROIDES*, Hodgson.

Resident, but rather a scarce bird in Ceylon. It has been found on the lower hills, near Kandy; and I have obtained specimens in immature and adult plumage in the low country near Colombo, and in the extreme south of the island. Although at first sight this Cuckoo may be readily mistaken for a King Crow, having the same general colour and remarkable shape of tail, it is not difficult to distinguish it when within a moderate distance. It usually perches lower and alights more frequently on the ground, besides having little of the Flycatcher-action so common among the *Dicruri*.

Bill black; inside of mouth deep orange; irides dark brown; feet black.

Ceylon, India, Burmah.



## 85. LAMPROCOCYX MACULATUS, Gmel.

This beautiful Emerald Cuckoo was first made known from Ceylon, and appears to be the one given by Kelaart and Layard in their Catalogue (1853) under the name of *Cuculus xanthorhynchos*, Horsf., a Malay species. I have seen no specimens of it; and it is undoubtedly rare.

Ceylon, India.

## 86. COCCYSTES JACOBINUS, Bodd.

*C. melanoleucos*, Gmel.; Jerdon, B. of Ind. no. 212.

Common in the north of the island. These birds are always numerous in the Aripo district, frequenting bushes and low trees, and usually perching on the highest branches. In December and January (the commencement of the breeding-season with many birds in Ceylon) they are very noisy and incessantly flying from one place to another, one or more males apparently chasing the female, and uttering their clamorous cries. Layard mentions finding a young Cuckoo of this species under the care of a pair of Mud-birds (*Malacocercus*); and, from the frequent battles I observed between this Cuckoo and a pair of *Malacocercus striatus* which were nesting in a low tree close to my house, I have no doubt that the Black-and-white Crested Cuckoo frequently lays its eggs in the nest of that common Babbler.

Bill black; irides red-brown; feet lead-colour.

Ceylon, India, Africa.

## 87. COCCYSTES COROMANDUS, Linn.

I believe this handsome Cuckoo is very scarce in Ceylon. I have only seen two specimens, both from the Kandy district.

Bill black; irides reddish brown; feet lead-colour.

Ceylon, India, Burmah, Malaya.

## 88. EUDYNAMIS HONORATA, Linn.

Formerly known as *E. orientalis*, Linn.

Layard says of this bird in Ceylon:—"Wherever Crows are found, there the Coël is found also." I have only seen this bird, however, during the N.E. monsoon, from November to April. During this period it is very common in the Aripo district; and I have also found it numerous near Colombo. After April, I have never met with the species until towards the end of the year. I believe it is a true migratory bird. Among the specimens I have shot in January and February is a young male in the spotted plumage, but having the top of the head rusty brown; in other respects the colours are the same as, but purer than, those in the female. These Cuckoos are very noisy in the morning and evening.

Bill dull green; irides crimson; feet slate-colour.

Ceylon, India, Burmah, N. Malaya, S. China.

## 89. ZANCILOSTOMUS VIRIDIROSTRIS, Jerdon.

This is a low-country species, and, so far as I know, not extending



to the south of the island. It is found abundantly throughout the year in the north; and I have occasionally met with it near Colombo. It is skulking in its habits, creeping rapidly through the low bushes, and rarely exposing itself when it has once been alarmed.

Bill apple-green; irides deep red, orbits cobalt; feet dark leaden. Ceylon, S. India.

90. *PHÆNICOPHÆUS PYRRHOCEPHALUS*, Forst.

This Cuckoo has hitherto been found only in Ceylon. It inhabits tree-jungle in the low country near the foot of the hills. One specimen, alive but injured, was brought to me by some natives who had caught it only a few miles from Colombo. I saw a second flying across a road in the Central Province, and followed it for some distance through the jungle, but failed to obtain it. Its flight was weak; but it moved rapidly through the trees, half flying and half hopping from branch to branch. Layard says the irides of this Cuckoo are white; but in the living bird (a male) I had they were brown, and they are marked as of that colour in specimens in Lord Walden's collection.

Bill light apple-green above, bluish green below; irides brown; orbital skin crimson; feet dark leaden.

Ceylon.

91. *TACCOCUA LESCHENAULTII*, Less.

I am indebted to Mr. Forbes Laurie for the opportunity of examining a male specimen of this fine Cuckoo, hitherto unknown in Ceylon. He tells me it was obtained in the Doombera valley (1800 feet), not far from Kandy, that it came into his hands immediately after it was shot, and he himself prepared the skin. As a S.-Indian species it is likely to occur in Ceylon; and the Doombera valley is a wild district, from which I have known many of the rare and peculiar Ceylon birds to have been obtained.

Bill red, tip yellow; irides reddish; feet lead-colour.

Ceylon, S. India.

92. *CENTROPUS RUFIPENNIS*, Illiger.

Common generally in the low country. It was very abundant at Aripo, feeding very much on the ground, where there was always a large supply of grasshoppers.

Bill black; irides red; feet black.

Ceylon, India, Burmah, Malaya.

93. *CENTROPUS CHLORORHYNCHUS*, Blyth.

Peculiar to Ceylon, and, I believe, almost confined to the lower hills in the Central district. I have only seen this bird alive on one occasion, and then in thick jungle under trees. It is either very scarce or escapes notice from its skulking habits. From *C. rufipennis* it may always be distinguished by its green bill, if



not by the very rich purple gloss over the anterior portion of its plumage.

Bill pale green ; irides red ; feet black.

Ceylon.

94. *NECTAROPHILA ZEYLONICA*, Linn.

Common in the low country. I have frequently seen it in the gardens at Colombo ; but have not met with it at Aripo. Layard speaks of it as abundant in the southern and midland districts.

Ceylon, India.

95. *NECTAROPHILA MINIMA*, Sykes.

I do not remember seeing this bird in the Aripo district, although Layard states that it is common in the north of the island. It is occasionally seen at Colombo.

Ceylon, S. India.

96. *ARACHNECHTHRA ASIATICA*, Latham.

This species was very common at Aripo, and was found there at all seasons. I have also seen it in the south. At a Government rest-house in the extreme south of the island, where I was staying in August 1869, a pair of these birds had a nest in the veranda ; it was fastened to the end of an iron rod hanging from the roof and once used for suspending a lamp. The birds showed very little fear, although I was for several days sitting within a few feet of the nest, engaged in the preparation of specimens. I have obtained this species at Nuwara Eliya in October.

Bill black ; irides red-brown ; feet black.

Ceylon, India, N. Burmah.

97. *ARACHNECHTHRA LOTENIA*, Linn.

This is a very common species at Colombo, and is said by Layard to be plentiful in the southern and midland districts. I have no note of its occurrence at Aripo. Some specimens have the bill very much curved.

Bill black ; irides brown ; feet black.

Ceylon, S. India.

98. *DICÆUM MINIMUM*, Tickell.

I have procured this little bird in all parts of the island ; and specimens obtained at Nuwara Eliya were precisely the same as those from Aripo and elsewhere.

Bill flesh-colour ; irides brown ; feet fleshy brown.

Ceylon, India, Burmah.

99. *PIPRISOMA AGILE*, Tickell.

Layard records having obtained a pair of these birds on the Central road.

Ceylon, India.



100. *DENDROPHILA FRONTALIS*, Horsf.

Layard speaks of this bird as "abundant about jack-trees," which are only found in the low country. Although I have known it killed in such parts of the island, I have always considered it a hill species, as it is one of the common birds at all seasons at Nuwara Eliya and on the upper hills. Jerdon states that in India it is most abundant on the Neilgherries—a situation corresponding in a remarkable manner with the higher hills in Ceylon, the birds and plants of the two ranges being in most respects the same.

These little Nuthatches appear to keep in small parties at all times of the year, and are very active in examining the branches of any trees they may happen to visit. The colours of this bird soon lose their brightness after death; and the peculiar delicacy of the tints can hardly be discovered in a cabinet specimen.

Bill coral-red; irides golden; feet yellow-brown.

Ceylon, India, Assam, Burmah, Malaya.

101. *UPUPA NIGRIPENNIS*, Gould.

Very abundant in the Aripo district during the winter months, and occasionally in the summer. Some of these birds are no doubt residents in Ceylon; but their numbers in the north are largely increased about October, either by migrants from India or from the east side of the island. Layard speaks of it under the name of *U. senegalensis*, Sw., and says he "shot young birds, not fully fledged, in August." This would agree with the breeding-time of the Hoopoe in Burmah, of which Jerdon says:—"I found it breeding in holes of trees in June and July." If Layard's birds, however, were bred in Ceylon, as might be supposed from his statement that they were not fully fledged, then there are two distinct breeding-seasons for this species in the north of the island, as in January 1870 I found, in my compound at Aripo, a nest of the Hoopoe in a hole in a small mustard-tree (*Salvadora persica*). I caught the old bird as it was leaving the nest; and after enlarging the hole, came down to three young birds, just hatched, and resting on a bed of rotten wood. These nestlings were quite naked, and their bills were barely a quarter of an inch long.

The Hoopoe was found by Layard on the east and south-east coasts, and once at Colombo. I have also had a specimen from the neighbourhood of Kandy.

The flight of this bird is easy and undulating; and its note is repeated whilst it is on the wing, as well as when perched on the top of a tall bush.

There is some variation in the colours and dimensions of the Hoopoes found in Ceylon, the tendency being towards the characters of the Burmese variety described by Jerdon. Of three specimens shot at Aripo, one has the bill at front 1.9 inch, the closed wing 5; first primary entirely black, chin whitish, and the feathers of the posterior half of the crest white between the black and rufous. The last characters have been regarded as specially belonging to



*U. epops*; but in this specimen there is no white spot on the first primary. This bird was killed in December. In a second specimen, shot in February, the bill is 2·25 inches, closed wing 5·1, the general colour of the bird very rufous, and a white spot on the left first primary, but only a very minute speck on the right. A third, a male killed in December, agrees generally with the Indian form, has no spot on the first primary, but has the bill at front 2·15 inches, and the closed wing 5·25. The last and largest of these specimens did not exceed 10·8 inches in length; and they may all doubtless be referred to *U. nigripennis*, if that form be really distinct from *U. epops*.

Bill black, base flesh-colour; irides brown; feet dark leaden.

Ceylon, India, Burmah?

#### 102. *LANIUS ERYTHRONOTUS*, Vigors.

Very common in the Aripo district and in other parts of Northern Ceylon. I have also seen it occasionally in the Cinnamon Gardens at Colombo; but it does not appear to visit the hills. A cup-shaped nest of this species was built in a thorn-bush close to my house at Aripo; but the young birds had left it before my arrival there in the beginning of April. In a subsequent year I obtained young birds able to fly as early as the middle of February, and older ones nearly full-grown in March. These young birds were all very rufous, with the head, upper back, and flanks closely barred, the lower part of the back more broadly marked, and the secondaries rufous with their centres dusky. Layard says "the young are fledged in June;" but they are out some months earlier than that in the Aripo district. These birds feed very much on dragonflies and grasshoppers.

Bill black; irides dark brown; feet black.

Ceylon, India, Central Asia.

#### 103. *LANIUS CRISTATUS*, Linn.

Referred to by Layard as *L. superciliosus*, Linn. This bird I have found common in the north, west, and central parts of Ceylon during the winter months. It remains about Aripo from October to April, and is tolerably common at Nuwara Eliya during the same period. Layard mentions their being particularly numerous at Hambantotte, on the south-east coast, but does not say at what time of the year. There is probably a migration of this species from the east to the west side of the island at the beginning of the N.E. monsoon, at which time no doubt many of these birds also come from India. A specimen obtained at Aripo in October is of a much richer brown than others I shot at Nuwara Eliya in February. These birds are fond of perching on the extreme top of a bush.

Blyth (*Ibis*, 1867, p. 304) refers the birds described by Layard to "*L. lucionensis*, Scopoli (?)," a race of *L. cristatus*, "distinguished by its prevalent ashy-brown hue." This character is not uncommon in Ceylon specimens which have old, worn plumage; but I have not seen it in newly moulted birds.

Bill dusky; irides dark brown; feet dark leaden.

Ceylon, India, Andamans, Malacca.



104. *TEPHRODORNIS PONDICERIANA*, Gmel.

A careful comparison of a series of *T. affinis*, Bl., from Ceylon, with a number of *T. pondiceriana* from India, has satisfied me that there is not sufficient ground for separating them specifically. The Ceylon birds appear to be smaller; but the depth of the general ashy brown of the upper surface varies in both, and to much the same extent. The supercilium also varies in distinctness in the birds from the two countries, but in those from India the maximum development is perhaps greater than in specimens from Ceylon.

These birds are common in the north and west of the island during the winter months, and probably migrate from the eastern side. They breed early in the year; and the young birds in their spotted plumage have been procured by Mr. Legge, in April, from the cinnamon-gardens at Colombo. Birds of the year are paler than in the following season. This may not have been known to Mr. Blyth when he described the Ceylon species as greyer than those from India.

Bill dusky; irides dull yellow; feet dusky lead-colour.

Ceylon, India, Assam, Upper Burmah.

105. *HEMIPUS PICATUS*, Sykes.

This bird is rare in the low country, and seems to be chiefly found on the upper hills. It is a common bird at Nuwara Eliya throughout the year, frequenting high bush jungle or low trees. Young birds have the colours less decided than adults.

Bill black; irides yellow; feet black.

Ceylon, S. India.

106. *VOLVOCIVORA SYKESII*, Strickl.

Generally distributed over the low country; it is resident in the Aripo district, and I have found it common near Colombo and in the extreme south. Although I have shot a great many of these birds, I have never obtained a female with any other than the barred under plumage and the grey head, and I cannot confirm Blyth's statement that the adult female has a black head and neck as in the male. The black in the young male first appears in spots on the top and sides of the head.

Bill black; irides brown; feet black.

Ceylon, India.

107. *GRAUCALUS LAYARDI*, Blyth.

*Graucalus pusillus*, Bl.

A smaller bird than the N. Indian *G. macei*, Less., with which it has been confounded. It differs also (Jerdon, 'Ibis,' 1872, p. 117) in having the under wing-coverts strongly barred, the abdominal bars absent in the adult male, and the outer tail-feathers only slightly tipped with white.

I have not seen this species alive; but it is occasionally found in



the Kandy district and, according to Layard, who speaks of it as *G. macei*, in the S. and W. provinces.

Ceylon, S. India.

108. *PERICROCOTUS FLAMMEUS*, Forster.

Widely distributed, but nowhere very common. I have not met with it at Aripo, but have obtained it near Colombo; and it is tolerably numerous in the cold season on the hills. I have seen it more abundantly at Nuwara Eliya than elsewhere. It is generally in pairs, and perches high up on the trees.

Bill black; irides brown; feet black.

Ceylon, India, Assam.

109. *PERICROCOTUS PEREGRINUS*, Linn.

Common all over the island. It is resident in the Aripo district, and is found at Nuwara Eliya in the cold season. It does not frequent trees so much as bush-jungle; and I have never observed it perching very high, as is the marked habit of the preceding species.

Bill black; irides brown; feet black.

Ceylon, India, Andamans, Burmah.

110. *BUCHANGA MINOR*, Blyth.

This bird has been separated from the common Indian species (*B. macrocerca*), which it resembles in colouring, but from which it differs in all its dimensions. My finest specimen of *B. minor* is 10·5 inches total length instead of 12; wing 5·25 instead of 5·75 or 6; and other parts in proportion. The tail of the Ceylon bird is always less deeply forked than in the Indian species; and the small white rictal spot is frequently absent. Whatever may be thought of the value of these differences, they are constant; and I have not heard of the larger *B. macrocerca* of India being found in Ceylon. *B. minor* is abundant in the north; it is very common at Aripo, and is the only species of Drongo Shrike I have seen there. It is also found about Colombo, but by no means commonly within my experience. Its place there and in the south is occupied by another species. None of the Drongo Shrikes in Ceylon go above the lower hills, and for the most part they are confined to the low country.

Bill black; irides red-brown; feet black.

Ceylon.

111. *BUCHANGA LONGICAUDATA*, A. Hay.

I have seen this species on the tops of the trees in forest-jungle between Kandy and Trincomalie, and shot one specimen in a small wood about sixteen miles from Colombo. Layard says it is common in the Jaffna peninsula, and that "it frequents open lands, and perches on the backs of cattle to seek for ticks, on which it feeds largely." There must surely be some mistake about the species to which Layard here refers. His account agrees precisely with the habits of *B. minor*; and Lord Walden, who first described *B. longicaudata*, from India, where it is well known, tells me it is strictly a



forest species, frequenting high trees, and is never seen on the backs of cattle.

Bill black; irides red-brown; feet black.

Ceylon, India.

112. *BUCHANGA CÆRULESCENS*, Linn.

Layard speaks of having procured one or two specimens of this species at Point Pedro, in the extreme north; but it is not otherwise known from Ceylon.

Ceylon, India.

113. *BUCHANGA LEUCOPYGIALIS*, Blyth.

Peculiar to Ceylon. Allied to *B. cærulescens*; but differing from that species in being smaller and having the dark grey of the breast continued (but gradually becoming paler) towards the vent, with the white confined to the under tail-coverts. In the immature bird the whole of the abdominal region is very dark grey, and the under tail-coverts have three or four broad dark bands on a paler ground.

This is the common species about Colombo and in the southern district. I have never seen it in the north.

Bill black; irides brown; feet black.

Ceylon.

114. *DISSEMURUS LOPHORHINUS*, Vieill.

Peculiar to Ceylon. *D. edoliiformis*, Blyth, is apparently the same species. The typical character consists in having the head subcrested, with the simple form of tail found in *Buchanga*. In Lord Walden's large collection of *Dicruri* from Ceylon, there are many examples showing an apparent gradation in the form of the tail between this species and *D. malabaricus*; but as the true *D. lophorhinus* is found in localities where the racket-tailed species is unknown, I shall keep them distinct, and in my notice of the next species refer to the apparent gradations between them.

*D. lophorhinus* is found on some of the lower hills, and in wild districts in the low country in the southern half of the island. It appears to be quite a jungle bird.

Bill black; irides brown; feet black.

Ceylon.

115. *DISSEMURUS MALABARICUS*, Scop.

This is no doubt the species referred to by Layard under *Edolius paradiseus*, Linn., as it has been obtained in abundance in the district where Layard procured his specimens. It is quite confined to the jungle, and frequents the forests in the northern and central parts of the island. An immature specimen I shot in very wild country between Kandy and Trincomalie has the outer tail-feathers three inches longer than the next; no part of the stem is bare; but the inner web is very much narrowed just on a level with the tip of the adjoining feather. Lord Walden has received many similar specimens and others with the long racket-feathers in different stages of



growth, one example showing a difference in the intermediate length of bare stem in the growing feathers on the two sides of the tail. The short tail-feathers in some specimens appear to me to be possibly a character of youth; but they are regarded by Lord Walden as individual variations; and the attention he has given to the *Dicruri* entitles his opinion to considerable weight.

Bill black; irides brown; feet black.

Ceylon, S. India.

#### 116. ARTAMUS FUSCUS, Vieill.

Generally distributed over the low country, but is locally abundant at certain seasons. It is very common at Aripo and in the neighbourhood of Colombo during the N.E. monsoon. I have always found it in small parties and easy of approach.

Bill pale blue; irides dark brown; feet dark slate.

Ceylon, India, Burmah.

#### 117. TCHITREA PARADISI, Linn.

Generally distributed over the low country, and, at certain seasons, not uncommon on the lower hills. It is, however, a great wanderer, and very uncertain in its movements. At Aripo they have at times been very numerous; and then I have not seen one for several weeks. They seem to be of a fearless disposition, and used sometimes to fly up under the roof of my veranda after spiders when I was standing within a few yards of them. I have procured specimens in all states of plumage at different times, and two examples showing the change from the red to the white feathers—one of them at Aripo, the other at Colombo, and both in January. Layard obtained one in February in which the change was far advanced. They are common about Kandy towards the end of the year.

Bill leaden blue; irides brown; feet pale blue.

Ceylon, India.

#### 118. MYIAGRA AZUREA, Bodd.

Widely distributed, according to Layard, who, however, speaks of this bird as *M. cærulea*, Vieill. I have only seen it from the western province, where it is locally not uncommon.

Ceylon, India to China, Malaya, Andamans?, Philippines.

#### 119. LEUCOCERCA AUREOLA, Less.

*Leucocerca albofrontata*, Frankl.

Layard records the occurrence of a *Leucocerca* in Ceylon which was described by Blyth as *L. compressirostris*, from its differing from the above species in having the bill more compressed. Mr. Blyth tells me, however, that he believes now it was only a variety and that it should come under the above heading. I have examined a Ceylon specimen of *L. compressirostris*; and the character of the bill is very decided, so much so as almost to justify the separation of the bird from the *Myiagrinae*, if in other respects it did not agree so closely with *L. albofrontata*. That species, however, has been received from



Ceylon; and *L. compressirostris* may perhaps best be considered a variety of it.

Ceylon, India.

*Leucocerca* —? Mr. Hugh Nevill (J. R. A. S., Cey. Br., 1867–70, pt. i. p. 138) records the occurrence in the country round Nuwara Eliya of a Flycatcher which he calls *Leucocerca fuscoventris*, Franklin. The characters he gives of the species are evidently taken from a description of *L. pectoralis*, Jerdon; and from what I know of the circumstances I believe I am quite justified in saying that Mr. Nevill never saw the Nuwara-Eliya bird except alive in the jungle.

It is not unlikely, however, that a *Leucocerca*, probably *L. pectoralis*, may be found on the Nuwara-Eliya hills, although it has not yet been clearly identified.

#### 120. MYIALESTES CINEREOCAPILLA, Vieill.

Resident and very common in the Nuwara Eliya district. It frequents the lower branches of trees and is very bold and familiar, so much so as to be rather a pest when one is collecting in the jungle, from its habit of following one about or flitting from branch to branch just in front of one. I believe that in Ceylon it is almost entirely confined to the upper hills.

Bill dusky; irides brown; feet fleshy brown.

Ceylon, India to Burmah and Tenasserim, China.

#### 121. ALSEONAX LATIROSTRIS, Raffles.

A winter visitor to Ceylon, arriving early in October at Aripo. It is common there until April, and is also found about Colombo at the same season. Its manners are precisely the same as those of the European *B. grisola*, to which species it is closely allied.

Bill black, base yellow; irides brown; feet black.

Ceylon, S. India, China.

#### 122. ALSEONAX TERRICOLOR, Hodg.

*Butalis muttui*, Layard, described by him from a single specimen obtained at Point Pedro, agrees with Hodgson's *A. terricolor* from India. It is very rare in Ceylon.

Ceylon, North and Central India.

#### 123. OCHROMELA NIGRORUFA, Jerdon.

Jerdon says of this Flycatcher that "it has hitherto (1862) only been found on the summit of the Neilgherries and highest mountains of Ceylon." I can find no record, however, of this species occurring in Ceylon except that by Layard, who says he saw a drawing made by Mr. E. L. Mitford from a specimen he obtained at Ratnapoora. This is in the low country and probably not a hundred feet above the level of the sea.

#### 124. EUMYIAS SORDIDA, Walden.

This species, distinguished and described by Lord Walden (Ann. Nat. Hist. 1870, p. 218), is very common at Nuwara Eliya. I have



observed it there at all seasons; but it appears to have been mistaken for *E. melanops*, Vigors, by both Kelaart and Layard, who evidently refer to it under that name in their catalogues. A specimen sent by Layard is in the British Museum, and is given in Gray's 'Hand-list' (4897) as *E. ceylonensis*, n. sp.?

The general colour of the head, back, and outer edges of the quill-feathers is a dark bluish grey; throat and breast more dingy, and becoming paler towards the vent; forehead and chin bright blue; wings and tail dusky.

Bill black; irides brown; feet black.

Ceylon.

#### 125. CYORNIS RUBECULOIDES, Vigors.

Probably only an occasional visitor to Ceylon. Layard records having obtained a few specimens in the north of the island in October 1851; and I have examined specimens from Ceylon in Lord Walden's collection. Examples of this species from Ceylon and Burmah differ from Indian birds in having the orange colouring of the breast running up the centre of the throat, a peculiarity pointed out to me by Lord Walden.

Ceylon, India, Burmah.

#### 126. CYORNIS JERDONI, G. R. Gray.

This species was at one time considered identical with *C. banyumas*, Horsf., from Java, and is given under that name by Jerdon in his 'Birds of India;' but it has been separated by Mr. G. R. Gray as distinct. It is a resident in Ceylon and not uncommon in the low country between Colombo and Kandy, but has not been recognized as being widely distributed. I have obtained specimens a few miles from Colombo in July. Mr. Legge describes the female as being brighter on the upper surface than the male, but this is not in accordance with what I have observed.

Bill black; irides brown; feet lavender (brown in dry skins).

S. India and Ceylon.

#### 127. ERYTHROSTERNA HYPERYTHRA, Cabanis. (Plate XVII.)

This Robin Flycatcher was described in 1866 by Cabanis (Journ. f. Orn. p. 391) from a specimen sent from Ceylon by my friend Mr. Nietner; and that example (in the Berlin Museum) was, I believe, the only one until now which had been brought to Europe. I was fortunate enough to obtain two specimens of this species at Nuwara Eliya in February 1870, and I have no doubt that it is not uncommon on the hills at that season. Mr. Nietner probably obtained his bird on his estate about 2000 feet below Nuwara Eliya; and further inquiries may perhaps lead to its discovery on the Neilgherries\*.

\* Since the above was written, a specimen of this Flycatcher has been sent home from Goona, Central India. It is in full breeding-plumage, and was supposed to be *E. parva*. It is very probable that these two species have been confounded when not in full plumage, and that *E. hyperythra* is not so rare or so local as appears to be the case at present.



The distinguishing characters of the species are the rich orange-brown of the throat and breast, and the black stripe running from the bill down the sides of the neck to the breast and terminating below the bend of the closed wing. The specimens I obtained were both males, adult and immature; and the above characters are distinct in both, but much more so in the older bird.

These birds frequented low thin jungle; and I did not hear them utter any note.

Bill dusky above, yellow beneath; irides dark brown; feet purplish brown.

Ceylon, Central India.

Jerdon mentions that *E. leucura* is found in Ceylon; but I cannot find any special record of its occurrence there. It may have been confounded with *E. hyperythra*.

#### 128. BRACHYPTERYX (?) PALLISERI, Blyth. (Plate XVIII.)

Peculiar to Ceylon. The generic position of this bird is not very clear. It was placed by Blyth doubtfully in *Brachypteryx*, but differs from the birds of that genus in the sexes being alike in colouring and in the well-developed tail. I believe it will require generic distinction; but for the present I shall leave it in *Brachypteryx*.

It is a species confined to the upper hills, and is by no means uncommon in the Nuwara Eliya district; but, from its habits, it is not an easy bird to watch or to obtain. It frequents the low brushwood in the true jungle, creeping about the stems of the underwood close to the ground, and may sometimes be seen busily examining the dead branches of some fallen tree. Frequently it betrays its close neighbourhood by its "cheep" once or twice repeated; and it will show itself for a moment within two or three yards of one; then it is lost again in the thick jungle. By giving up a good deal of time I succeeded in obtaining a few specimens; but I have often been out for many hours without being able to get a shot, although I have occasionally heard the bird close to me. It will sometimes show itself on a jungle-path; but it then keeps close to the side, turning over the dead leaves in search of insects, and disappearing on the slightest alarm. When on the ground it often jerks its tail up after the manner of the Robins; but I have not observed this habit when it has been on the stems of the jungle plants or creeping about the dry sticks. The sexes are alike in colouring. I have one specimen which on dissection proved to be undoubtedly a male; and it could not be distinguished by any external character from the female. Two other birds, of different sex and evidently young, were also alike, and differed from the adults only in the absence of the rusty throat and dark grey cheeks, and in having the tail shorter. I have been unable to ascertain any thing of the nesting-habits of this species; and the bird itself is exceedingly rare in collections.

The whole upper surface is of a dark olive-brown, the wings, rump, and tail being of a richer brown tint; chin and throat pale rusty, beneath the eye and the ear-coverts dark greyish; the underparts pale olive, becoming brown at the flanks, vent, and under tail-coverts.



Bill dusky above, dark grey below; irides pale buff; feet dark flesh.

Ceylon (upper hills).

129. *ARRENGA BLIGHI*, n. sp. (Plate XIX.)

In the adult, or perhaps nearly adult, male the whole head, nape, and throat pure black; back, wing-coverts, and breast black strongly glossed with indigo; carpal joint dark smalt-blue; wings, tail, rump, flanks, and abdomen dusky brown, the two last slightly rufous. The upper tail-coverts, rump, and flanks are tinged with blue; and it is not improbable that in an older bird these parts may become of the same colour as the back and breast. In the young the whole bird is brown, darker on the upper surface and more rufous below, the feathers of the forehead, throat, and breast centred with yellow-brown, and there is an indication of blue on the carpal joint.

The dimensions of the adult male are:—length 8 inches, wing 4·4, tail 3·5, tarsus 1·4, bill at front 0·6.

Bill black; irides greyish; feet black.

An adult female, shot by Mr. Bligh, but almost knocked to pieces, had very much the character of a young bird of the same sex I obtained at Nuwara Eliya (fig. 2); and the wing-spot was brighter, but not of so deep a blue as in the male.

The only example of this new species of *Myiophonus* I saw in Ceylon was the immature bird I obtained at Nuwara Eliya in July 1870; and the tinge of blue on the wing led Mr. Samuel Bligh of Ceylon to the opinion that it was the young of a species he had shot on the hills two or three years before, and which had been sent with other skins to Mr. Master of Norwich. By the kindness of that gentleman I have been able to examine his specimen and compare it with the one I myself obtained. There is no doubt of their belonging to the same species; and as it has hitherto been unknown I have named it after my friend Mr. Bligh, who procured this first specimen of what is entirely a new form in the island.

Some credit is due to Mr. Edward Blyth for his remarks on the absence of certain birds on the Ceylon hills. He says ('Ibis,' 1867, p. 312) "That *Myiophonus horsfieldi* (or a specialized representative of this bird) has not been observed in the island is worthy of notice; but I have before expressed an opinion that the higher regions of Ceylon have not yet been sufficiently explored." At the time Mr. Blyth wrote this the first specimen of the Ceylon *Myiophonus* was probably on its way to England; and its true character has only now been recognized. Its nearest ally is *A. cyanea*, Horsf., from Java.

The habits of the Ceylon bird correspond, so far as is known, with those of the other *Myiophoni*. The young bird I procured at Nuwara Eliya was killed on a low branch of a jungle tree close to a little mountain-stream; and Mr. Bligh, who obtained his specimens at an elevation of between 4000 and 5000 feet, told me he had never met with the bird excepting in the immediate neighbourhood of water-courses. He writes me that although he has seen this species several



times it is very difficult to obtain. The bird frequently perches on a rock in the midst of some mountain-torrent, but is very impatient of observation. On these occasions it "gives utterance to a peculiarly long-drawn, plaintive though loud whistling note; at the same time the body is dipped and the tail slightly raised." It soon seeks shelter under the dense jungle foliage.

### 130. PITTA BRACHYURA, Linn.

Generally distributed in Ceylon during the winter months, and at that time very abundant at Aripo. Although most of these birds seen in Ceylon are probably visitors from India coming in October, I have reason to think some of them are residents, as I have frequently heard and more than once seen them at Nuwara Eliya in August. My house at Aripo was surrounded by *Suriya* trees, the branches of many of them touching the roof of the veranda; and to these trees the Pittas used to come every evening shortly before sunset, perching about six or eight feet from the ground and continually repeating their cry of "A-vitch-i-a" (the name given to the bird by the Singhalese), which was frequently followed by a low hissing scream. On being alarmed by my too close approach they would fly direct to the hedge about thirty yards distant and hide themselves under the darkest and thickest part of it. A frequent attitude of this bird when perched on a stout branch of a tree was with the head and body stretched up to the full height, the legs straight, and the tail turned upwards.

Bill orange, tip dusky; irides brown; feet flesh-colour.  
Ceylon, India.

### 131. GEOCICHLA LAYARDI, Walden.

Peculiar to Ceylon. A single example of this Thrush was sent to Lord Walden in a collection of birds from the island. From what I heard in Ceylon from the person who made the collection I have no doubt this bird was obtained on the hills on the south-east side of the island, a part of the country which has not yet been properly examined and is likely to produce more novelties. This bird is described as more nearly allied to *G. citrina* of North and Central India than to *G. cyanota* of Malabar, with the orange colour of the underparts brighter and richer than in *G. citrina*, but not nearly so deep as in *G. rubecula* of Java.

The colours of this specimen are rich orange on the head, neck, and underparts, bluish grey above, and a white spot on the wing.  
Ceylon.

### 132. TURDULUS WARDII, Jerdon.

Generally a rare bird in Ceylon; but Mr. Laurie tells me it is not uncommon during the north-east monsoon in some of the hill-forests. I have seen specimens collected by that gentleman and others from the Kandy district, but have not met with the bird alive.

Ceylon, India.



133. *MERULA KINNISI* (Kelaart), Blyth.

Peculiar to Ceylon, and, I believe, confined to the upper hills. It is very common at Nuwara Eliya, frequenting alike the edges of the jungle and the gardens of the English houses, and often building in the stables and outhouses. It has the habits generally of the English Blackbird; but its song is by no means so fine.

The male has the whole upper surface black with a bluish-grey tinge, the underparts more dingy; the female has the upper colour less intense, and is dark ashy brown below. Young birds have the head and back brown, with the throat and breast mottled, the feathers being pale-centred and with dark brown tips.

Bill bright orange (adult), yellowish brown (young); irides brown; orbits yellow; feet yellow.

Ceylon.

134. *OREOCINCLA NILGIRIENSIS*, Blyth.

This handsome long-billed Thrush was described by Layard under the name of *Zoothera imbricata* from a specimen received from Mr. Thwaites, who probably obtained it on the hills. It has since been recognized as the above species. I have examined two skins sent home by Mr. Bligh, and Layard's specimen now in the British Museum; and the scale-like appearance of their plumage, arising from the black border to each feather, is well marked.

"Bill corneous; legs brown" (*Layard*).

Ceylon; Neilgherries.

135. *OREOCINCLA SPILOPTERA*, Blyth.

Peculiar to Ceylon. This is quite a jungle bird and not very uncommon in suitable places on the hills. Many specimens have been procured in wild country not far from Kandy, and in the forest-land adjoining the coffee estates between 2000 and 5000 feet high. I have not met with it at Nuwara Eliya.

Bill black; irides brown; feet pale brown.

Ceylon.

136. *PYCTORHIS SINENSIS*, Gmel.

Layard observed this bird in widely separated localities in the low country, but does not speak of it as numerous. I have seen a specimen in the possession of Mr. Legge, R.A., at Colombo, which I believe he told me was killed near his house; and I have seen others from the Kandy country.

Ceylon, India to Burmah and China?

137. *ALCIPPE NIGRIFRONS*, Blyth.

Peculiar to Ceylon. This little bird is well distinguished from the allied species *A. atriceps* by the greater part of the head being brown, the black being confined to the forehead, and a broad streak through the eye to the ear instead of covering the whole top of the head. I have not seen this bird in the north of Ceylon; and Layard does



not say where he discovered it ; but it is abundant in the central and probably in the southern districts. It is, however, somewhat migratory within the island, and it is difficult to say to what cause its irregular movements are due. I have shot it both near Colombo and at Nuwara Eliya in January, and have found it abundant at the latter place in July, August, and September ; then it has entirely disappeared. It is an amusing little bird, usually found in small parties and frequenting underwood and low thick bushes, or creeping among the stems of the taller jungle-plants, occasionally coming to the edge of a path and betraying its presence by an angry hissing note, evidently intended to warn off intruders.

Bill dusky above, pale flesh-colour beneath ; irides golden ; feet purplish flesh.

Ceylon.

### 138. DUMETIA ALBOGULARIS, Blyth.

This species is said by Layard to be confined to the vicinity of Colombo ; and although it is unlikely to be so purely local, I certainly never saw the bird alive until I became a resident close to the cinnamon-gardens in which he observed it. Like the following species it will probably be found in bush jungle in the interior as well as in the immediate neighbourhood of Colombo.

Ceylon, S. India.

### 139. DRYMOCATAPHUS FUSCICAPILLUS, Blyth.

Peculiar to Ceylon and rarely met with. I only know of three\* specimens having been obtained—two of them by Layard in Colombo and in the central road leading from Kandy northwards, and one (a male) by myself also from the latter part of the island. I found this bird among thick underwood in forest-jungle by the side of the road on which I was travelling ; and it was perched within two feet of the ground when I had my first fair view of it as with outstretched neck and swelling throat it poured forth a torrent of babbling notes.

I have restored this bird to its original position in the genus *Drymocataphus*, as its bill does not agree in form with that of *Pellorneum*, and the fifth quill-feather is the longest, the fourth and sixth being equal and slightly shorter. The colour of the back, wings, and tail dark olive-brown, the last tipped rufous ; wings and tail in some lights showing distinct transverse striæ ; crown rich dark brown, the feathers slightly pale-shafted ; lores, cheeks, sides of neck, and all the underparts pale rufous brown, the breast being rather darker.

Bill dusky above, flesh-colour below ; irides red ; orbits yellow ; feet pale flesh.

Ceylon.

### 140. POMATORHINUS MELANURUS, Blyth.

Peculiar to Ceylon ; rather local in its distribution, but generally numerous where it is found. It is very abundant at all times of the

\* I have since seen a fourth, which was procured a few years ago by Mr. Bligh from the hills.



year at Nuwara Eliya and in the surrounding district, frequenting the primitive jungle with which the upper hills are covered. It is also found occasionally in wild country near Kandy, and was first seen by Layard "in low, scrubby, and almost impenetrable brushwood" a few miles from Colombo. It was probably not far from this last locality that I also met with it, in the low country, a wild district of no great extent, to which I have referred in my notice of *Harpactes fasciatus*. Like its congeners, however, this Scimitar-bill is essentially a hill bird. It creeps about underwood and the lower branches of trees, half opening and closing its wings, and assuming various kinds of strange attitudes. It is at all seasons noisy; and just about the pairing-time in February the cries of a party of these birds remind one more of a concert of Cats than any thing else. It is to this species the name of Gamut-bird is often applied, from the powerful notes of the male beginning very low and running up the scale; they have a very striking sound when heard amid the silence of the deep jungle.

The colour of the sexes is alike. The back, wings, flanks, vent, and under tail-coverts rich olive-brown with a rufous tinge, especially on the flanks; from the base of the upper mandible to the nape black, extending to the mixed olive-brown and black on the top of the head: throat, breast, middle of abdomen, and a conspicuous supercilium pure silky white; tail blackish brown. The young bird is much more rufous generally, and has the ear-coverts and the sides of the neck and breast quite rusty.

Lord Walden has a series of specimens of *Pomatorhinus* the localities of which are not very intelligible on the labels; but the birds were probably obtained in the south or south-east of the island. All these have the upper surface quite rufous, extending also to the tail. This colouring is not found in one of the many specimens I have from Nuwara Eliya, and is so marked as almost to justify a specific distinction.

Bill yellow, with the base dusky above; irides dark red; feet lead-colour.

Ceylon.

#### 141. GARRULAX CINEREIFRONS, Blyth.

Peculiar to Ceylon. This species is confined to the southern half of the island, frequenting the lower hills, and, according to Layard, "it much resembles the *Malacocerci*, hunting in small parties and incessantly calling to each other." It is not uncommon in the Kandy district and in the hilly country between that and Galle. I have examined a great number of specimens of this species, and have found them agree very closely with each other; but they differ so materially in dimensions from those given by Blyth that I can only suppose he had but one example before him, and that an immature bird. This impression is confirmed by the specific name *cinereifrons*, given by him, and agreeing with his description "forehead and cheeks pale ashy;" whereas the birds I have examined have the whole top of the head ashy, that colour often extending over the nape, as well as



the cheeks, which are paler than the rest of the head; chin albescent, becoming rufous on the throat; in other respects the colours agree with Blyth's description. The dimensions of a specimen I obtained at Kandy, and which is not at all unnaturally stretched out, but fairly represents an adult bird, measures fully 10 inches instead of 8·5; the other comparative dimensions are:—wing 4·75, 4·5; tail 4·5, 4; bill to gape 1·3, 1·25; tarsus 1·5, 1·25.

Bill black; irides buff; feet dusky.

Ceylon.

#### 142. MALACOCERCUS STRIATUS, Swains.

A comparison of specimens of *M. striatus* I obtained in Ceylon with *M. malabaricus* in the Calcutta Museum left me in great doubt as to the reason for separating them specifically, and I cannot but think they will ultimately be included under the same name. The depth of the striæ in *M. striatus* varies with age; in a well-grown young bird there is not a trace of striæ on the tertiaries, and they are very indistinct on the tail. In a fully adult bird now before me the striation exactly agrees with Jerdon's description of that character in *M. malabaricus*: "the tertiaries are but very obscurely striated, but the tail is distinctly so." The distinctive character of *M. striatus* has hitherto been shown by comparing it with *M. terricolor*; but it should have been placed by the side of the Malabar species.

The Ceylon bird is universally distributed over the low country, frequenting alike the jungle, half-cultivated ground, and the gardens and compounds in Colombo. Its manners are the same as those of the common Indian species. I have found it nesting at Aripo in January.

Young birds are slightly rufous.

Bill pale yellow; irides pale buff; feet pale yellow.

Ceylon, S. India?

The only record I can find of the occurrence of *M. griseus*, Gmel., in Ceylon is in the 'Appendix' to Kelaart's 'Prodromus Faunæ Zeylanicæ' (p. 45), where, in a report by Mr. Blyth on a collection of Ceylon Mammals, Birds, Reptiles, and Fishes, and, I presume, made to the Asiatic Society of Bengal, the following appears among the list of birds:—

"*Malacocercus griseus* (Lath.), var.—Resembling the species of S. India, except that the head is concolorous with the rest of the upper parts."

I have neither seen nor heard of the true *M. griseus* in Ceylon.

#### 143. LAYARDIA RUFESCENS, Blyth.

Peculiar to Ceylon, and tolerably common in the wilder parts of the low country in the southern half of the island. It was formerly considered a hill species; but I believe it only visits the upper hills during the cold season. I have only found it at Nuwara Eliya at the beginning of the year; but it is at all times to be met with a few miles from Colombo where there is jungly or half-cultivated land.

PROC. ZOOLOGICAL SOCIETY.—1872, No. XXIX.



It keeps in small parties, and has generally the habits of the other *Malacocerci*.

Bill dull orange ; irides white ; feet yellow.

Ceylon.

144. *HYPsipetes ganeesa*, Sykes.

As *H. neilgherriensis*, Jerdon, is now united with *H. ganeesa*, Sykes, the Ceylon birds will come under the latter title.

This species in Ceylon is, I believe, confined to the hills, and is most abundant at a moderate elevation. I have only seen it at Nuwara Eliya in February ; but it is tolerably common in jungle from the Kandy country to about 5000 feet. I have generally found it in small parties on rather low trees.

Ceylon, S. India.

145. *CRINIGER ICTERICUS*, Strickland.

Layard says of this bird that it "abounds in the mountain zone." This probably means the lower ranges, as he tells me he has never visited the Nuwara-Eliya district, and he does not profess to know the hill birds. Kelaart, on the other hand, who specially collected the birds of the upper hills, says it is "a common species in the low country." I have no doubt Layard was right in suggesting that Kelaart mistook the common *Ixos luteolus* for this species ; and this is confirmed by Mr. Legge's observation that *Criniger ictericus* "is strictly a jungle bird" (J. R. A. S., Ceylon Branch, 1870-71, p. 43). Mr. Legge, however, whose knowledge of the low country at the time he wrote was confined to the western province, says "Kelaart wrote correctly of this bird ;" but "a strictly jungle bird" can hardly be described as common in a district principally consisting of paddy-fields and cultivated land.

I have only obtained this bird once in the neighbourhood of Colombo, among trees in a native village ; it is most numerous in forest country on the lower hills, as is the case with this species in India.

Bill black ; irides red ; feet dark leaden.

Ceylon, Malabar.

146. *IXOS LUTEOLUS*, Less.

This bird, the *Pycnonotus flavirictus* of Strickland, is one of the commonest species in the low country. It is equally abundant at Aripo and Colombo wherever there are low bushes, and has a hurried twittering song of a few notes, loud and frequently repeated.

Bill black ; irides red ; feet blue-black.

Ceylon, South and Central India.

147. *KELAARTIA PENICILLATA*, Blyth.

Very abundant at Nuwara Eliya and on the upper hills, frequenting low bushes and thin jungle.

The general colour of this bird is dark olive-green above and greenish yellow below, brighter yellow on the throat, middle of



abdomen, and under tail-coverts; head black in front and shading into olive-green at the nape, with the feathers tipped paler or white; chin and a narrow vertical stripe on each side of the forehead white; lores and cheeks black, paling to leaden grey behind, with a yellow spot below the ear and a tuft of bright yellow feathers springing from immediately behind the eye and directed backwards. These tufts stand out from the sides of the head when the bird is alive, and add much to its generally handsome appearance.

This species was described from Ceylon specimens; but is believed by Jerdon to be "identical with one procured from the Mysore country below the Neilgherries, which was accidentally destroyed," but from which a coloured sketch was made.

Bill black; irides red-brown; feet leaden.  
Ceylon, S. India?

148. *RUBIGULA MELANICTERA*, Gmel.

Peculiar to Ceylon, and tolerably common in the low country and lower hills of the central and southern portions of the island. I have obtained specimens near Colombo and close to Kandy.

The colour of the upper surface is olive-brown, and of the under parts bright yellow, with the flanks tinged with dull olive; top and sides of the head black; quills brownish black, with the outer edge olive, and tail dingy black, with all but the central feathers tipped white.

Bill black; irides red; feet purplish black.  
Ceylon.

149. *PYCNONOTUS HÆMORRHOUS*, Gmel.

Very common all over the low country, and less so on the lower hills. I have never seen it at Nuwara Eliya or above 5000 feet, and I am inclined to think it is only a seasonal visitor to that elevation. I have found this species breeding in December at Aripo; its cup-shaped nest was placed under the eaves of my bath-house out of doors, and supported by the sticks of which the rough framework was constructed.

Bill black; irides dark brown; feet leaden black.  
Ceylon, South and Central India.

150. *PHYLLORNIS JERDONI*, Blyth.

Common in the low country. I have obtained it on several occasions at Aripo, near Colombo, and quite in the south. It generally keeps among the upper branches of the trees.

Bill slate; irides brown; feet lavender.  
Ceylon, India.

151. *PHYLLORNIS MALABARICUS*, Lath.

Recorded by Layard and Kelaart from the hills; and I have seen a specimen obtained by Mr. Laurie. I believe it is rather rare in Ceylon.

Ceylon, South and Central India.



152. *IORA ZEYLONICA*, Gmel.

This species is well known in the low country ; it is very abundant at Aripo at all seasons, and almost as common about Colombo. It breeds at Aripo at the end of the year ; and I have obtained it in November with hardly a trace of green on the black back. Its notes are very much varied ; and some of them sound as if uttered at a considerable distance when the bird is really within a few yards.

I regret that I did not know whilst I was in Ceylon of the question as to *I. typhia*, Linn., being found in the south of India and Ceylon. As I brought home no male specimens of *Iora* which were not in such a state of plumage as to leave a doubt about their belonging to *I. zeylonica*, I shall not include *I. typhia* among the Ceylon species ; but I have a very strong impression, partly based on my recollection of a pair of birds with dull green backs which for several days frequented some shrubs close to a house where I was staying, a few miles from Colombo, that *I. typhia* is found in Ceylon. I had no doubt of it at the time, as the male of *I. zeylonica* should then (February), according to my observations, have the back nearly or entirely black.

In case this paper should fall into the hands of any one collecting in Ceylon, but who is not familiar with the distinctive characters of the two species of *Iora*, I may mention that the females are at all times practically alike. In the breeding-season the male of *I. zeylonica* has the back entirely black or, more frequently, black and green irregularly mixed, the colours being in patches and not generally blending with each other ; at the same season the male of *I. typhia* has the back wholly green, contrasting with the black wings, which in both species have two white bars. A further distinction is said to exist in the colour of the irides (this would hold good at all seasons), those of *I. zeylonica* being grey and those of *I. typhia* light hazel ; I can answer for the former being correct.

Bill slate ; irides grey ; feet dull leaden.

Ceylon, S. India.

153. *IRENA PUELLA*, Latham.

Layard and Kelaart have each recorded an example of this species, both from near Kandy.

Specimens of this bird from Ceylon are much desired for comparison with those from India. The male has the whole upper parts and under tail-coverts bright cobalt-blue ; wings, tail, and lower plumage deep velvet-black. The female is of a dull, slightly mottled Antwerp blue throughout. (*Jerdon.*)

Ceylon, Malabar, Assam, Arracan, Burmah.

154. *ORIOLOUS INDICUS*, Jerd.

I include this species of Oriole on the authority of Layard, who speaks of a pair of these birds having been shot near Colombo, and coming under his notice.

Ceylon, India.



155. *ORIOLOUS CEYLONENSIS*, Bonaparte.

Generally distributed in the low country. I have met with it commonly at Aripo, Colombo, and in the south; but I have no reason to think it ascends above the lower hills. The young bird has the back pale dirty yellow, purer on the rump; top of the head brownish black, becoming streaked on the cheeks and strongly so on the throat and under neck; quills margined externally with whitish, and the colours generally very much less pure than in the adult. The bill in the young is black.

Bill deep flesh-colour; irides red; feet leaden.

Ceylon, S. India.

156. *COPSYCHUS SAULARIS*, Linn.

Abundant in the low country, and rarely found far from native villages or the houses of English residents. The familiarity of the "Magpie Robin" makes it a general favourite; and whether when perching on the roof of the house (a frequent station for it when singing) or furiously attacking some intruding rival, there is always something attractive in this showy and well-known species. During the last hour before sunset these birds become very noisy and frequent fights take place between the cocks, two or three of them going through a sort of tournament before the hen bird which has taken up her quarters in the neighbourhood. It is at this time the cocks put themselves in such strange attitudes, turning back the tail till it almost touches the head, as Layard mentions; but Jerdon says he has never observed these performances, which from my own observation I should say are regularly gone through every afternoon; the birds frequently utter a harsh kind of scream; and this goes on until the sun disappears and the quickly following darkness puts an end to the proceedings.

Good specimens of this Robin are very difficult to obtain at Colombo, unless immediately after moulting; as the birds soon become discoloured with the red soil, and the tails rapidly worn out at the end.

Females of this species from Ceylon have the back darker than those from Burmah, and perhaps from India generally, but they do not differ from a Madras specimen in the British Museum.

The young birds are greyish brown above, with the throat and breast mottled with dark brown on a paler ground, and the bill dusky.

Bill black; irides brown; feet dark leaden.

Ceylon, India, Arracan, Tenasserim, S. China, Hainan.

157. *KITTACINCLA MACRURA*, Gmel.

This bird is confined to wild jungly districts in the low country and on the lower hills. In such localities it is numerous and its fine song may be constantly heard in the morning and evening. It is abundant in the wilder parts of the northern road from Kandy; and I have also heard it occasionally in a piece of thick jungle close to



Kandy itself. It usually perches low ; and from its habit of frequenting dense jungle, it is often difficult to obtain sight of.

Bill black ; irides brown ; feet flesh-colour.

Ceylon, India, Assam, Burmah, Malacca, Hainan.

158. *THAMNOBIA FULICATA*, Linn.

Common about houses and outbuildings, and, I believe, generally distributed through the lower parts of the country. I have seen them more numerous in the north than elsewhere ; and they were always about my house at Aripo, frequently coming into the veranda, and generally very tame.

Bill black ; irides brown ; feet black.

Ceylon, S. India.

159. *PRATINCOLA CAPRATA*, Linn.

Layard and Kelaart both mention having obtained this species on the lower hills ; but I have never met with it, either alive or as a skin.

Ceylon, India, Burmah, Malaya, Philippines.

160. *PRATINCOLA ATRATA*, Blyth.

Very common at Nuwara Eliya and on the upper hills. It frequents gardens rather than jungle ; and the top of a rhododendron bush is a favourite station for the male, which always chooses a conspicuous position when it sings its short Robin-like song. Young males at first have the general brown plumage and rufous rump of the female ; the change to the pure black and white of the adult male is very gradual, the quills and rump being the last to assume the mature colours.

Bill black ; irides brown ; feet black.

Ceylon hills, Neilgherries.

161. *LARVIVORA CYANA*, Hodgson.

This bird is a winter visitor to Ceylon. Layard obtained specimens in October in the extreme north ; and I procured adult and immature examples of both sexes at Nuwara Eliya in January, February, and March. It was at that time tolerably common on the hills ; but I have not met with it at any other season.

The female is olive-brown above ; underparts rufous, paler on the throat and centre of abdomen ; under tail-coverts white. These particulars, taken from one of my Ceylon specimens, agree with Hodgson's last description of the colours in the female.

Bill dusky ; irides brown ; feet flesh-colour.

Ceylon, India (generally on the hills).

162. *CYANECULA SUECICA*, Linn.

Layard obtained this species in March in one of the coffee-districts. I have not met with it.

Ceylon, India, N. and W. Asia, N. Europe.



163. *ACROCEPHALUS DUMETORUM*, Blyth.

This is apparently the bird given by Layard as *Phyllopneuste montanus*, Blyth.

Generally distributed; it is a winter visitor and numerous in Ceylon at that season. I have killed it at Aripo, Colombo, and Nuwara Eliya. All my specimens have the greenish shade on the upper surface mentioned by Blyth as found in the birds from Ceylon.

Bill dusky above, pale flesh below; irides brownish yellow; feet in different specimens pale brown to purplish flesh.

Ceylon, India, Nepal, Assam.

164. *ORTHOTOMUS LONGICAUDA*, Gmel.

Common in all parts of the island, but especially frequenting gardens and the neighbourhood of habitations. It is as abundant at Nuwara Eliya as at Aripo or other parts of the low country. I have examined many of these birds from different localities, and have found them to agree in all respects with Jerdon's description of this species, except in the length of the tail; this in Ceylon birds I have never found to exceed  $2\frac{1}{4}$  inches.

Bill dusky flesh; irides yellow; feet flesh.

Ceylon, India to Burmah, S. China.

165. *PRINIA SOCIALIS*, Sykes.

Layard found this species in the extreme north; and I believe Mr. Legge discovered it nesting in a patch of Guinea grass close to his house at Colombo. It will probably be found in suitable situations in other parts of the island.

Ceylon, S. India.

166. *CISTICOLA SCHÆNICOLA*, Bonap.

*Cisticola homalura*, Blyth?

I place these two species together as it is difficult to speak of them separately, in consequence of the confusion existing between them, if they are really distinct. Layard says of "*C. cursitans*, Blyth" ( $= C. schœnicola$ , Bonap.) that it "is much less common than *C. homalura*; and though found in the same locality, it frequents trees and jungle." This bird surely cannot be a *Cisticola*. Kelaart says of *C. cursitans*:—"frequents the grass-plains; very common at Trincomalie." I can confirm this statement and say precisely the same of it at Colombo; it is common there wherever there is a patch of long grass.

*C. homalura* was discovered by Layard in paddy-fields near Galle; he "subsequently found it sparingly about Colombo, and abundantly in fields of gingelle (*Sesamum orientale*) at Pt. Pedro." Kelaart says it "is found in great abundance on Horton Plains and Nuwara Ellia," these last localities resembling each other in being elevated grass-plains surrounded by forest-jungle.

I am almost ashamed to think of the number of specimens of *Cisticola* I have shot at Nuwara Eliya in the hope of getting one of



*C. homalura*, which Blyth says ('Ibis,' 1867, p. 302) "differs from *C. schœnicola* in having a stouter bill, the whole upper parts much darker, and the tail almost even, except that its outermost feathers are .25 inch shorter than the next;" but, except in some considerable variation occasionally in the depth of the general rufous tint, there was nothing to distinguish them from the grass-frequenting species at Colombo. Mr. Layard tells me that the fine collection of Ceylon birds he brought to England is now in such a state as to be useless for scientific purposes; and as I can obtain no specimens of *C. homalura* for examination, I must regard that species as very doubtful until further evidence is procured from the localities whence Layard obtained his birds.

*C. schœnicola* from Ceylon agrees with the European bird in size, and is larger than the Indian representative; it has, however, the same decided markings as the latter form, and they are even more conspicuous. The dimensions given by Jerdon are greater than those of any of the Indian specimens I have examined.

Bill dusky above, flesh below; irides pale yellow; feet flesh-colour.

Ceylon, India to Europe, Africa, China, Hainan, Formosa.

#### 167. *DRYMOIPUS INORNATUS*, Sykes.

The difficulty in determining the species of *Drymoipus* is so well known that it may prevent additional confusion if I mention that the three species included in this list of Ceylon birds have been compared with specimens in the British Museum, and satisfactorily identified with the species there labelled with the names I have given. The identification of at least one of the two Colombo species by myself and Mr. Legge whilst I was in Ceylon was not correct; and it is uncertain to which of them Mr. Legge's observations (J. R. A. S., C. B., 1870-71, p. 50) refer.

I believe *D. inornatus* is not uncommon about Colombo; but the only specimen I brought to England came from Kandy, and agreed with those in the British Museum in having the lores, throat, and cheeks whitish, the whole under surface and flanks very light, with a dull yellowish tinge, and a rather broad subterminal dusky band of uniform tint on the under surface of the tail-feathers. The bill is rather slight and black, with the base of the under mandible abruptly pale (dried skin). The wing exactly 2 inches. Jerdon says of this species, "in no case does the wing ever come up to 2 inches, more generally  $1\frac{3}{4}$ ." I cannot think, however, there is any doubt about this specimen being *D. inornatus*. Layard says the eggs of this species are "verditer, with purplish blotches and wavy lines;" Mr. Legge gives "ground-colour clear blue-green, clouded here and there, or blotched mostly towards the obtuse end, with sepia." It is doubtful to which species either of these gentlemen refers.

#### 168. *DRYMOIPUS JERDONI*, Blyth.

The common Ceylon species, of which I have obtained specimens close to Colombo, agrees perfectly with *D. jerdoni*, Blyth, in



the British Museum, where there is a specimen named and sent by Dr. Jerdon himself. In the 'Birds of India,' vol. ii. p. 180, Jerdon mentions that Blyth described this species from specimens he sent him from Southern India; but he afterwards absorbed it into *D. longicaudatus* in the belief that the specimen he described was in imperfect plumage. Jerdon further says:—"It appears to me very similar to some Ceylon birds which Mr. Blyth doubtfully considered identical with *D. inornatus*."

My Ceylon birds are greyish brown on the upper surface, rather paler on the head, cheeks, and neck; lores pale and much less conspicuous than in *D. inornatus*; under surface pale fulvous, and flanks rather dusky; the upper surface of the tail-feathers distinctly striated, the striæ showing as faint narrow bars on the under surface, which has a narrow dark subterminal band, generally darker in the centre, and giving the appearance of a spot. In fresh specimens the bill is dusky above, fleshy below; irides pale yellow; feet flesh-colour.

Length 5·5 inches, wing 2·3, tail 2·5, tarsus ·8, bill at front ·4.

Ceylon, S. India.

#### 169. *DRYMOIPIUS VALIDUS*, Blyth.

This species, at first called *D. robustus* by Blyth, is peculiar to Ceylon, and, according to Layard (who discovered it), rather a rare bird. Mr. Legge and I were both mistaken in believing it common about Colombo, as I now find I did not see the species in Ceylon. A specimen in Lord Walden's collection, agreeing with another in the British Museum, has the bill entirely black, stouter and considerably deeper than I have seen in any other Ceylon species; top of the head, lores, and general upper surface dark greyish brown; beneath whitish, with a pale fulvous tinge; cheeks, sides of the breast, and flanks dusky. Length 6 inches, wing 2·4, tarsus 1, bill at front ·5.

The dry specimen has the bill black; tarsus yellow-brown (probably flesh-colour when alive); irides "light red-brown" (Layard). Ceylon.

#### 170. *PHYLLOSCOPUS NITIDUS*, Lath.

This bird is common at Nuwara Eliya in the cold season; and I have seen it also at Aripo.

Bill dusky above, flesh below; irides dark brown; feet pale brown. Ceylon, India.

#### 171. *PHYLLOSCOPUS VIRIDANUS*, Blyth.

Recorded by Layard, who also gives *Phyllopneuste montanus*, Blyth, which is probably a synonym of *Acrocephalus dumetorum*, Blyth.

#### 172. *SYLVIA AFFINIS*, Blyth.

I obtained one specimen of this species at Aripo in December. Layard also appears to have only met with it on one occasion.

Bill, base slate, tip dusky; irides pale yellow; feet dark leaden. In this species and very many others Jerdon has apparently given



the colour of the bill and legs from their appearance in dried specimens.

Ceylon, Central India.

173. *MOTACILLA MADERASPATENSIS*, Gmel.

Layard mentions having seen one specimen in a private collection in Ceylon.

Ceylon, India.

174. *CALOBATES SULPHUREA*, Bechst.

I have obtained this bird at Nuwara Eliya in the beginning of the cold season; it is better known on the hills than in the low country. Asia to Australia, Africa, Europe.

175. *BUDYTES VIRIDIS*, Gmel.

This is the common Wagtail in Ceylon, appearing with other migratory birds in October.

Ceylon, India, "N. Africa, S.E. Europe, and W. Asia" (*Jerdon*).

176. *LIMONIDROMUS INDICUS*, Gmel.

I have only seen this bird on wild jungle-roads between Kandy and Trincomalie; but Layard has apparently met with it in other localities.

Ceylon, India, Arracan, Burmah, and part of Malaya, China.

177. *CORYDALLA RICHARDI*, Vieill.

This species is numerous in winter on the "Galle face"—the esplanade at Colombo, and a great place of resort for Pipits, Wag-tails, and small Sand-Plovers at that season. It is no doubt, as Layard states, widely distributed; but I do not think it is a resident in Ceylon.

Ceylon, India, and Asia generally, Africa, Europe.

178. *CORYDALLA RUFULA*, Vieill.

Resident and very common in Ceylon; I have found it at Aripo, Colombo, and Nuwara Eliya; and I believe it is generally distributed throughout the island.

Bill dark brown above, yellowish below; irides brown; feet light fleshy brown.

Ceylon, India, Assam, Burmah.

179. *CORYDALLA STRIOLATA*, Blyth.

This bird is also common at Colombo in the winter. I have compared and identified specimens of this and *C. richardi* from Ceylon with birds in the Indian Museum at Calcutta.

Ceylon, India, China.

180. *ZOSTEROPS PALPEBROSUS*, Temm. (Plate XX. fig. 1.)

Common in the central and southern parts of Ceylon, but only ascending the hills to about 2000 feet. It frequents trees and



flowering shrubs, and, Mr. Legge says, is often to be seen on the tulip-trees in the principal street of the Fort at Colombo. It is common about Kandy and the surrounding district; but I have never met with it in the north or on the upper hills. Specimens of this *Zosterops* from the low country in Ceylon vary somewhat in size, but have been identified in England and Calcutta with *Z. palpebrosus*, and agree with Jerdon's description of that species except in being generally smaller and in the colour of the bill and legs. He says, "Bill blackish, horny at the base beneath; legs reddish horny;" but I find in freshly-killed birds the following colours:—

Bill dark leaden, paler at the base beneath; irides light brown; legs and feet lavender.

Ceylon, India, Assam, Arracan, Tenasserim.

181. *ZOSTEROPS CEYLONENSIS*, n. sp. (Plate XX. fig. 2.)

Upper surface dark olive-green, deeper on the head and paler on the upper tail-coverts; a circle of small white feathers round the eye; lores and below the eye dusky, but not very conspicuous; chin, throat, and centre of breast greenish yellow, shading at the sides of the neck and breast into the colour of the back, and giving the appearance of an incomplete pectoral band; the rest of the underparts bluish white, darkest on the flanks, and sometimes tinged in the centre with yellow; under tail-coverts yellow; quills and tail dusky brown, both margined externally with olive-green, and the latter faintly marked with transverse striæ. Sexes alike.

Length 4·75 inches, wing 2·4, tail 1·8, bill at front ·5, tarsus ·7.

Bill dark leaden above, paler below; irides light brown; feet lavender.

This is at all seasons one of the commonest birds at Nuwara Eliya and on the upper hills. It is, I have no doubt, the one recorded by Kelaart as *Z. annulosus*, Swainson, an African species. Layard, in speaking of this bird in his 'Notes on the Ornithology of Ceylon,' says: "Dr. Kelaart writes, 'we fear that the Nuwara Eliya *Zosterops* is wrongly identified; it is of a darker green than the common *Z. palpebrosus*.'" He then adds, "I, however, much doubt the distinctness of this and the preceding species." A comparison of the two birds, however, leaves no doubt that there is a marked difference between them, both in colour and in the form of the bill. The bird from the Ceylon hills cannot be identified with any recognized species; and Mr. A. O. Hume, to whom I showed specimens of it when I was at Calcutta, told me he had never seen it in any of his many collections from the Neilgherries, a district (as I have before mentioned) agreeing closely in character and productions with the Ceylon hills. Mr. W. T. Blanford, in a paper on the Birds of Western India (J. A. S. B. 1869, vol. xxxviii. p. 170), says, in speaking of *Z. palpebrosus*, "the Nilgiri race is a little larger and appears to be a little darker in colour." He gives as the measurements of a specimen, "beak ·4, wing 2·2, tail 1·75, tarsus ·7," and says "the black lores appear more developed in the Nilgiri bird." These observations evidently refer to *Z. palpebrosus*; but it appeared to me desirable to



mention them in my account of *Z. ceylonensis* for the purpose of showing the difference between the hill species of the two countries.

I believe Dr. Jerdon is under the impression that he has seen *Z. ceylonensis* in India ; but he has no record of it.

*Z. ceylonensis* differs somewhat in habits from *Z. palpebrosus*. It frequents hedges and bush-jungle rather than trees, clinging Tit-like to the stems, and often covering its forehead with pollen from the flowers which it busily examines for insects. As these birds are very common and constantly flying in small parties from bush to bush, uttering their lively chirp, they attract attention ; and the little "White-eye" is familiar to most Europeans who visit Nuwara Eliya. In the winter the males associate in flocks of fifteen or twenty ; and it is then rare to find a female in their company. I believe the latter are for the time solitary, as, with one exception, the numerous specimens I have shot from different flocks have proved to be males. The breeding-season is probably about April or May ; but I have been unable to obtain any particulars of their nesting.

The distinction between the two species of *Zosterops* found in Ceylon will be readily seen on reference to Plate XX.

#### 182. PARUS CINEREUS, Vieill.

Very abundant at Nuwara Eliya and on the upper hills at all seasons, and found occasionally on the western coast, around Colombo and not far from Galle. Layard says it is "not uncommon throughout the island," but I have never seen it in the Aripo district or in the extreme south. Like many hill birds it is often met with near Kandy ; but I expect its appearance about Colombo and in some other parts of the low country is exceptional, as when found there it is by no means numerous. It has the usual habits of the Titmouse family.

Bill black ; irides black ; feet leaden.

Ceylon, India (except Bengal), Malaya.

#### 183. CORVUS LEVAILLANTI, Less.

*Corvus culminatus*, Sykes.

General in the low country, and especially frequenting native villages and the more uncultivated districts in the interior. It is rare at Colombo compared with *C. splendens*, and was not so numerous as that species at Aripo. I believe Crows are unknown on the upper hills ; but I have heard of their having been occasionally seen for a day or two on coffee-estates 3000 or 4000 feet high.

Bill black ; irides dark brown ; feet black.

Ceylon, India to the Malay peninsula.

#### 184. CORVUS SPLENDENS, Vieill.

This well-known bird is much more numerous on the coast generally than inland, and is found in great abundance in all the large towns, but is not met with in native villages so much as the last species. It was common at Aripo ; and at Colombo it is very abundant, not confining itself to the shore, but boarding the vessels as soon as



they are anchored in the harbour or roadstead, paying frequent visits while they remain there, and only reluctantly leaving them at their departure when they are two or three miles away. It is unnecessary to say more of the well-known inquisitive, thievish habits of these birds than that in Ceylon they fully keep up the character they have obtained elsewhere. From the comparative localization of this bird in the larger towns in the south-west of Ceylon, Mr. Hugh Nevill has stated (Journ. Roy. Asiat. Soc., Ceylon Branch, 1870-71, p. 33) that "there is no doubt it is not indigenous to the south of the island, having been introduced by the Dutch at their various stations as a propagator of cinnamon, the seeds of which it rejects uninjured." I have been unable to discover on what evidence this statement has been made. This Crow has certainly been in Ceylon long enough to spread over every part of the island if its habits or inclinations had led it to do so; but on both sides of the island it is comparatively local; and whilst on a coasting voyage from Ceylon to Calcutta, and calling at numerous places on my way, I found on the Indian coast the same localization of this bird in the larger ports as is the case in Ceylon.

The Ceylon birds are smaller than those in India, and, according to Blyth, are darker, but I have not had an opportunity of comparing a sufficient number of specimens from the two countries to be able to judge on this point. Jerdon says nothing of the neck changing from ashy to a dull fawn-colour in old birds in India; but this is the case in Ceylon. The young birds are very dark on the neck; and these may possibly have been the subjects of Blyth's observations. Specimens of this Crow from Ceylon and India are now, however, in the Gardens of the Society, and will afford ready means of comparison of any changes that may take place.

Bill black; irides brown; feet black.

Ceylon, India, Assam, Burmah?

#### 185. *CISSA ORNATA*, Wagler.

Peculiar to Ceylon. This remarkably handsome species has attracted some attention since it was described by Blyth as *C. puella* from specimens forwarded by Layard; but it had been previously made known by Wagler. It is, so far as is known, essentially a hill bird, found most abundantly at about 5000 feet and upwards, but at certain seasons descending as low as 1500 feet. This is about the elevation of Kandy; and the jungles in the immediate neighbourhood of that city, nearly in the centre of the island, appear to be the lower limit of the range of this and many other hill species. In the cold season, which is only really perceptible on the hills, these birds are numerous at Nuwara Eliya, frequenting the dense bushes growing under the trees in forest-jungle. They are very noisy, continually uttering a harsh Jay-like scream, both when perched and flying. There is consequently little difficulty in finding them out when they are in the neighbourhood; but from their keeping so much to the dense jungle I have on several occasions worked my way quietly through the bushes to within a few yards of the birds without being



able to get sight of them. The specimens I obtained were apparently not quite mature, as the blue of the underparts was not uniformly developed; otherwise they were in good feather, and enable me to give a description of the species.

Whole head and neck rich deep chestnut; back, tail, and underparts (in adults) bright cobalt-blue, the tail-feathers tipped and more or less margined externally with white; quills light chestnut on the outer webs, black on the inner.

Bill red (adult), tipped black (young); irides light brown; feet coral-red.

Ceylon hills.

186. *ACRIDOTHERES TRISTIS*, Linn.

Very common in the low country, and generally distributed. They were very numerous at Aripo; and a young bird brought to me by some natives soon became tame enough to be allowed its liberty in the house, sometimes escaping through the window to the adjoining trees, but always allowing itself to be caught, or going into its cage when held up to it. It became rather troublesome at last from its fondness for standing on the top of my head or perching on my hand when I was writing or engaged in some other work at the table.

Ceylon specimens are much darker than those obtained in India.

Bill yellow; orbits yellow; irides dark brown; feet pale yellow.

Ceylon, India, Assam, Burmah.

187. *TEMENUCHUS PAGODARUM*, Gmel.

Obtained by Layard in the north of the island, and by Kelaart at Trincomalie.

Ceylon, India.

188. *TEMENUCHUS SENEX*, Temm.

Peculiar to Ceylon; described by Layard as *T. albofrontatus*, as it was believed to be new; it has since been recognized as *T. senex*, Temm., erroneously described by Bonaparte as from Bengal. Several specimens have been received by Lord Walden, of which, however, only one has the head entirely grey, the true character of *T. senex*. Layard gives the following description of his bird, which is now in the British Museum:—

“General colour of back, tail, and wings black with a green gloss; forehead albescent; hinder feathers of crest brownish black with albescent shafts; general colour of breast, throat, vent, and under tail-coverts albescent, the shafts of the feathers on the throat shining white.”

It is, I believe, from the lower hills, and appears to be rather a local species.

Ceylon.

189. *PASTOR ROSEUS*, Linn.

Layard “found large flocks of these birds” quite at the north of the island in July, but did not see them afterwards. They have also



been obtained at Putlam, on the west coast ; and I have little doubt that I saw a flock at Aripo in 1866, but I could not get near them. It is rather remarkable that this bird is not better known in Ceylon, as in India, according to Jerdon, it is most abundant in the south and south-west.

Ceylon and India westward.

190. *EULABES RELIGIOSA*, Linn.

Recorded by Layard as common on the west coast. I have never met with it at Aripo, and believe it is more frequently seen in the south. There are many Ceylon specimens in Lord Walden's collection, most probably procured in the south-east of the island.

Ceylon, South India.

191. *EULABES PTILOGENYS*, Blyth.

This well-marked species of hill Myna is peculiar to Ceylon, and is found in flocks on the upper hills chiefly, but sometimes met with in the neighbourhood of Kandy. It frequents the tops of the trees ; and at Nuwara Eliya, where it is often numerous, I have found it wild and difficult of approach. I have heard, however, of large numbers having been killed on some of the coffee-estates in the early morning or evening. Its call is constantly repeated when on the wing, and sometimes when perched on the tops of the trees.

This species may be readily distinguished by the yellow lappets at the back of the head, and the absence of any naked skin about the eye and cheeks.

Bill deep orange, base black ; irides brown, lappets yellow ; feet dull yellow.

Ceylon.

192. *PLOCEUS BAYA*, Blyth.

This bird, called by Layard *P. philippinus*, is said by him to be migratory and to breed in June. It was, however, generally to be found at Aripo ; and there they used to build their curious nests in December on the trees close to my house. A young bird was brought to me in February which was just ready to leave the nest. I have never seen the nest of this species in any other than ordinary branching trees ; but Layard says it builds on palms and other trees indiscriminately.

Ceylon, India, Assam, Burmah, Malaya.

193. *PLOCEUS STRIATUS*, Blyth.

I have not met with this species at Aripo or on the west side of Ceylon ; and Layard, who found it on the east side, thinks it is confined to that part of the island. It is rather remarkable, however, that this bird should not change its quarters according to the season and, like many other species, migrate from one side to the other at the change of the monsoons.

This is the species most probably given by Layard under *P. manyar*, Horsf., which is a Javan bird.

Ceylon, North and Central India, Burmah, parts of Malaya.



194. *MUNIA MALACCA*, Linn.

195. *MUNIA RUBRONIGRA*, Hodgs.

196. *MUNIA UNDULATA*, Latham.

197. *MUNIA STRIATA*, Linn.

198. *MUNIA MALABARICA*, Linn.

With the exception of *M. rubronigra*, which I have not seen, and was recorded by Layard only from Galle, the above species are more or less abundant in the low country—*M. undulata* and *M. malabarica* being the most numerous, and the former perhaps the most widely distributed. I have seen many nests of *M. undulata* at Aripo and near Colombo, and have often watched the birds biting off the grass-stems and taking them to the nest, which has been generally a large structure, sometimes placed near the end of a branch, but more commonly in a thick bush.

These species are more or less distributed through India and the neighbouring countries eastward of it.

199. *MUNIA KELAARTI*, Blyth.

Peculiar to Ceylon, and confined to the upper hills. It is abundant at Nuwara Eliya at all seasons, frequenting the gardens and cultivated ground, and may often be seen on the roads feeding, like the Sparrows, on what it can find there. I have specimens in all stages of plumage. The adult bird may be distinguished from *M. pectoralis*, Jerdon, with which it was at first confused, by its having the rump and underparts, from the breast downwards, brownish black, with each feather centred, barred, and margined with white, producing a mottled effect; the under tail-coverts are only centred white; and the extremity of the upper tail-coverts is tinged with glistening yellow. Young birds have the throat speckled brown and white, and the underparts faintly mottled with two shades of light yellowish brown.

Bill lead-colour, very dark in adults; irides brown; feet leaden. Ceylon.

200. *ESTRElda AMANDAVA*, Linn.

I have seen specimens of this bird which were procured by Mr. Legge from a grass-field adjoining his house at Colombo. It had not been previously observed in Ceylon; and it may be, as Mr. Legge suggests as possible (J. R. A. S., C. B., 1870-71, p. 53, note), that some of the many birds of this species imported into Ceylon have escaped from confinement and become acclimatized. The occurrence of *Munia rubronigra* (a North-Indian species) only about Galle may perhaps be accounted for in the same manner, if no mistake was made in its identification.

Ceylon, India, Assam, Burmah.

201. *PASSER INDICUS*, Jard. & Selby.

Found in Ceylon wherever there are human habitations. It is



abundant at Nuwara Eliya, but I was told by old residents that they remembered the time when the now common Sparrows and Musquitos were unknown at that elevation.

Ceylon, India, eastward to Siam.

202. *MIRAFRA AFFINIS*, Jerdon.

I have found this species common at Aripo; and Layard has recorded it from the north also. I am not sure that it does not also occur at Colombo.

Bill dusky above, pale brown below; irides brown; feet fleshy brown.

Ceylon, South India, Upper Burmah.

203. *PYRRHULAUDA GRISEA*, Scop.

Confined to the northern part of the island. Layard believed it was migratory; but I have seen it at Aripo at all seasons, in pairs during the summer, and in flocks during the winter months.

Bill pale brown; irides brown; feet fleshy.

Ceylon, India westward to Arabia.

204. *ALAUDA GULGULA*, Frankl.

Very common in the low country; but I have no recollection of seeing it on the hills. It has, however, been recorded, I believe, from the upper country by Kelaart. It was abundant at Aripo.

Bill dusky above, paler below; irides brown; feet fleshy brown.

Ceylon, India.

205. *CROCOPUS CHLOROGASTER*, Blyth.

I have obtained this Pigeon near Aripo; and it is said by Layard to be confined to the north of the island.

Ceylon, South and Central India.

206. *OSMOTRERON BICINCTA*, Jerdon.

This species is also found at times in wild jungle south of Aripo. I have likewise met with it a few miles from Coombo; but it is recorded as more numerous further south.

Ceylon, India eastward to Tenasserim.

207. *OSMOTRERON POMPADOURA*, Gmel.

The description of this species given by Gmelin was from a drawing of a Ceylon bird. Layard believed it to be a variety of *O. malabarica*, Jerdon; and Blyth has since given it the name of *flavogularis*; but the difference between Blyth's species and *O. pompadoura* can only be traced in the under tail-coverts, and there is a variation in this difference. *O. pompadoura* and *O. flavogularis* agree precisely in differing from *O. malabarica* in having the head less grey and the throat more yellow, and in not having the under tail-coverts cinnamon; this colour, however, Mr. Blyth tells me is only found in the male of *O. malabarica*. Specimens of *O. pompa-*



*doura* and *flavo-gularis* in Lord Walden's collection are undistinguishable, except in having the under tail-coverts green margined with white, entirely white, or white margined with yellow. Lord Walden's opinion that these differences are due to season or age appears to me likely to be correct; if not, the number of species founded on the colour of these particular feathers will have to be increased.

Ceylon, South India.

208. *CARPOPHAGA SYLVATICA*, Tickell.

Recorded by Layard as mostly found on the mountain-zone. He mentions it under the name of *C. pusilla*, Blyth; but the difference in size of this Pigeon from Ceylon and parts of India is not generally recognized as of specific value.

Ceylon, India to Burmah, and Hainan.

209. *ALSOCOMUS PUNICEUS*, Tickell.

This Pigeon, known to the Singhalese by a name literally translated "Season Pigeon," is recorded by Layard only as a rare visitor; and, according to the natives, "it appears during the fruiting of the cinnamon-trees." I have never seen it.

Ceylon, Eastern side of Central India, Assam, Arrakan, and Tenasserim.

210. *PALUMBUS TORRINGTONIÆ*, Kelaart.

Peculiar to Ceylon. It is found in great abundance on the hills, but changes its locality according to the season and the time at which the fruit of particular trees ripens. I have found it numerous at Nuwara Eliya at the end and beginning of the year; and it is occasionally found there at other times. It is allied to *P. elphinstonei*, Sykes, but differs essentially from it in having the back and wings dark slaty, and the underparts strongly vinaceous. It is known on the hills as the "Blue Pigeon."

Bill dusky, tip pale green; irides dark yellow; feet fleshy red.

Ceylon.

*Macropygia macroura* (Gmelin). With reference to the occurrence of this species in Ceylon, as stated by Bonaparte, Lord Walden has been good enough to send me the following note, with permission to make use of it:—

"The titles *Columba macroura*, Gmel. (1788), and *Columba mace-rona*, L. S. Müller (1776), were founded on the *Tourocco* of Buffon (Hist. Nat. Ois. ii. p. 553, and Pl. Enl. 329). Buffon figured this Pigeon from a Senegal example, presented by Adanson under the name of *Tourterelle à large queue du Sénégal*. But he afterwards (Hist. Nat.) substituted for Adanson's title that of *Tourocco*, because, as he says, while Adanson's bird possessed many of the characters of the European Turtledove, it carried its tail like 'le Hocco' (*Crax*). *Tourocco* may therefore be translated Turtledove-Curassow. Buffon is most circumstantial in his account of the locality whence his bird was obtained; and the fact that the specimen bore a title given



by Adanson strongly corroborates the Senegal origin. Yet Bonaparte (Consp. ii. p. 57) says 'ex Ceylon, nec Senegal.' The Prince was also (*l. c.*), I believe, the first who referred *C. macroura*, Gm., to the genus *Macropygia*. Still it is doubtful whether he ever saw an example of the bird, and the diagnosis given by him of the species only contains the prominent characters discernible in the plate quoted."

211. *COLUMBA INTERMEDIA*, Strickl.

There are two stations on the Ceylon coast which "Rock-Pigeons" are known to frequent. The principal one is Pigeon Island, a large mass of isolated rocks well known on the east coast, and about eighteen miles north of Trincomalie. I have visited this locality; and I have no doubt that Pigeons, probably of this species, are found there at a particular season of the year, according to the general report of the natives on the adjoining mainland; but I did not see any when I was there. Layard mentions their having been killed about fifty miles inland from Trincomalie. The other station is off Berberyn, not far from Galle.

Ceylon, India to Burmah.

212. *TURTUR RUPICOLA*, Pall.

Layard records having shot a young bird of *T. orientalis*, Lath. This may be the above species; but I am disposed to think his identification doubtful, as his only specimen was a young bird.

213. *TURTUR SURATENSIS*, Gmel.

Very common in the low country, and abundant at Aripo.  
Ceylon, India.

214. *TURTUR RISORIA*, Linn.

Very numerous in the north, and, I believe, not uncommon throughout the low country.  
Ceylon, India.

215. *CHALCOPHAPS INDICA*, Linn.

This handsome Dove is found in all parts of the island except the north. I have met with it in cultivated districts near Colombo and in the extreme south, on the road through the forest between Kandy and Trincomalie, and at Nuwara Eliya, where at the end of the year it frequents the jungle in great numbers. It has a low rapid flight, and a peculiar moaning coo, more like the note of some Owls than that of a Dove.

Ceylon, India, eastward to Tenasserim.

216. *PAVO CRISTATUS*, Linn.

Common in all jungly districts within a moderate distance of the coast. So far as my observations and inquiries have gone, it is unknown in the hill-country; and it is more numerous in the eastern



and northern parts of the island than in the more cultivated south and west.

Ceylon, India.

217. *GALLUS STANLEYI*, Gray.

The Ceylon Jungle-fowl is remarkable not only for being peculiar to the island, but also for being common in all parts of it where the country is uncultivated and there is jungle of a moderate height. Although especially abundant in the low country, it is often very numerous even on the upper hills, and is attracted to the particular localities where the "*nilloo*," the native name for some species of *Strobilanthes* growing at 5000 feet and upwards, is at the time in seed. I have entirely failed to discover that any thing is known among botanists of the seeds of the *Acanthaceæ* possessing narcotic or other poisonous properties; but it is well known that the Jungle-fowl after feeding for a time among the *nilloo* become partially blind or stupified, so that they may frequently be knocked down with a stick. This stupefaction is generally attributed to the *nilloo*-seeds, which are so largely eaten by these birds; but in the absence of any known poisonous properties in these seeds, it appears possible that the birds may really suffer from devouring some fungus or other plant found in the damp woods where the *nilloo* grows.

At daybreak the crow of the Jungle-cock is first heard; and for an hour or two after sunrise, if the birds are at all numerous, they may be heard challenging each other on all sides. On these occasions a successful shot may sometimes be obtained by remaining perfectly still between two birds which are challenging and gradually approaching each other. Some of the native hunters are very expert in calling the Jungle-cocks, by beating on a loose fold of their cloth, so as to produce an imitation of the sound of a bird's wings just as it is alighting: no time must be lost with the gun on these occasions, as the cocks discover the deception the moment they get sight of you, and instantly run off with drooping tails like Pheasants. It is not difficult in favourable jungle to approach a calling bird within easy shot; and under these circumstances I have generally found the cock strutting up and down a low horizontal branch of a tree, raising and lowering its head, and every now and then giving utterance to its peculiar crow, which has been likened to the sound of "George Joyce." When the bird is tolerably close, the syllable "ek" is heard preceding those two sounds, which are so familiar to persons who have been wandering in the jungles of Ceylon. In some of the wilder jungle-roads, a cock and hen may sometimes be seen feeding together; but generally the hens are very shy, and not many of them are killed.

Mr. Layard tells me that there is no doubt about this Jungle-fowl sometimes breeding with the domestic poultry in the native villages. I have seen young Jungle-fowl, which had been hatched under domestic hens, running about with the other chickens; but they were always rather wild and invariably roosted out of doors; and those which were not sooner or later killed by some accident,



ultimately took to the jungle. Some others, which, however, were reared in an aviary at Colombo by my friend Dr. Boake, became quite tame, and were in good feather when he kindly allowed me to send them to London for the Society's Gardens; but they all died when they were almost within sight of England.

Mr. Blyth can hardly be correct in his description of the head and appendages in this species. He says (*Ibis*, 1867, p. 307), "The cock has a yellow comb with a red edge, and the cheeks and wattles (as I remember them in the living bird) are chiefly yellow." His description of the colour of the comb is approximately correct, as the extent of the yellow varies in different specimens; but I am too familiar with the appearance of the living or freshly killed bird to have any doubt about the cheeks and wattles being red. These parts assume a dark livid appearance a few hours after death; but the yellow in the comb remains, and is evident even in old dry skins. The size of the comb and wattles varies, and probably depends on age.

The following details were taken from a fine adult cock I killed at Aripo, and were noted down on the spot:—

Bill brown, front of the lower mandible pale yellow; irides buff; comb, wattles, and naked skin about the head purplish red, the comb having a large wing-shaped spot of yellow occupying the middle of the posterior half, very bright at its origin immediately over the eye, and shading off at its margin into the colour of the comb; feet and legs pale yellow.

Ceylon.

## 218. *GALLOPERDIX BICALCARATA*, Forst.

Peculiar to Ceylon; abundant on many parts of the hills, and frequenting also jungly places in the low parts of the southern half of the island. During the winter months it is numerous in the coffee-districts and upper hills, and is trapped in large numbers by the natives. It is skulking in its habits and difficult to flush, usually seeking concealment in the thicker parts of the jungle when it is disturbed. They bear confinement well in Ceylon; but some specimens I brought to England, although apparently strong and well on their arrival, all died within three days after the ship entered the Thames.

Bill red ♂, dusky ♀; irides brown; feet fleshy red.

Ceylon.

## 219. *FRANCOLINUS PICTUS*, Jard. & Selby.

The occurrence of this species, said to have been well identified, was noticed three or four years ago in one of the Colombo newspapers. I did not see the specimens, and I cannot now give the precise date or particulars of where they were obtained.

Ceylon, Central India.

## 220. *ORTYGORNIS PONTICERIANA*, Gmel.

Common in the north of Ceylon, and found also in the cinnamon-gardens at Colombo. These birds may have escaped from confine-



ment, as large numbers of them are brought alive to the Colombo market from Tuticorin on the Indian coast; Mr. Legge, however, has also seen the bird at Galle. This species is indigenous in the north, and is always very abundant at Aripo. The large compound surrounding my house at that place was virtually nothing but a considerable piece of jungle fenced in, and was frequented by many kinds of wild animals and birds. Partridges were very numerous there; and they might be seen or heard at all hours of the day, and often within a few yards of the house. They roosted in low bushes.

Bill dusky; irides brown; feet dull red.

Ceylon, South, Central, and North-west India, Persia?

221. *PERDICULA ASIATICA*, Lath.

Layard mentions having seen a pair of these birds which were caught alive near Colombo. He speaks of it under the name of *P. argoondah*, Sykes.

Ceylon, South India.

222. *EXCALFACTORIA CHINENSIS*, Linn.

I have seen this bird from Kandy and the cinnamon-gardens at Colombo; and Layard says it is common in the south.

Ceylon, India, eastward to China, Malaya, Australia.

223. *TURNIX TAIGOOR*, Sykes.

Common in all parts of the low country. I have found its eggs at Aripo in February.

Bill lead-colour; irides pale yellow; feet pale leaden.

Ceylon, India.

224. *CURSORIUS COROMANDELICUS*, Gmel.

I believe the Indian Courser is resident in the north of Ceylon, as I have seen it in almost every month of the year at Aripo. It is more numerous, however, in the winter months, being then in small parties of six or eight. Its flight is heavy and flapping, like that of the Lapwings; but it runs lightly and fast; and when separated from its companions, I have more than once seen it running along behind the bund of a dry paddy-field, with head lowered and wings trailing on the ground, presenting a most curious appearance, as the colour of the back resembled that of the dry mud, and there was nothing to attract attention but the drooping black primaries. Layard appears to have occasionally met with this bird, but only in April.

Bill black; irides dark brown; feet cream-colour.

Ceylon, Central and West India.

225. *CHARADRIUS FULVUS*, Gmel.

*Charadrius longipes*, Temm., apud Jerdon.

The Ceylon birds have the ash-coloured axillary plume characteristic of this species; they are migratory, appearing at Aripo in August, many of them then having some remains of the black



breeding-plumage. Throughout the winter they are abundant in the north, and are occasionally seen as far south as Colombo, frequenting the esplanade with some of the smaller Plovers.

Bill black; irides brown; feet dark leaden.

Ceylon, India, Eastern Asia to Australia, and Polynesia.

226. *ÆGIALITES MONGOLICUS*, Pall.

Mr. Edmund Harting, in a series of exhaustive papers "On rare or little known Limicolæ" ('Ibis' 1870), has worked out the synonymy of this species, among others, and identified the bird given by Jerdon under *Æ. pyrrhothorax*, Temm., with that described by Pallas. It is doubtless the one mentioned by Layard as *Hiaticula leschenaultii*, Less., as I have no reason to think the much larger *Æ. geoffroyii*, Wagler, is found in Ceylon.

*Æ. mongolicus* is a winter visitor to Ceylon, and is then very abundant on the coast, commonly associating with *Æ. cantianus*. All the specimens I have examined have been in winter dress.

Bill black; irides dark brown; legs grey, feet dark grey.

Asia to North Australia.

227. *ÆGIALITES CANTIANUS*, Lath.

Mr. Harting has been good enough to examine my specimens of this and the preceding species, and he tells me that a small Plover which I had been unable to identify is the young of *Æ. cantianus*. I obtained specimens of this species in different states of plumage; but the greater number of these birds found in Ceylon are young ones, and apparently diminutives of *Æ. mongolicus*. I have occasionally got specimens in nearly, if not quite, full plumage.

Bill black; irides dark brown; feet dark grey, legs paler (in the young).

Europe, Asia.

228. *ÆGIALITES DUBIUS*, Scop.

*Ægialites philippensis*, Scop., apud Jerdon.

This well-known little Plover is common in Ceylon, and, I believe, resident there, as it is certainly found during a great part of the year at Aripo. Although associating to some extent with the other Sand-Plovers, it does not always keep with the party, but wanders off to some distance when feeding. It is particularly fond of standing on any little natural elevation of the ground or heap of rubbish on the beach.

Bill black; irides dark brown; feet yellow.

Ceylon, India, eastward to China and Japan.

229. *CHETTUSIA GREGARIA*, Pall.

I have identified a single specimen of this Plover shot by Mr. Bligh on the Galle face at Colombo. It has not been before observed in Ceylon.

Ceylon, parts of India, West Asia, and South-east Europe.



230. *LOBIVANELLUS INDICUS*, Bodd.

*Lobivanellus goensis*, Gmel., apud Jerdon.

Found in all open parts of the low country, and generally in pairs.  
Resident in Ceylon.

Ceylon, India.

231. *SARCIOPHORUS MALABARICUS*, Bodd.

*Sarciophorus bilobus*, Gmel., apud Jerdon.

Distribution much the same as that of the last species, but it is more numerous in the north. It was always abundant at Aripo, and was found in large flocks during winter. Jerdon, in his description of this species, has omitted to mention that the chin and upper part of the throat are dull black. This appears as soon as the young are well able to fly, and remains at all seasons. It is present in all the Indian specimens I have seen.

Bill yellow, tip black; irides pale yellow; wattles yellow; feet yellow.

Ceylon, India.

232. *ESACUS RECURVIROSTRIS*, Cuv.

I have only seen this bird occasionally in the Aripo district. It was usually in pairs on the banks of the Aripo river. I have shot this bird in August, from which it would appear to be a resident.

Bill greenish yellow, tip black; irides pale yellow; feet yellow.

Ceylon, India.

233. *ÆDICNEMUS CREPITANS*, Temm.

Common in the north at all seasons. I have also flushed it in the cinnamon-gardens at Colombo.

Asia, N. Africa, Europe.

234. *STREPSILAS INTERPRES*, Linn.

I obtained one specimen in August on the coast a few miles north of Aripo. Layard also met with it in the north, and once at Colombo. It is rather a scarce bird in Ceylon.

World-wide distribution.

235. *DROMAS ARDEOLA*, Paykul.

I have never seen this remarkable bird; but specimens were obtained by Layard—at sea, with one exception. He follows Blyth in placing it near the Terns.

Ceylon, India, Red Sea.

236. *HÆMATOPUS OSTRALEGUS*, Linn.

Layard records having seen one or two of these birds near Jaffna. Ceylon, Indian and European coasts.

237. *SCOLOPAX RUSTICULA*, Linn.

The occasional appearance of the Woodcock on the Ceylon hills



has been reported on "sportsman's authority;" and it is now confirmed by Mr. S. Bligh, who writes to me that he has just examined a specimen quite recently killed at Nuwara Eliya.

238. *GALLINAGO NEMORICOLA*, Hodgson.

239. *GALLINAGO STENURA*, Temm.

240. *GALLINAGO SCOLOPACINA*, Bonap.

241. *GALLINAGO GALLINULA*, Linn.

Of these four reputed Ceylon species *G. stenura* appears to be the only one which has been positively identified. It is the Common Snipe of sportsmen; and I believe there are but few persons in the island who are aware of the peculiarity in the tail-feathers by which it can be at once distinguished from *G. scolopacina*, which it is generally believed to be. *G. stenura* is found all over the island in the winter months; and although of course much more abundant in paddy-growing districts, it is also numerous in swampy plains on the upper hills. *G. nemoricola* was recorded from Nuwara Eliya by Mr. Hugh Nevill as new to the island (J. R. A. S., C. B., 1867-70, p. 138); but although he, I believe, examined the specimen, the skin was not preserved, and he himself told me that he identified the bird, after he had left the hills, by the coloured figure in Jerdon's 'Illust. Ind. Ornith.,' from which work he has evidently taken his description of the species. Although neither Layard nor Kelaart mentions this bird, Jerdon speaks of it as being found on the "elevated regions of Southern India and Ceylon," but does not give any authority. In the case of *G. gallinula*, Layard thought that "sportsman's authority" might be trusted, as the "Jack" would not be easily confounded with the other Indian Snipes, and he had been informed by a person likely to be acquainted with it that it was not uncommon in the north a few years previously. None of these species is unlikely to occur in Ceylon; but, except in the case of *G. stenura*, the evidence in their favour is not quite as clear as could be wished.

242. *RHYNCHÆA BENGALENSIS*, Linn.

Not uncommon in the low country during the winter. Layard says some remain to breed, "the season of incubation being from May to July." He tells me that he obtained many eggs of this species. Jerdon also gives June and July as the breeding-time of this bird in India. It apparently varies, however, as a bird caught near Colombo, and sent alive to me on the 31st of December, was found to have laid an egg in the basket in which it was packed. This egg has been identified by Mr. Layard as that of the Painted Snipe, although its ground-colour is rather paler than usual.

Ceylon, India, Burmah to S. China, Africa.

243. *LIMOSA ÆGOCEPHALA*, Linn.

Recorded by Layard. I have not met with it.

Ceylon, parts of continental Asia and Europe.



244. *TEREKIA CINEREA*, Gmel.

I obtained one specimen in winter plumage, out of a flock of five, in April 1869; they were in a small swamp near the sea at Aripo. It appears to be new to Ceylon.

Bill dusky, base yellow; irides brown; feet pale orange.  
Europe, Asia to Australia.

245. *NUMENIUS ARQUATA*, Linn.?

The Ceylon Curlew requires further examination; it may prove to be *N. lineatus*, Cuv., which Mr. Blyth tells me is commonly found in India.

246. *NUMENIUS PHÆOPUS*, Linn.

This and the preceding species are common on many parts of the coast, and were often found at Aripo, but never in flocks.

Europe, Africa, Asia.

247. *TRINGA SUBARQUATA*, Gmel.

I have obtained this bird in May at Aripo with the breeding-plumage far advanced.

248. *TRINGA MINUTA*, Leisler.

This was the common Stint on the shore at Aripo, yet it appears doubtful whether it is found in India (Jerdon, *Birds of India*, App. p. 875).

Bill black; irides brown; feet leaden-black.

249. *TRINGA SALINA*, Pall.

*Tringa subminuta*, Midd.

I obtained two of this species at Aripo, in January 1870, for the first time. It is new to Ceylon, although Blyth, as quoted by Jerdon (*ut supra*), states that it is the common Little Stint of India. My specimens of these two Stints have been carefully examined and identified for me in Eugland.

Bill black; irides brown; feet dull olive.

250. *TRINGA PLATYRHYNCHA*, Temm.

This appears to be rare in Ceylon. Layard only obtained one two specimens quite in the north.

251. *ACTITIS GLAREOLA*, Gmel.

Exceedingly abundant in all wet places. I have counted twenty round a small pool in my compound at Aripo during the rains.

252. *ACTITIS OCHROPUS*, Linn.

Numerous, but less so than the last species.

253. *ACTITIS HYPOLEUCOS*, Linn.

Very common in all parts of the low country, and less so on the



hills. I have seen it as high as Nuwara Eliya in February. It is probably resident in Ceylon.

254. *TOTANUS GLOTTIS*, Linn.

255. *TOTANUS STAGNATILIS*, Bechst.

Both very common at Aripo, and generally so in the low country.

256. *TOTANUS FUSCUS*, Linn.

257. *TOTANUS CALIDRIS*, Linn.

These species were considered common by Layard ; but I have not seen them.

258. *HIMANTOPUS AUTUMNALIS*, Hass.

*Himantopus candidus*, Bonn.

Not uncommon at Aripo during the rains.

259. *RECURVIROSTRA AVOCETTA*, Linn.

Two of these birds, killed near Jaffna, are recorded by Layard.

Almost all these small Waders are, I believe, winter visitors to Ceylon after breeding in Northern Europe or Asia.

260. *HYDROPHASIANUS CHIRURGUS*, Scop.

Very common in the neighbourhood of Colombo. Beautiful specimens in various states of plumage are sometimes brought in for sale by the Singhalese, who walk through the flooded marshes and wait patiently, with the water often above their waists, till they can make sure of a successful shot. I have not seen this bird in the north ; but, as Layard mentions, it may be sometimes observed walking on the lotus-leaves in the lake at Colombo.

Bill bluish, tip green ; irides red-brown ; feet leaden.

Ceylon, India, China.

261. *PORPHYRIO POLIOCEPHALUS*, Lath.

Common in suitable situations, but shy and fond of concealment. They are numerous in the neighbourhood of Colombo.

Ceylon, India.

262. *GALLICREX CRISTATUS*, Lath.

This bird is also common about Colombo and in marshes in the south.

Ceylon, parts of India, Burmah, Malaya, China.

263. *GALLINULA CHLOROPUS*, Linn.

Layard met with one specimen of this bird in the north ; but I have not heard of any others, although it appears to be general in India.

Europe, Asia, Africa.



264. *GALLINULA PHÆNICURA*, Forster.

This well-known species was first described and characteristically figured by Forster (1781) from a Ceylon specimen. It is very common in suitable situations throughout the low country; but I am not aware that it is found on the hills, except near their foot.

In what appears to be rather an old bird the upper part of the back is irregularly barred with grey, and the chestnut is confined to the sides of the rump and under tail-coverts. This was a male, and when shot was in company with a presumed female and a small black chick. Jerdon does not mention the characteristic white face and forehead in his description of this species; and I observe in the British Museum a specimen from Malaya labelled *G. phœnicura* in which the white is confined to the underparts. It appears to have rather a stouter bill, and may be a distinct race or species, possibly the one from which Jerdon took his description, in which he says "irides blood-red, legs green." These characters do not agree with the following in true *G. phœnicura* from Ceylon:—

Bill green, ridge dull red; irides brown; legs and feet light yellow-brown.

Ceylon, India to Malaya, S. China, Formosa.

265. *PORZANA PYGMÆA*, Naum.

Layard records having obtained one specimen.

Ceylon, India, China, Japan.

266. *PORZANA FUSCA*, Linn.

Recorded by Layard as rare.

Ceylon, India, E. Asia.

267. *RALLINA CEYLONICA*, Gmel.

This bird arrives in Ceylon in October, just at the change of the monsoon, and takes refuge in the first place of concealment it can find, often entering the houses and hiding amongst the furniture. I have caught the bird under these circumstances at the hotel at Colombo. Although this Rail is only a winter visitor to Ceylon, specimens of it from India appear to be rare, and the North-Indian race has been separated by Blyth under the name of *R. amauroptera*. The distribution of the true *R. ceylonica* appears to be uncertain.

Bill dusky above, green below; irides red-brown ("carmine, with an inner circle of yellow," Layard); feet leaden brown.

Ceylon, S. India.

268. *RALLUS STRIATUS*, Linn.

Ceylon, India, Burmah to Malaya, Formosa.

269. *RALLUS INDICUS*, Blyth.

Ceylon, India, Tientsin.

These two species have both been recorded by Layard, but are said to be rare. *R. indicus* is very close to *R. aquaticus* of Europe, but



has been separated by Blyth; I have not had an opportunity of comparing them.

270. *LEPTOPTILOS JAVANICA*, Horsf.

I have seen this Stork close to Aripo and a few miles from Trincomalie, on both occasions in small parties. I believe it is a winter visitor and that it is only found in the northern half of the island, although by no means uncommon in particular districts.

Ceylon, India, Burmah to part of Malaya, Hainan.

271. *MYCTERIA AUSTRALIS*, Shaw.

Layard mentions having seen this bird near Jaffna; but I have never met with it.

Ceylon, India to Australia.

272. *CICONIA EPISCOPUS*, Bodd.

*Ciconia leucocephala*, Gmel.

Described by Layard as common in swampy lands; and although I have not met with the bird, it appears to be well known in suitable situations.

Ceylon, India to Malaya.

273. *ARDEA CINEREA*, Linn.

This Heron, considered by Layard to be very rare, is not at all uncommon in Ceylon. I have seen it in many parts of the island, and have had an opportunity of examining young birds on more than one occasion.

Asia, Africa, Europe.

274. *ARDEA PURPUREA*, Linn.

More common than the last species; it is very numerous in the south, and breeds near the Amblangodde Lake, a few miles from Galle.

Asia, Africa, Europe.

275. *HERODIAS ALBA*, Linn.

276. *HERODIAS EGRETTOIDES*, Temm.

*Ardea intermedia*, Wagler, apud Layard.

277. *HERODIAS GARZETTA*, Linn.

278. *DEMIEGRETTE ASHA*, Sykes.

279. *BUPHUS COROMANDUS*, Bodd.

These five species are all said by Layard to be common, and to breed in Ceylon. I have no doubt he is quite correct. Egrets of different kinds are abundant in the swamps throughout the island; but as I brought home no specimens with me, I cannot be sure of the correctness of my identifications. I have occasionally seen spe-



cimens of one species at Nuwara Eliya, in October, with a black bill and greenish feet, probably *H. garzetta*; but these birds are mostly found in the low country. They appear to be all widely distributed.

280. *ARDEOLA GRAYII*, Sykes.

*Ardeola leucoptera*, Bodd., apud Jerdon.

Exceedingly common in Ceylon.

281. *BUTORIDES JAVANICA*, Horsf.

Common, and resident in the north. I have occasionally seen it near Colombo.

Bill black above, yellow below; irides yellow; feet yellow-green. Ceylon, India, Burmah to Malaya, China.

282. *ARDETTA FLAVICOLLIS*, Lath.

283. *ARDETTA CINNAMOMEA*, Gmel.

284. *ARDETTA SINENSIS*, Gmel.

Of these three species *A. cinnamomea* is the commonest. They appear to be confined to the southern half of the island, and are all found in the neighbourhood of Colombo. They range from India more or less eastward.

285. *NYCTICORAX GRISEUS*, Linn.

Not uncommon in suitable places.

Asia, Africa, Europe.

286. *GOISACHIVS MELANOLOPHUS*, Raffles.

I was fortunate in getting a specimen of this purely eastern Bittern at Aripo, in November 1866; it was hiding among some low bushes a few yards from my house, and was a female in immature plumage. My shot disabled but did not kill it, and it struck at me furiously with its bill as I endeavoured to extricate it from among the thorns, the neck-feathers being erected in true Bittern fashion. Layard first observed this species in Ceylon, and obtained two or three specimens near Colombo. It is remarkable that this common Malay species should not yet have been observed in India, as the birds obtained by Layard and myself (no others are recorded from Ceylon) were all found on the west coast, and my specimen was from that part of the coast where migrants from India generally appear first.

Ceylon, Malaya, Japan, Philippines.

287. *TANTALUS LEUCOCEPHALUS*, Forster.

Ceylon, India, Burmah, Amoy.

288. *PLATALEA LEUCORODIA*, Linn.

Asia, Africa, Europe.



289. *ANASTOMUS OSCITANS*, Bodd.

Ceylon, India.

290. *THRESKIORNIS MELANOCEPHALUS*, Linn.

Ceylon, India, Burmah, Arrakan, China?

These four species are resident in Ceylon, and abundant in many localities, but, I believe, more numerous in the south and south-east than elsewhere, except, perhaps, the last one, which Layard says is common in the north and north-west. I have only seen it on two occasions near Aripo. *T. leucocephalus* was first described from a Ceylon specimen.

291. *FALCINELLUS IGNEUS*, S. G. Gmel.

Not uncommon near Aripo, and apparently confined to the north. World-wide distribution.

292. *PHÆNICOPTERUS ANTIQUORUM*, Temm.*Phœnicopterus roseus*, Pall., apud Jerdon.

I have seen this species occasionally at Aripo in October and November; they were in flocks of from twenty to thirty, and presented a remarkable appearance as they flew in single file, with outstretched head and legs. Layard speaks of having seen them in very large numbers on the north and east coasts.

Parts of Asia, Africa, and Europe.

293. *SARKIDIORNIS MELANONOTUS*, Forster.

Ceylon, India, Burmah.

294. *ANSERELLA COROMANDELIANA*, Gmel.

Ceylon, India, Burmah, Malaya.

295. *DENDROCYGNA JAVANICA*, Horsf.*Dendrocygna arcuata*, Cuv.

Ceylon, India, Burmah, Malaya.

296. *SPATULA CLYPEATA*, Linn.

Asia, Europe.

297. *ANAS PÆCILORHYNCHA*, Penn.

Ceylon, India, Burmah.

298. *DAFILA ACUTA*, Linn.

Asia, Europe.

299. *QUERQUEDULA CRECCA*, Linn.

Asia, Europe.

300. *QUERQUEDULA CIRCIA*, Linn.

Asia, Africa, Europe.



These eight species are all recorded by Layard; and *S. melanotus* was first obtained in Ceylon. The only one commonly distributed throughout the island is *D. javanica*; it breeds in many localities, and is known among Europeans as the "Teal." *A. coromandeliana* is tolerably numerous in the north and east, and, I am told, breeds near Battacaloe; it is sometimes found near Colombo. The others are mostly found in the north, where, according to Layard, *Q. crecca* and *circia* are very abundant in winter. Layard also mentions having seen on several occasions, through a telescope, what he believed to be *Branta rufina*, Pall.; but that species has not been yet identified from Ceylon.

301. *PODICEPS PHILIPPENSIS*, Bonn.

Very common on all large pieces of water, and often associated in flocks. I have counted thirty-eight together on the Colombo lake.

Bill black, tip white, base of lower mandible dull green; irides dark yellow; legs and feet blackish green in front, black below (♀ killed in July).

Ceylon, India, China, Formosa, Hainan.

302. *THALASSIDROMA* —?

A species of Stormy Petrel is often seen in Colombo harbour and on the west coast in the bad weather during the south-west monsoon, but no specimen of it has yet been obtained; it has appeared to me to be entirely black, with the exception of the white rump.

303. *CROICOCEPHALUS ICHTHYAËTUS*, Pall.

Layard mentions having seen a pair of these birds at Pt. Pedro after a severe storm. It appears to be only occasionally seen on the Indian coast.

304. *XEMA BRUNNICEPHALA*, Jerdon.

This is the only true Gull commonly found in Ceylon. It is very abundant in the north, and is seen at times on all parts of the coast.

Ceylon, India, Pekin?

305. *SYLOCHELIDON CASPIA*, Lath.

306. *GELOCHELIDON ANGLICA*, Montagu.

307. *HYDROCHELIDON LEUCOPAREIA*, Natt.

*Hydrochelidon indica*, Stephens.

These species I have found common in Ceylon, and I have no doubt of their being resident there. *S. caspia* may be seen at all times of the year, almost invariably in pairs, flying along the shore just outside the line of beach. I have shot *G. anglica* in April, July, and December, but have not met with one in the full breeding-plumage.

308. *SEENA AURANTIA*, Gray.

Said by Layard to be common. I have not actually identified



this species, but believe I have often seen it near the Aripo pearl-banks.

309. *STERNA MELANOGASTER*, Temm.

I have frequently seen this Tern near Aripo, and occasionally at Colombo. Layard found it common on some of the inland lakes as well as on the coast.

310. *STERNA NIGRA*, Linn.

*Sterna leucoptera*, Temm.

I shot one of a pair of these birds in May 1866. They were flying about over a small tank, not very far from the shore, about six miles from Aripo, and were in rather imperfect plumage, the head and neck being speckled. The characters of the species, however, were unmistakable. My specimen is now in the Colombo Museum. This is the only occasion of this Tern having been recognized in Ceylon; and it has only been recently added by Mr. Hume to the Indian avifauna.

Ceylon, India, China, North Africa, South Europe.

311. *STERNA GRACILIS*, Gould?

I include, with some doubt, under this heading a Tern shot in July 1869, on the Colombo beach; others of the same kind were killed at the time; and they were all in rather immature plumage. This specimen has been examined by Mr. Howard Saunders and Mr. Gould, and is believed by those gentlemen to be *S. gracilis*, and in that case a visitor to Ceylon during the Australian winter. It had the bill reddish black, irides black, and feet dull fleshy red. *S. gracilis* is allied to *S. hirundo*, but has the bill slighter, the upper tail-coverts grey as on the back and tail, and the whole under surface white.

312. *STERNULA SINENSIS*, Gmel.?

There is, I think, some doubt about the species to which Layard refers under the name of *S. minuta*, and which he speaks of as frequenting the inland lakes, though "most common on tanks and still waters near the sea-shore." I have never succeeded in obtaining a specimen of true *S. minuta*; and Mr. Legge, who has collected many of the Ceylon Terns, has been equally unsuccessful; but we have both frequently met with a small species in winter dress which may have been mistaken for it. This bird agrees in measurements and general colouring with *S. sinensis*, Gmel. (*S. sumatrana*, Raffles), and differs from *S. minuta* in having a black bill and the shaft of the first primary white. It was also collected by Mr. Jesse during the late Abyssinian expedition.

313. *THALASSEUS CRISTATUS*, Stephens.

This Tern is not uncommon on the west coast during summer. I have identified a specimen killed on the beach at Colombo in company with smaller species.



314. *THALASSEUS MEDIUS*, Horsf.

*Thalasseus bengalensis*, Less., apud Jerdon.

A Tern apparently of this species is very common.

315. *ONYCHOPRION ANÆSTHETUS*, Scop.

I do not know this Tern; but Layard mentions having obtained three specimens.

316. *PHAËTON RUBRICAUDA*, Bodd.

During my annual cruises on the Ceylon coast, I have seen this bird sufficiently near to identify it with certainty, as it hovered over the vessel. All the Tropic birds I have seen there, however, have had white tails; and, as I find among my notes mention of one instance of the bill being red, I conclude that bird was an immature example of the above species. I am very confident I have also seen the yellow-billed species, *P. flavirostris*, Brandt, but I have no special record of the colour of the bill. I shall therefore only call the attention of future observers to that species.

317. *SULA FIBER*, Linn.

In February and March 1868 I had many opportunities of watching a pair of Boobies which frequented the neighbourhood of the Aripo pearl-banks, about ten miles from the land. They used often to perch on a large iron buoy close to my usual anchorage at night. I only saw them during that one season; and they have not been otherwise recorded.

318. *ATTAGEN MINOR*, Gmelin.

*Attagen ariel*, Gould.

Frigate-birds have been killed in several localities on the west coast; and I have observed them on many occasions at Aripo during the strength of the south-west monsoon. They were generally in parties of five or six, and at a considerable height above the shore. Their action, as they hung as it were against the gale, slowly swaying, first on one side, then on the other, strongly reminded me of the behaviour of a large paper kite when it has mounted high in the air. Without any perceptible movement of their partially extended wings, these birds remained as if suspended in the air, but very slowly working against the wind, and gradually advancing along the line of beach. Layard mentions these birds under the name of *A. ariel*, Gould, a species from the Australian seas, but which also has been recorded by Swinhoe from Amoy.

*A. aquilus*, Linn., is found in the Indian seas; and it is not unlikely that some of the Frigate-birds seen on the Ceylon coast may belong to that species.

319. *PELECANUS PHILIPPENSIS*, Gmel.

I have seen Pelicans near Trincomalie, and at the entrance to Kokeley Lake, on the north-east coast. Their breeding-stations













Gould, John. 1872. "March 5, 1872." *Proceedings of the Zoological Society of London* 1872, 355–492. <https://doi.org/10.1111/j.1469-7998.1872.tb00490.x>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/90880>

**DOI:** <https://doi.org/10.1111/j.1469-7998.1872.tb00490.x>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/73724>

**Holding Institution**

Natural History Museum Library, London

**Sponsored by**

Natural History Museum Library, London

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.