

Synonym: Eremomela griseoflava perimacha Oberholser, Proc. Biol. Soc. Wash., 33, 1920: Asbestos Mts.

Breast pale grey, belly somewhat brighter yellow than in *E. i. saturator*.

Range: The Orange River basin from the Vaal confluence downstream, Gordonia, and most of Griqualand West, southern S. W. Africa. Intergrades with the next race between 22° and 23°S in South West Africa.

3. *Eremomela icteropygialis sharpei* Reichenow

Eremomela flaviventris sharpei Reichenow, Orn. Monatsb., 13, 1905: 25—Windhoek

Breast almost white; belly decidedly brighter than in *E. i. icteropygialis*.

Range: Northern South West Africa, the whole of Bechuanaland Protectorate except the north-east corner, and the Kalahari Gemsbok Park area of the northern Cape.

4. *Eremomela icteropygialis polioxantha* Sharpe

Eremomela polioxantha Sharpe, Cat. Bds. B.M., 7, 1883: 160—Swaziland.

Breast much as in *E. i. icteropygialis*, belly much richer yellow than in any of the preceding.

Range: as outlined in par. 3 above.

References:

- Acocks, J. H. P. (1953), 'Veld Types of South Africa,' *Bot. Surv. S. Afr. Mem. No.* 28.
 Clancey, P. A. (1959), 'Miscellaneous Taxonomic Notes on African Birds, XIII (h), *Durb. Mus. Novit.*, 5: 204–6 (1962), 'Comments on the Typical Race of *Eremomela icteropygialis* (Lafresnaye), *Bull. B.O.C.*, 82: 44–5.
 Forbes, V. S. (1958), 'The Limits of Le Vaillant's Travels in South Africa,' *The Ostrich*, 29: 79–85.
 Grant, C. H. B. (1957), 'Levaillant's Travels in South Africa,' *The Ostrich*, 28: 83–97.
 Macdonald, J. D. (1957), *A Contribution to the Ornithology of Western South Africa*. London.
 White, C. M. N. (1961), Notes on *Eremomela icteropygialis* (Lafresnaye), *Bull. B.O.C.*, 81: 90–2.
 Winterbottom, J. M. (1958), 'Distribution of *Eremomela icteropygialis* Lafresnaye,' *Bull. B.O.C.*, 78: 132–3.
 Winterbottom, J. M. (1960), 'Southern Limit of the Range of the Swallow-tailed Bee-eater *Merops hirundineus* Licht.,' *The Ostrich*, 31: 27.
 Winterbottom, J. M. and Courtenay-Latimer, M. (1961), 'A List of the Birds of Little Namaqualand,' *S. Afr. Avif. Ser.*, 1.

A new name for *Bradypterus barratti major* (Roberts)

by S. A. PARKER

Received 6th January, 1962

If both *Caffrillas barratti* (Sharpe) and *Dumeticola major* Brooks are placed in the genus *Bradypterus*, following Delacour (*Ibis*, 1942, p. 512), then the name *Bradypterus barratti major* (Roberts) becomes a junior secondary homonym. I therefore propose:—

Bradypterus barratti lysis, nom. nov. for *Caffrillas barratti major* Roberts, Ann. Transvaal Mus., viii, p. 234, 1922—Wakkerstroom, Transvaal.

= *Bradypterus barratti major*, not

Dumeticola major Brooks, Journ. As. Soc. Bengal, 41, p. 77, 1872—Kashmir

= *Bradypterus major major*.



Parker, S. A. 1962. "A new name for *Bradypterus barratti* major (Roberts)." *Bulletin of the British Ornithologists' Club* 82, 122–122.

View This Item Online: <https://www.biodiversitylibrary.org/item/126998>

Permalink: <https://www.biodiversitylibrary.org/partpdf/77898>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: British Ornithologists' Club

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.