

Miocene mollusks from the Kumano Group of the Ukui area, southeastern part of the Kii Peninsula, southwestern Honshu, Japan

YUTAKA HONDA¹, SEISUKE USHIRO² and SAYURI MORITANI³

¹Department of Earth Sciences, Faculty of Education, Mie University, Tsu 514-8507, Japan

²Shingu Senior High School attached to Kinki University, Shingu 4966, Shingu, Wakayama 647-0081, Japan

³Yasusaka 1010, Nakagawara-cho, Sumoto, Hyogo 656-0004, Japan

Received 19 December 1996 ; Revised manuscript accepted 5 March 1998

Abstract. The Shimosato Formation comprises the lower part of the Kumano Group of the southeastern Kii Peninsula, southwestern Honshu. It yields such mollusks as *Saccella miensis* (Araki), *Cyclocardia siogamensis* (Nomura), *Macoma (Macoma) izurensis* (Yokoyama), *Cultellus izumoensis* Yokoyama, *Dosinia (Phacosoma) nomurai* (Otuka), *Periploma (Aelga) mitsuganoense* Araki, *Thracia watanabei* Itoigawa and Shibata, *Turritella (Hataiella) sagai* Kotaka, and *Fulgoraria (Musashia) yanagidaniensis* Araki. The *Periploma*–*Saccella*–*Cyclocardia* and *Turritella*–*Dosinia* assemblages of the Shimosato Formation are inferred to represent lower sublittoral to bathyal and sublittoral environments, respectively.

The fauna of the Shimosato Formation is comparable with the subtropical Akeyo Fauna of late Early Miocene age, based on the occurrence of diagnostic species such as *Dosinia (Phacosoma) kawagensis* Araki, *Thracia watanabei* Itoigawa and Shibata, *Turritella sagai*, and *Fulgoraria yanagidaniensis*. The assemblages of the middle part of the Shikiya Formation and the lower part of the Mitsuno Formation are comparable with those of the tropical Kurosedani (subtropical Kadonosawa) Fauna of latest Early to earliest Middle Miocene age.

Key words : Assemblages, fauna, Kumano Group, Miocene, mollusks, systematic description

Introduction

The Miocene Kumano and Owase Groups of the Nankai Geologic Province are exposed in the southeastern part of the Kii Peninsula, southwestern Honshu (Figure 1). The Kumano Group rests upon the Shimanto Supergroup with clino-unconformity or is faulted against it and is intruded by the Kumano Acidic Rocks. The age of the Kumano Acidic Rocks is between 14 and 15 Ma (Chijiwa, 1988). The Kumano Group consists largely of mudstone and sandstone from 1,500 to 4,000 m thick and is divided into three lithologic units, in ascending order: the Shimosato (Onuma), Shikiya (Koguchi) and Mitsuno Formations (Hisatomi, 1984) (Figure 2).

The contemporaneous Setouchi Miocene Series (upper Lower to lower Middle Miocene) of the Setouchi Geologic Province is extensively distributed in central and western Honshu, to the north of the Kii Peninsula (Figure 1). The molluscan paleontology of the Setouchi strata has been studied in detail (*vide* Itoigawa and Shibata, 1992). For example, Itoigawa (1987, 1988) named and defined the subtropical Akeyo Fauna (late Early Miocene; ca. 18 to 16 Ma) based on mollusks of the Akeyo Formation of the Mizunami Group. Similar fauna has also been recorded from: the Ichishi, Ayugawa and Tsuzuki Groups of the Eastern Setou-

chi Miocene Series in southwestern Honshu; the Kurami Group in central Honshu; the upper part of the Kunugidaira Formation, and the Nakayama and Yotsuyaku Formations in northeastern Honshu (Ozawa *et al.*, 1995; Matsubara, 1995a).

On the other hand, the tropical Kurosedani Fauna (Tsuda, 1960) has been recorded from southwestern Japan, and the contemporaneous subtropical Kadonosawa Fauna has been recorded from northeastern Honshu and southern Hokkaido, northern Japan (Itoigawa, 1988). Both faunas are restricted within the N. 8 zone of Blow (1969) of latest Early to earliest Middle Miocene age (Ogasawara and Noda, 1996). The Akeyo Fauna differs from the Kurosedani Fauna by lacking the tropical mollusks of the latter (Itoigawa, 1988). It also differs from the Kadonosawa Fauna by lacking the characteristic species of the latter: *Anadara (Hataiarca) kakehataensis* Hatai and Nisiyama, *Vasticardium ogurai* (Otuka), *Glycydonta itoigawae* (Tsuda), *Tateiwaia tateiwaia* (Makiyama), *T. yamanarii* (Makiyama), and *Zeuxis minoensis* (Itoigawa) (Itoigawa, 1988).

A Miocene fauna has been reported from several places in the Kii Peninsula (Mizuno, 1953, 1957; Tanai and Mizuno, 1954; Katto and Masuda, 1978; Katto *et al.*, 1976, 1980; Ujihara and Shibata, 1982; Chijiwa and Tomita, 1985;



Figure 1. Distribution of the Miocene strata of the Nankai and Setouchi Geologic Provinces, central and southwestern Honshu (redrawn from: Itoigawa and Shibata, 1992; Editorial Committee of KINKI, ed., Part 6 of Regional Geology of Japan, 1987).

Honda, 1992). However, much data are required to reconstruct the Miocene paleoenvironments in the Nankai Geologic Province.

In 1976 and 1977, one of us (SU) collected several hundred fossil mollusks from a construction site at Ukui Junior High School (Loc. 2; Figures 3, 4), as well as several tens of specimens from an outcrop on a wave-cut terrace at Loc. 7 on Kitsune-jima (Figure 3). Recently, one of us (YH) obtained more specimens from additional localities in the Ukui area. All of these localities are assigned to the Shimosato Formation and together they contain mollusks such as *Saccella miensis* (Araki), *Cyclocardia siogamensis* (Nomura), *Macoma (Macoma) izurensis* (Yokoyama), *Cultellus izumoensis* Yokoyama, *Dosinia (Phacosoma) nomurai* (Otuka), *Periploma (Aelga) mitsuganoense* Araki, *Thracia watanabei* Itoigawa and Shibata, *Turritella (Hataiella) sagai* Kotaka, and *Fulgoraria (Musashia) yanagidaniensis* Araki (Table 1).

The purpose of this study is to record the molluscan assemblages of the Shimosato Formation, and to describe their characteristic species, including those representative of the Akeyo Fauna. We also wish to clarify the molluscan faunal succession of the Kumano Group of the Nankai Geologic Province of southwestern Japan.

Outline of geology and molluscan fauna

The upper Muro Group (Upper Oligocene to Lower Miocene) of the Shimanto Supergroup and the Shimosato

Formation of the Kumano Group are exposed in the Ukui area. The Muro Group consists largely of interbedded gray sandstone and black shale. The Shimosato Formation, on the other hand, is made up of massive, gray, fine- or very fine-grained sandstone, and sandy siltstone. In a wave-cut platform on Kitsune-jima (Loc. 7 in Figure 3), the Shimosato Formation is made up of well-sorted, well-jointed, pale gray, tuffaceous, fine-grained sandstone. The fine-grained sandstone at this locality frequently contains sand-pipes. The Shimosato Formation strikes about N50°E and dips about 10° to 20°SE, and is approximately 200 m thick. Conglomerate of the lowermost Shimosato Formation, which attains a thickness of 30 to 80 cm, rests upon black shale of the uppermost Muro Group with clino-unconformity.

The Kumano Acidic Rocks also crop out in this area. They consist largely of granite porphyry, and intrude into the upper Muro and Kumano Groups. A vertical pyroclastic dike, which is closely related to the Kumano Acidic Rocks (Suzuki, 1976), is exposed in two small cuts at Ukui Junior High School. This dike intrudes into very fine-grained sandstone of the Shimosato Formation, varies in width from approximately 0.5 to 5 m, and trends about N80°E. Figure 3 is a map showing the location of fossil localities in the Shimosato Formation, and Figure 4 shows fossil localities in columnar sections.

Mollusks reported previously from the Shimosato Formation by Ujihara and Shibata (1982) include *Neilonella* cf. *N. isensis* Shibata, *Saccella miensis*, *Portlandia (Portlandella)*




Figure 3. Map showing fossil localities in the Shimosato Formation. 1-7, fossil localities.




Figure 4. Columnar sections of the Shimosato Formation. 1-7, fossil localities.

Table 1. Occurrences of fossil mollusks in the Shimosato Formation. VA, very abundant (20 or more individuals); A, abundant (10 to 19 individuals); C, common (5 to 9 individuals); F, few (2-4 individuals); R, rare (one individual). One individual is defined herein as consisting of more than half of a separated valve or an articulated pair of bivalves, and more than half of a gastropod specimen. See also Figure 5 and Tables 2, 3.

SPECIES	LOCALITY						
	1	2	3	4	5	6	7
Bivalves:							
<i>Acila</i> cf. <i>A. (Acila) submirabilis</i> Makiyama					R	F	R
<i>Acila</i> sp.							
<i>Ennucula</i> sp.		A					
<i>Saccella miensis</i> (Araki)		VA				R	
<i>Solamen</i> sp.		F					
<i>Septifer</i> ? sp.		F					
<i>Chlamys</i> cf. <i>C. iwamurensis</i> Itoigawa		R					
<i>Cycladicama</i> sp.		R					
<i>Cyclocardia siogamensis</i> (Nomura)		VA					C
<i>Cyclocardia</i> sp.		C					
<i>Lucinoma</i> sp.		F	R				
<i>Macra</i> sp.		R					
<i>Macoma (Macoma) izurensis</i> (Yokoyama)		A					
<i>Angulus</i> cf. <i>A. kagayamensis</i> Ogasawara and Tanai		F					
<i>Nitidotellina</i> sp.		R					
<i>Cultellus izumoensis</i> Yokoyama		A					
<i>Disinia (Phacosoma) kawagensis</i> Araki							F
<i>Dosinia (Phacosoma) nomurai</i> (Otuka)		R					A
<i>Dosinia</i> sp.		R					C
<i>Paphia</i> sp.		R					
<i>Glaucanome</i> sp.		F					
<i>Mya</i> sp.		A					
<i>Anisocorbula</i> sp.		R					
" <i>Teredo</i> " sp.			C				
<i>Periploma (Aelga) mitsuganoense</i> Araki		VA		R			
<i>Periploma (Aelga)</i> sp.		VA					
<i>Thracia watanabei</i> Itoigawa and Shibata		F				R	
<i>Thracia</i> sp.					R		
Gastropods:							
<i>Turritella (Hataiella) sagai</i> Kotaka		A					A
<i>Turritella</i> sp.		R					F
<i>Bittium</i> sp.	F	A					
<i>Euspira</i> sp.		R					
<i>Ancistrolepis</i> sp.		A					
<i>Fulgoraria (Musashia) yanagidaniensis</i> Araki		C					
<i>Eocylichna</i> sp.		C					

southwestern Honshu.

Molluscan assemblages and environments

Approximately 400 specimens were collected from 7 localities in the Shimosato Formation. These specimens comprise 22 genera and 28 species of bivalves and 6 genera and 7 species of gastropods (Table 1). Mollusks occur sporadically in the fine- and very fine-grained sandstone and sandy siltstone of the Shimosato Formation. Almost all specimens are complete or nearly so, but all except those from Locality 7 lack original shell material. Two molluscan assemblages

are recognized herein: the *Periploma-Saccella-Cyclocardia* (Loc. 2) and *Turritella-Dosinia* assemblages (Loc. 7). Mollusks from Localities 1 and 3-6 are not used for recognition of assemblages, because each yielded fewer than 10 specimens. Table 2 summarizes the molluscan assemblages and characteristic and associated species.

The *Periploma-Saccella-Cyclocardia* assemblage

This assemblage occurs in beds of sandy siltstone and overlying very fine-grained sandstone at Locality 2. Both beds are up to about 10 m thick. The assemblage is characterized by the dominant occurrence of *Periploma* (*P.*

Table 2. Molluscan assemblages of the Shimosato Formation. vfs : very fine sandstone ; sdy slt : sandy siltstone ; fs : fine sandstone.

MOLLUSCAN ASSEMBLAGE	CHARACTERISTIC SPECIES	ASSOCIATED SPECIES	LITHOLOGY	LOC.
<i>Periploma-Saccella-Cyclocardia</i> Ass.	<i>Periploma mitsuganoense</i> <i>Saccella miensis</i> <i>Cyclocardia siogamensis</i>	<i>Ancistrolepis</i> sp. <i>Bittium</i> sp. <i>Ennucula</i> sp. <i>Mya</i> sp. <i>Cultellus izumoensis</i> <i>T. sagai</i> <i>Macoma izurensis</i>	vfs sdy slt	2
<i>Turritella-Dosinia</i> Assemblage	<i>Turritella sagai</i> <i>Dosinia nomurai</i> <i>D. kawagensis</i>	<i>C. siogamensis</i> <i>Acila</i> cf. <i>A. submilabilis</i>	fs	7

mitsuganoense : Figures 5-12, 5-13, and *P. sp.* : Figure 5-8), *Saccella miensis* (Figure 5-1), and *Cyclocardia* (*C. siogamensis* : Figure 5-14, and *C. sp.*), but also contains *Ancistrolepis* sp., *Bittium* sp., *Ennucula* sp. (Figure 5-2), *Mya* sp. (Figure 5-10), *Cultellus izumoensis* (Figure 5-11), *Turritella sagai* (Figure 5-9), and *Macoma izurensis* (Figure 5-7 ; Table 2). This assemblage typically contains many extant genera that live in upper sublittoral to bathyal depths (N_1 -B) off Japan and neighboring countries, including *Saccella*, *Cyclocardia* and *Turritella* (Table 3). However, it also includes *Periploma*, *Ancistrolepis* and *Musashia*, which live in lower sublittoral to bathyal depths (N_3 -B, Table 3). Consequently, the *Periploma-Saccella-Cyclocardia* assemblage is inferred to represent a lower sublittoral to bathyal environment.

Comparison : The Onuma Formation of the Kumano Group contains a number of species known from this assemblage, including *Saccella miensis*, *Cyclocardia siogamensis*, *Macoma izurensis*, and *Cultellus izumoensis* (Chijiwa and Tomita, 1985). This assemblage is therefore correlative with those of the Onuma Formation, which represent lower sublittoral environments (Chijiwa and Tomita, 1985).

In addition, the *Periploma-Saccella-Cyclocardia* assemblage contains many species in common with the *Periploma-Acila* and *Macoma-Lucinoma* assemblages in the Eastern Setouchi Miocene Series of central and southwestern Honshu, including *Saccella miensis*, *Periploma* sp., *M. izurensis*, *Cultellus izumoensis*, and *Cyclocardia siogamensis* (Shibata, 1978). This assemblage is therefore also correlative with the *Periploma-Acila* and *Macoma-Lucinoma* assemblages, which represent upper sublittoral and sublittoral environments, respectively (Shibata, 1978).

The *Turritella-Dosinia* assemblage


This assemblage occurs in gray, fine-grained sandstone at Locality 7 and is characterized by the dominant occurrence of *Turritella* (*T. sagai* : Figure 5-9, and *T. sp.*) and *Dosinia* (*D. nomurai* : Figures 5-15, 5-19, *D. kawagensis* : Figure 5-16, and *D. sp.*) (Table 2). *Cyclocardia siogamensis* and *Acila* cf. *A. submirabilis* (Figure 5-6) are also present in this assemblage (Table 2). The presence of many intact specimens in this assemblage implies that there was little or no post-mortem transportation. Modern *Turritella*, *Cyclocardia* and

Table 3. Modern bathymetric distribution of extant northwestern Pacific bivalves and gastropods in the Shimosato molluscan fauna (Higo and Goto, 1993). Depth ; T, tidal zone ; N_1 , low-water line to 30 m ; N_2 , 30-60 m ; N_3 , 60-120 m ; N_4 , 120-200 m ; B, over 200 m deep (Taki and Oyama, 1954, partly revised).

GENERA (SUBGENERA)	DEPTH					
	T	N ₁	N ₂	N ₃	N ₄	B
<i>Acila</i> (<i>Acila</i>)						
<i>Ennucula</i>						
<i>Saccella</i>						
<i>Solamen</i>						
<i>Chlamys</i>						
<i>Cycladicama</i>						
<i>Cyclocardia</i>						
<i>Lucinoma</i>						
<i>Mactra</i>						
<i>Macoma</i>						
<i>Angulus</i>						
<i>Nitidotellina</i>						
<i>Phacosoma</i>						
<i>Paphia</i>						
<i>Glaucanome</i>						
<i>Mya</i>						
<i>Anisocorbula</i>						
<i>Periploma</i>						
<i>Thracia</i>						
<i>Turritella</i>						
<i>Bittium</i>						
<i>Euspira</i>						
<i>Ancistrolepis</i>						
<i>Musashia</i>						
<i>Eocylichna</i>						

Acila species live in upper sublittoral to bathyal depths (N_1 (N_2)-B), and *Dosinia* (*Phacosoma*) dwells in the tidal to lower sublittoral zone (T- N_4) (Table 3). The *Turritella-Dosinia* assemblage is therefore inferred to represent a sublittoral environment.

Comparison : This assemblage contains *Turritella sagai*



and *Dosinia nomurai* in common with the *Turritella*-*Lucinoma* assemblage of the Tomikusa Group (upper Lower Miocene) in the Eastern Setouchi Miocene Series of central Honshu (Shibata, 1978). This assemblage is therefore correlative with the *Turritella*-*Lucinoma* assemblage, which represents an upper sublittoral environment (Shibata, 1978).

Discussion

Honda (1992) noted that two types of molluscan assemblage are present in the Kumano and Owase Groups. The first type has been recorded from the Shimosato, Shikiya, and Onuma Formations within the lower and middle parts of the Kumano Group (Mizuno, 1953; Tanai and Mizuno, 1954; Ujihara and Shibata, 1982; Chijiwa and Tomita, 1985); and the Yukunoura Formation of the Owase Group (Honda, 1992). The type is characterized by *Portlandia watasei*, *Cyclocardia siogamensis*, *Lucinoma* sp., *Macoma optiva* (*M. izurensis*), *Cultellus izumoensis*, and *Turritella sagai* (*T. kadonosawaensis*, *T. shataii*, *T. kiiensis*).

The *Periploma*-*Saccella*-*Cyclocardia* and *Turritella*-*Dosinia* assemblages of the Shimosato Formation contain many species in common with those of the first type, including *Cyclocardia siogamensis*, *Macoma izurensis*, *Cultellus izumoensis*, and *Turritella sagai* (Table 2). Both assemblages therefore belong to the first type. In addition, the Shimosato Formation contains *Dosinia kawagensis*, *Thracia watanabei*, *Turritella sagai*, and *Fulgoraria yanagidaniensis* (Table 1), which characterize the fauna of the Akeyo Formation. The fauna of the Shimosato Formation is therefore comparable with the Akeyo Fauna of late Early Miocene age (ca. 18 to 16 Ma; Itoigawa, 1987, 1988) (Figure 2).

On the other hand, the second type of molluscan assemblage has been reported from the middle part of the Shikiya and the lower part of the Mitsuno Formation within the middle and upper parts of the Kumano Group (Mizuno, 1953; Tanai and Mizuno, 1954; Katto and Masuda, 1978; Katto *et al.*, 1980). The type is characterized by many warm-water mollusks such as *Anadara* cf. *A. ogawai*, *Cucullaea toyamaensis*, *Crassatellites pauxillus*, *C. nanus*, *Vasticardium ogurai*, *Mikadotrochus* sp., and *Conus tokunagai*.

Among these species, *Cucullaea toyamaensis* and *Conus tokunagai* characterize the Kurosedani Fauna of southwestern Japan. *Conus tokunagai* is also characteristic in the Kadonosawa Fauna of northern Japan. The middle part of the Shikiya Formation yields *Cucullaea toyamaensis* and *Conus tokunagai* (Katto and Masuda, 1978), in association with larger foraminifers such as *Lepidocyclus japonica* and

Miogyopsina sp. (Nishimura and Miyake, 1973). In addition, *Conus tokunagai* has been reported from the lower part of the Mitsuno Formation (Mizuno, 1953; Tanai and Mizuno, 1954; Katto *et al.*, 1980).

Accordingly, the assemblages of the middle part of the Shikiya Formation and the lower part of the Mitsuno Formation are comparable with those of the Kurosedani (Kadonosawa) Fauna (Figure 2). The occurrence of the Kurosedani (Kadonosawa) Fauna corresponds to the warm marine climatic episode during the latest Early to earliest Middle Miocene period (ca. 16–15 Ma) (Chinzei, 1986).

Such a different stratigraphic occurrence of the molluscan assemblages suggests a warming of climate and not merely a change in the sedimentary environment. The warming of climate is also recognized in the transition from the subtropical Akeyo to tropical Kurosedani Fauna in the Mizunami Group of the Eastern Setouchi Miocene Series (Itoigawa, 1988, 1989).

A similar scenario has also been proposed for another taxonomic group in the Kumano Group. For instance, Nishimura and Miyake (1973) reported larger foraminifers such as *Lepidocyclus japonica* and *Miogyopsina* sp. from the middle part of the Shikiya Formation. Using planktonic foraminifers, Ibaraki (1990) assigned the *Lepidocyclus*-*Miogyopsina*-bearing horizon to the N8.b zone of Blow (1969). From the occurrence of *Orbulina universa* d'Orbigny (Ikebe *et al.*, 1975), the uppermost part of the Shikiya Formation and the Mitsuno Formation correlate with the N.9 zone of Blow (1969) (Hisatomi, 1987) (Figure 2).

As was discussed in the aforementioned lines, the molluscan assemblages of the lower part of the Mitsuno Formation (Mizuno, 1953; Tanai and Mizuno, 1954; Katto *et al.*, 1980) are comparable with those of the Kurosedani (Kadonosawa) Fauna. The fauna, however, is restricted within the N.8 zone of Blow (1969) of latest Early to earliest Middle Miocene age (Ogasawara and Noda, 1996). Further studies are needed to clarify the molluscan faunal succession of the Kumano Group of the Nankai Geologic Province of southwestern Japan.

Summary of faunal discussion

1) The *Periploma*-*Saccella*-*Cyclocardia* and *Turritella*-*Dosinia* assemblages of the Shimosato Formation are inferred to represent lower sublittoral to bathyal and sublittoral environments, respectively.

2) The fauna of the Shimosato Formation is comparable with the subtropical Akeyo Fauna of late Early Miocene age,

Figure 5. 1. *Saccella miensis* (Araki). Loc. 2, MES* 1015. 2. *Ennucula* sp. x2, Loc. 2, MES 1016. 3. 5a-b. *Thracia watanabei* Itoigawa and Shibata. 3; Loc. 2, MES 1017. 5a-b; Loc. 6, MES 1018. 4a-b. *Glaucanome* sp. Loc. 2, MES 1019. 6. *Acila* cf. *A. (Acila) submirabilis* Makiyama. x1.5, Loc. 7, MES 1020. 7. *Macoma (Macoma) izurensis* (Yokoyama). Loc. 2, MES 1021. 8. *Periploma (Aelga)* sp. Loc. 2, MES 1022. 9. *Turritella (Hataiella) sagai* Kotaka. Loc. 7, MES 1023. 10a-c. *Mya* sp. Loc. 2, MES 1024. 11. *Cultellus izumoensis* Yokoyama. Loc. 2, MES 1025. 12a-b, 13. *Periploma (Aelga) mitsuganoense* Araki. 12a-b. Loc. 2, MES 1026. 13; Loc. 2, MES 1027. 14. *Cyclocardia siogamensis* (Nomura). x1.5, Loc. 2, MES 1028. 15, 19. *Dosinia (Phacosoma) nomurai* (Otuka). 15; Loc. 2, MES 1029. 19; Loc. 7, MES 1030. 16. *Dosinia (Phacosoma) kawagensis* Araki. Loc. 7, MES 1031. 17. *Fulgoraria (Musashia) yanagidaniensis* Araki. Loc. 2, MES 1032. 18. *Chlamys* cf. *C. iwamurensis* Itoigawa. Loc. 2, MES 1033. All figures natural size unless otherwise stated. *Abbreviation for the Department of Earth Sciences, Faculty of Education, Mie University.

based on the occurrence of diagnostic species such as *Dosinia kawagensis*, *Thracia watanabei*, *Turritella sagai*, and *Fulgoraria yanagidaniensis*.

3) The assemblages of the middle part of the Shikiya Formation and the lower part of the Mitsuno Formation are comparable with those of the tropical Kurosedani (subtropical Kadonosawa) Fauna of latest Early to earliest Middle Miocene age.

Systematic notes on some important species

Phylum Mollusca
Class Bivalvia
Family Veneridae
Subfamily Dosiniinae
Genus *Dosinia* Scopoli, 1777
Subgenus *Phacosoma* Jukes-Browne, 1912

Dosinia (Phacosoma) kawagensis Araki, 1960

Figure 5-16

Dosinia nomurai Otuka; Itoigawa, 1956, pl. 2, fig. 3 (*non* Otuka, 1934).

Dosinia japonica kawagensis Araki, 1960, p. 95, pl. 7, fig. 3.

Dosinia (Phacosoma) kawagensis Araki; Masuda, 1963, p. 22, pl. 4, figs. 1-8 (fig. 1, reproduced from Araki, 1960); Yoon, 1979, p. 15, pl. 2, figs. 7, 9, 10; Matsubara, 1995b, p. 330, pl. 4, figs. 9-11.

Dosinia kawagensis Araki; Hayashi and Miura, 1973, pl. 2, fig. 12; Hayashi, 1973, pl. 1, figs. 2, 3; Ishida *et al.*, 1980, pl. 3, fig. 12.

Dosinorbis kawagensis (Araki); Itoigawa in Itoigawa *et al.*, 1974, p. 87, pl. 23, figs. 1-6.

Phacosoma kawagensis (Araki); Itoigawa *et al.*, 1981 (1982, p. 80), pl. 14, figs. 6a-b; Shibata and Ina, 1983, p. 48, pl. 5, fig. 12; Itoigawa and Shibata, 1986, pl. 16, fig. 13; Muramatsu, 1992, pl. 49, fig. 10.

Dosinia (Phacosoma) chikuzenensis nomurai Otuka; Hayashi, 1988, p. 5, pl. 1, figs. 2, 3 (reproduced from illustration of *Dosinia kawagensis* in Hayashi (1973)).

Remarks.—Two incomplete closed valves were obtained from fine-grained sandstone of the Shimosato Formation. The present species was originally described by Araki (1960) (as *D. japonica kawagensis*) from the Kaisekizan Formation (upper Lower Miocene; *vide* Yoshida, 1991; Shibata, 1967) of the Ichishi Group, Mie Prefecture, southwestern Honshu. Itoigawa (1956) had earlier cited "*D. nomurai*" from the Tsuzuki Group (upper Lower Miocene) of Kyoto Prefecture, but Masuda (1963) later assigned this occurrence to *D. kawagensis*.

The present species resembles *D. (Phacosoma) nomurai* Otuka, 1934, which was originally described from the Kadonosawa Formation (uppermost Lower to lowermost Middle Miocene; Irizuki and Matsubara, 1994) of Iwate Prefecture, northeastern Honshu. However, *D. (P.) kawagensis* differs by having a more anteriorly expanded anterodorsal margin. The present species also resembles *D. (P.) japonica* (Reeve), which is living in Japan, but differs from the latter by having a smaller and somewhat more inequilateral shell.

The present species has also been recorded from the

upper Lower to lowermost Middle Miocene formations of Honshu and Korea. These include the Akeyo Formation of the Mizunami Group, and other strata of the Eastern Setouchi Miocene Series in central and southwestern Honshu (Itoigawa *et al.*, 1981); the Shiode, Ajiri, and Yotsuyaku Formations of northeastern Honshu (Masuda, 1963; Matsubara, 1995b); and the Sinhyeon Formation of Korea (Yoon, 1979).

Occurrence.—As characteristic species of *Turritella-Dosinia* ass. at Locality 7.

Family Periplomatidae
Genus *Periploma* Schumacher, 1817
Subgenus *Aelga* Slodkewitsch, 1935

Periploma (Aelga) mitsuganoense Araki, 1959

Figures 5-12a-b, 13

Periploma mitsuganoense Araki, 1959, p. 163, pl. 18, figs. 2a-b; Araki, 1960, p. 85, pl. 5, figs. 13a-b (reproduced from Araki, 1959); Shibata in Itoigawa *et al.*, 1974, p. 108, pl. 34, figs. 12-17; Ishida *et al.*, 1980, pl. 4, fig. 21; Itoigawa *et al.*, 1981 (1982, p. 115), pl. 22, figs. 14a-b; Itoigawa and Shibata, 1986, pl. 17, fig. 13; *non* Hayashi, 1973, pl. 5, fig. 1.

Remarks.—Approximately 80 slightly deformed specimens were collected from sandy siltstone of the Shimosato Formation. Araki (1959) based this species on specimens from the Kaisekizan Formation (upper Lower Miocene) of the Ichishi Group, Mie Prefecture, southwestern Honshu. It closely resembles *Periploma (Aelga) besshoense* (Yokoyama, 1924), which was originally described from the Asagai Formation (Lower Oligocene) of the Joban coal field, northeastern Honshu, but is distinguished from the latter in having a more inequilateral shell. *Periploma owasense* Suzuki (1934, p. 348, text-fig. 4), described from the Owase Group (upper Lower to lowermost Middle Miocene) of Mie Prefecture, southeastern Kii Peninsula, is allocated here to *Periploma* sp. indet., owing to the poor preservation of the specimen. *Periploma mitsuganoense* has also been recorded from the Akeyo Formation of the Mizunami Group, and other strata of the Eastern Setouchi Miocene Series (Itoigawa *et al.*, 1981).

Occurrence.—As characteristic species of *Periploma-Saccula-Cyclocardia* ass. at Locality 2 and also from Loc. 4.

Family Thraciidae
Genus *Thracia* Leach, 1824

Thracia watanabei Itoigawa and Shibata, 1975

Figures 5-3, 5a-b

Thracia sp. (n. sp.); Itoigawa in Itoigawa *et al.*, 1974, p. 108, pl. 34, figs. 10-11.

Thracia watanabei Itoigawa and Shibata, 1975, p. 31, pl. 8, figs. 21-24; Itoigawa *et al.*, 1981 (1982, p. 116), pl. 22, fig. 16.

Remarks.—Only four specimens were obtained from gray, very fine-grained sandstone at two localities in the Shimosato Formation. The present species was proposed by Itoigawa and Shibata (1975) from the Yamanouchi Mem-

ber of the Akeyo Formation of the Mizunami Group. *Thracia watanabei* has also been recorded from the Tomikusa, Iwamura and Ayugawa Groups of the Eastern Setouchi Miocene Series in central and southwestern Honshu (Itoigawa *et al.*, 1981).

Occurrence.—Localities 2, 6.

Class Gastropoda
Family Turritellidae
Genus *Turritella* Lamarck, 1799
Subgenus *Hataiella* Kotaka, 1959

Turritella (Hataiella) sagai Kotaka, 1951

Figure 5-9

Turritella s-hataii sagai Kotaka, 1951, p. 87, pl. 12, figs. 13-17.

Turritella (Hataiella) sagai Kotaka; Kotaka, 1959, p. 89, pl. 9, figs. 6-8, 10, 12, 18; Yoon, 1979, p. 21, pl. 5, fig. 11; Marincovich and Kase, 1986, p. 61, pl. 2, figs. F-H (in part); Honda, 1992, pl. 58, fig. 8.

Non Turritella (Hataiella) sagai Kotaka; Marincovich and Kase, 1986, pl. 2, figs. A-E (in part); Marincovich, 1988, p. 15, pl. 3, figs. 3, 5-9 (3, 5-7, 9; reproduced from Marincovich and Kase, 1986, p. 61, pl. 2, figs. A-E); Zhidkova and Sal'nikov, eds., 1992, p. 248, pl. 46, figs. 3a-b.

Turritella sagai Kotaka; Shibata *in* Itoigawa *et al.*, 1974, p. 132, pl. 40, figs. 11, 12; Ishida *et al.*, 1980, pl. 5, figs. 10, 11; Itoigawa *et al.*, 1981 (1982, p. 164), pl. 28, figs. 13-14; Itoigawa and Shibata, 1986, pl. 17, fig. 2; Muramatsu, 1992, pl. 50, fig. 9.

Remarks.—A total of 26 specimens are to hand. The present species was originally described by Kotaka (1951, p. 87, pl. 12, figs. 13-17) from the Togari Member of the Akeyo Formation of the Mizunami Group. As noted by Kotaka (1951), *T. (Hataiella) sagai* resembles *T. (H.) shataii* Nomura, originally described from the "Shiogama" Formation (Lower Miocene; Ishii *et al.*, 1982, p. 14, 20) of Miyagi Prefecture, northeastern Honshu. The present species also resembles *T. (H.) belogolovaensis* Ilyina, originally described from the Kuluven Formation (Lower Miocene) of western Kamchatka (*vide* Titova, 1994, p. 7), but it differs from the latter in having a more slender shell.

Marincovich and Kase (1986, p. 61, pl. 2, figs. A-H) cited *Turritella sagai* from the Bear Lake Formation (lowermost Middle Miocene, in part) of southwestern Alaska (figs. A-E), and the Akebihara Sandstone of the Ayugawa Group (upper Lower Miocene) of southwestern Honshu (figs. F-H). Specimens from the Bear Lake Formation are assigned here to *T. belogolovaensis*, as was done by Titova (1994). Zhidkova and Sal'nikov, eds. (1992, p. 248, pl. 46, figs. 3a-b) cited *T. (H.) sagai* from the Neveliskaya Formation (Lower Miocene) of southern Sakhalin, and this occurrence is also assigned here to *T. belogolovaensis*.

Turritella sagai has been recorded from the Akeyo Formation of the Mizunami Group and other strata of the Eastern Setouchi Miocene Series (Itoigawa *et al.*, 1981), and the Hwabongri Formation (lower Middle Miocene) of Korea (Yoon, 1979).

Occurrence.—As characteristic species of *Turritella-Dosinia* ass. at Loc. 7, and as associated species of *Peri-*

ploma-Sacella-Cyclocardia ass. at Loc. 2.

Family Volutidae
Subfamily Fulgorariinae
Genus *Fulgoraria* Schumacher, 1817
Subgenus *Musashia* Hayashi, 1960

Fulgoraria (Musashia) yanagidaniensis Araki, 1959
Figure 5-17

Fulgoraria hirasei yanagidaniensis Araki, 1959, p. 165, pl. 18, fig. 6; Araki, 1960, p. 104, pl. 8, fig. 5 (reproduced from Araki, 1959).

Musashia (Neopsephaea ?) yanagidaniensis (Araki); Shikama, 1967, p. 115, text-fig. 20, pl. 13, figs. 5-8.

Musashia (Neopsephaea) yanagidaniensis (Araki); Shikama, 1967, pl. 14, fig. 1 (same specimen as Shikama, 1967, pl. 13, fig. 8).

Musashia yanagidaniensis (Araki); Hayashi, 1973, pl. 3, fig. 8; Shibata and Ina, 1983, p. 62, pl. 9, fig. 6.

Psephaea ? yanagidaniensis (Araki); Shibata *in* Itoigawa *et al.*, 1974, p. 167, pl. 51, figs. 1-3, pl. 52, figs. 1, 2, pl. 53, figs. 1-3, pl. 54, figs. 5a-c.

Fulgoraria yanagidaniensis (Araki); Ishida *et al.*, 1980, pl. 6, figs. 34a-b.

Musashia (s. s.) yanagidaniensis (Araki); Itoigawa *et al.*, 1981.

Musashia ? yanagidaniensis (Araki); Itoigawa *et al.*, 1982, p. 239, pl. 40, figs. 1-2.

Fulgoraria (Psephaea ?) cf. ashiyaensis Shikama; Hayashi, 1988, p. 7, pl. 3, fig. 8 (reproduced from *Musashia yanagidaniensis*; Hayashi, 1973).

Remarks.—Seven specimens were collected from gray, very fine-grained sandstone at one locality in the Shimosato Formation. Araki (1959) proposed the present species under the name of *Fulgoraria hirasei yanagidaniensis* from the Kaisekizan Formation (upper Lower Miocene) of the Ichishi Group. The present species has also been recorded from the Akeyo Formation of the Mizunami Group and other strata of the Eastern Setouchi Miocene Series (Itoigawa *et al.*, 1981).

Occurrence.—Locality 2.

Acknowledgments

We wish to express our deep gratitude to Tamio Kotaka (Emeritus Professor of Tohoku University) for his continuous encouragement of our study of the Cenozoic molluscan faunas of Japan, and to Kazuo Okamoto (Emeritus Professor of Hiroshima University) for providing the opportunity to study fossil mollusks from the Shimosato Formation of the Ukui area. We would like to thank Kenshiro Ogasawara of the University of Tsukuba for valuable comments on the manuscript. Thanks are expressed to Louie Marincovich, Jr. of Earth Science Editing for critical review of the manuscript. Thanks are also expressed to Paul Callomon of Elle Scientific Publications for proofreading an early draft of the manuscript.

References

Araki, Y., 1959: On some Miocene Mollusca from Mie

- Prefecture, Japan. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 36, p. 161-167, pl. 18.
- Araki, Y., 1960: Geology, paleontology and sedimentary structures (including problematica) of the Tertiary formations developed in the environs of Tsu City, Mie Prefecture, Japan. *Bulletin of the Liberal Arts Department, Mie University, Special Volume*, no. 1, p. 1-118, pls. 1-11.
- Blow, W.H., 1969: Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. In Brönnimann, P. and Renz, H.H. eds., *Proceedings of the First International Conference on Planktonic Microfossils*, Geneva, 1967, Vol. 1, p. 119-422, pls. 1-54, E.J. Brill, Leiden, The Netherlands.
- Chijiwa, K., 1988: Post-Shimanto sedimentation and organic metamorphism: an example of the Miocene Kumano Group, Kii Peninsula. *Modern Geology*, vol. 12, p. 363-387.
- Chijiwa, K. and Tomita, S., 1985: On the Onuma Formation of the Kumano Group (A study of the Tertiary formations of the Kumano coal-field in the Kii Peninsula, Southwest Japan, Part 3). *Memoirs of the Faculty of Science, Kyushu University, Ser. D, Geology*, vol. 25, no. 3, p. 319-336, pl. 41.
- Chinzei, K., 1986: Faunal succession and geographic distribution of Neogene molluscan faunas in Japan. *Palaeontological Society of Japan, Special Papers*, no. 29, p. 17-32.
- Editorial Committee of KINKI, ed., 1987: *Regional Geology of Japan Part 6 KINKI*, 297 p. Kyoritsu Shuppan Co., Tokyo. (in Japanese)
- Hayashi, T., 1973: Fossils from the environs of Horaijisan. *Bulletin of the Horaijisan Natural Science Museum*, p. 10-17, incl. pls. 1-6. (in Japanese)
- Hayashi, T., 1988: Reexamination of fossils from the environs of Horaijisan. *Bulletin of the Horaijisan Natural Science Museum*, no. 18, p. 3-14, incl. pls. 1-6. (in Japanese)
- Hayashi, T. and Miura, Y., 1973: The Cenozoic sediments in the southern part of Okazaki City, central Japan. *The Bulletin of Aichi University of Education*, vol. 22, *Natural Science*, p. 133-149, incl. pls. 1-2. (in Japanese with English abstract)
- Higo, S. and Goto, Y., 1993: *A Systematic List of Molluscan Shells from the Japanese Is. and the Adjacent Area*, 879 p. Elle Scientific Publications, Yao, Osaka. (in Japanese)
- Hisatomi, K., 1984: Sedimentary environment and basin analysis of the Miocene Kumano Group in the Kii Peninsula, Southwest Japan. *Memoirs of the Faculty of Science, Kyoto University, Series of Geology and Mineralogy*, vol. 50, nos. 1-2, p. 1-65, pls. 1-4.
- Hisatomi, K., 1987: The Kumano Group. In Editorial Committee of KINKI ed., *Regional Geology of Japan Part 6 KINKI*, p. 128-131. Kyoritsu Shuppan Co., Tokyo. (in Japanese)
- Hisatomi, K. and Miyake, Y., 1981: Upheaval movement and igneous activity in the Shionomisaki area, Kii Peninsula, Southwest Japan. *The Journal of the Geological Society of Japan*, vol. 87, no. 10, p. 629-639. (in Japanese with English abstract)
- Honda, Y., 1992: A Miocene molluscan fauna from the Owase Group, Kii Peninsula, Southwest Honshu. *Bulletin of the Mizunami Fossil Museum*, no. 19, p. 413-425, pl. 58.
- Ibaraki, M., 1990: Geologic ages of *Lepidocyclina* in Japan with implications in the northward drift of the Izu Peninsula. In Tsuchi, R. ed., *Pacific Neogene Events, Their Timing, Nature and Interrelationship*, p. 137-149. University of Tokyo Press.
- Ikebe, N., Chiji, M. and Morozumi, Y., 1975: *Lepidocyclina* horizon in the Miocene Kumano Group in reference to planktonic foraminiferal biostratigraphy. *Bulletin of the Osaka Museum of Natural History*, no. 29, p. 81-89, pls. 20-21. (in Japanese with English abstract)
- Irizuki, T. and Matsubara, T., 1994: Vertical changes of depositional environments of the Lower to Middle Miocene Kadonosawa Formation based on analysis of fossil ostracode faunas. *The Journal of the Geological Society of Japan*, vol. 100, no. 2, p. 136-149. (in Japanese with English abstract)
- Ishida, S., Okamura, Y. and Matsuoka, C., 1980: Stratigraphy and fossils of the Ayugawa Group. In Branch of Geomorphology and Geology, Nature Study Society of Shiga Prefecture ed., *Land and Life in Shiga, Separate Volume for Geomorphology and Geology and Geological Map of Shiga Prefecture 1: 100,000 in Scale*, p. 251-299, incl. pls. 1-6. Foundation of Nature Conservation in Shiga Prefecture. (in Japanese with English summary)
- Ishii, T., Yanagisawa, Y., Yamaguchi, S., Sangawa, A. and Matsuno, K., 1982: *Geology of the Matsushima District. Quadrangle Series, Scale 1: 50,000*, 121 p. Geological Survey of Japan. (in Japanese with English abstract)
- Itoigawa, J., 1956: Molluscan fauna of the Tsuzuki Group in Kyoto Prefecture, Japan. *Memoirs of the College of Science, University of Kyoto, Ser. B (Geology and Mineralogy)*, vol. 23, no. 2, p. 179-192, pls. 1-3.
- Itoigawa, J., 1987: Early to Middle Miocene molluscan faunas in Southwest Japan and Korea. *Geological Reports of Shimane University*, no. 6, p. 29-38. (in Japanese with English abstract)
- Itoigawa, J., 1988: The Miocene Kadonosawa fauna of Japan. In Grant-Mackie, J.A., Masuda, K. and Ogasawara, K. eds., *Professor Tamio Kotaka Commemorative Volume on Molluscan Paleontology, Special Publication, Saito Ho-on Kai Museum of Natural History*, no. 2, p. 397-403.
- Itoigawa, J., 1989: Tropical spike in early Middle Miocene (ca. 16 MA) of Southwest Japan. In Gengwu, L., Tsuchi, R. and Qibin, L. eds., *Proceedings of International Symposium of Pacific Neogene Continental and Marine Events*, p. 19-26. National Working Group of China for IGCP-246, Nanjing University Press, Nanjing.
- Itoigawa, J. and Shibata, H., 1975: New Miocene pelecypods from the Mizunami Group, Mizunami City, central Japan. *Bulletin of the Mizunami Fossil Museum*, no. 2, p. 15-34, pls. 6-8.
- Itoigawa, J. and Shibata, H., 1986: Molluscan fauna of the Setouchi Miocene Series, Southwest Japan. *Palaeontological Society of Japan, Special Papers*, no. 29, p. 149-159, pls. 16-17.
- Itoigawa, J. and Shibata, H., 1992: Miocene paleogeography of the Setouchi Geologic Province, Japan, a revision. *Bulletin of the Mizunami Fossil Museum*, no. 19, p. 1-12. (in Japanese with English abstract)

- Itoigawa, J., Shibata, H. and Nishimoto, H., 1974 : Molluscan fossils from the Mizunami Group. *Bulletin of the Mizunami Fossil Museum*, no. 1, p. 43-203, pls. 1-63. (in Japanese)
- Itoigawa, J., Shibata, H., Nishimoto, H., and Okumura, K., 1981 : Miocene fossils of the Mizunami Group, central Japan. 2. Molluscs. *Monograph of the Mizunami Fossil Museum*, no. 3-A, p. 1-53, pls. 1-52. (in Japanese)
- Itoigawa, J., Shibata, H., Nishimoto, H., and Okumura, K., 1982 : Miocene fossils of the Mizunami Group, central Japan. 2. Molluscs (Continued). *Monograph of the Mizunami Fossil Museum*, no. 3-B, p. 1-330. (in Japanese)
- Katto, J. and Masuda, K., 1978 : Tertiary Mollusca from the southern part of Kii Peninsula, Wakayama Prefecture, Southwest Japan. *Research Reports of the Kochi University*, vol. 27, *Natural Science*, p. 97-111, pls. 1-5.
- Katto, J., Masuda, K. and Sako, Y., 1976 : A note on the so-called Upper Muro Group, Wakayama Prefecture. *Research Reports of the Kochi University*, vol. 24, *Natural Science*, no. 15, p. 1-10, pls. 1-4. (in Japanese with English abstract)
- Katto, J., Masuda, K., Taira, A., and Sako, Y., 1980 : On the molluscan fauna and lithofacies of the Kumano Group at the Komugi area, Nachikatsuura Town, Wakayama Prefecture. In, Taira, A. and Tashiro, M. eds., *Selected Papers in Honor of Prof. Jiro Katto, Geology and Paleontology of the Shimanto Belt*, p. 101-108, pls. 14-15. Rinyakosaikai Press, Kochi. (in Japanese with English abstract)
- Kotaka, T., 1951 : Recent *Turritella* of Japan. *Short Papers from the Institute of Geology and Paleontology, Tohoku University*, no. 3, p. 70-90, pls. 11-12.
- Kotaka, T., 1959 : The Cenozoic Turritellidae of Japan. *The Science Reports of the Tohoku University, Second Series (Geology)*, vol. 31, no. 2, p. 1-135, pls. 1-15.
- Marincovich, L., Jr., 1988 : Recognition of an earliest Middle Miocene warm-water event in a southwestern Alaskan molluscan fauna. In, Grant-Mackie, J.A., Masuda, K. and Ogasawara, K. eds., *Professor Tamio Kotaka Commemorative Volume on Molluscan Paleontology, Special Publication, Saito Ho-on Kai Museum of Natural History*, no. 2, p. 1-24, pls. 1-3.
- Marincovich, L., Jr. and Kase, T., 1986 : An occurrence of *Turritella (Hataiella) sagai* in Alaska : implications for the age of the Bear Lake Formation. *Bulletin of the National Science Museum, Tokyo, Ser. C*, vol. 12, no. 2, p. 61-66.
- Masuda, K., 1963 : Three Miocene *Dosinia* from near Sendai City, Miyagi Prefecture, Northeast Honshu, Japan, with reference to some Miocene *Dosinia*. *Saito Ho-on Kai Museum of Natural History, Research Bulletin*, no. 32, p. 18-29, incl. pls. 3-4.
- Matsubara, T., 1995a : Fossil Mollusca of the Lower Miocene Yotsuyaku Formation in the Ninohe district, Iwate Prefecture, Northeast Japan. Part 1. General consideration of the fauna. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 180, p. 303-320.
- Matsubara, T., 1995b : Fossil Mollusca of the Lower Miocene Yotsuyaku Formation in the Ninohe district, Iwate Prefecture, Northeast Japan. Part 2 (1). Bivalvia and Scaphopoda. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 180, p. 321-335.
- Mizuno, A., 1953 : Notes on the Miocene molluscs from the Kumano Group in the south-eastern Kii Peninsula, Japan. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 9, p. 9-18.
- Mizuno, A., 1957 : *Explanatory text of the geological map of Japan. Scale 1 : 50,000, Nachi*, 40 p. Geological Survey of Japan. (in Japanese with English abstract)
- Muramatsu, K., 1992 : Geology and molluscan assemblages of the Ayugawa Group, Shiga Prefecture, Japan. *Bulletin of the Mizunami Fossil Museum*, no. 19, p. 301-317, pls. 48-50. (in Japanese with English abstract)
- Nishimura, A. and Miyake, Y., 1973 : Occurrences of *Lepidocyclina* and *Miogyopsina* from the Kumano Group. *Collected Papers from Symposium on the Shimanto Geosyncline*, p. 37-38. Published by Wakayama University, Wakayama. (in Japanese)
- Ogasawara, K. and Noda, H., 1996 : Miocene *Hataiarca* (Mollusca, Bivalvia) invasion event in the Japanese Islands from a viewpoint of Indo-Pacific connection. In, Noda, H. and Sashida, K. eds., *Professor Hisayoshi Igo Commemorative Volume on Geology and Paleontology of Japan and Southeast Asia*, p. 133-139.
- Otuka, Y., 1934 : Tertiary structures of the northwestern end of the Kitakami mountainland, Iwate Prefecture, Japan. *Bulletin of the Earthquake Research Institute*, vol. 12, part 3, p. 566-638, pls. 44-51.
- Otuka, Y., 1939 : Tertiary crustal deformations in Japan (with short remarks on Tertiary paleogeography). *Jubilee Publication in the Commemoration of Prof. H. Yabe, M.I.A. Sixtieth Birthday*, vol. 1, p. 481-519.
- Ozawa, T., Inoue, K., Tomita, S., Tanaka, T., and Nobuhara, T., 1995 : An outline of the Neogene warm-water molluscan faunas in Japan. *Fossils*, no. 58, p. 20-27. (in Japanese with English abstract)
- Saeki, H. and Koto, J., 1972 : Geology and ore deposits of the central Kii Peninsula. *Mining Geology*, vol. 22, no. 7, p. 437-447. (in Japanese with English abstract)
- Shibata, H., 1967 : Geology of the Miocene Ichishi Group. *The Journal of the Geological Society of Japan*, vol. 73, no. 7, p. 337-346. (in Japanese with English abstract)
- Shibata, H., 1978 : Molluscan paleoecology of the Miocene First Setouchi Series in the eastern part of the Setouchi geologic province, Japan. *Bulletin of the Mizunami Fossil Museum*, no. 5, p. 23-110.
- Shibata, H. and Ina, H., 1983 : Mollusks and plants from the Shidara Group (Miocene), central Japan. *Monograph of the Mizunami Fossil Museum*, no. 4, p. 1-89, pls. 1-24. (in Japanese)
- Shikama, T., 1967 : System and evolution of Japanese fulgorarid Gastropoda. *Science Reports of the Yokohama National University, Sec. 2, Biological and Geological Sciences*, no. 13, p. 23-132, pls. 1-17.
- Suzuki, H., 1976 : The Muro Group and the Kumano Group of the Ukui district in the Kii Peninsula, Southwest Japan. *The Science and Engineering Review of Doshisha University*, vol. 17, no. 2, p. 126-137. (in Japanese with English abstract)
- Suzuki, K., 1934 : On the geological age of the Miyai Coal-bearing Tertiary in the southeastern Kii Peninsula. *Chikyū (The Globe)*, vol. 21, no. 5, p. 329-364. (in

- Japanese)
- Taki, I. and Oyama, K., 1954: Matajiro Yokoyama's the Pliocene and later faunas from the Kwanto region in Japan. *Palaeontological Society of Japan, Special Papers*, no. 2, p. 1-68, pls. 1-49.
- Tanai, T. and Mizuno, A., 1954: Geological structure in the vicinity of the Kumano coal field in southeastern Kii Peninsula. *The Journal of the Geological Society of Japan*, vol. 60, no. 700, p. 28-39. (in Japanese with English abstract)
- Titova, L.V., 1994: Revision of the Neogene Turritellidae (Mollusca: Gastropoda) from Kamchatka. *Ruthenica*, vol. 4, no. 1, p. 1-20.
- Tsuda, K., 1960: Paleo-ecology of the Kurosedani fauna. *Journal of the Faculty of Science, Niigata University*, Ser. 2, vol. 3, no. 4, p. 171-203.
- Ujihara, A. and Shibata, H., 1982: Molluscs and elasmobranchs from the Miocene Kumano Group in the southern part of the Kii Peninsula, Japan. *Bulletin of the Mizunami Fossil Museum*, no. 9, p. 25-33, pls. 5-6. (in Japanese with English abstract)
- Yokoyama, M., 1924: Molluscan remains from the lowest part of the Jōban coal-field. *Journal of the College of Science, Tokyo Imperial University*, vol. 45, p. 1-22, pls. 1-5.
- Yoon, S., 1979: The Tertiary deposits of the Ulsan Basin. No. 2. Molluscan Fossils. *The Journal of the Geological Society of Korea*, vol. 15, no. 1, p. 1-36, incl. pls. 1-5.
- Yoshida, F., 1991: Planktonic foraminifera from the Ichishi, Fujiwara, and Morozaki groups in the Eastern Setouchi Geologic Province, central Japan. *Bulletin of the Mizunami Fossil Museum*, no. 18, p. 19-31, pls. 1-2.
- Zhidkova, L., S. and Sal'nikov, B.A. eds., 1992: *Reference section of Paleogene-Neogene deposits of south-east Sakhalin (Makarov Section)*, 357 p., 68 pls. All-Russia Petroleum Scientific-Research Geological-Exploration Institute (VNIGRI), St.-Petersburg. (in Russian)

Ajiri 網尻, Akebihara 山女原, Akeyo 明世, Asagai 浅貝, Ayugawa 鮎川, Ichishi 一志, Iwamura 岩村, Kadosawa 門ノ沢, Kaisekizan 貝石山, Kitsune-jima キツネ島, Kurosedani 黒瀬谷, Koguchi 小口, Kumano 熊野, Kunugidaira 櫛平, Kurami 倉真, Mitsuno 三津野, Muro 牟婁, Nachikatsuura 那智勝浦, Nakayama 中山, Nankai 南海, Onuma 大沼, Osone 大曾根, Owase 尾鷲, Semotobana 瀬元鼻, Setouchi 瀬戸内, Shikiya 敷屋, Shimosato 下里, Shiode 塩手, Shiogama 塩釜, Tanabe 田辺, Togari 戸狩, Tomikusa 富草, Tsuzuki 綴喜, Uematsu 植松, Ukui 宇久井, Yamanouchi 山野内, Yotsuyaku 四ツ役, Yukunoura 行野浦



Honda, Yutaka, Ushiro, Seisuke, and Moritani, Sayuri. 1998. "Miocene mollusks from the Kumano Group of the Ukui area, southeastern part of the Kii Peninsula, southwestern Honshu, Japan." *Paleontological research* 2, 12–24.

View This Item Online: <https://www.biodiversitylibrary.org/item/124737>

Permalink: <https://www.biodiversitylibrary.org/partpdf/82927>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.