Recent Literature.

the Lower California bird receiving the new name castaneiceps. Seven species and ten subspecies of Golden Warblers are now recognized only two of which (*D. æstiva* and *D. bryanti castaneiceps*) pertain to the fauna of North America, as defined in the new 'A. O. U. Check List,' the remainder being West Indian and Central American.—J. A. A.

Ridgway on the Type Specimen of Buteo oxypterus Cassin.* — A reexamination of the type specimens of *Buteo oxypterus* Cassin, preserved in the Museum of the Philadelphia Academy of Natural Sciences, has led M1. Ridgway to refer *B. oxypterus* to *B. swainsoni* as a pure synonym, the type of oxypterus proving to be unquestionably a young *B. swainsoni*. "Upon the whole," says Mr. Ridgway, "I cannot see the slightest reason for recognizing '*B. oxypterus*' even as a local race of swainsoni," He also adds that the "specimen described in 'History of North American Birds' (III, p. 266) as the melanistic adult of '*Buteo swainsoni* var. oxypterus is not *B. swainsoni* at all, but *B. fuliginosus* Scl." Comparative diagnoses are given of these two species.—J. A. A.

Ridgway's List of Emended Names of North American Birds.[†]—The names "represent new or hitherto unpublished combinations" adopted by the A. O. U. 'Committee on Classification and Nomenclature of North American Birds.' "They are here presented in order that the first known use of such combinations may be cited among the references which the committee has decided to give under each species, viz., the first pertinent binomial or trinomial appellation, and the first use of the name as adopted in the new list now being prepared by the committee." The changes, some 77 in number, relate chiefly to the generic part of the name, and result largely from the depression of formerly current genera to the rank of subgenera, partly from the change of status of the form in question from specific to subspecific rank, or the reverse, and in small part from actual change of names, found necessary on synonymic grounds. Comparatively few are therefore of a very 'radical' character, and are mainly the following, with which are given their current equivalents.

Parus cinctus obtectus (Cab.) = P. cinctus. Sylvania microcephala Ridgw. = Myiodioctes minutus. Pyranga rubra cooperi = P. æstiva cooperi. Loxia curvirostra stricklandi Ridgw. = L. c. mexicana. Ammodramus savannarum passerinus = Coturniculus passerinus. Melospiza georgiana (Lath.) = M. palustris. Corvus corax sinuatus (Wagl.) = C. c. carnivorus. Aphelocoma sieberii arizonæ = A. sordida arizonæ. Zenaidura macroura = Z. carolinensis.

* Remarks on the Type Specimen of Buteo oxypterus, Cassin. By Robert Ridgway, Proc. U. S. Nat. Mus., VIII, pp. 75-77. Published April 22, 1885.

† Some Emended Names of North American Birds. By Robert Ridgway, Proc. U. S. Nat. Mus., VIII, pp, 354, 355. Published Sept. 2, 1885.

1885.]



Ridgway, Robert. 1885. "Ridgway on the Type Specimen of Buteo Oxypterus Cassin." *The Auk* 2, 371–371. <u>https://doi.org/10.2307/4625305</u>.

View This Item Online: https://doi.org/10.2307/4625305 Permalink: https://www.biodiversitylibrary.org/partpdf/86921

Holding Institution Smithsonian Libraries and Archives

Sponsored by Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.