

Chaetogaster limnaei

(Oligochaeta : Naididae)

Inhabiting the Mantle Cavity of the Asiatic Clam,
Corbicula fluminea, in Barkley Lake, Kentucky

BY

JAMES B. SICKEL AND MARK B. LYLES

Department of Biological Sciences, Murray State University, Murray, Kentucky 42071

DURING A TWO YEAR SAMPLING program in which the Asiatic clam, *Corbicula fluminea* (Müller, 1774), was being collected from Barkley Lake (Cumberland River), Trigg County, Kentucky, a naidid oligochaete, *Chaetogaster limnaei* von Baer, 1827, was discovered living on the gills within the mantle cavity of the clams during a brief two month period in late spring. ENG (1976) first reported *C. limnaei* in the Asiatic clam from California and indicated that infestation was seasonal with the highest prevalence (87%) occurring from March through May. He noted no evidence of parasitism and reported a low intensity of symbionts within their host (several worms per clam).

Chaetogaster limnaei was first noticed in *Corbicula* from Barkley Lake (Cumberland River Mile 67.0) on 21 May 1980 when 4 clams were being examined to determine their reproductive condition and were found to contain many worms. The clams were in poor condition, evidenced by their light weight and emaciated appearance. Clams were collected again on 4 June, and one was found to contain between 40 and 50 worms crawling on the gills, foot and mantle surface. Dissection of the clam revealed no worms internally, but there were a number of unidentified hyaline spheres in the gonads along with deteriorating clam eggs. Those clams in poorest physical condition, which generally had a high intensity of *C. limnaei*, possessed nodules or calcareous concretions on the inner shell surface producing a chalky appearance. Clams kept in aquaria died within a few days and floated to the surface. No *C. limnaei* were found in dead clams or in aquarium water putrefied by dead clams. Clams were collected again on 6, 17, and 24 June and the prevalence of infection was greater than 80% with the highest intensity in one clam being 167 worms. On 15 August many floaters, floating dead clams with closed shells, were observed in Barkley

Lake. Twelve live and 8 dead *Corbicula* were examined and no *C. limnaei* were found. Over half of the clams in the bed at Cumberland River Mile 67.0 were dead. Most of those still alive showed signs of stress—emaciated body, separating gill filaments, and rough inner surface of shell.

The effect of *Chaetogaster limnaei* on *Corbicula* is uncertain. In one clam, worms were observed between the mantle and shell. However, mantle tissue of clams in poor condition was very thin and easily torn. It is not known whether the worms penetrated the mantle on their own or crawled through the lesions caused by some other agent.

Chaetogaster limnaei has been characterized as either parasitic or commensal on snails (BRINKHURST & JAMIESON, 1971). GRUFFYDD (1965) provided evidence that two forms of *Chaetogaster limnaei* exist, one commensal on *Lymnaea pereger*, *C. limnaei limnaei*, and the other *C. limnaei vaghini* Gruffydd, 1965, parasitic in the kidney of *L. pereger*. GAMBLE & FRIED (1976) reported *C. l. limnaei* feeding on subepithelial tissue of *Physa acuta*. *Chaetogaster limnaei* has been reported also from fingernail clams, *Sphaerium* (BARBOUR, 1977) and mussels, Unionidae, (COKER *et al.*, 1921).


Although more *Chaetogaster limnaei* were found in clams in poor condition than in healthy clams, there was no conclusive evidence indicating parasitism. The relationship between *C. limnaei* and *Corbicula fluminea* is uncertain and needs further investigation.

ACKNOWLEDGMENTS

The authors thank Jarl K. Hiltunen, Great Lakes Fishery Laboratory, for confirming the identification of *Chaetogaster limnaei* and Murray State University students Mark A. Young and Michael L. Russell for assistance in collecting and examining clams.

Literature Cited

- BARBOUR, MICHAEL T.
1977. *Chaetogaster limnaei limnaei* (Oligochaeta: Naididae) inhabiting the mantle cavity of the pill clam *Sphaerium*. Trans. Amer. Microsc. Soc. 96 (1): 141 - 142
- BRINKHURST, R. O. & B. G. M. JAMIESON
1971. Aquatic Oligochaeta of the world. Univ. Toronto Press; 860 pp.
- COKER, R. E., A. F. SHIRA, H. W. CLARK & A. D. HOWARD
1921. Natural history and propagation of fresh-water mussels. Bull. U. S. Bur. Fish., 37: 77 - 181
- ENG, LARRY L.
1976. A note on the occurrence of a symbiotic oligochaete, *Chaetogaster limnaei*, in the mantle cavity of the Asiatic clam, *Corbicula manilensis*. The Veliger 19 (2): 208 (1 October 1976)
- GAMBLE, H. RAY & BERNARD FRIED
1976. Experimental evidence for parasitism in the relationship between *Chaetogaster limnaei limnaei* (Oligochaeta) and *Physa acuta* (Gastropoda). The Veliger 18 (4): 393-395; 1 plt. (1 April 1976)
- GRUFFYDD, LL. D.
1965. Evidence for the existence of a new subspecies of *Chaetogaster limnaei* (Oligochaeta) in Britain. Journ. Zool. 146: 175 - 196





Sickel, Jb and Lyles, Mb. 1981. "CHAETOGASTER-LIMNAEI (OLIGOCHAETA, NAIDIDAE) INHABITING THE MANTLE CAVITY OF THE ASIATIC CLAM, CORBICULA-FLUMINEA, IN BARKLEY LAKE, KENTUCKY." *The veliger* 23, 361–362.

View This Item Online: <https://www.biodiversitylibrary.org/item/134165>

Permalink: <https://www.biodiversitylibrary.org/partpdf/93727>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: California Malacozoological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.