A CLASSIFIED CATALOGUE OF AMERICAN LAND SHELLS, WITH LOCALITIES.

BY HENRY A. PILSBRY.

Almost a decade has elapsed since the publication of the "Check List of American Land Shells" (Proc. Acad. N. S. Phila., 1889), and the frequent calls for copies of that list since the reprint edition was exhausted, as well as the progress of the science during that period, have induced the editors of The Nautilus to offer a new list covering the same ground.

In the nomenclature of this list we have admitted changes shown to be necessary; and in the classification have frequently forsaken the old roads for paths lately "blazed out," and as yet not much travelled.

Additions and corrections of any kind will be gratefully received.

Class GASTROPODA.

Subclass STREPTONEURA.

Order PROSOBRANCHIATA.

Suborder Tænioglossa.

Family CYCLOSTOMATIDÆ.

Genus CHOANOPOMA Pfeiffer, 1847.

Subgenus Ctenopoma Shuttleworth, Pfr., 1856.

1. Choanopoma Bahamense Shuttl. Key West, Fla. (Dr. W. H. Rush). Also Bahamas.

Genus CHONDROPOMA Pfeiffer, 1847.

2. CHONDROPOMA DENTATUM (Say). Key West and Naples, near Gordon's Pass, Fla. Also Cuba.

Family TRUNCATELLIDÆ.

Genus TRUNCATELLA Risso, 1826.

- 3. Truncatella caribæensis 'Sowb.' Rve. Key Largo and Anclote Key, Fla. Also West Indies.
- 4. TRUNCATELLA CARIBÆENSIS PULCHELLA Pfr. Marco and Cedar Keys, Fla. Also Bermuda, Cuba, Yucatan, etc.

- 5. TRUNCATELLA BILABIATA Pfr. Key West, Key Largo, Biscayne Bay, Micco, Indian River, Furguson's Pass, Marco, Cedar Keys, Fla. Also Bermuda, Cuba, etc.
- 6. Truncatella stimpsoni Stearns. False Bay, near San Diego, California.
 - 7. TRUNCATELLA CALIFORNICA Pfr. San Diego, California.

[Note.—T. subcylindrica "Gray" has been reported from Florida. The identity of the species is doubtful, and the occurrence in Florida of the form so-called by Pfeiffer, equally so].

Suborder Rhipidoglossa.

Family HELICINIDÆ.

Genus HELICINA Lamarck, 1801.

- 8. Helicina Chrysocheila Binney. Texas, near mouth of the Rio Grande. Also State of Tamaulipas, Mexico.
- 9. Helicina orbiculata Say. St. Simon's I., Ga., Chattanooga, Tenn., Eureka Springs, Ark., Stone Co., Mo., southward to southern Florida, and southwest to the Rio Grande; type locality near mouth of the St. John's River, Florida. [H. Hanleyana Pfr. is a synonym].

9a. HELICINA ORBICULATA TROPICA 'Jan' Pfr. Texas, mainly south and southwest. An ill defined race.

10. Helicina occulta Say. Western Pa. to Minn., south to Tenn. Distribution markedly discontinuous and local. Allegheny Co., Pa.; Athens, Ill.; near Iowa City and Eldora, Iowa; Winona, Minn.; Sheboygan, Whitefish Bay near Milwaukee, and near Du Pere, Wis.; Lexington and at Natural Bridge, Va.; Harriman and South Pittsburg, Tenn. An abundant fossil of the Post-Pliocene Loess formation in eastern and central Iowa, the adjacent parts of Ill., western Ia. and E. Neb.; also fossil at New Harmony, Ind., the type locality.

[Note.—H. subglobulosa Poey, a Cuban species introduced into United States works on the evidence of specimens found at Key Biscayne and Fort Dallas, Fla., many years ago, has not yet been shown to be actually living in Florida].

Subclass EUTHYNEURA.

Order PULMONATA.

Suborder Stylommatophora.

(Superfamily HOLOPODA).

Family HELICIDÆ.

Subfamily Helicinæ (vel Belogona).

(Belogona Siphonadenia).

Genus HELIX Linné, 1758.

Section Helicogena Férussac.

11. Helix Aspersa Müll. Charleston, S. C.; New Orleans; Santa Barbara, Santa Clara and San Jose, Cal. Imported from Europe.

Section Tachea 'Leach,' Turton, 1831.

- 12. Helix Nemoralis Linn. Burlington, N. J.; Lexington, Va.; ? Baraboo, Sauk Co., Wis. Imported from northern Europe.
- 13. Helix hortensis Müll. Labrador; Anticosti I.; Barachois, Gaspéregion; Nova Scotia; Casco Bay, Me.: Eagle, House, Kettle, and Nantucket Is., Mass.; also New Bedford, Marblehead, Manchester and Magnolia, Mass.

Genus HYGROMIA Risso, 1826.

Section Fruticicola Held, 1837.

- 14. HYGROMIA HISPIDA (L.). Halifax, N. S.; Montreal. A species of northern Europe, imported.
- 15. HYGROMIA RUFESCENS (Penn.). Quebec and Levis, Quebec, Canada. A species of northern Europe, imported.

Genus HELICELLA Férussac.

Section Trochula Schlüter, 1838.

16. Helicella terrestris (Penn.). Charleston, S. C. A native of southern Europe.

Subgenus Cochlicella Risso, 1826.

17. Helicella ventricosa (Drap.). Sullivan's I., S. C. Imported from Europe.

Subgenus Theba Risso, 1826.

18. Helicella cantiana (Montagu). Quebec. A native of northern and central Europe.

(Belogona Euadenia).

Genus GLYPTOSTOMA Bland & Binney, 1873.

19. GLYPTOSTOMA NEWBERRYANUM (W. G. Binn.). Los Angeles, Cal., to Todos Santos Bay, Lower Cal., and about 40 miles inland.

Genus LYSINOE H. & A. Adams, 1855.

20. Lysinoe humboldtiana (Val.). Altuda, Texas. Also Mexico.

Genus EPIPHRAGMOPHORA Doring, 1875.

Subgenus Monadenia Pilsbry, 1895.

- 21. EPIPHRAGMOPHORA FIDELIS (Gray). Humboldt and Shasta Co., Cal., to Vancouver I.
 - 21a. EPIPHRAGMOPHORA FIDELIS, f. flava Hemph.
 - 21b. EPIPHRAGMOPHORA FIDELIS f. minor.
- 21c. EPIPHRAGMOPHORA FIDELIS SUBCARINATA Hemph. Humboldt Co., Cal. This is directly intermediate between *fidelis* and *infumata*.
- 21d. EPIPHRAGMOPHORA FIDELIS INFUMATA (Gld.). Coast counties of Cal. from Siskiyou to Alameda.
- 22. EPIPHRAGMOPHORA MORMONUM (Pfr.) Shasta Co. to Tulare and Santa Barbara Co., Cal.
- 23. EPIPHRAGMOPHORA HILLEBRANDI (Newc.) Calaveras, Tuolumne and Mariposa Cos., Cal.
- 24. EPIPHRAGMOPHORA CIRCUMCARINATA (Stearns). Near Turloch, Stanilaus Co., Cal.

Subgenus Helminthoglypta Ancey.

- 25. Epiphragmophora dupetithouarsi (Desh.). Monterey, Cal.
- 26. EPIPHRAGMOPHORA SEQUOICOLA (Coop.). Santa Cruz Co., Cal.
- 27. EPIPHRAGMOPHORA AYRESIANA (Newc.). Santa Cruz, San Miguel, San Clemente and Santa Rosa Islands.
- 28. EPIPHRAGMOPHORA TRASKI (Newc.). Los Angeles to Ft. Tejon and to San Luis Obispo, Cal.
- 28a. EPIPHRAGMOPHORA TRASKI PROLES Hemphill. Fraser's Mills, Tulare Co., Cal.

(To be Continued.)



Pilsbry, Henry Augustus. 1897. "A classified catalogue of American land shells, with localities." *The Nautilus* 11, 45–48.

View This Item Online: https://www.biodiversitylibrary.org/item/86834

Permalink: https://www.biodiversitylibrary.org/partpdf/95119

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.