

A New Species of *Terebra* from the Moçambique Channel

BY

R. A. WHITNEY

953 East Prairie Street, Decatur, Illinois 62521

(1 Plate)

A SPECIES OF *Terebra*, which has not been described previously, was received from Moçambique. The specimens were obtained through a supplier in the village of Mossuril, Moçambique, who got them from native fishermen. They were trawled in the Moçambique channel, where the fauna is semi-tropical and tropical from the influence of the southward flowing Moçambique Current. A strong Indo-Pacific component is present in this area, but as the present species has little resemblance to any previously described species, it may be that it is endemic to the Moçambique Channel.

With the advent of SCUBA diving and fishermen extending their operational range for trawling, this area may soon produce more taxa to be added to the faunal record.

Terebra lillianae Whitney, spec. nov.

(Figures 1 to 4)

Description: Size medium; color shiny white with butterscotch brown banding on the anterior $\frac{1}{3}$ of whorls; whorls slightly convex with convex white double subsutural bands marked by impressed suture and deep subsutural grooves forming the 2 subsutural bands; nucleus of $1\frac{1}{2}$ dome-shaped glassy whorls; first postnuclear whorl translucent with almost straight ribs about equal to interspaces forming slight nodes toward the apical end and no noticeable subsutural band or spiral sculpture; in next 6 whorls axial ribs much wider than interspaces, forming rounded nodes with double subsutural band dividing whorls into approximate thirds; in next 7 whorls ribs become progressively wider spaced, with one inconspicuous spiral groove on anterior portion of whorl crossing ribs; subsutural bands now have a pronounced nodular appearance; ribs axially join lower subsutural band by elongated rib extension; penultimate whorl with weak spiral cords and axial ribs numbering 20, becoming more nodose on subsutural bands than on anterior portion of

whorl; body whorl of medium length with butterscotch brown stripe and divided by spiral groove at periphery; anterior to periphery spiral grooves continue to cross axial ribs with somewhat cancellate appearance; aperture semi-quadrate; outer lip sturdy, white within; with 2 brown bands showing through from outside markings; columella white, twisted, broad, with 2 extremely weak plications; anterior canal broad, very twisted; length 42.2 mm; diameter 9.1 mm; 16 whorls plus nucleus.

Holotype: Los Angeles County Museum of Natural History, Type Collection No. 1724

Type Locality: The holotype and 2 paratypes were collected during 1974 off Mossuril, Moçambique, $14^{\circ}55'S$ latitude, $40^{\circ}41'E$ longitude.

Paratypes: One paratype, length 41.8 mm, width 8.0 mm in the R. A. Whitney collection, no. 66; the second paratype, length 41.0 mm, width 8.0 mm, is in the Douglas and Sherry Welker collection no. 2, in Decatur, Illinois.

Largest Specimen Examined: The holotype.

Discussion: This species shows little variation among the specimens examined. It has a striking and consistent pattern in both sculpture and coloration. The sculpture is extremely nodose on the 2 subsutural bands, only slightly less so on the anterior portion of the whorl, giving the shell a file-like appearance. The 2 subsutural bands are shiny white with the butterscotch brown color of the whorls slightly impinging on the subsutural band. The main difference exists in the number of spiral grooves on the anterior portion of early whorls. Among the specimens examined, the spiral grooves vary in number from 1 to 4.

Terebra lillianae has little resemblance to other species of *Terebra*; however, some superficial resemblance to other species should be considered in making identifications. The Japanese species, *T. torquata* Adams & Reeve, 1850, has the nodular appearance on the 2 subsutural

bands, but the anterior portion of whorls is crossed with spiral cords, giving a cancellate appearance, whereas in *T. lillianae* the anterior portion of whorls is quite nodose. *Terebra cracilenta* Li, 1930, is quite nodose, but has only one subsutural band, and is a uniform pale flesh color, whereas *T. lillianae* has 2 subsutural bands and is a striking striped bicolor. *Terebra hancocki* Bratcher & Burch, 1970, might also be compared, although this species is consistently broader and the coloration is a shiny pale beige with irregular blotches of reddish brown.

The new species is named in honor of Mrs. Lillian Whitney of Decatur, Illinois, in appreciation for her encouragement of, and assistance in, the author's study of Terebridae.

Literature Cited

- BRATCHER, TWILA
1973. Six *Terebra* look-alikes. Hawaiian Shell News XXI (4), n. ser. 160: 1, 3 (April 1973)
- BRATCHER, TWILA & ROBERT DONALD BURCH
1970. Five new species of *Terebra* from the eastern Pacific. The Veliger 12 (3): 295-300; plt. 44 (1 January 1970)
- BURCH, ROBERT DONALD
1964. Notes on the Terebridae of the Philippine Islands (Mollusca: Gastropoda). The Veliger 6 (4): 210-218 (1 April 1964)
1965. New terebrid species from the Indo-Pacific Ocean and from the Gulf of Mexico, with new locality records and provisional list of species collected in Western Australia and at Sabah, Malaysia. The Veliger 7 (4): 241-253; plt. 31 (1 April 1965)
- CATE, JEAN MCCREERY & ROBERT DONALD BURCH
1964. Mitridae and Terebridae (Mollusca: Gastropoda) of Malaita, Fiji, and Bileau Island, New Guinea. The Veliger 6 (3): 139-147; 1 map (1 January 1964)
- CERNOHORSKY, WALTER OLIVER
1969. List of type specimens of Terebridae in the British Museum (Natural History). 11 (3): 210-222 (January 1969)
- CERNOHORSKY, WALTER OLIVER & ALBERT JENNINGS
1966. The Terebridae of Fiji (Mollusca: Gastropoda). The Veliger 9 (1): 37-67; pls. 4-7; 13 text figs. (1 July 1966)
- KEEN, A. MYRA
1971. Sea shells of tropical West America: marine mollusks from Baja California to Peru. Stanford Univ. Press, Stanford, Calif. i-xiv+1066 pp.; ca. 4000 figs.; 22 color pls. (1 September 1971)
- SMITH, EDGAR ALBERT
1873. Remarks on a few species belonging to the family Terebridae and descriptions of several new forms in the collection of the British Museum. Ann. Mag. Nat. Hist. (4)
- WEAVER, CLIFTON STOKES
1960-1961. Hawaiian marine mollusks; the genera *Terebra* and *Hastula*. Insert in Hawaiian Shell News 1 (1-9): 1-36; pls. 1-9 (1 January 1960 to 1 January 1961)

Explanation of Figures 1 to 4

Terebra lillianae Whitney, spec. nov.

Figure 1: Holotype, Los Angeles County Museum of Natural History type no. 1724

Figure 2: Paratypes; Whitney specimen on left. Welker specimen on right × 2.6

Figure 3: Enlargement of protoconch and early whorls, Paratype in Whitney Collection × 5

Figure 4: Enlargement of Aperture of Holotype × 5.5



Whitney, R. A. 1976. "A NEW SPECIES OF TEREBRA FROM THE MOZAMBIQUE CHANNEL." *The veliger* 18, 381–382.

View This Item Online: <https://www.biodiversitylibrary.org/item/137755>

Permalink: <https://www.biodiversitylibrary.org/partpdf/97610>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: California Malacozoological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.