

Anatomy, Ecology and Distribution of the Volutidae and Volutomitridae of the Southern Indian Ocean

(Gastropoda: Prosobranchia)

BY

PATRICK M. ARNAUD

Station Marine d'Endoume, 13007 Marseille, France

AND

JEAN-JACQUES VAN MOL

Université Libre de Bruxelles, 50 Avenue F. D. Roosevelt, 1050 - Bruxelles, Belgique

(1 Plate; 10 Text figures)

THREE RECENT BENTHIC SURVEYS have been done with the M/S *Marion-Dufresne* in the subantarctic waters of the Indian Ocean. During cruise MD.03 (1974) several stations were made (including 50 macrobenthic samples) off Kerguelen, Heard and Crozet Islands between shallow water and 4200 m. Cruise MD.04 (1975) concentrated on the shelf around Kerguelen Islands, with 231 macrobenthic samples, and cruise MD.08 was mostly concerned with the Marion-Prince Edward shelf (49 macrobenthic samples) and the Crozet shelf (99 macrobenthic samples).

We collected 3 species of volutid and volutomitrid gastropods at 65 stations, in 73 samples, by trawls, dredges and grabs (Table 1): *Provocator pulcher* Watson, 1882; *Volutomitra curta* (Strebel, 1908) and *V. fragillima* Watson, 1882 (Figures 1 to 3). Study of the plentiful specimens provides the first thorough insight into these 2 allied neogastropod families in this area. Many faunistic and biogeographic data are recorded and the resulting distribution of the 3 species obtained is mapped (Figures 7, 8). Our anatomical studies, made with the help of D. van Weert, have led to discussion of several of their characteristics and assessment of their positions in the 2 families involved. The egg capsule of the volutid *Provocator pulcher* is also described and figured for the first time.

Family Volutidae

Provocator pulcher Watson, 1822

Provocator pulcher WATSON, 1882: 330, 331; 1886: 260, pl. 13, fig. 5; CARCELLES, 1947: 6; POWELL, 1960: 156; WEAVER & DUPONT, 1970: 123, fig. 53 F-G; CLOVER, 1975a: 10, 1 photo; 1975b: 1, 1 photo
Provocator provocator SOWERBY, 1887: 305, pl. 18, fig. 176; SMITH, 1942: 62, 63, pl. 13, fig. 95
Zidona (Provocator) pulchra WENZ, 1943: 1350, fig. 3822

Specimens (empty shells recorded in parentheses):

- Kerguelen Islands

MD.03: 7-CP4, 3; 11-CP7, (2); 13-CP9, 3; 17-CB5, 12 (3); 21-CP14, 5 (2); and 2 egg capsules (incl. 1 empty); 24-CB6, 5 (1)

MD.04: 7-CP13, 1 (2); 26-CP61, 2; 34-DC88, (1); 35-DC89, 1; 38-CP92, 1; 39-DC93, (1); 45-DR106, (2); 70-CP169, 1 egg capsule; 95-DC233, 1 egg capsule; 118-CP284, 1 (1)

- Crozet Islands

MD.08: 42-CP197, (1); 75-CP303, (3); 78-CP319, (2)

Distribution (Figures 7, 8):

The type locality is Kerguelen Islands, West Christmas Harbour, 48°43'S; 69°15'E, 105 fathoms [190 m], volcanic mud. Previously known only by empty shells (WATSON, 1882) and by shells from unknown stations off Kerguelen (CLOVER, 1975a, 1975b), the species is here re-




Figure 7

Present records of Volutidae and Volutomitridae off Kerguelen Islands and Heard Island: *Provocator pulcher* (1: living; 2: dead; 3: egg capsule), *Volutomitra fragillima* (4: living; 5: dead) and *Volutomitra curta* (6: living; 7: dead)




Figure 13

Provocator pulcher: valve of Leiblein; A, frontal section; B, transverse section A. Oes. - anterior oesophagus Gr. - non-ciliated groove M. Oes. - mid-oesophagus

When crossing the nerve ring, the mid-oesophagus has a diameter of about 1.5 mm. Its walls are thin. The muscular layer is very weak at its beginning, the fibers have the same disposition as in the anterior part. The external layer of circular muscle fibers is thicker. The inner epithelium forms a few low ridges. At a short distance from the beginning, the gland of Leiblein (Figure 11) begins with swellings that appear on the dorsal wall of the oesophagus. The gland is progressively differentiated as a tubular, elongate organ coiled upon itself in a compact mass surrounding the oesophagus. The lumen is alveolar with muscular compartments. The gland is composed of 3 distinct parts. The first is lined with tall columnar gland cells with few mucocytes; the muscular layer is very thin. In the next part, the wall has a thick muscular layer and bears longitudinal and transverse ridges. The glandular epithelium is composed of a higher percentage of mucocytes. The dead end of the gland forms a sac 2 to 2.5 mm wide. Its wall is very thin and lined in most of its parts by a flattened epithelium. There are a few scattered low ridges of gland cells comprising some mucocytes. The bad fixation of the tissues in this part of the digestive tract prevented better interpretation.

Unidentified food remnants have been found in this part of the alimentary canal. Behind the gland of Leiblein, the posterior part of the mid-oesophagus has a diameter of 1.9 mm; its internal wall bears many longitudinal ridges.

The posterior oesophagus is separated from the preceding part by a constriction. Its section increases rapidly to a diameter of 2 to 2.5 mm. The muscular layer is very thin, the epithelium consists of ciliated cells and mucocytes. There are digitated internal ridges.

The U-shaped stomach is a very simple sac with an internally plicated wall. The ridges of the dorsal part of the posterior oesophagus reach the single digestive gland aperture. The ventral ridges continue in the intestine without interruption.

The intestine runs from the stomach along the right pallial wall. Its diameter is 2 to 3 mm. The thin muscular layer is lined with an epithelium which is thickened into longitudinal ridges of columnar, ciliated gland cells with numerous dark granules. The mucocytes, scattered at the beginning, are much more abundant towards the end. The anal gland is a branching tubule derived from an invagination of the renal wall. Its cells are ciliated and granular.

Discussion: With these observations, it is possible to determine the systematic position of the genus *Provocator*. This genus has been classified in the subfamily Zidoninae by CLENCH & TURNER (1964) on the basis of the morpho-

logy of the radula. This placement is corroborated by the following observations:

- uniseriate radula with tricuspid teeth;
- accessory salivary glands loosely bound around moderately compact salivary glands;
- lobes at the bases of the siphon of equal development;
- absence of operculum.

It should be added that the genus *Provocator* could be considered more primitive than *Alcithoe* by the fact that its gland of Leiblein is less detached from the oesophageal wall than in the latter genus (compare the anatomical description of *Alcithoe arabica* by PONDER, 1971).

On the specific level, the very close similarity should be noted between the radular teeth of *Provocator pulcher* (Figure 12A) and *P. corderoi* (cf. plt. 107 in CLENCH & TURNER, 1964); the teeth of the latter species are somewhat more slender, with more arcuate bases.

Family Volutomitridae

Volutomitra curta (Strebel, 1908)

Paradmete curta STREBEL, 1908: 23, plt. 3, figs. 34a-e; POWELL, 1951: 166; CARCELLES, 1953: 196; POWELL, 1958: 198; 1960: 157

Paradmete longicauda STREBEL, 1908: 24, plt. 3, figs. 36a-b; POWELL, 1951: 165; CARCELLES, 1953: 196, plt. 3, fig. 70

Volutomitra (Paradmete) curta. CERNOHORSKY, 1970: 103, plt. 13, figs. 11-13

Specimens (all from Kerguelen Islands):

MD.03: 3-CP2, 6 (1); 3-DS1, 1; 6-CP3, 3; 17-CB5, 70 (9); 21-CP14, 1; 24-CB6, 2

MD.04: 15-DC37, 1; 17-DC39, 1; 24-DR58, 1 (2); 38-CP92, 1; 47-DC108, 1; 82-CP196, (1); 113-DC269, (1); 115-DC275, 1

Distribution: The type locality is Shag Rock Bank, 53° 34'S, 43° 23'W, 160 m, gravel and sand, bottom temperature +2.05°C. It was known from Shag Rock Bank,

South Georgia Island, Enderby Land, Mackenzie Sea. First recorded here (cf. Figure 7) from Kerguelen Islands, a major northward extension of range of this volutomitrid. Not obtained at Crozet Islands or Marion/Prince Edward Islands during our surveys.

Bathymetric Range: Was known living from 75 to 300 m. Thus it is of interest to point out that this species was obtained living off Kerguelen Islands from 120 m to as deep as 650 m (present material).

Ecology: Rather similar to that of *Volutomitra fragillima* (the 2 species are frequently collected together) but with a lower temperature-tolerance which would account for its absence from both the Crozet and Marion/Prince Edward shelves.

Anatomy (Figures 14, 15): Animal rather small; maximum height of the shell: 28.6 mm. Tentacles cylindrical, bearing eyes on their posterior side. Operculum present.

The Alimentary Canal:

The pleurembolic proboscis is relatively long. The wall of the proboscis sheet is transversely plicate. The strong retractor muscles connect its sides to the lateral walls of the body. The ventral wall of the buccal cavity is protected by a chitinous shield. The accessory salivary gland is an unpaired organ as in other Volutomitridae. This gland is quite small and entirely concealed in the proboscis. Its thin secretory duct lies just behind the chitinous shield of the buccal floor and opens at the mouth aperture. Histologically the gland consists of a tube lined with cubic epithelium, the cells possessing large nuclei, and an outer layer of circular muscle fibers. Few gland cells are situated in its terminal part. The radula is triseriate. The central tooth is quite large with a single thin lanceolate cusp and 2 long basal processes (Figure 12C). The lateral teeth are small and short.

The salivary glands are small; the secretory tubules are composed of one type of cell. The salivary ducts enter the wall of the oesophagus in front of the valve of Leiblein. They open into the buccal cavity close to the junction of

Explanation of Figures 1 to 6

Figure 1: *Provocator pulcher*, height 85 mm, from Kerguelen Islands, MD. 03-24-CB6

Figure 2: *Volutomitra curta*, height 15.6 mm, from Kerguelen Islands, MD. 03-17-CB5

Figure 3: *Volutomitra fragillima*, height 15.4 mm, from Kerguelen Islands, MD. 04-F51-DC119

Figure 4: Egg capsule of *Provocator pulcher* on a valve of *Malletia gigantea* measuring 47 × 28 mm; Kerguelen Islands, MD 04-H95-DC233

Figure 5: Two egg capsules of *Provocator pulcher* on an egg capsule of skate, Kerguelen Islands, MD. 03-21-CP14

Figure 6: The same capsules, enlarged




Figure 14

Volutomitra curta: anterior part of the digestive tract. Abbreviations as for Figure 11. E - eye M - muscular bundles

the oesophagus. The ducts are densely ciliated, the cilia having a forward orientation.

The anterior oesophagus is relatively narrow. There are 2 lateral ciliated ridges. Ventrally the epithelium is composed of flattened, nonciliated cells. Gland cells filled with red granules are in all this part of the digestive tract.

The mid-oesophagus begins with the valve of Leiblein. The tract is rather long and coiled upon itself, a strong muscular bundle is attached between 2 loops. The valve of Leiblein, situated well in front of the nerve ring, is quite typical. The ventral non-ciliated groove of the anterior oesophagus persists as a deep slit into the fold of the valve. This groove, ventral in the anterior part of the valve, moves progressively to the right and becomes dor-

sal posteriorly. This evidence of torsion thus occupies the same level as in the Volutidae. Two bundles of longitudinal muscle fibers are developed into the wall of the valve, below the ridges lining the non-ciliated groove. The histological organisation of the valve corresponds to the one observed in *Provocator*, except the 2 conspicuous muscular bundles present in its walls. The mid-oesophagus possesses a ciliated groove which corresponds to the pretorsional dorsal alimentary tract of the anterior oesophagus. At its beginning, the wall of the mid-oesophagus is thin; later on, it becomes thicker by increase of the internal layer of circular muscle fibers. The numerous mucocytes and the red-stained cells that constitute the dorsal epithelium of the first part of the mid-oesophagus are pro-




Figure 15

Volutomitra curta: transverse section through proboscis.

B. C. - buccal cavity Car. - subradular cartilage D. A. Sal. Gl. - common duct of accessory salivary glands Rad. - radula
Oe. - oesophagus Pr. - proboscis Pr. S. - proboscis sheet

gressively replaced by gland cells of irregular shape and filled with brown granules. These latter elements are restricted to the dorsal part of the lumen and are separated from the ventral ciliated groove by 2 low ciliated ridges. Further on the dorsal part is completely isolated and forms the gland of Leiblein. At its posterior end, the gland emerges out of the wall of the mid-oesophagus as a small

translucent vesicle. Its histology seems to be uniform.

The transition from the mid-oesophagus to the posterior oesophagus is visible externally as a rapid decrease in the external diameter of the duct. Nevertheless, the size of the lumen remains constant, as there is a considerable diminution of the muscular layers of the wall. The lumen is lined with ciliated epithelium.

THE VELIGER is open to original papers pertaining to any problem concerned with mollusks.

This is meant to make facilities available for publication of original articles from a wide field of endeavor. Papers dealing with anatomical, cytological, distributional, ecological, histological, morphological, physiological, taxonomic, etc., aspects of marine, freshwater or terrestrial mollusks from any region, will be considered. Even topics only indirectly concerned with mollusks may be acceptable. In the unlikely event that space considerations make limitations necessary, papers dealing with mollusks from the Pacific region will be given priority. However, in this case the term "Pacific region" is to be most liberally interpreted.

It is the editorial policy to preserve the individualistic writing style of the author; therefore any editorial changes in a manuscript will be submitted to the author for his approval, before going to press.

Short articles containing descriptions of new species or lesser taxa will be given preferential treatment in the speed of publication provided that arrangements have been made by the author for depositing the holotype with a recognized public Museum. Museum numbers of the type specimens must be included in the manuscript. Type localities must be defined as accurately as possible, with geographical longitudes and latitudes added.

Short original papers, not exceeding 500 words, will be published in the column "NOTES & NEWS"; in this column will also appear notices of meetings of the American Malacological Union, as well as news items which are deemed of interest to our subscribers in general. Articles on "METHODS & TECHNIQUES" will be considered for publication in another column, provided that the information is complete and techniques and methods are capable of duplication by anyone carefully following the description given. Such articles should be mainly original and deal with collecting, preparing, maintaining, studying, photographing, etc., of mollusks or other invertebrates. A third column, entitled "INFORMATION DESK," will contain articles dealing with any problem pertaining to collecting, identifying, etc., in short, problems encountered by our readers. In contrast to other contributions, articles in this column do not necessarily contain new and original materials. Questions to the editor, which can be answered in this column, are invited. The column "BOOKS, PERIODICALS, PAMPHLETS" will attempt to bring reviews of new publications to the attention of our readers. Also, new timely articles may be listed by title only, if this is deemed expedient.

Manuscripts should be typed in final form on a high grade white paper, 8½" by 11", double spaced and accompanied by a carbon copy.

A pamphlet with detailed suggestions for preparing manuscripts intended for publication in **THE VELIGER** is available to authors upon request. A self-addressed envelope, sufficiently large to accommodate the pamphlet (which measures 5½" by 8½"), with double first class postage, should be sent with the request to the Editor.

EDITORIAL BOARD

DR. DONALD P. ABBOTT, *Professor of Biology*
Hopkins Marine Station of Stanford University

DR. WARREN O. ADDICOTT, *Research Geologist*, U. S.
Geological Survey, Menlo Park, California, and
Consulting Professor of Paleontology, Stanford
University

DR. HANS BERTSCH, *Curator of Marine*
Invertebrates
San Diego Museum of Natural History

DR. JERRY DONOHUE, *Professor of Chemistry*
University of Pennsylvania, Philadelphia, and
Research Associate in the Allan Hancock Foundation
University of Southern California, Los Angeles

DR. J. WYATT DURHAM, *Professor of Paleontology*
Emeritus
University of California, Berkeley, California

DR. CADET HAND, *Professor of Zoology and*
Director, Bodega Marine Laboratory
University of California, Berkeley, California

DR. CAROLE S. HICKMAN, *Assistant Professor of*
Paleontology
University of California, Berkeley, California

DR. A. MYRA KEEN, *Professor of Paleontology and*
Curator of Malacology, *Emeritus*
Stanford University, Stanford, California

DR. VICTOR LOOSANOFF, *Senior Biologist*, *Emeritus*
U. S. National Marine Fisheries Service

DR. JOHN MCGOWAN, *Professor of Oceanography*
Scripps Institution of Oceanography, La Jolla
University of California at San Diego

DR. FRANK A. PITELKA, *Professor of Zoology*
University of California, Berkeley, California

DR. ROBERT ROBERTSON, *Pilsbry Chair of Malacology*
Department of Malacology
Academy of Natural Sciences of Philadelphia

DR. PETER U. RODDA,
Chairman and Curator, Department of Geology
California Academy of Sciences, San Francisco

DR. CLYDE F. E. ROPER, *Curator*
Department of Invertebrate Zoology (Mollusca)
National Museum of Natural History
Washington, D. C.

DR. JUDITH TERRY SMITH, *Visiting Scholar*
Department of Geology, Stanford University
Stanford, California

DR. RALPH I. SMITH, *Professor of Zoology*
University of California, Berkeley, California

DR. CHARLES R. STASEK,
Bodega Bay Institute
Bodega Bay, California

DR. T. E. THOMPSON, *Reader in Zoology*
University of Bristol, England


EDITOR-IN-CHIEF

DR. RUDOLF STOHLER, *Research Zoologist*, *Emeritus*
University of California, Berkeley, California

ASSOCIATE EDITOR

MRS. JEAN M. CATE
Rancho Santa Fe, California


594.05
V43
Moll.



ISSN 0042-3211

THE VELIGER

A Quarterly published by
CALIFORNIA MALACOOLOGICAL SOCIETY, INC.
Berkeley, California



VOLUME 22

APRIL 1, 1980

NUMBER 4

CONTENTS

Egg Masses of Mollusca from the Mediterranean Waters of Israel and Notes on
Reproduction of the Freshwater Species *Theodoxus jordani* and *Melanoides
tuberculata*. (14 Plates; 1 Text figure)

AL. BARASH & Z. ZENZIPER 299

Aligena laterodentata, New Species, from the Caribbean Coast of Honduras (Bival-
via : Leptonacea). (5 Text figures)

HAROLD W. HARRY, JOSEPH C. BRITTON & RODNEY NUNLEY 318

The Lamellariidae of the North Eastern Pacific. (2 Plates; 15 Text figures)

DAVID W. BEHRENS 323

Predation by the Prosobranch Mollusk *Lamellaria diegoensis* on *Cystodytes lobatus*,
a Colonial Ascidian. (1 Text figure)

GRETCHEN LAMBERT 340

Stone Boring Marine Bivalves from Monterey Bay, California. (1 Text figure)

E. C. HADERLIE 345

Genetic Variation in Sympatric Sibling Species of *Littorina*.

N. P. WILKINS & D. O'REGAN 355

Range Extension and Notes on the Feeding of the Nudibranch *Okenia cupella*.

PHILLIP NIMESKERN, JR. 360

Two New Molluscan Species (Gastropoda : Muricidae) from the Tropical Eastern
Pacific. (1 Plate)

LEROY H. POORMAN 361

[Continued on Inside Front Cover]

Distributed free to Members of the California Malacozoological Society, Inc.
Subscriptions (by Volume only) payable in advance to Calif. Malacozool. Soc., Inc.

Volume 23: \$37.50 plus \$1.50 for postage (U. S. A. only)

For ALL foreign countries: Swiss Francs 60.- plus SF 7.- for postage

Single copies this issue \$22.00. Postage additional.


Send subscription orders to California Malacozoological Society, Inc.

1584 Milvia Street, Berkeley, CA 94709, U. S. A.

Address all other correspondence to Dr. R. STÖHLER, Editor, Department of Zoology
University of California, Berkeley, CA 94720

CONTENTS — Continued

On the Taxonomic Position, the Species and the Paleocological Significance of the Genera <i>Eubora</i> , <i>Toxosoma</i> and <i>Littoridina</i> (?) in the Pliocene Pebas Forma- tion of the Upper Amazon Region (Gastropoda : Prosobranchia). (17 Text figures)	
DIETRICH KADOLSKY	364
Gastropod Mimicry by Another Pleustid Amphipod in Central California. (1 Plate)	
JOHN W. CARTER & DAVID W. BEHRENS	376
Early Development of <i>Mytilopsis leucophaeata</i> (Bivalvia : Dreissenacea). (1 Plate)	
SCOTT E. SIDDALL	378
Fish Predation on Pulmonate Limpets. (2 Text figures)	
SUSAN BLACKFORD COOK	380
Intermediate Forms and Range Extension of <i>Pedicularia californica</i> and <i>Pedicularia</i> <i>ovuliformis</i> . (3 Text figures)	
ROBERT W. SCHMIEDER	382
<i>Helix aperta</i> Introduced in Richmond, California (Mollusca : Pulmonata). (1 Text figure)	
BARRY ROTH & DUSTIN D. CHIVERS	385
<i>Spiroglyphus</i> and <i>Stoa</i> , Taxonomic Problems in the Vermetidae. (1 Text figure)	
A. MYRA KEEN	388
NOTES & NEWS	392
Soviet Contributions to Malacology in 1978. KENNETH J. BOSS & MORRIS K. JACOBSON	
Sea Star Predation on Rock-Boring Bivalves. E. C. HADERLIE	
BOOKS, PERIODICALS & PAMPHLETS	404



Note: The various taxa above species are indicated by the use of different type styles as shown by the following examples, and by increasing indentation.

ORDER, Suborder, DIVISION, Subdivision, SECTION,
SUPERFAMILY, FAMILY, Subfamily, *Genus*, (*Subgenus*)
New Taxa

Egg Masses of Mollusca from Mediterranean Waters of Israel and Notes on Reproduction of the Freshwater Species

Theodoxus jordani and *Melanoides tuberculata*

BY

AL. BARASH AND Z. ZENZIPER

(14 Plates; 1 Text figure)

INTRODUCTION

A SERIES OF PAPERS on the molluscan fauna of the Mediterranean coasts of Israel has been published during the last decades. But only a few observations deal with the reproduction of Mollusca in this area (BARASH & DANIN, 1973; RAHAT, 1973). The purpose of the present work is to give a preliminary report on egg masses of marine molluscs found in the intertidal and infralittoral zones of the Mediterranean Israel and Sinai Peninsula. The Mediterranean coast, along which the material discussed here was collected, extends from Rosh-Haniqra in the North (Lebanon border) to Bardawil (Sinai Peninsula), near the northern entrance of the Suez Canal (see map).

Most egg masses were collected during field trips and expeditions, some were brought after months and even years of having been preserved in alcohol; accurate counting and measuring thus were often beyond our reach. In several cases, adult parent animals were put into aquaria, where they deposited their spawn, thus confirming the identity of the egg masses found in the sea. These included: *Cerithium scabridum*, *C. rupestre*, *Columbella rustica*, *Bulla striata*, *Aplysia fasciata*, *Bursatella leachi savigniana*, *Elysia timida*. The spawn of *Rhinoclavis kochi* was found only in the aquarium.

Egg masses were first sent for identification in 1966 to the late Professor Gunnar Thorson at the Zoological Museum of the University of Copenhagen. Unfortunately, his death interrupted his great scientific work and the material was returned to us containing identified egg masses of 6 species. Dr. Klaus Bandel, Friedrich-Wilhelms University, Bonn, identified egg masses of 5 species, and Dr. Nellie B. Eales, University of Reading, identified the egg masses of 2 species of Aplysiidae.

The greatest part of the material is kept in the collection of the Tel-Aviv University, Department of Zoology. The egg masses of *Cerithium rupestre* and *Elysia timida* are in the collection of the Hebrew University of Jerusalem, Department of Zoology.

The localities in which living adult specimens of the species treated in this paper were found are indicated; the vertical distribution of the species is also given. These data should be useful for future investigations of spawn of these species in the area.

LIST OF SPECIES CONSIDERED

GASTROPODA

PROSOBRANCHIA

MESOGASTROPODA

CERITHIIDAE

Cerithium scabridum Philippi, 1849*Cerithium rupestre* Risso, 1826*Rhinoclavis kochi* (Philippi, 1848)

JANTHINIDAE

Janthina ? nitens Menke, 1828

NATICIDAE

4 types, species not determined

CASSIDIDAE

Cassidaria echinophora (Linnaeus,
1758)*Semicassis undulata* (Gmelin, 1791)




Figure 33

Collecting Localities of Molluscan Egg Masses at the Coasts of Israel and the Sinai Peninsula

TONNIDAE

Tonna galea (Linnaeus, 1758)

NEOGASTROPODA**MURICIDAE**

Trunculariopsis trunculus (Linnaeus, 1758)

Murex brandaris Linnaeus, 1758

THAIDIDAE

Thais haemastoma (Linnaeus, 1767)

Thais carinifera (Lamarck, 1822)

COLUMBELLIDAE

Columbella rustica (Linnaeus, 1758)

BUCCINIDAE

Euthria cornea (Linnaeus, 1758)

Pisania striata (Gmelin, 1791)

NASSARIIDAE

Sphaeronassa mutabilis (Linnaeus, 1758)

FASCIOLARIIDAE

Fasciolaria lignaria (Linnaeus, 1758)

CONIDAE

Conus ventricosus Gmelin, 1791

OPISTHOBRANCHIA**CEPHALASPIDEA**

(Bullomorpha)

BULLIDAE

Bulla striata Bruguière, 1789




Figure 1




Figure 2



Figure 3



Arnaud, Patrick M. and Van Mol, Jean-Jacques. 1979. "ANATOMY ECOLOGY AND DISTRIBUTION OF THE VOLUTIDAE AND VOLUTOMITRIDAE GASTROPODA PROSOBRANCHIA OF THE SOUTHERN INDIAN OCEAN." *The veliger* 22, 19–31.

View This Item Online: <https://www.biodiversitylibrary.org/item/134937>

Permalink: <https://www.biodiversitylibrary.org/partpdf/97674>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: California Malacozoological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.